

# Oral health in a group of sugar-cane chewers

S Harris and P Cleaton-Jones

Dental Research Unit of the South African Medical Research Council and the University of the Witwatersrand, Jan Smuts Avenue, Johannesburg, South Africa, 2001.

*Keywords: Caries; epidemiology; periodontal disease; sugar-cane*

## SUMMARY

The oral health of 98 sugar-cane chewers was investigated. There were progressive increases in D M F T values and Ramfjord's P D I scores as age increased but degree of attrition remained constant. The investigation showed that simple oral hygiene procedures were followed, 4-5 Kg of sugar-cane were chewed per day (= 500 g sucrose), and yet the D M F T values were low.

## OPSOMMING

Die mondigesondheid van 98 suikerrietkouers is ondersoek. 'n Progressiewe vermeerdering in D M F T-waardes en Ramfjord se P D I-tellings het plaasgevind namate die ouderdom toegeneem het. Die graad van attrisie het egter konstant gebly. Hierdie ondersoek het getoon dat eenvoudige mondhigiëne toegepas is, dat daar 4-5 Kg suikerriet per dag (= 500 g sukrose) gekou is, en tog het die D M F T-waardes laag gebly.

## INTRODUCTION

The prevalence of dental caries is lower among populations eating an unrefined diet than those ingesting highly refined carbohydrates. This has been shown in various groups such as South African Bantu (Oranje, Noriskin and Osborn 1935; Staz 1938), Maya and Navajo Indians, Hollanders and Jamaicans (Steggerda and Hill 1936), Indians (Ramfjord 1961), Nigerians (Sheiham 1967), and Finns (Ainamo and Alvesalo 1970).

In 1952 Dreizen and Spies questioned the generally held belief that although refined carbohydrates might be conducive to dental caries, the same foods, if ingested in the raw or natural state, were protective against this disease. They referred to two opposing studies. Osborn and Noriskin (1937) had reported that there was a low prevalence of dental caries among Bantu labourers in the sugar-cane fields of South Africa; in contrast Steggerda and Hill (1936) had reported a high caries prevalence in sugar-cane chewers on the island of Jamaica. Dreizen and Spies then investigated 147 sugar-cane chewers in Cuba 141 of whom were white and 6 negro, and observed a high caries prevalence.

More recently, Künzel *et al* (1973) examined 81 Cuban sugar-cane farm workers and observed high prevalences of both dental caries and periodontal disease. They concluded that the hypotheses which assumed

mechanical cleaning effect of sugar-cane, leading to caries reduction, must be rejected.

The present study was undertaken to examine the dental caries prevalence in Bantu sugar-cane chewers in South Africa, some 40 years after the initial study of Osborn and Noriskin (1937).

## MATERIALS AND METHODS

The survey was carried out on a group of 98 rural Bantu, all the labourers in the region, living and working on sugar farms in the Pongola district, a subtropical region of the South-eastern Transvaal. All ate a typical Bantu diet consisting of maize meal supplemented by legumes and meat (Walker 1977).

The following data was recorded.

1. Personal details: age, sex, length of residence in area
2. Heights (cm) mass (Kg)
3. Oral hygiene methods
4. DMFT index
5. Plaque index
6. Ramfjord's periodontal disease index (PDI)
7. Amount of dental attrition
8. Sugar-cane chewing habits

Forty-seven of the subjects studied were unsure of their age so the classification of age groups suggested by van Reenen (1964) was used. This comprises children, ado-

escents, young adults, middle-aged and old-aged individuals. The criteria for these are shown in Table 1.

Table 1. Details of subjects examined

Age Group (van Reenen 1964)	Males	Females
Old-age (> 60 yrs)	6	1
Middle-aged (40-60 yrs)	7	11
Young adult (+ 3rd molars)	23	15
Adolescent (+ 2nd molars)	26	9
Child (- 2nd molars)	0	0
Total	62	36

All examinations were carried out in natural light using mouth mirror, probe, periodontal probe and disclosing solution. Caries was diagnosed whenever a sharp dental probe 'caught' in a pit, fissure or suspicious area, and any teeth lost due to trauma were regarded as being sound.

The plaque index used was a modified version of the Silness and Løe and Quigley and Hein indices (Glickman 1972), as used in the department of conservative dentistry, University of the Witwatersrand. The six teeth used by Ramfjord (1959) for his periodontal disease index were scored as follows:

1. The teeth were stained with a disclosing solution
2. Each of the six teeth were divided into 4 areas, mesial, buccal, distal and lingual
3. Each of the four areas were scored from 0-3 according to the following criteria:
  - (a) 0 = No plaque
  - (b) 1 = Plaque covering up to 1/3 of the crown of the tooth
  - (c) 2 = Plaque covering from 1/3 to 2/3 of the crown of the tooth
  - (d) 3 = Plaque covering whole of the crown of the tooth

The scores around each tooth were totalled and divided by four to determine the mean plaque index for each tooth. These indices were then recorded.

The periodontal disease index was determined according to Ramfjord (1959).

The criteria used for the attrition index were those suggested by van Reenen (1964):

1. The occlusal surfaces of the six teeth were examined
2. The scores were allocated as follows:
  - (a) 0 = No attrition of enamel
  - (b) 1 = Attrition of enamel
  - (c) 2 = Attrition of enamel with exposed dentine
  - (d) Devitalized tooth and pulp chamber exposed

The observations were transferred to punch cards and analysed in an IBM 370/158 computer using the Statistical Package for the Social Sciences (Nie *et al* 1975).

In addition water samples from all sources of drinking water, namely boreholes, rivers and canals, were collected for measurement of their fluoride content. The fluoride analyses were performed using a combined fluoride ion selective electrode (Orion Research, Model 96-09) coupled to a digital pH meter (Orion Research, Model 701).

## RESULTS

Almost twice as many males as females were examined. No children were investigated and there were only one in the old-age group. The majority of subjects fell within the adolescent and young adult groups (Table 1). They were a stable population having lived in the area for a mean of  $17.2 \pm 11.8$  years.

The traditional method of chewing the sugar-cane practised by all the subjects. The outer hard skin is stripped off with the teeth to expose the fleshy part of the cane containing fibre and sucrose (Fig. 1). In 1974 the mean composition of this flesh was 14.2 per cent fibre, 13.5 per cent sucrose with the remainder being organic and inorganic impurities, cellulose and water. The mean pH of the sugar-cane juice was 5.2 (Stellenbosch Milling Research Institute 1974 - personal communication).



Fig. 1. The outer hard skin is stripped off using the teeth.




Fig. 2. A substantial bite is taken of the inner, hard, fibre bundle containing the sucrose juice.

A substantial bite of the inner fleshy fibre bundle containing the sugar-cane juice was then taken (Fig. 2) to produce a bolus of about 1-4 cm. This was then tentatively chewed and sucked until most of

Table II. Details of dental caries in the subjects (means  $\pm$  standard deviations)

Index	Sex	Age group				
		Adolescent	Young adult	Middle-aged	Old-aged	All ages
" subjects caries free	Male	17,3	8,2	1,0	0	25,5
	Female	6,1	4,1	1,0	0	11,2
Decayed teeth	Male	0,7 $\pm$ 1,2	1,6 $\pm$ 2,0	1,6 $\pm$ 1,6	2,0 $\pm$ 2,1	1,3 $\pm$ 1,7
	Female	0,9 $\pm$ 1,7	1,7 $\pm$ 2,0	1,6 $\pm$ 2,0	5,0 $\pm$ 0	1,6 $\pm$ 1,9
Missing teeth	Male	0,1 $\pm$ 0,3	0,7 $\pm$ 1,3	5,4 $\pm$ 5,9	5,8 $\pm$ 4,7	1,5 $\pm$ 3,3
	Female	0,2 $\pm$ 0,7	2,5 $\pm$ 4,2	3,7 $\pm$ 4,1	5,0 $\pm$ 0	2,4 $\pm$ 3,8
DMFT	Male	0,8 $\pm$ 1,3	2,4 $\pm$ 2,6	7,0 $\pm$ 6,5	7,8 $\pm$ 4,9	2,8 $\pm$ 4,0
	Female	1,1 $\pm$ 2,3	4,2 $\pm$ 4,7	5,4 $\pm$ 4,6	10,0 $\pm$ 0	4,0 $\pm$ 4,6

Table III. Plaque index, Ramfjords PDI and attrition indices (means  $\pm$  standard deviations)

Index	Sex	Age group				
		Adolescent	Young adult	Middle-aged	Old-aged	All ages
Plaque index (max = 3)	Male	1,4 $\pm$ 0,4	1,4 $\pm$ 0,5	1,6 $\pm$ 0,6	1,6 $\pm$ 0,5	1,5 $\pm$ 0,5
	Female	1,4 $\pm$ 0,3	1,5 $\pm$ 0,3	1,6 $\pm$ 0,5	2,0 $\pm$ 0	1,5 $\pm$ 0,4
Ramfjord PDI (max = 6)	Male	0,9 $\pm$ 0,7	1,8 $\pm$ 1,1	3,3 $\pm$ 1,3	2,4 $\pm$ 1,1	1,6 $\pm$ 1,2
	Female	0,4 $\pm$ 0,7	1,3 $\pm$ 0,8	3,3 $\pm$ 1,0	3,1 $\pm$ 0	1,7 $\pm$ 1,4
Attrition index (max = 3)	Male	1,2 $\pm$ 0,2	1,2 $\pm$ 0,1	1,3 $\pm$ 0,2	1,2 $\pm$ 0,2	1,2 $\pm$ 0,2
	Female	1,1 $\pm$ 0,1	1,2 $\pm$ 0,2	1,2 $\pm$ 0,2	1,0 $\pm$ 0	1,2 $\pm$ 0,2

sweetness had been extracted after which the chewed bolus was discarded and another bite taken.

It proved impossible to record accurately the frequency of sugar-cane chewing which varied considerably from day to day; the labourers chew the cane only during tea and lunch breaks and in off-duty hours. The amount of cane chewed varied with age with the younger individuals chewing more than the older. Usually, however, about 5 metres of cane was chewed per day which is equivalent to 4-5 Kg of sugar-cane and approximately 500 g of sucrose.

In spite of the high sucrose intake the heights and mass of the subjects fell within the norms accepted by the International Biological Programme and the Metropolitan Life Insurance Company (Walker 1977 - personal communication).

Dental caries observations are listed in Table II. Of all the subjects 36,7 per cent were caries free, the greatest number being in the adolescent and old age groups. A greater number of males were free of caries than were females but this difference was not statistically significant. There was an increase in caries with aging, the lowest prevalence was in the adolescent group. No significant differences were found between males and females in this nor any other section of the study. In the adolescents and young adults there were more carious than missing teeth while the opposite was noted in the middle-aged and old-aged groups. The tooth loss was due to both caries and periodontal disease. It was not possible to determine which condition was responsible for the greater loss. No restored teeth were seen.

When the plaque index was calculated this was found to be constant throughout all the groups (Table III).

The values of Ramfjord's periodontal index showed an increase with age (Table III). The value in the middle-

aged group was more than double those in the younger individuals, which difference was statistically significant ( $p < 0,5$ ). Careful questioning revealed that a reduced ability to chew the hard sugar-cane due to pain and mobile teeth was associated with this increase in the periodontal index.

Attrition was seen in all the age groups. The amount of attrition did not increase with age.

The various oral cleansing habits used are listed in Table IV. These were rudimentary and mostly consisted of rinsing the mouth with water. Traditional sand and ash cleansing were used by only 11 per cent of the sample. Statistical analyses revealed no significant differences in any of the indices between groups using different cleansing habits.

The water supply in the area was found to be fluoride deficient with a mean value of 0,088 ppm.

Table IV. Oral cleansing habits in the 98 subjects

Method	Agent		Frequency per day	
Nil	1	Nil	2	0 - 1
Finger	12	Soap	2	1 - 36
Toothbrush	20	Ash	2	2 - 20
Water rinse only	65	Sand	9	3 - 34
		Toothpaste	17	
		Water only	66	

## DISCUSSION

The low prevalence of dental caries in this study confirmed initial clinical impressions. The findings of Oranje and Noriskin (1937) almost 40 years ago have thus been corroborated. Unfortunately, due to differing terminology, the absolute DMFT values obtained in the present study cannot be directly compared to the results in the investigations of Oranje and Noriskin (1937) and Steggerda and Hill (1936).

Dreizen and Spies (1952) and Künzel *et al* (1973), however, employed similar terminology. They recorded mean DMFT values of 15.1 and 16.5 respectively in Cuban sugar-cane chewers. These values are very much higher than the very low DMFT of 3.2 noted in the present study, which must include teeth lost due to periodontal disease. Age groups and numbers in the present study and those of Dreizen and Spies (1952) and Künzel *et al* (1973) were similar, as was the method of sugar-cane chewing. All of Künzel *et al*'s subjects and all but 6 of Dreizen and Spies' subjects were white, however, in contrast to the Bantu in the present study. It has been reported elsewhere that there are differences in caries prevalences in different racial groups e.g. whites and negroes in the United States (Kelly *et al* 1967; Littleton *et al* 1970; Rowe *et al* 1976), whites, negroes and other groups in South Africa (Staz 1938; Retief *et al* 1975). It is therefore conceivable that this variation in caries prevalence may also be present in the sugar-cane chewers. If so, then the DMFT values of 15.1 recorded by Dreizen and Spies (1952) and 16.5 of Künzel *et al* (1973) are similar to those in other non-sugar-cane chewing whites, while the DMFT findings in the present study in the Bantu sugar-cane chewers, are similar to the finding in Bantu non-sugar-cane chewers (Staz 1938; Retief *et al* 1975).

Thus, bearing in mind the low numbers examined, one may suggest that caries prevalence is no worse in Black habitual sugar-cane chewers than in Black non-sugar-cane chewers, in spite of an enormous daily sucrose intake of over 500 g.

The reason for this apparent contradiction of the generally accepted dogma is not clear. The abrasive action of the sugar-cane probably plays little role since the amount of plaque present is considerable. Osborn *et al* (1937) have suggested, following *in vitro* studies on tooth decalcification, that there is a protective agent in crude sugar-cane juice which is removed during processing. This hypothesis however, is not supported by the studies of Dreizen and Spies (1952) and Künzel *et al* (1973).

Use of the teeth to tear and chew the hard sugar-cane stalks over a prolonged period was associated with both attrition and periodontal disease. The attrition, although marked, is less than that seen in Kalahari bushmen living in a sandy environment (van Reenen 1964). No mention of the degree of attrition has been made in the other studies of sugar-cane chewers. The effect of the attrition on dental caries coupled with the lack of information on the frequency of sugar-cane ingestion, the possibility of an extraneous fluoride source and the possible absence of a cariogenic flora must temper the apparent rejection of the dogma.

The amount of periodontal disease seen, does not appear excessively high. The PDI values in our study

closely resemble those of Künzel *et al* (1973) as well as those reported in the rural Finnish population by Ainamo and Alvesalo (1970). It is probably a "normal" increment with advancing age. It is not possible to say whether the periodontal disease occurred at an earlier age in the present study, due to the inaccuracy of the age recordings.

Finally, the normal values for mass seen in this study, in spite of a high daily sucrose intake, are probably related to the heavy manual labour involved in cutting the sugar-cane.

## ACKNOWLEDGEMENTS

We gratefully acknowledge the assistance of Mr. P. Fatti of the Department of Applied Mathematics, I. A. R. P. Walker of the M. R. C. Human Biochemistry Research Unit, Miss B. Slack, Mrs. H. Ball and Mrs. Wilton-Cox of the Dental Research Unit and especially Miss E. Joubert.

## REFERENCES

- Ainamo J. & Alvesalo L. (1970) Caries prevalence in a Finnish population. *Acta Odontologica Scandinavica*, **28**, 271-281.
- Dreizen S. & Spies T.D. (1952) The incidence of dental caries in habitual sugar cane chewers. *Journal of the American Dental Association*, **45**, 193-200.
- Glickman I. *Clinical Periodontology*: Saunders, Philadelphia, 1967.
- Kelly, J.E., van Kirk L.E. & Garst C.C. (1967) Decayed, missing filled teeth in adults, United States 1960-1962. *National Center for Health Statistics Ser. 11*, No. 23, U.S. Dept. Health, Education and Welfare, Washington.
- Künzel Von W., Borroto R.C., Lanier S. & Soto F. (1973) Auswirkungen habituellen Zuckerrohrkanens auf Kariesbefall Parodontalzustand kubanischer Zuckerrohrarbeiter. *Deutsche Stomatologie*, **23**, 554-561.
- Littleton W., Kakehashi S. & Fitzgerald R.J. (1970) Study of differences in the occurrence of dental caries in Caucasian Negro children. *Journal of Dental Research*, **49**, 742-751.
- Nie N.H., Hull C.H., Jenkins J.G., Steinbrenner K. & Bent D. (1975) *Statistical Package for the Social Sciences*. 2nd ed. McGraw Hill, New York.
- Oranje P., Noriskin J.N. & Osborn T.W.B. (1935) The effect of upon dental caries in the South African Bantu. *South African Journal of Medical Sciences*, **1**, 57-62.
- Osborn T.W.B. & Noriskin J.N. (1937) The relationship between diet and caries in the South African Bantu. *Journal of Dental Research*, **16**, 431-441.
- Osborn T.W.B., Noriskin J.N. & Staz J. (1937) A comparison of crude and refined sugar and cereals in their ability to promote *in vitro* decalcification of teeth. *Journal of Dental Research*, **16**, 165-171.
- Ramfjord S.P. (1959) Indices for prevalence and incidence of periodontal disease. *Journal of Periodontology*, **30**, 95-112.
- Ramfjord S.P. (1961) Periodontal studies of boys 11-17 years old in Bombay, India. *Journal of Periodontology*, **32**, 237-248.
- Retief D.H., Cleaton-Jones P.E. & Walker A.R.P. (1975) Dental caries and sugar intake in South African pupils of 16 to 17 in four ethnic groups. *British Dental Journal*, **138**, 463-469.
- Rowe N.H., Garn S.M., Clark D.C. & Guire K.E. (1976) The effect of age, sex, race and economic status on dental caries experience of the permanent dentition. *Pediatrics*, **57**, 457-466.
- Sheiham A. (1967) The prevalence of dental caries in Negro populations. *British Dental Journal*, **123**, 145-148.
- Staz J. (1938) Dental caries in South Africa. *South African Journal of Medical Sciences*, **3**, supplement, 1-63.
- Steggerda M. & Hill T.J. (1936) Incidence of dental caries among Maya and Navajo Indians. *Journal of Dental Research*, **15**, 23.
- van Reenen J.F. (1964) Dentition, jaws and palates of the Kalahari Bushmen. *Journal of the Dental Association of South Africa*, **19**, 1-15, 38-44.
- Walker A.R.P. (1977) Health implications of fibre depleted sugar-cane. *South African Medical Journal*. (In Press).