

REFERENCES

Ada L, Canning C, Westwood P 1990 'The patient as an active learner' in Ada L and Canning C (eds) Key issues in Neurological Physiotherapy, Butterworth-Heinemann, Oxford

Andriacci T P, Ogle J A, Galante J O 1977 Walking speed as a basis for normal and abnormal gait measures. *Journal of Biomechanics* 10: 262-268

Bernstein M L, Despars J A, Singh N P, Avalos K, Stansbury D W, Light R W 1994 Reanalysis of the 12-min walk in patients with chronic obstructive pulmonary disease. *Chest* 105: 163-167

Bradsater M E, de Bruin H, Gowland C, Clark B M 1983 Hemiplegic gait: Analysis of temporal variables. *Archives of Physical Medicine and Rehabilitation* 64: 583-587

Butland R J A, Pang J, Gross E R, Woodcock A A, Geddes D M 1982 Two, six, and twelve minute walking tests in respiratory disease. *British Medical Journal* 284: 1604-1608

Carr J H, Shepherd R B 1998 *Neurological Rehabilitation optimizing motor performance*. Butterworth Heinemann, Oxford

Carr J H, Shepherd R B 2003 *Stroke Rehabilitation - guidelines for exercise and training to optimize motor skill*. Butterworth Heinemann, Elsevier Science Limited

Census 2001: <http://www.statssa.gov.za/census01>

Chiou I L, Burnett C N 1985 Values of activities of daily living: a survey of stroke and their home therapists. *Archives of Physical Medicine and Rehabilitation* 65: 901-906

Cole B, Finch E, Gowland C, Mayo N 1994 Physical Rehabilitation Outcome Measures. Basmajian J (editor). Journal of the Canadian Physiotherapy Association (Publisher)

Collin C, Wade D T, Davies S, Horne V 1988 The Barthel ADL Index: A reliability study. International Disability Studies 10: 61-63

Connelly D M, Stevenson T J, Vandervoort A A 1996 Between - and within - rater reliability of walking tests in a frail elderly population Physiotherapy Canada 48: 47-51

Connor M D, Thorogood M, Casserly B, Dobson C, Warlow C P 2004 Prevalence of Stroke Survivors in Rural South Africa; Results from the Southern African Stroke Prevention Initiative (SASPI) Agincourt Field Site. Stroke 35: 627-632

Dawes H, Bateman A, Culpan J, Scott O M, Roach N K, Wade D T 2003 Heart rate as a measure of exercise testing after acquired brain injury. Physiotherapy 89(10): 570-574

Dean C M, Richards C L, Malouin F 2001 Walking speed over 10 metres overestimates locomotor capacity after stroke. Clinical Rehabilitation 15: 415-421

D'Olhaberriague L, Litvan I, Mitsias P, Mansbach H H 1996 A reappraisal of reliability and validity studies in stroke. Stroke 27: 2331-2336

Duncan P W, Badke M B Eds 1987 Stroke rehabilitation: The recovery of motor control. Chicago, Illinois. Year Book Medical Publishers

Duncan P, Richards L, Wallace D, Stoker-Yates J, Pohl P, Luchies C, Ogle A, Studenski S 1998 A Randomized controlled pilot study of a home-based exercise program for individuals with mild and moderate stroke. Stroke 29: 2055-2060

Duncan P W, Jorgensen H S, Wade D T 2000 Outcome measures in acute stroke trials. Stroke 31: 1429

Eales C J, Stewart A V 1996 The exercise capacity of three socio-economic groups of elderly hypertensive patients. *Physiotherapy Research International* 1: 255-264

Eng J J, Chu K S, Dawson A S, Kim C M, Hepburn K E 2002 Functional walk tests in individuals with stroke: relation to perceived exertion and myocardial exertion. *Stroke* 33: 756-761

Enright P L, Sherrill D L 1998 Reference equations for the six-minute walk in healthy adults. *American Journal of Respiratory Critical Care Medicine* 158: 1384-1387

Fiatarone M A, Marks E C, Ryan N D 1990 High intensity strength training on nonagenarians. *Journal of the American Medical Association* 263: 3029-3034

Friedman P J, Richmond D E, Baskett J J 1988 A prospective trial of serial gait speed as a measure of rehabilitation in the elderly. *Age Ageing* 17: 227-235

Friedman P 1990 Gait recovery after hemiplegic stroke. *International Journal of Disability Studies* 12: 119-122

Goldie P A, Matyas T A, Evans O M 1996 Deficit and change in gait velocity during rehabilitation after stroke. *Archives of Physical Medicine and Rehabilitation* 77: 1074-1082

Granger C V, Albrecht G L, Hamilton B B 1979 Outcome of Comprehensive Medical Rehabilitation: Measurement by PULSES Profile and the Barthel Index. *Archives of Physical Medicine and Rehabilitation* 60: 145-154

Grimby G 1983 On the energy cost of achieving mobility. *Scandinavian Journal of Rehabilitation Medicine Supplement* 9: 49-54

Hale L A, Eales C J 1998 Recovery of walking function in stroke patients after minimal rehabilitation. *Physiotherapy Research International* 3(3): 194-205

Hale L A, Eales C J, Fritz V U 1998 The Soweto Stroke Questionnaire. *South African Journal of Physiotherapy* 54(4): 16-20

Hale L A, Eales C J, Stewart A, Fritz V U 1999 The problems experienced by black stroke patients in Soweto, South Africa. *South African Journal of Physiotherapy* 55(2): 12-17

Hale L A, Wallner P J 1996 The challenge of service provision in South Africa for patients with hemiplegia. *Physiotherapy* 82: 156-158

Hill K D, Ellis P S, Bernhardt J, Maggs P, Hull S 1997 Balance and mobility outcomes for stroke patients: a comprehensive audit. *Australian Journal of Physiotherapy* 43: 173-180

Holden M K, Gill K M, Magliozzi M R 1986 Gait assessment for neurologically impaired patients: standards for outcome assessment. *Physical Therapy* 66: 1530-1539

http://www.safrika.info/plan_trip/holiday/cities/soweto.htm

Jorgensen H S, Nakayama H, Pedersen P M, Kammersgaard L, Raashou H O, Olsen T S 1995a Outcome and time course of recovery in Stroke. Part 11: Time course of recovery. The Copenhagen Stroke Study. *Archives of Physical Medicine and Rehabilitation* 76: 406-412

Jorgensen H S, Nakayama H, Raashou H O, Olsen T S 1995b The recovery of walking function in stroke patients: the Copenhagen Stroke Study. *Archives of Physical Medicine and Rehabilitation* 76: 27-32

Kalra L, Evans A, Perez I, Knapp M, Swift C, Donaldson N 2005 A randomized controlled comparison of alternative strategies in stroke care. *Health Technology Assessment* 9(18):

Krebs D E, Jette A M, Assmann S F 1998 Moderate exercise improves stability in disabled elders. *Archives of Physical Medicine and Rehabilitation* 79: 1489-1495

King J L, Guarracini M, Leniham L, Freeman D, Gagas B, Boston A, Bates E, Nori S 1989. Adaptive exercise testing for patients with hemiparesis. *Journal of Cardiopulmonary Rehabilitation* 9: 237-242

Lerner-Frankiel M B, Vargas S, Brown M, Krusell L, Schoheberger W 1986 Functional community ambulation: What are your criteria? *Clinical Management in Physical Therapy* 6(2): 12-15

Macko R F, Katzel L I, Yataco A, Tretter L D, DeSouza C A, Dengel D R, Smith G V, Silver K H 1997 Low-velocity graded treadmill stress testing in hemiparetic stroke patients. *Stroke* 28: 988-992

Mahoney F I, Barthel D W 1965 Functional evaluation: The Barthel Index. *Maryland State Medical Journal* 14: 61-65

Mbeki T M. 1997. White Paper on an Integrated National Disability Strategy. Office of the Deputy President.

McArdle W D, Katch F I, Katch V L 1996 *Exercise Physiology. Energy, nutrition and human performance* 4th Edition. Baltimore Md: Williams and Wilkins

Meijer R, van Limbeek J, Peusens G, Rulkens M, Dankoor K, Vermeulen M, de Haan R J 2005 The Stroke Unit Discharge Guideline, a prognostic framework for the discharge outcome from the hospital stroke unit. A prospective study. *Clinical Rehabilitation* 19(7): 770-778

Moreland J D, Goldsmith C H, Huijbregts M P, Anderson R E, Prentice D M, Brunton K B, O'Brien M A, Torresin W D 2003 Progressive resistance strengthening exercises after stroke: a single-blind randomized controlled trial. *Archives of Physical Medicine and Rehabilitation* 84(10): 1433-1440

Murray M P, Kory R C, Sepic S B 1970 Walking pattern of normal women. *Archives of Physical Medicine and Rehabilitation* 51: 637-650

Nazzal M, Sa'adah M A, Al-ansari D, Al-awadi O, Eyadah A A, Al-kadiri M A, Inshasi J, Trebinjac S, Khader S 2001 Stroke rehabilitation: application and analysis of the modified Barthel index in an Arab community. *Disability and Rehabilitation* 23: 36-42

Olsen T S 1990 Arm and leg paresis as outcome predictors in stroke rehabilitation. *Stroke* 21: 247-251

Olney S J, Monga T H, Costigan P A 1986 Mechanical energy of walking of stroke patients. *Archives of Physical Medicine and Rehabilitation* 67: 92-98

Olney S J, Griffin M P, McBride I D 1994 Temporal, kinematic and kinetic variables related to gait speed in subjects with hemiplegia: A regression approach. *Physical Therapy* 74: 872-885

Olney S J, Richards C 1996 Hemiparetic gait following stroke. Part 1 Characteristics. *Gait Posture* 4: 136-148

Pedersen P M, Jorgensen H S, Nakayama H, Raaschou H O, Olsen T S 1997 Comprehensive assessment of activities of daily living in stroke. The Copenhagen Stroke Study. Archives of Physical Medicine and Rehabilitation 78: 161-165

Perry J, Garrett M, Gronley J K, Mulroy S J 1995 Classification of walking handicap in the stroke population 26: 982-989

Potempa K, Lopez M, Braun L T, Szidon J P, Fogg L, Tincknell T 1995 Physiological outcomes of aerobic exercise training in hemiparetic stroke patients. Stroke 26: 101-105

Roberts L, Counsell C 1998 Assessment of clinical outcomes in acute clinical trials. Stroke 29: 986-991

Robinett C S, Vondran M A 1988 Functional ambulation velocity and distance requirements in rural and urban communities. Physical Therapy 68: 1371-1373

Rossier P, Wade D T 2001 Validity and reliability comparison of 4 mobility measures in patients presenting with neurologic impairment. Archives of Physical Medicine and Rehabilitation 82: 9-13

Rothstein J M 1985 Measurement in Physical Therapy. Churchill Livingstone, New York

SAMA – Neurological Association of South Africa. Stroke working group 2000 Stroke Therapy Clinical Guideline. South African Medical Journal 90(3):

Scurba F C, Slivka W A 1998 Six-Minute walk testing. Seminars in Respiratory and Critical Care medicine 19: 383-392

Shah S, Vanclay F, Cooper B 1989 Improving the sensitivity of the Barthel Index for Stroke rehabilitation. Journal of Clinical Epidemiology 42(8): 703-709

Silver K H C, Macko R F, Forrester L W 2000 Effects of treadmill training on gait velocity, cadence, and gait symmetry in chronic hemiparetic stroke:a preliminary report. *Neurorehabilitation and Neural Repair* 14: 65-71

Skillbeck C E, Wade D T, Hewer R L, Wood V A 1983 Recovery after stroke. *Journal of Neurology, Neurosurgery and Psychiatry* 46: 5-8

Stadler A 1979 "Birds in the cornfields" in Bozzoli B. (editor) *Labour, Townships and Protest: Studies in the Social History of the Witwatersrand*. Ravan Press

Steyn K, Fourie J, Bradshaw D 1992 The impact of chronic diseases of lifestyle and their major risk factors on mortality in South Africa. *South African Medical Journal* 82: 227-231

Sulter G, Steen C, De Keyser J 1999 Use of the Barthel Index and Modified Rankin Scale in acute stroke trials. *Stroke* 31: 1538-1541

Teixeira-Salmela L, Olney S J, Nadeau S 1999 Muscle strengthening and physical conditioning to reduce impairment and disability in chronic stroke survivors. *Archives of Physical Medicine and Rehabilitation* 80: 1211-1218

The Economist 2004 *Pocket World in Figures*. 2005 Edition. Profile Books. London

von Schroeder H P, Coutts R D, Lyden P D 1995 Gait parameters following stroke: a practical assessment. *Journal of Rehabilitation Research Development* 32: 25-31

Wade D T, Wood V A, Heller A, Maggs J Hewer R L 1987 Walking after stroke: Measurement and recovery over the first three months. *Scandinavian Journal of Rehabilitation Medicine* 19: 25-30

Wade D T, Hewer R L 1987 Functional abilities after stroke: Measurement, natural history and prognosis. *Journal of Neurology, Neurosurgery and Psychiatry* 50: 177-182

Wade D T 1992 *Measurement in Neurological Rehabilitation*. Oxford. Oxford University press

Wallner P J, Stewart A V 1994 A community approach to physiotherapy. *South African Journal of Physiotherapy* 50(3): 54

Watson M 2002 Refining the Ten-metre Walking Test for Use with Neurologically Impaired People. *Physiotherapy* 88(7): 386-397

Wilkinson P R, Wolfe C D, Warburton F G, Rudd A G, Howard R S, Ross-Russell R W, Beech R 1997 Longer term quality of life and outcome in stroke patients: Is the Barthel Index alone an adequate measure of outcome? *Quality Health Care* 6(3): 125-30

World Health Organization MONICA 1988 The World Health Organisation MONICA Project. *Journal of Clinical Epidemiology* 41: 105-114

World Health Organization 2001 *The International Classification of Functioning, Disability and Health*. World Health Organization. Geneva