

https://dspace.org/

Cardiac and cerebellar histomorphology and inositol 145trisphosphate Ip3r perturbations in adult xenopus laevis following atrazine exposure

Jaclyn Johnson; P Nkomozepi; P Opute; Ejikeme Mbajiorgu

https://hdl.handle.net/10539/35394
WOS

Downloaded from a DSpace repository. https://dspace.org/

applied sciences

MDPI

Article

Cardiac and Cerebellar Histomorphology and Inositol 1,4,5-Trisphosphate (IP_3R) Perturbations in Adult *Xenopus laevis* Following Atrazine Exposure

Jaclyn Asouzu Johnson [1,*], Pilani Nkomozepi [2], Prosper Opute [3] and Ejikeme Felix Mbajiorgu [1]

[1] School of Anatomical Sciences, University of the Witwatersrand, Johannesburg 2193, South Africa; ejikeme.mbajiorgu@wits.ac.za
[2] Department of Human Anatomy and Physiology, University of Johannesburg, Johannesburg 2006, South Africa; pilanin@uj.ac.za
[3] Department of Animal and Environmental Biology, University of Benin, Benin City 1154, Nigeria; prosper.opute@uniben.edu
* Correspondence: jaclyn.johnson@wits.ac.za; Tel.: +277-34-694-543

Abstract: Despite several reports on the endocrine-disrupting ability of atrazine in amphibian models, few studies have investigated atrazine toxicity in the heart and cerebellum. This study investigated the effect of atrazine on the unique Ca^{2+} channel-dependent receptor (Inositol 1,4,5-trisphosphate; IP_3R) in the heart and the cerebellum of adult male *Xenopus laevis* and documented the associated histomorphology changes implicated in cardiac and cerebellar function. Sixty adult male African clawed frogs (*Xenopus laevis*) were exposed to atrazine (0 µg/L (control), 0.01 µg/L, 200 µg/L, and 500 µg/L) for 90 days. Thereafter, heart and cerebellar sections were processed with routine histological stains (heart) or Cresyl violet (brain), and IP_3R histochemical localization was carried out on both organs. The histomorphology measurements revealed a significant decrease in the mean percentage area fraction of atrial (0.01 µg/L and 200 µg/L) and ventricular myocytes (200 µg/L) with an increased area fraction of interstitial space, while a significant decrease in Purkinje cells was observed in all atrazine groups ($p < 0.008$, 0.001, and 0.0001). Cardiac IP_3R was successfully localized, and its mean expression was significantly increased (atrium) or decreased (cerebellum) in all atrazine-exposed groups, suggesting that atrazine may adversely impair cerebellar plasticity and optimal functioning of the heart due to possible disturbances of calcium release, and may also induce several associated cardiac and neural pathophysiologies in all atrazine concentrations, especially at 500 µg/L.

Keywords: atrazine; Purkinje; cerebellum; myocytes; toxicology; IP_3Rs

check for updates

Citation: Asouzu Johnson, J.; Nkomozepi, P.; Opute, P.; Mbajiorgu, E.F. Cardiac and Cerebellar Histomorphology and Inositol 1,4,5-Trisphosphate (IP_3R) Perturbations in Adult *Xenopus laevis* Following Atrazine Exposure. *Appl. Sci.* **2021**, *11*, 10006. https://doi.org/10.3390/app112110006

Academic Editors: Panagiotis Berillis and Božidar Rašković

Received: 22 September 2021
Accepted: 20 October 2021
Published: 26 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

Atrazine is designed specifically to impair the physiology of weeds [1]; however, the physical and chemical properties of atrazine, such as its half-life, slow degradation [2], and solubility in different mediums, contribute to its persistence in the environment and, consequently, its contamination of ground, surface, and drinking water [3–5]. This suggests a constant presence of atrazine and its degradation products in the environment and, therefore, in drinking water and aquatic and terrestrial environments [3], with inevitable pathological consequences.

The effects of atrazine exposure have been widely reported such as impaired development in fish [6] and perturbed immunology and growth in *Xenopus laevis* [7,8]. However, most information on atrazine-induced effects in adult animal models have focused mostly on its endocrine-disrupting abilities, particularly in *Xenopus laevis* gonadal morphology [9–11]. Furthermore, atrazine and its metabolites can bio-accumulate in aquatic microbiota with adverse aquatic ecosystem implications [12] and can induce several organ

Please note that certain pages of this article have been removed in order to reduce the file size so that the PDF can be uploaded on the system (the system has a limit of 1MB for files and several PDF files are larger than this).

The first and last pages of each paper (with full bibliographic details and affiliations) are included.

If the entire unredacted paper is required, this can be emailed directly to whomever requires them by contacting Prof. Paul R. Manger on Paul.Manger@wits.ac.za

study, IP_3Rs were successfully localized in the atrium and ventricles of frog hearts, and the results showed a significant increase in atrial IP_3Rs and a non-significant increase in ventricular IP_3Rs following atrazine treatment, which may be related to the differences in function of atrial and ventricular myocytes [43,44]. Additionally, elevated IP_3R expression is reputably observed in cardiac ventricular hypertrophy, heart failure, atrial fibrillation, and dilated cardiomyopathy [45,46], which mirrors our histomorphometry observations. These elevated cardiac IP_3R levels are also suggested to compensate for waning RYRs in human heart diseases [45,47].

The observed micromorphological changes in the heart and cerebellum following atrazine treatment and IP_3R perturbations in both organs (peripheral and central toxicity) indicates atrazine's potential to disrupt cardiac and cerebellar excitability with attendant dysfunctions. These histopathological observations together suggest impaired motor agility, mild ventricular cardiomyopathy, and progressed atrial arrythmias, all of which severely impair the ability of *Xenopus* to thrive in its environment. The results further emphasize that caution should be applied to atrazine use and application and regulatory guidelines should be enforced, as significant organ toxicities are evident at the environmentally relevant concentration (0.01 µg/L), which may have grave consequences in aquatic animal life conservation, the ecosystem, and general environmental health.

Author Contributions: Conceptualization, J.A.J. and E.F.M.; methodology, J.A.J. and E.F.M.; software, P.N.; validation, J.A.J. and E.F.M.; formal analysis, J.A.J., E.F.M. and P.N.; investigation, J.A.J. and E.F.M.; resources, J.A.J. and E.F.M.; data curation, J.A.J.; writing—original draft preparation, J.A.J. and E.F.M.; writing—review and editing, J.A.J., E.F.M. and P.O.; visualization, J.A.J., E.F.M. and P.O.; supervision, E.F.M.; project administration, J.A.J. and E.F.M.; funding acquisition, J.A.J., E.F.M. and P.N. All authors have read and agreed to the published version of the manuscript.

Funding: This work was partially supported by a Faculty Research Committee Individual Grant awarded to Jaclyn Asouzu Johnson in 2015 by the University of Witwatersrand, Faculty of Health Sciences (grant number; 00125484211015121101512110500498 5).

Institutional Review Board Statement: This study was conducted according to the guidelines of the approved standard procedures of the University of the Witwatersrand Animals Ethics Research Committee (2014/32/D) and the Gauteng Province Nature Conservation (CPF6 0115 (2015) and CPF6 0120, 2016).

Data Availability Statement: The supporting data reported in this study will be made available publicly during the course of publication, if this manuscript is accepted.

Acknowledgments: The authors especially acknowledge Hasiena Ali for her hands-on assistance at various stages of this research, and Ihunwo, Cornelius Rimayi, and Lynette Sena for their amazing collaboration and inspiration.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Bai, X.; Sun, C.; Xie, J.; Song, H.; Zhu, Q.; Su, Y.; Qian, H.; Fu, Z. Effects of atrazine on photosynthesis and defense response and the underlying mechanisms in Phaeodactylum tricornutum. *Environ. Sci. Pollut. Res.* **2015**, *22*, 17499–17507. [CrossRef]
2. Ribaudo, M.; Bouzaher, A. Atrazine: Environmental Characteristics and Economics of Management. 1994. Available online: http://www.ers.usda.gov/publications/pub-details/?pubid=40594 (accessed on 21 September 2021).
3. Barchanska, H.; Sajdak, M.; Szczypka, K.; Swientek, A.; Tworek, M.; Kurek, M. Atrazine, triketone herbicides, and their degradation products in sediment, soil and surface water samples in Poland. *Environ. Sci. Pollut. Res.* **2016**, *24*, 644–658. [CrossRef] [PubMed]
4. Dabrowski, J.M.; Shadung, J.M.; Wepener, V. Prioritizing agricultural pesticides used in South Africa based on their environmental mobility and potential human health effects. *Environ. Int.* **2014**, *62*, 31–40. [CrossRef]
5. Tiryaki, O.; Temur, C. The fate of pesticide in the environment. *J. Biol. Env. Sci.* **2010**, *4*, 29–38.
6. Scahill, J.L. Effects of atrazine on embryonic development of fathead minnows (Pimephales promelas) and Xenopus laevis. *BIOS* **2008**, *79*, 139–149. [CrossRef]
7. Langerveld, A.J.; Celestine, R.; Zaya, R.; Mihalko, D.; Ide, C.F. Chronic exposure to high levels of atrazine alters expression of genes that regulate immune and growth-related functions in developing Xenopus laevis tadpoles. *Environ. Res.* **2009**, *109*, 379–389. [CrossRef]

8. Sifkarovski, J.; Grayfer, L.; Andino, F.D.J.; Lawrence, B.P.; Robert, J. Negative effects of low dose atrazine exposure on the development of effective immunity to FV3 in Xenopus laevis. *Dev. Comp. Immunol.* **2014**, *47*, 52–58. [CrossRef]
9. Hayes, T.B.; Case, P.; Chui, S.; Chung, D.; Haeffele, C.; Haston, K.; Lee, M.; Mai, V.P.; Marjuoa, Y.; Parker, J.; et al. Pesticide Mixtures, Endocrine Disruption, and Amphibian Declines: Are We Underestimating the Impact? *Environ. Health Perspect.* **2006**, *114* (Suppl. 1), 40–50. [CrossRef] [PubMed]
10. Jooste, A.M.; Du Preez, L.H.; Carr, J.A.; Giesy, J.P.; Gross, T.S.; Kendall, R.J. Gonadal development of larval male Xenopus laevis exposed to atrazine in outdoor microcosms. *Environ. Sci. Technol.* **2005**, *39*, 5255–5261. [CrossRef] [PubMed]
11. Rimayi, C.; Odusanya, D.; Weiss, J.; de Boer, J.; Chimuka, L.; Mbajiorgu, F. Effects of environmentally relevant sub-chronic atrazine concentrations on African clawed frog (Xenopus laevis) survival, growth and male gonad development. *Aquat. Toxicol.* **2018**, *199*, 1–11. [CrossRef] [PubMed]
12. Kabra, A.N.; Ji, M.-K.; Choi, J.; Kim, J.R.; Govindwar, S.; Jeon, B.-H. Toxicity of atrazine and its bioaccumulation and biodegradation in a green microalga, Chlamydomonas mexicana. *Environ. Sci. Pollut. Res.* **2014**, *21*, 12270–12278. [CrossRef] [PubMed]
13. Lin, J.; Zhao, H.-S.; Qin, L.; Li, X.-N.; Zhang, C.; Xia, J.; Li, J.-L. Atrazine Triggers Mitochondrial Dysfunction and Oxidative Stress in Quail (Coturnix C. coturnix) Cerebrum via Activating Xenobiotic-Sensing Nuclear Receptors and Modulating Cytochrome P450 Systems. *J. Agric. Food Chem.* **2018**, *66*, 6402–6413. [CrossRef]
14. Podda, M.V.; Deriu, F.; Solinas, A.; Demontis, M.P.; Varoni, M.V.; Spissu, A.; Anania, V.; Tolu, E. Effect of atrazine Administration on Spontaneous and Evoked Cerebellar Activity in the RAT. *Pharmacol. Res.* **1997**, *36*, 199–202. [CrossRef]
15. Xia, J.; Qin, L.; Du, Z.-H.; Lin, J.; Li, X.-N.; Li, J.-L. Performance of a novel atrazine-induced cerebellar toxicity in quail (Coturnix C. coturnix): Activating PXR/CAR pathway responses and disrupting cytochrome P450 homeostasis. *Chemosphere* **2017**, *171*, 259–264. [CrossRef]
16. Lin, Z.; Dodd, C.A.; Xiao, S.; Krishna, S.; Ye, X.; Filipov, N.M. Gestational and Lactational Exposure to Atrazine via the Drinking Water Causes Specific Behavioral Deficits and Selectively Alters Monoaminergic Systems in C57BL/6 Mouse Dams, Juvenile and Adult Offspring. *Toxicol. Sci.* **2014**, *141*, 90–102. [CrossRef] [PubMed]
17. Asouzu Johnson, J.; Ihunwo, A.; Chimuka, L.; Mbajiorgu, E.F. Cardiotoxicity in African clawed frog (Xenopus laevis) subchronically exposed to environmentally relevant atrazine concentrations: Implications for species survival. *Aquat. Toxicol.* **2019**, *213*, 105218. Available online: https://www.sciencedirect.com/science/article/pii/S0166445X18308099 (accessed on 16 September 2021). [CrossRef] [PubMed]
18. Amir, S.; Shah, S.; Mamoulakis, C.; Docea, A.; Kalantzi, O.-I.; Zachariou, A.; Calina, D.; Carvalho, F.; Sofikitis, N.; Makrigiannakis, A.; et al. Endocrine Disruptors Acting on Estrogen and Androgen Pathways Cause Reproductive Disorders through Multiple Mechanisms: A Review. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1464. [CrossRef]
19. Spealman, C.R. The action of ions on the frog heart. *Am. J. Physiol. Content* **1940**, *130*, 729–738. [CrossRef]
20. Fijorek, K.; Püsküllüoğlu, M.; Tomaszewska, D.; Tomaszewski, R.; Glinka, A.; Polak, S. Serum potassium, sodium and calcium levels in healthy individuals—literature review and data analysis. *Folia Med. Crac.* **2014**, *54*, 53–70.
21. Philipson, K.D.; Nicoll, D.A. Sodium-Calcium Exchange: A Molecular Perspective. *Annu. Rev. Physiol.* **2000**, *62*, 111–133. [CrossRef]
22. Bers, D.M.; Barry, W.H.; Despa, S. Intracellular Na+ regulation in cardiac myocytes. *Cardiovasc. Res.* **2003**, *57*, 897–912. [CrossRef]
23. Eisner, D.A.; Caldwell, J.L.; Kistamas, K.; Trafford, A.W. Calcium and Excitation-Contraction Coupling in the Heart. *Circ. Res.* **2017**, *121*, 181–195. [CrossRef] [PubMed]
24. Ghosh, A.; Greenberg, M.E. Calcium signaling in neurons: Molecular mechanisms and cellular consequences. *Science* **1995**, *268*, 239–247. [CrossRef] [PubMed]
25. Dellis, O.; Dedos, S.G.; Tovey, S.C.; Rahman, T.U.; Dubel, S.J.; Taylor, C.W. Ca 2+ Entry Through Plasma Membrane IP 3 Receptors. *Science* **2006**, *313*, 229–233. [CrossRef] [PubMed]
26. Dellis, O.; Rossi, A.M.; Dedos, S.G.; Taylor, C. Counting Functional Inositol 1,4,5-Trisphosphate Receptors into the Plasma Membrane. *J. Biol. Chem.* **2008**, *283*, 751–755. [CrossRef]
27. Koulen, P.; Janowitz, T.; Johnston, L.D.; Ehrlich, B.E. Conservation of localization patterns of IP3 receptor type 1 in cerebellar Purkinje cells across vertebrate species. *J. Neurosci. Res.* **2000**, *61*, 493–499. Available online: https://onlinelibrary.wiley.com/doi/abs/10.1002/1097-4547%2820000901%2961%3A5%3C493%3A%3AAID-JNR3%3E3.0.CO%3B2-9 (accessed on 21 September 2021). [CrossRef]
28. Ando, H.; Hirose, M.; Mikoshiba, K. Aberrant IP3 receptor activities revealed by comprehensive analysis of pathological mutations causing spinocerebellar ataxia 29. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 12259–12264. [CrossRef]
29. Garcia, M.I.; Boehning, D. Cardiac inositol 1,4,5-trisphosphate receptors. *Biochim. Biophys. Acta (BBA)—Bioenerg.* **2017**, *1864*, 907–914. [CrossRef]
30. Rohr, J.R.; McCoy, K.A. A qualitative meta-analysis reveals consistent effects of atrazine on freshwater fish and amphibians. *Environ. Health Perspect.* **2010**, *118*, 20. [CrossRef] [PubMed]
31. Farruggia, F.T.; Rossmeisl, C.M.; Hetrick, J.A.; Biscoe, M.; Branch, M.E.R., III. *Refined Ecological Risk Assessment for Atrazine*; US Environmental Protection Agency, Office of Pesticide Programs: Washington, DC, USA, 2016.
32. Mitrut, R.; Stepan, A.; Pirici, D. Histopathological Aspects of the Myocardium in Dilated Cardiomyopathy. *Curr. Health Sci. J.* **2018**, *44*, 243–249. [CrossRef]

33. Marshall, L.; Vivien, C.; Girardot, F.; Péricard, L.; Demeneix, B.A.; Coen, L.; Chai, N. Persistent fibrosis, hypertrophy and sarcomere disorganisation after endoscopy-guided heart resection in adult Xenopus. *PLoS ONE* **2017**, *12*, e0173418. [CrossRef] [PubMed]
34. Guo, C.; Witter, L.; Rudolph, S.; Elliott, H.L.; Ennis, K.A.; Regehr, W.G. Purkinje Cells Directly Inhibit Granule Cells in Specialized Regions of the Cerebellar Cortex. *Neuron* **2016**, *91*, 1330–1341. [CrossRef]
35. Ralcewicz, T.A.; Persaud, T. Purkinje and granule cells distribution in the cerebellum of the rat following prenatal exposure to low dose ionizing radiation. *Exp. Toxicol. Pathol.* **1994**, *46*, 443–452. [CrossRef]
36. Manzano, A.S.; Herrel, A.; Fabre, A.-C.; Abdala, V. Variation in brain anatomy in frogs and its possible bearing on their locomotor ecology. *J. Anat.* **2017**, *231*, 38–58. [CrossRef]
37. Redmond, L.; Ghosh, A. Regulation of dendritic development by calcium signaling. *Cell Calcium* **2005**, *37*, 411–416. [CrossRef]
38. Nanou, E.; Catterall, W.A. Calcium Channels, Synaptic Plasticity, and Neuropsychiatric Disease. *Neuron* **2018**, *98*, 466–481. [CrossRef]
39. Berridge, M.J.; Bootman, M.; Roderick, H. Calcium signalling: Dynamics, homeostasis and remodelling. *Nat. Rev. Mol. Cell Biol.* **2003**, *4*, 517–529. [CrossRef]
40. Rossi, M.L.; Prigioni, I.; Gioglio, L.; Rubbini, G.; Russo, G.; Martini, M.; Farinelli, F.; Rispoli, G.; Fesce, R. IP3 receptor in the hair cells of frog semicircular canal and its possible functional role. *Eur. J. Neurosci.* **2006**, *23*, 1775–1783. [CrossRef]
41. Hisatsune, C.; Mikoshiba, K. IP3receptor mutations and brain diseases in human and rodents. *J. Neurochem.* **2017**, *141*, 790–807. [CrossRef]
42. Fill, M.; Copello, J.A. Ryanodine Receptor Calcium Release Channels. *Physiol. Rev.* **2002**, *82*, 893–922. [CrossRef] [PubMed]
43. Tijskens, P.; Meissner, G.; Franzini-Armstrong, C. Location of Ryanodine and Dihydropyridine Receptors in Frog Myocardium. *Biophys. J.* **2003**, *84*, 1079–1092. [CrossRef]
44. Lipp, P.; Laine, M.; Tovey, S.C.; Burrell, K.M.; Berridge, M.J.; Li, W.; Bootman, M.D. Functional InsP3 receptors that may modulate excitation–contraction coupling in the heart. *Curr. Biol.* **2000**, *10*, 939–942. [CrossRef]
45. Signore, S.; Sorrentino, A.; Ferreira-Martins, J.; Kannappan, R.; Shafaie, M.; Del Ben, F.; Isobe, K.; Arranto, C.; Wybieralska, E.; Webster, A.; et al. Inositol 1,4,5-Trisphosphate Receptors and Human Left Ventricular Myocytes. *Circulation* **2013**, *128*, 1286–1297. [CrossRef] [PubMed]
46. Go, L.O.; Moschella, M.C.; Watras, J.; Handa, K.K.; Fyfe, B.S.; Marks, A.R. Differential regulation of two types of intracellular calcium release channels during end-stage heart failure. *J. Clin. Investig.* **1995**, *95*, 888–894. [CrossRef] [PubMed]
47. Harzheim, D.; Movassagh, M.; Foo, R.S.-Y.; Ritter, O.; Tashfeen, A.; Conway, S.J.; Bootman, M.D.; Roderick, H.L. Increased InsP3Rs in the junctional sarcoplasmic reticulum augment Ca2+ transients and arrhythmias associated with cardiac hypertrophy. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 11406–11411. [CrossRef] [PubMed]