

https://dspace.org/

A comparison between a mobility programme alone and a standard physiotherapy rehabilitation approach, on the functional outcome of patients following primary total hip arthroplasty.

Naidoo, Umantha

2007-04-10T07:51:44Z
http://hdl.handle.net/10539/2350

Downloaded from a DSpace repository. https://dspace.org/

TABLE OF CONTENTS

Page

Declaration i

Dedication ii

Abstract iii

Acknowledgements iv

List of Tables vii

List of Abbreviations viii

Chapter 1 Introduction 1

Chapter 2 Literature Review:

1. Introduction 5
2. Primary THA as a treatment tool: 6
   - 2.1. The indications for THA
   - 2.2. Types of THA Prosthesis
   - 2.3. Postoperative complications:
     - 2.3.1. Early complications 9
     - 2.3.2. Late complications 11

3. The postoperative physiotherapy management of THA patients:
   - 3.1. Education of THA patients 12
   - 3.2. Mobility regimens and exercise therapy as part of the postoperative rehabilitation process 13

4. Outcome measures: Evidence for use of chosen assessment tools used to evaluate outcome measures:
4.1. Visual Analogue Scale (VAS) 20
4.2. Iowa Level of Assistance (ILOA) Scale 21
4.3. Goniometry as a tool for measuring hip range of motion 22

Chapter 3 Method 24

Chapter 4 Results 35

Chapter 5 Discussion 45

Chapter 6 Conclusion 51

References 52

Appendix A: Ethical clearance 57
Appendix B: Patient information sheet 58
Appendix C: Patient consent form 59
Appendix D: ILOA Scale 60
Appendix E: Pre-operative patient data collection sheet 61
Appendix F: Post-operative patient data collection sheet 62
Appendix G: Instructional Handout to Patients 63
Appendix H: Instructional Handout to Physiotherapists 64