

DOES THE USE OF MICROPLEGIA SIGNIFICANTLY IMPROVE OUTCOMES IN CONGENITAL HEART SURGERY?

Sharmel Bhika^{1,*}, Geoffrey Candy² Krubin Naidoo³

1. Department of Cardiothoracic Surgery, Charlotte Maxeke Johannesburg Academic Hospital, Johannesburg, South Africa
2. Department of Surgery, University of Witwatersrand, Johannesburg, South Africa
3. Department of Cardiothoracic Surgery, Nelson Mandela Children's Hospital, Johannesburg, South Africa

* Corresponding author: Department of Cardiothoracic Surgery, Charlotte Maxeke Johannesburg Academic Hospital, Jubilee Road, Johannesburg, 2196, South Africa. Tel: +27 82 485 9923; e-mail: sharmel@vodamail.co.za (S. Bhika).

Clinical registration: International Standard Randomised Controlled Trial Number (ISRCTN) registry: 12983759

VISUAL ABSTRACT

KEY QUESTION

Does the use of Microplegia significantly improve outcomes in congenital heart surgery?



KEY FINDINGS

Microplegia showed lower biochemical markers; less inotropic support required and quicker myocardial recovery.




TAKE-HOME MESSAGE

Significantly improved short-term postoperative outcomes were found in all clinical and biochemical parameters measured in the Microplegia sample group.

49

CENTRAL IMAGE

50



51

52

ABSTRACT

Objectives: The aim of this prospective randomized study is to compare short-term postoperative outcomes of elective congenital cardiac patients undergoing corrective cardiac surgery. A comparison of clinical and biochemical findings were assessed between two cardiac arresting agents, namely all-blood Microplegia and crystalloid St Thomas' Hospital Cardioplegic II solution.

Methods: Forty-two patients were prospectively randomised prior to corrective cardiac surgery to two techniques of myocardial protection: group I tepid (28°C) Microplegia (MPS) and group II cold (4°C) St Thomas' Hospital Cardioplegic Solution II (STS) crystalloid cardioplegia. Perioperative data was prospectively collected. Endpoints included serial measurements of Troponin I, haemoglobin/ haematocrit and lactate; time taken for spontaneous return to normal sinus rhythm; and need for postoperative inotropic support.

Results: Surgery resulted in increased immediate post-aortic unclamping myocardial lactate and troponin levels in both groups, however, significantly lower lactate (2.36 ± 0.76 vs 4.19 ± 1.09 mmol/l, $p < 0.0001$) and Troponin I levels (8.18 ± 3.68 vs 22.88 ± 13.39 ng/ml, $p < 0.0001$) were observed in Group I vs Group II. Improved myocardial contractility as measured by quicker spontaneous return to normal sinus rhythm and reduced perioperative requirement for inotropic support as well as reduced need for blood transfusion was evident in Group I.

Conclusions: Microplegia showed significantly less myocardial injury, haemodilution and reduced need for inotropic support in the first 24 hours post congenital cardiac surgery.

Keywords: myocardial protection; cardioplegia; cardiopulmonary bypass; congenital cardiac surgery

81

82 **INTRODUCTION**

83 Myocardial protection is critical during the surgical repair of almost all congenital heart
84 surgery requiring cardiopulmonary bypass. The goal of myocardial protection is to minimize
85 myocardial metabolism during ischaemia thus preserving myocardial function in the relaxed
86 arrested heart, whilst facilitating cardiac repair in a bloodless surgical field. The basic
87 principles of myocardial protection include: rapid diastolic cardiac arrest, hypothermia to
88 decrease myocardial oxygen consumption, and avoidance of myocardial oedema [1].

89

90 Innovations in techniques of myocardial preservation have allowed improvements in
91 myocardial protection and subsequent outcomes following congenital heart surgery. Low
92 cardiac output syndrome incurred as a result of perioperative myocardial injury is still the
93 commonest cause of morbidity and mortality in cardiac surgery [2]. Currently, many options
94 may be considered when choosing a myocardial preservation strategy. Arresting the heart to
95 provide myocardial preservation from ischaemia and reperfusion can be achieved using a
96 diversity of cardioplegic solutions.

97

98 Changes in the myocardial metabolic status occur during the vulnerable ischaemic period and
99 upon reperfusion following removal of the aortic cross-clamp. It is this metabolic status that
100 largely determines the functional recovery of the myocardium [3]. Myocardial tissue acidosis
101 resulting in lactate production is one of the most sensitive markers of inadequate myocardial
102 preservation [4]. It is well established that the persistence of peripheral lactate release during
103 cardiac reperfusion is an independent predictor of postoperative low cardiac output syndrome
104 [5]. Further techniques that have been validated in assessing the degree of myocardial injury
105 are measurements of specific metabolites or biochemical markers, such as cardiac troponin
106 and serum creatine kinase MB isoenzyme (CK-MB) [6-8]. This is done by direct catheter
107 sampling from the coronary sinus (venous drainage of the myocardium) post removal of the

aortic cross-clamp. This evaluation of myocardial metabolism during cardiac surgery, therefore allows the investigator to quantify the degree of physiologic damage. Serum troponin I has been recognised to be a highly sensitive and specific marker of myocardial injury, and thereby reflects the adequacy of intraoperative myocardial protection [4,8,9]. Furthermore, postoperative troponin I prognosticates long-term cardiovascular events at follow-up [10,11]. Both lactate and troponin I release, can therefore explain the efficacy of myocardial preservation [4].

A cornerstone of cardiac surgical practice has been cold crystalloid potassium-based cardioplegia which was introduced in the early 1950's. This early cardioplegic technique to initiate and maintain cardiac arrest during heart surgery was soon followed by the use of intermittent cold blood cardioplegia [12]. Subsequently, blood cardioplegia has become the most popular choice amongst surgeons in the United States of America (USA) [13] and is being used increasingly in both Europe and the United Kingdom. All-blood, potassium-enriched cardioplegia, called microplegia or miniplegia, was originally described by Menasche *et al* in 1996 as a safe and effective alternative to blood or crystalloid cardioplegia [14]. Conventionally, blood cardioplegia is composed of a blood:crystalloid mixture (commonly in the ratio of 4:1 or 8:1), additives, and a potassium-rich solution. Microplegia is reconstituted by combining small volumes of concentrated additives (66:1) to blood from the cardiopulmonary bypass circuit, including a potassium-rich solution, thereby allowing a large volume of blood cardioplegia to be delivered with relatively minimal amounts of crystalloid. In our practice, we observed upon release of the aortic crossclamp and reperfusion of the ischaemic heart, a quicker recovery of myocardial activity and decreased requirement for post-operative inotropic support following Microplegia. This observation initiated the current investigation to formally compare short-term patient outcomes of these two cardioplegic solutions used routinely in our centre.

134

135 The aim of this randomised study was to determine short-term patient outcomes using two
136 different compositions of cardioplegia used in our institution during congenital cardiac
137 surgery. The study was to answer the question: Does myocardial protection with microplegia
138 in paediatric congenital open heart surgery significantly reduce the perioperative myocardial
139 insult and improve the postoperative haemodynamic (need for inotropic support) status. The
140 objectives were to measure myocardial ischaemic injury as reflected by time taken for
141 spontaneous return to normal sinus rhythm and the need for postoperative inotropic support
142 and the need for postoperative blood transfusion as reflected by postoperative haemoglobin
143 measurements.

144

145 **MATERIALS AND METHODS**

146 **Patients and Study design**

147 The study was approved by the Human Research Ethics Committee (Medical) of the
148 University of Witwatersrand, Johannesburg, South Africa and conducted at the Charlotte
149 Maxeke Johannesburg Academic Hospital, South Africa. Preoperative consent for the study
150 and procedure was obtained from the parents / legal guardians of the patients for surgery in a
151 written and signed document.

152

153 **Patients:** Patients referred from surrounding cardiology clinics for elective corrective
154 congenital heart surgery, participated in the study.

155 **Inclusion criteria:** Patients of either gender and of body weight 5-20kg, with one of the
156 three most commonly seen congenital heart defects, namely Tetralogy of Fallot, ventricular
157 septal defect or atrioventricular septal defects, of varying complexity, were included.

158 **Exclusion criteria:** Neonates, patients with interstitial lung disease or with poor preoperative
159 ventricular function (EF < 50%), inherent bleeding disorders (eg Di George Syndrome,

haemophiliacs, Von Willebrands disease), more than one cardiopulmonary bypass run, or failure to separate from cardiopulmonary bypass / Extra Corporeal Life Support (ECLS), preoperative liver disease and redo surgery were excluded from the study. Neonates were also excluded from the study.

Study design:

During 2016, a pilot study of forty-two elective paediatric congenital cardiac patients were enrolled in a single-centre randomised study comparing outcomes using two different cardioplegia solutions, each administered at different temperatures. The two randomised groups of cardioplegia were as follows:

1. Microplegia (MPS); n=21, delivered to the aortic root with the Quest MPS system (Integrated Myocardial Protection System-2, Quest Medical Inc; Allen TX, US). Ratio blood to crystalloid > 60:1. Temperature: tepid, 28°C
2. St Thomas' Hospital Cardioplegic Solution II (STS); n=21, Adcock Ingram Critical Care (Pty) Ltd; Johannesburg. Crystalloid cardioplegia. Temperature: cold, 4°C.

Both cardioplegia solutions are used in our current practice in congenital cardiac surgery. See Table 2 for cardioplegia solution composition.

Computer randomisation and cardioplegia solution assignment was done by an unbiased person with no involvement or interest in the outcome of the study. Only one consultant surgeon and one registrar operated on these cases to keep the operating technique constant. The surgeons were blinded to the cardioplegia solution used for the operation until administration of the initial dose after the aorta was cross-clamped. At this point, the different colour of the Microplegia (sanguinous) and crystalloid (asanguinous) solutions would identify the solution being used. With the exception of the perfusionists and surgeons, all other staff (Anaesthetists, biochemists, cardiologists, nursing staff, intensivists) dealing with the biological sampling, perioperative care, and/or collection and analysis of data were

blinded to the randomisation assignment of the patient. Perioperative data was collected on a study datasheet by a nonbiased, blinded member of the surgical team.

Operative Technique

Both anaesthesia and surgery employed routine standard procedures to correct these defects in all cases. Surgery was performed through a median sternotomy. The standard management of cardiopulmonary bypass (CPB) was used - Aorto bi-caval cannulation, systemic cooling, snaring of the cavae, aortic crossclamping followed by antegrade administration of cardioplegia. The CPB circuit included a Medtronic phosphorylcholine-coated paediatric/infant custom pack tubing set, Terumo roller pump and a hollow-fiber membrane, coated oxygenator incorporating a 40 micron filter. Heparin was administered at a dose of 300 IU/kg to attain a target activated clotting time of 400 seconds or above prior to initiating CPB. For this study the extracorporeal circuit was always primed with a fresh frozen plasma to leucodepleted homologous packed red cell solution reconstituted in a 1:1 ratio and 100mg/kg IU of heparin. A non-pulsatile CPB flow was established at 2.4 L/min/m². Patients were systemically cooled to mild-moderate hypothermia (28-32 °C) at the time of cardioplegic arrest. Cardiac arrest was accomplished by antegrade administration of the hyperkalaemic cardioplegia solutions via a cannula placed in the ascending aorta, proximal to both the aortic crossclamp and the aortic arterial cannula. The cardioplegia was delivered at an induction arresting dose of 30 ml/kg and maintenance doses of 10 ml/kg given repeatedly at 20 minute intervals at an average aortic root pressure of 80-100 mmHg. The volume of the doses for both cardioplegia solutions being compared were the same, only the compositions and temperatures differed. As each maintenance dose of cardioplegia was administered, topical ice slush was applied to the anterior heart surface.

Study End points

End points included:

1. Serial (x5) biochemical measurements: Lactate + Troponin I + Haemoglobin level

blood sampling:

- In theatre preoperatively (T0);
- In theatre 10 minutes post removal of aortic crossclamp (T1);
- 6 hours post crossclamp removal (T2);
- 12 hours post crossclamp removal (T3);
- 24 hours post aortic unclamping (T4).

The lactate and haemoglobin levels were measured from arterial blood gases and the troponin I levels were measured with iStat (Point of care) cartridges.

2. The time of intraoperative spontaneous recovery to normal sinus rhythm (SRNSR) after unclamping of the aorta. The perfusionists had timers running from the time of unclamping and observations for sinus rhythm were made by the minute.

3. The requirement for inotropic support in the form of intravenous infusions was categorized as:

Low: Dobutamine $\leq 2,5\mu\text{g/kg/min}$

Medium: Dobutamine $> 2,5\mu\text{g/kg/min} \leq 5\mu\text{g/kg/min}$

High: Dobutamine $> 5\mu\text{g/kg/min}$ and/or addition of adrenaline and/or phenylephrine.

The attending anaesthetists practice is to introduce inotropic support routinely post unclamping of the aorta, irrespective of whether or not the patient requires additional support of cardiac contractility. The study therefore assessed the need for inotropic support as follows:

- On admission to ICU (T1)
- At 10am the following morning (T2).

Coronary Sinus Sampling

Coronary sinus blood (venous drainage of the myocardium) sampling was done through the right atrium (the right atrium was routinely opened during the corrective procedures in this trial). This venous blood was aspirated 10 minutes after the aorta was unclamped, allowing the heart to eject and perfuse the coronaries with oxygenated blood while flushing residual cardioplegia from the coronary system. Aspiration was done with an Argyle Polyurethane Umbilical catheter size 5Fr and this sample specimen was analysed for lactate and troponin I levels which should reflect metabolic parameters for myocardial injury during the reperfusion state.

Data Handling and Statistical Comparison

Data was recorded in EXCEL (Microsoft) and analysed using Statistica V8. Data has been reported in tables and graphs as frequency (n) and as means ($\pm$ standard deviation or standard error of the mean) or median and range. Frequencies in each group were compared with a Chi-squared test with a Fischer Exact test if $n < 5$. Comparison between the study groups was made using a t-test if the data was normally distributed. Statistical differences between groups where the data was not normally distributed used a Mann Whitney test. Serial measurements were compared using repeated measures ANOVA. A p-value of < 0.05 was regarded as being statistically significant.

RESULTS

Patients

Table 1 shows the demographic and perioperative data. No significant differences were observed between any of the variables. Post-randomisation, no patient was excluded from the study. All patients enrolled completed the study and there were no deaths. None of the patients had post-operative renal failure or a severe cardiac event requiring resuscitation. None of the patients had residual lesions detected after the initial repair on trans-oesophageal echocardiography in either group, and no patient required cardiopulmonary bypass to be re-instituted intraoperatively.

Biochemical results

The troponin I levels collected at several sampling times are presented in Figure 2. The preoperative levels were not elevated in both groups. Surgery was associated with significant perioperative troponin leak in both groups ($p < 0.0001$ for both groups) which is reflected by increased T1 values. However, coronary sinus sampling (T1) showed significantly lower mean troponin I ($\pm$ SD) values with MPS compared to STS (8.18 ± 3.68 vs 22.88 ± 13.39 ng/ml, $p < 0.0001$; Fig. 2.). The levels remained significantly lower throughout the 24 hour postoperative time-course with a gradual reduction in troponin levels seen in both groups. Troponin I level was elevated 10 minutes (T1) and 6 hours (T2) post aortic unclamping for both groups (Fig. 2.). The troponin I level for the STS group was significantly higher (highest reading for one patient was >50 ng/ml) than the MPS group (highest reading for one patient was 15.3 ng/ml) at these times ($p < 0.0001$). T3 and T4 show gradual decreases in these serial troponin measurements which were not significantly different from T1 and T2.

Surgery resulted in increased immediate post aortic crossclamp removal myocardial lactate levels (T1) in both groups. Furthermore, significant differences were shown in higher mean

lactate levels ($\pm$ SD) for the STS group (2.36 ± 0.76 vs 4.19 ± 1.09 mmol/l, $p < 0.0001$). This significant difference continued during the postoperative (first 24 hours) period (1.074 ± 0.56 vs 1.89 ± 0.54 mmol/l, $p < 0.0001$; Fig. 3.).

Preoperative haemoglobin levels readings (T0) were not significantly different (Fig. 4.). Haemoglobin levels dropped significantly for both groups between preoperative haemoglobin levels and T1 readings (specimens taken 10min post aortic unclamping) and in the STS group showed significantly lower haemoglobin values (9.6 ± 1.0 vs 8.2 ± 0.06 g/dl, $p < 0.0001$) suggesting possible haemodilution in this group and need for blood transfusion.

Spontaneous return to normal sinus rhythm

Following reperfusion, the time of spontaneous return to normal sinus rhythm (SRNSR) in the MPS group was significantly quicker than those of the STS group (1.14 ± 0.35 vs 3.33 ± 0.57 seconds, $p = < 0.0001$) (Fig.5.). RNSR in < 60 seconds occurred in 85% of the MPS group vs 0% of the STS group for this time interval. These results suggesting that the immediate myocardial protective effect of MPS solution is superior to that of St. Thomas crystalloid cardioplegia. Only 1 out of the 42 patients in the study was paced until day 1 postoperatively, this patient was in the STS group which contributed no statistical difference in the comparison ($p = 0.8$).

Haemodynamic results

The need for inotropic support for the first postoperative night proved to be more prevalent in the STS group ($p = 0.0022$), with the majority of cases in the STS group showing the need for inotropic support $> \text{Dobutamine } 2.5 \mu\text{g/kg/min}$ even on day 1 post op review (T2). The trend for both groups demonstrated a reduced need in inotropic support from T1 to T2. (Fig. 6.)

DISCUSSION

Currently, extrapolations from findings of comparative studies in myocardial protective strategies using different cardioplegia solutions in adult and animal population groups have erroneously been ascribed to the paediatric population. However critical differences in subcellular structural and functional organization as well as metabolic substrate utilization differentiate the adult from the paediatric models. Even today, in congenital heart surgery requiring CPB, the single most common cause of postoperative morbidity and mortality is perioperative myocardial injury [2]. To our knowledge there are no prospective clinical studies published comparing the efficacy of traditional crystalloid cardioplegia to all-blood Microplegia in paediatric congenital cardiac surgery.

In neonates, the enhanced tolerance of the immature myocardium to hypoxia is partially explained by the biochemical differences in energy production between immature and mature myocardium. As well as long chain fatty acids (predominant source), the adult heart also uses lipids, carbohydrates and amino acids to fuel metabolism, in contrast, the foetal heart relies more on anaerobic metabolism. At present, the bulk of evidence available points to the neonatal myocardium being considerably more resistant to ischaemia than the mature myocardium [2,15]. Neonatal patients were therefore excluded in this study.

Most European centers use crystalloid cardioplegia, while most North American surgeons use blood cardioplegia to provide additional oxygen substrate. In a North American multi-institutional survey, 86% of surgeons used blood-based cardioplegia (of which 5% used Microplegia) as their preferred cardioplegia regime in paediatric cardiac surgery, whilst only 14% of surgeons in northern USA used crystalloid cardioplegia solutions [16]. Microplegia relies on a precision pump to administer specific volumes of additives within the cardioplegia solution while minimizing the quantity of crystalloid delivered to the patient. Administration of less crystalloid should avoid volume overload and minimize haemodilution, which may

lead to decreased blood product use. Additionally, the increased number of red blood cells, thus haemoglobin molecules present in MPS, should improve oxygen delivery. The combination of lower crystalloid volume and improved oxygen delivery should, in theory, lead to reduced myocardial oedema and more rapid and improved ventricular function [14]. There is clear evidence that myocardial oedema negatively impacts both systolic and diastolic ventricular function. A previous study in Yorkshire pigs showed that the detrimental consequences of haemodilution, volume overload and blood transfusions may also be avoided with Microplegia [17]. Starr et al. [18] further showed that a lower osmolarity cardioplegia solution is associated with higher myocardial water content and impaired diastolic filling. One goal of any myocardial protection strategy should be to minimize myocardial oedema. Although interstitial myocardial water content was not measured in this study, the detrimental effects of myocardial oedema, mainly delayed intraoperative spontaneous return to normal sinus rhythm and an increased need for postoperative inotropic support showed significantly superior results in the MPS group. Intraoperative spontaneous return to normal sinus rhythm is typically considered an indicator of poor myocardial preservation. Additionally, low cardiac output syndrome is frequent during the first hours post cardiac surgery, however, this occurrence reflects a relatively delayed degree of myocardial recovery. The minimization of the ischaemic insult resulted in improved 24-hour postoperative ventricular function as evidenced by the significantly lower inotropic requirement and lactate measurements in the MPS group.

Postoperative lactate levels, particularly the measurement taken ten minutes after removing the aortic crossclamp, correlates with ischaemic myocardial acidosis that occurs in the arrested heart. The significantly elevated lactate levels of the STS group in comparison with the MPS group shows clear evidence of the resulting superior myocardial preservation of the latter.

374

375 The significant difference in haemoglobin levels at T1 (10 minutes post aortic crossclamp
376 removal) and T2 (6 hours post crossclamp removal) demonstrates that the STS group of
377 patients likely underwent more significant perioperative haemodilution as compared with the
378 MPS group. T3 and T4 haemoglobin levels of STS were higher than MPS and this could
379 have either been due to excessive blood transfusion or over-diuresis of the STS group.
380 Consequently, there may be a tendency for increased blood transfusion in the STS group
381 owing to the immediate postoperative reduced haemoglobin levels.

382

383 Cardiac troponin I, a regulatory protein unique to myocardium, reflects the quality of
384 perioperative myocardial preservation [19]. Mildly elevated perioperative troponin I levels
385 are of no prognostic significance and are expected owing to intraoperative cardiac
386 manipulation and surgical trauma to cardiac tissue. Onorati et al [20] following sampling
387 from the coronary sinus during reperfusion, demonstrated the predictive role of troponin I on
388 hospital outcome as well as postoperative left ventricular recovery post adult coronary
389 surgery. Even though this study was on patients with unstable angina, our results reflect
390 similar findings of significantly reduced perioperative myocardial damage in patients
391 subjected to MPS.

392

393 Hypothermia during cardiopulmonary bypass exerts its protective effect by multiple
394 mechanisms; the most obvious mechanism being the reduction of both metabolic rate and
395 oxygen consumption[21]. The optimal cardioplegia temperature for myocardial protection is
396 controversial but has been previously [studied](#). It is well established, however that crystalloid
397 cardioplegia should be used at temperatures lower than the tepid (27- 29°C) temperature used
398 in some studies which show inferior results. In canine models Magovern et al. [22]

demonstrated equivocal outcomes in myocardial protection regardless of whether blood cardioplegia at 20°C or crystalloid cardioplegia at 4°C was administered, however, reduced recovery was observed with crystalloid at 20°C versus blood at 4°C. Comparisons between warm blood cardioplegia (37°C) against tepid blood cardioplegia show no clear advantage of the one over the other. Buckberg and colleague's results [23] suggested that there are very slight differences in oxygen consumption when the temperature is reduced from 37°C to 29°C. The relationship, however, between myocardial wall tension and myocardial oxygen consumption (MvO_2) in the context of hypothermia is considered to be optimal at 28°C [23]. Hayashida et al. [24] determined the ideal blood cardioplegia temperature and concluded that antegrade cardioplegia administered at a tepid (29°C) temperature was associated with reduced myocardial injury resulting in optimal myocardial preservation. Similarly other studies have confirmed the benefit of tepid blood cardioplegia in comparison with cold blood cardioplegia, with regard to reduced myocardial damage, a quicker recovery of left ventricular function and even improved clinical outcomes [24-28]. In our study, the MPS group which showed results of superior myocardial preservation may have confirmed or even optimized the benefits of MPS when use at a tepid temperature.

Patients with congenital heart disease often have to adapt with the effects of volume overload, cyanosis or ventricular hypertrophy. Once subjected to cardiac surgery using cardiopulmonary bypass and ischaemic cardiac arrest, these pathologies become more concerning and thus, an optimal myocardial protective strategy would need to be chosen with consideration. The conclusion that the subendocardium [29] and hypertrophied ventricles [9] may be more predisposed to ischemic injury implies that the imbalance between myocellular energy and oxygen supply relative to consumption is the predominant mechanism of post-ischaemic myocardial dysfunction. It is, therefore, imperative that especially these patients would benefit from any refinements in myocardial preservation regarding recovery of

ventricular contractility, optimal cardioplegic temperature and prevention of ischaemic injury.

Study limitations

The time of assessment and data analysis only reflects outcomes of short-term myocardial recovery. The trend of myocardial recovery after 24 hours was seen in all patients subjected to both cardioplegia solutions. A longer term analysis is required to determine conclusively if any cardioplegia solution really shows significant benefit.

The fact that two different temperatures of cardioplegia were used, may have introduced a confounding variable. It is difficult to determine if temperature or cardioplegia solution beneficially affected the study end-points.

It remains to be determined if the reduced haemodilution with MPS further translates into the reduction of myocardial oedema upon coronary reperfusion.

Peri-operative echocardiography would more accurately determine left and right ventricular systolic and diastolic function.

Although blinded to the study groups, the different anaesthesiology teams varied according to their practice preference which may have introduced variation in the administration of fluid management, transfusion triggers observed and thresholds for introducing and/or escalating inotropes. These may have introduced confounding variables into the study.

In summary, all-blood MPS resulting in the administration of less crystalloid, may have decreased myocardial oedema to permit more rapid recovery of ventricular function in surgery to correct congenital heart defects in paediatric patients. The detrimental consequences of haemodilution, volume overload and potential blood transfusions may also be avoided. The key findings indicate that serum troponin, lactate and haemoglobin measurements intraoperatively, 6 hours, 12 hours and 24 hours post aortic crossclamp

removal in the MPS sample group were all significantly lower than crystalloid St Thomas Hospital's Cardioplegic Solution II. Furthermore, the MPS group required less inotropic support and myocardial recovery to spontaneous normal sinus rhythm was significantly quicker. This comparative study has clearly shown significant short-term (over 24 hours) clinical and biochemical superior outcomes in our MPS group. These convincing results has directed these investigating surgeons towards the routine use of Microplegia in their cardioplegic myocardial protection strategy in paediatric cardiac surgery.

Acknowledgements

Cossni Medical Pty Ltd (Distributors of Quest Medical MPS) sponsored the Argyle Polyurethane Umbilical size 5Fr catheters used for coronary sinus blood sampling.

Funding

None declared.

Conflict of Interest

None declared.

FIGURES




Figure 1. Methodology for the prospective study comparing STS to MPS in congenital cardiac surgery.





Figure 2. Time course of troponin I in the peri- and post-operative repair of congenital heart defects. Differences ($p < 0.0001$) are between patients randomised to MPS (solid line) or STS (-----). Times T0: in theatre preoperatively (T0); T1: in theatre 10 minutes post removal of aortic crossclamp; T2: 6 hours post crossclamp removal; T3: 12 hours post crossclamp removal; T4: 24 hours post aortic unclamping. Data as means $\pm$ SEM.

504



505

506

507

508


Figure 3. Time course of lactate in the peri- and post-operative repair of congenital heart defects. Differences ($p < 0.0001$) are between patients randomized to MPS (solid line) or STS (-----). Times as given in Fig 2.

509

510

511

512




513

514

515 **Figure 4.** Time course of haemoglobin in the peri- and post-operative repair of congenital
 516 heart defects. Differences ($p < 0.0001$) are between patients randomised to MPS (solid line) or
 517 STS (-----). Times as given in Fig2.

518

519
520



521
522

523 **Figure 5.** Time take to achieve sinus rhythm following cardioplegia with MPS compared to
524 STS solution. Median values shown as circles and the lines as the range.

525
526




Figure 6. Requirement for inotropic support for the first postoperative night (T1) and day 1 post op review (T2) following cardioplegia with MPS and STS solutions (see text for further detail). Data as median and range.

TABLES

Table 1 – Patient demographic and perioperative variables. Data as mean $\pm$ STD and as median [range] for skewed data.

Variable	All (n=42)	Group 1 Tepid MPS (n=21)	Group 2 Cold STS (n=21)	Statistical difference between groups p value
Weight (kg)	10.1 $\pm$ 4.0	10.0 $\pm$ 4.0	10.1 $\pm$ 4.0	0.93
Gender (male/female)	21/21 (50% female)	10/11	11/10	0.76
Body surface area (m ²)	0.50 $\pm$ 0.16	0.48 $\pm$ 0.15	0.51 $\pm$ 0.16	0.56
Aortic cross clamp time (min)	72.9 $\pm$ 46.0 60 [19-229]	69.0 $\pm$ 44.4 57 [31-229]	76.8 $\pm$ 48.2 65 [19-228]	0.41
CPB time (min)	118.6 $\pm$ 55.1 100.5 [48.0-316]	115.3 $\pm$ 49.4 107 [66-292]	121.9 $\pm$ 61.4 100 [48-316]	0.89
Volume of cardioplegia (ml)	673 $\pm$ 512 550 [165-3300]	613 $\pm$ 335 520 [165 – 1790]	732 $\pm$ 641 600 [270-3300]	0.82
Diagnosis: 1. Tetralogy of Fallot (n) 2. Ventricular septal defect (n) 3. Atrioventricular septal defect (n)	7 31 4	2 18 1	5 13 3	0.21

Abbreviations : CPB - Cardiopulmonary bypass; MPS – Microplegia; STS - St Thomas' Hospital Cardioplegic Solution II.

547 **Table 2.** Preparation/Composition of cardioplegia solutions being compared.

		Induction dosage	Maintenance dosage
Administered volume	Solution composition	30ml/kg bolus	10ml/kg given at 20min intervals
St Thomas Solution (n=21) Temperature: 4°C			
Composition	KCl (mmol/L)	16	16
	NaHCO ₃ (mmol/L)	10	10
	MgCl (mmol/L)	16	16
	CaCl ₂ (mmol/L)	1,2	1,2
Microplegia (n=21) Temperature: 28°C			
Composition	KCl (mEq/L)	30	15
Additives (50ml)	(ml/L)	15	7
	MgSO ₄	5g/10ml	5g/10ml
	Lidocaine 2%	50mg/5ml	50mg/5ml
Blood priming solution: 35ml			

548

549

550

551

552

553

554

REFERENCES

1. Buckberg GD. A proposed “solution” to the cardioplegic controversy. *J Thorac Cardiovasc Surg.* 1979;77:803–815.
2. Castaneda AR, Jonas RA, Mayer JE, Hanley FL. Myocardial preservation in the immature heart. Ch. 3. In: *Cardiac surgery of the neonate and infant*, ed by Castaneda AR, Jonas RA, Mayer JE, Hanley FL. Published by WB Saunders Philadelphia, 1994 pp.41-54.
3. Khabbaz KR, Zankoul F, Warner KG. Intraoperative metabolic monitoring of the heart: online measurement of myocardial tissue pH. *Ann Thorac Surg* 2001;72:2227–34.
4. Onorati F, Renzulli A, De Feo M, Santarpino G, Gregorio R, Biondi A, *et al.* Does antegrade cardioplegia alone provide adequate myocardial protection in patients with left stem main disease? *J Thorac Cardiovasc Surg* 2003;126(5):1345–51.
5. Rao V, Ivanov J, Weisel R, Cohen G, Borger MA, Mickle DA. Lactate release during reperfusion predicts low cardiac output syndrome after coronary bypass surgery. *Ann Thorac Surg* 2001;71(6):1925–30.
6. Siouffi SY, Kwasnik EM, Khouri SF. Methods for the metabolic quantification of regional myocardial ischemia. *J Surg Res* 1987;43:360 –78.
7. Vijay P, Szekely L, Aufiero TX, Sharp TG. Coronary sinus adrenomedullin rises in response to myocardial injury. *Clin Sci* 1999;96:415–20.
8. Koh TW, Hooper J, Kemp M, Ferdinand FD, Gibson DG, Pepper JR. Intraoperative release of troponin T in coronary venous and arterial blood and its relation to recovery of left ventricular function and oxidative metabolism following coronary artery surgery. *Heart* 1998;80(4):341-8.
9. Hottenrott CE, Buckberg GD, Bugyi HI, Kurkji HJ, Towers B, Maloney JV Jr. Myocardial ischemia in hypertrophied ventricles following ventricular fibrillation during

cardiopulmonary bypass. *Bull Soc Int Chir* 1974;33:261-8.

10. Mair J, Larue C, Mair P, Balogh D, Calzolari C, Puschendorf B. Use of cardiac troponin I to diagnose perioperative myocardial infarction in coronary artery bypass grafting. *Clin Chem* 1994;40:2066–70.

11. Fellahi JL, Gué X, Richomme X, Monier E, Guillou L, Riou B. Short and long-term prognostic value of postoperative cardiac Troponin I concentration in patients undergoing coronary artery bypass grafting. *Anesthesiology* 2003;99(2):270–4.

12. Follette DM, Mulder DG, Maloney JV, Buckberg GD. Advantages of blood cardioplegia over continuous coronary perfusion or intermittent ischemia. *J Thorac Cardiovasc Surg* 1978;76:604-19.

13. Robinson LA, Schwarz GD, Goddard DB, Fleming WH, Galbraith TA. Myocardial protection for acquired heart disease surgery: results of a national survey. *Ann Thorac Surg*. 1995;59(2):361-72.

14. Menasche P. Blood cardioplegia: do we still need to dilute? *Ann Thoracic Surg* 1996;62:957-960.

15. Hammon JW Jr. Myocardial protection in the immature heart. *Ann Thoracic Surg* 1995;60(3):839-42.

16. Kotani, Y, Tweddell J, Gruber P, Pizarro C, Austin EH, Woods EK, *et al.* Current cardioplegia practice in paediatric cardiac surgery: A north American multi-institutional survey. *Ann Thorac Surg*. 2013;96:923-9.

17. McCann UG II, Lutz CJ, Picone AL, Searles B, Gatto LA, Dilip KA, *et al.* 2006. Whole blood cardioplegia (minicardioplegia) reduces myocardial edema after ischemic injury and cardiopulmonary bypass. *J Extra Corpor Technol.* 38:14–21.

18. Starr JP, Jia C, Amirhamzeh MMR, Rabkin DG, Hart JP, Hsu DT, *et al.* Coronary perfusate composition influences diastolic properties, myocardial water content, and histologic characteristics of the rat left ventricle. *Ann Thorac Surg* 1999;68:925–30.
19. Nesher N, Alghamdi AA, Singh SK, Sever JY, Christakis GT, Goldman BS, *et al.* Troponin after cardiac surgery: a predictor or a phenomenon? *Ann Thoracic Surg* 2008;85(4):1348-54.
20. Onorati F, Cristodoro L, Caroleo S, Esposito A, Amantea B, Santangelo E, *et al.* Troponin and lactate from the coronary sinus predict cardiac complications after myocardial revascularization. *Ann Thoracic Surg* 2007;83(3):1016-23.
21. Melrose DG, Dreyer B, Bentall HH, Baker JB. Elective cardiac arrest. *Lancet* 1955;269(6879):21-2
22. Magovern GJ Jr, Flaherty JT, Gott VL, Bulkley BH, Gardner TJ. Failure of blood cardioplegia to protect myocardium at lower temperatures. *Circulation* 1982;66 (Suppl. I):I60-7.
23. Buckberg GD, Brazier JR, Nelson RL, Goldstein SM, McConnell DH, Cooper N. Studies of the effects of hypothermia on regional myocardial blood flow and metabolism during cardiopulmonary bypass. The adequately perfused beating, fibrillating and arrested heart. *J Thorac Cardiovasc Surg* 1977;73(1):87–94.
24. Hayashida N, Ikonomidis JS, Weisel RD, Shirai T, Ivanov J, Carson SM, *et al.* The optimal cardioplegic temperature. *Ann Thorac Surg* 1994;58(4):961–71.
25. Fan Y, Zhang AM, Xiao YB, Wang YG, Hetzer R. Warm versus cold cardioplegia for heart surgery: a meta-analysis. *Eur J Cardiothorac Surg* 2010;37(4):912–9.
26. Abah U, Garfjeld Roberts P, Ishaq M, De Silva R. Is cold or warm blood cardioplegia superior for myocardial protection? *Interact CardioVasc Thorac Surg* 2012;14(6):848–55.^[17]_[SEP]

27. Durandy Y, Hulin S. Intermittent warm blood cardioplegia in the surgical treatment of congenital heart disease: clinical experience with 1400 cases. *J Thorac Cardiovasc Surg* 2007;133(1):241–6.
28. Badak MI, Guncun U, Discigil B, Boga M, Ozkisacik EA, Alayunt EA. Myocardium utilizes more oxygen and glucose during tepid blood cardioplegic infusion in arrested heart. *Int Heart J* 2005;46(2):219–29.
29. Najafi H, Henson D, Dye WS, Javid H, Hunter JA, Callaghan R, *et al.* Left ventricular hemorrhagic necrosis. *Ann Thorac Surg* 1969;7(6):550-61.