

LIST OF TABLES

TABLE 1.1	Summary of potential foodborne pathogens associated with foodborne illness.....	18
TABLE 1.2	Summary of predominant bacterial pathogens associated with various food commodities (Mossel <i>et al.</i> , 1995; ICMSF, 1996; Jay <i>et al.</i> , 2005).....	19
TABLE 2.1	Description of RTE foods, environmental swabs and cleaning tools.....	36
TABLE 2.2	Culture media, times, temperatures and plating techniques used in the microbiological survey of RTE foods, associated preparation surfaces and cleaning tools sampled from four retail delicatessens.....	37
TABLE 2.3	Mean bacterial counts obtained for RTE foods (n=77) sampled from four retail delicatessens	38
TABLE 2.4	Incidences of <i>Listeria (L.) monocytogenes</i> and <i>Salmonella</i> spp. in RTE foods (n=77) sampled from four retail delicatessens before and after preliminary incubation (PI)	39
TABLE 3.1	Mean aerobic bacterial, coliform, <i>E. coli</i> , <i>S. aureus</i> and <i>B. cereus</i> counts from RTE filled baguettes (n=35) and assorted salads (n=35) sampled from four retail delicatessens.....	71
TABLE 3.2	Incidences of <i>L. monocytogenes</i> and <i>Salmonella</i> spp. in RTE filled baguettes (n=35) and assorted salads (n=35) sampled from four retail delicatessens.....	72

TABLE 3.3	Microbiological guidelines suggested for RTE filled baguettes and assorted salads from the retailer participating in this study and the South African Department of Health. In addition the percentage of RTE samples (n=70) complying with these recommended guidelines.....	73
TABLE 3.4	Percentage distribution of aerobic bacteria associated with RTE filled baguettes and assorted salads (280 isolates) and associated preparation surfaces (368 isolates) sampled from four retail delicatessens.....	74
TABLE 3.5	Percentage distribution of 280 predominant isolates from aerobic plate counts (APC) of RTE filled baguettes and assorted salads sampled from four retail delicatessens.....	75
TABLE 4.1	Incidence of <i>Salmonella</i> spp. and <i>Listeria monocytogenes</i> on floor mops (n=25), cleaning cloths (n=39) and disposable plastic gloves (n=20) sampled from four retail delicatessens during the preparation of filled baguettes and assorted salads.....	97
TABLE 4.2	Percentage distribution of predominant microbial isolates from floor mops (100 isolates), cleaning cloths (156 isolates) and disposable plastic gloves (80 isolates) sampled from four retail delicatessens during the preparation of filled baguettes and assorted salads	98