

https://dspace.org/

Cooling methods to treat capture-induced hyperthermia in blesbok (Damaliscus dorcas phillipsi)

Sawicka, Joanna

2011-12-07
http://hdl.handle.net/10539/10884

Downloaded from a DSpace repository. https://dspace.org/

COOLING METHODS TO TREAT CAPTURE-INDUCED HYPERTHERMIA IN BLESBOK *(DAMALISCUS DORCAS PHILLIPSI)*

Joanna Sawicka

A dissertation submitted to the Faculty of Science, University of the Witwatersrand, Johannesburg, in fulfilment of the requirements for the degree of Master of Science.

Johannesburg, 2010

DECLARATION

I declare that this dissertation is my own work, with all assistance acknowledged. This dissertation is being submitted for the degree of Master of Science at the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at any other University.

…….

(**JOANNA SAWICKA**)

…… day of ……. 2010

I certify that the procedures used in this dissertation were approved by the Animal Ethics Screening Committee of the University of the Witwatersrand (AESC number 2008/34/04).

DEDICATION

I dedicate this dissertation to my family (both immediate, and to all my friends who have become my family) who supported me through all the difficult and rewarding times that went into making this dissertation possible. Thank you to my dad for all his encouraging words, my sister for all her help with grocery shopping and to my mom who helped me pack bakkies. The years of my dissertation have been incredibly stressful and I would not have made it without you all. A special mention and thank you to Allan Stacey for all his help when computer technology was an obstacle to be overcome. And thank you to my partner Alexei Krassnokutski, who has understood and supported me throughout many tribulations and frustrations. Lastly, I would also like to dedicate this dissertation to Charles Vermeulen, who will always be missed.

PRESENTATIONS AND PUBLICATIONS

The following presentations are offered in support of this dissertation

Sawicka JA, Fuller A, Fick L, Hetem R, Meyer LCR. 2009. The effectiveness of different cooling methods at alleviating capture-induced hyperthermia in blesbok *(Damaliscus dorcas phillipsi)* (oral presentation). At the South African Wildlife Management Association (SAWMA), Black Mountain, Free State, 13-16 September.

Sawicka JA, Fuller A, Fick L, Hetem R, Meyer LCR. 2009. The effectiveness of different cooling methods at alleviating capture-induced hyperthermia in blesbok *(Damaliscus dorcas phillipsi)* (poster presentation). At the University if the Witwatersrand's Postgraduate Symposium, Johannesburg, 20-21 October.

ABSTRACT

Wild animals are captured for management, health, translocation and research purposes. Capture is an unnaturally stressful event, which may result in morbidity or mortality. An attributing cause of the morbidity and mortality is capture-induced hyperthermia; the larger the magnitude and the longer the duration of this captured-induced hyperthermia, the greater the likely risk to the animal. The most common practice currently used in the field to lower body temperature is to douse hyperthermic animals with water. However, the water used is often at ambient temperatures and its efficacy is not known. We investigated whether this method and alternative methods are effective at lowering the body temperature of hyperthermic animals. To achieve these aims we implanted 19 blesbok with miniature temperature-sensitive data loggers in their abdomens and into their subcutaneous layers (at the sites of the flank, groin, lower neck and upper neck). The loggers continuously recorded core body temperatures of the blesbok throughout the study period at an interval of six minutes. We successfully retrieved complete data sets from 12 blesbok. The animals were captured on six separate occasions using a technique which elicited hyperthermia. Five animals were cooled by dousing with water of different temperatures (4°C, 17°C, 28°C) and fanning after dousing with 28°C water, in random order. Seven animals were cooled by ice packs, spraying a fine mist spray, intravenous (IV) infusion of one litre of 4°C water and 28°C water-dousing. Through the use of our continuous logging of body temperature we established the normal body temperature of the blesbok, which displayed a regular 24 hour body temperature pattern. The average daily body temperature of the blesbok was 38.8°C ± 0.4°C, with a minimum body temperature of 37.9°C ± 0.1°C and a maximum body temperature of 39.4°C ± 0.1°C. The body temperature after capture was as

high as 41°C-42°C, which was significantly higher than the normal body temperature (Student's t-test, $P < 0.05$). The animals were cooled once they were immobilised and the start of cooling was denoted as time zero. In the control (no active cooling) intervention the body temperature decreased to only about 40°C. Dousing animals with water, irrespective of its temperature, resulted in significant cooling ($P < 0.05$) of the animals, as indicated by their minimum body temperature reached, change in body temperature and rate of cooling. The water-dousing interventions decreased the body temperature to about 38°C after an hour, which was significantly lower than the control (RM-ANOVA, $P < 0.05$) but there was no significant difference in the minimum body temperature reached between the different water temperatures or by the addition of fanning (RM-ANOVA, $P > 0.05$). The water-dousing interventions cooled the animals more quickly than did the control (RM-ANOVA, $P < 0.05$), and the coldest water (4°) cooled the animals quicker than did the 28°C water-dousing (RM-ANOVA, $P < 0.05$). The core body temperature minus the subcutaneous temperature was calculated, and revealed a peak difference of about 3.5°C after the 4°C water-dousing. Ice-packs also resulted in significant cooling ($P < 0.05$) of the animals, as depicted by their minimum body temperature reached, change in body temperature and rate of cooling. The ice-packs lowered the body temperature to a minimum of about 38°C, which was significantly lower than the control (RM-ANOVA, $P < 0.05$). The ice-packs also cooled the animals significantly faster than did the control, intravenous infusion and mist spray (RM-ANOVA, $P < 0.05$) but cooled as quickly as the 28°C water-dousing (RM-ANOVA, $P > 0.05$). The core body temperature minus the subcutaneous temperature for the ice-packs peaked at a difference of about 3°C. The IV infusion and mist spray were ineffective cooling methods and did not significantly ($P > 0.05$) alter the minimum body temperature or rate of cooling. Even though the IV infusion caused a significant reduction in body temperature by

1°C, the cooling effect from the IV infusion was short-lived because the minimum body temperature reached after the intravenous infusion and mist spray was ultimately similar to the body temperature seen in animals receiving the control (RM-ANOVA $P > 0.05$). Also, the intravenous infusion and mist-spray cooled as slowly as did the control (RM-ANOVA $P > 0.05$). Therefore, water-dousing in this study was the most effective and practical method to cool hyperthermic blesbok. Although all the water temperatures (4°C, 17°C and 28°C) that we tested were effective, the coldest water (4°C) cooled the animals quickest. The addition of fanning to the 28°C water-dousing did not increase cooling. Ice-packs were also effective but may be not as easy to use as the water-dousing method as ice-packs are large and need to be kept frozen, and therefore are cumbersome for use in the field.

ACKNOWLEDGMENTS

I would like to thank my supervisors, Dr. Linda Fick, Dr Leith Meyer and Prof. Andrea Fuller, for their valued guidance and invaluable advice. I would also like to thank the remainder of the Brain Function Research Group and the staff of the Central Animal Service for their assistance in surgery and for providing crucial volunteers for collecting data during long days in the field.

Thank you to our collaborators, the National Zoological Gardens, for the use of their facilities and staff to make this project possible.

Lastly, thank you to the National Research Foundation for financially supporting this study.

TABLE OF CONTENTS **PAGE**

DECLARATION ii

DEDICATION iii

PUBLICATIONS AND PRESENTATION iv

ABSTRACT v

ACKNOWLEDGEMENTS viii

TABLE OF CONTENTS ix

LIST OF FIGURES xii

LIST OF TABLES xv

1. INTRODUCTION 1

1.1 Capture of game animals 2

1.2 Capture-induced hyperthermia 6

1.2.1 Pathophysiology of hyperthermia 6

1.2.2 Consequences of hyperthermia for animal welfare 10

1.2.3 Incidence and magnitude 11

1.2.4 Causes of capture-induced hyperthermia 17

1.3 Cooling of hyperthermic animals 23

1.3.1 Principles of heat loss 23

1.3.2 Cooling methods used for humans 29

1.3.3 Cooling methods used for domesticated animals 38

1.3.4 Cooling methods used for game animals 42

1.4 Dissertation aims 45

2. MATERIALS AND METHODS 47

2.1 Subjects and study site 48

2.2 Surgery 49

2.2.1 Anaesthesia 49

2.2.2 Implantation and removal of data loggers 49

2.3 Body temperature measurement 54

2.3.1 Core body temperature 54

2.3.2 Subcutaneous temperature 55

2.4 Experimental procedure 55

2.5 Microclimate measurements 64

2.6 Data analysis 64

3. RESULTS 68

3.1 Normal body temperature rhythm 69

3.2 The body temperature change due to capture 72

3.3 Mean body temperatures for the experimental interventions 75

3.4 Peak body temperature 77

3.5 Minimum body temperature 79

3.6 Amplitude of cooling 81

3.7 Thermal response indices 83

3.8 Rate of cooling 85

3.9 The temperature difference between the core and subcutaneous temperature 87

3.10 The temperature difference between core and ear skin temperature 90

3.11 Core abdominal temperature versus rectal temperature 92

3.12 Environmental conditions 92

4. DISCUSSION 96

5. CONCLUSION 119

6. REFERENCES 131

LIST OF FIGURES

Figure 1.1 25

An overview of how animals lose and gain heat to and from the environment.

Figure 2.1 50

The typical size of the Stowaway miniature data logger, waxed (left) and unwaxed (right), which measured core body temperature intra-abdominally.

Figure 2.2 51

The typical size of the iButton miniature data logger unit, waxed (left) and unwaxed (right), which measured subcutaneous body temperature.

Figure 2.3 53

An illustration of the sites of data logger implantation in the blesbok.

Figure 2.4 57

Schematic representation of the boma and chase sequence.

Figure 2.5 63

A flow diagram showing the order of the experimental procedures.

Figure 3.1 70

The body temperature of one blesbok over seven days.

Figure 3.2 71

The typical mean body temperature rhythm of all the blesbok shown as temperature over 24 hours.

Figure 3.3 73

The body temperature of one blesbok over eleven days that include two capture days..

Figure 3.4 74

The body temperature of one blesbok before capture, during capture, during experimental interventions and after experimental interventions.

Figure 3.5 76

The mean body temperature of the blesbok for each experimental intervention over time.

Figure 3.6 78

The peak body temperatures for each experimental intervention.

Figure 3.7 80

The minimum temperatures reached within an hour from the start of cooling.

Figure 3.8 82

The magnitude of the decrease in temperature within 30 minutes (panels A and C) and 1 hour (panels B and D).

Figure 3.9 84

The TRI values within 30 minutes and within one hour from the start of cooling for the water dousing and alternative groups.

Figure 3.10 86

The rate of cooling over 30 minutes by each experimental intervention

Figure 3.11 88

The temperature difference between the core and the periphery over time.

Figure 3.12 91

The temperature difference between the core and ear skin over time.

Figure 3.13 93

The body temperature discrepancy between abdominal temperature and rectal temperature.

LIST OF TABLES

Table 1.1 12

An overview of hyperthermia and the mortalities associated with capture.

Table 1.2 18

An overview of species that are reported to be susceptible to hyperthermia, capture myopathy or stress

Table 1.3 30

An overview of cooling techniques that have been tested for the purpose of cooling humans

Table 2.1. 60

A summary of animal body masses and cooling interventions for both the water-dousing group and the alternative methods group.

Table 3.1 94

The environmental conditions for both groups.