

ASSESSMENT OF CURRENT PRACTICES OF INTRAHOSPITAL TRANSPORTATION IN CRITICALLY ILL PATIENTS AT AN ACADEMIC HOSPITAL IN SOUTH AFRICA


Bhavic Mawjee

A research report submitted to the Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, in the partial fulfilment of the requirements for the degree of Master of Medicine in the branch of Anaesthesiology.

Johannesburg, 2023

Declaration

I, Bhavic Mawjee, declare that this Research Report is my own, unaided work. It is being submitted for the Degree of Master of Medicine at the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at any other University. The article is prepared according to the Author Guidelines for Journal of Critical Care (Section 2) and may differ from the rest of the document.

A handwritten signature in black ink, appearing to read 'Bhavic Mawjee', is written over a horizontal line.

(Signature of candidate)

On this day of 31 October 2023

Abstract

Purpose:

There is a paucity of guidelines for safe intrahospital transportation(IHT) in a resource-limited setting. The aim of this study was to describe the practices of IHT of critically ill patients perioperatively by anaesthetists.

Methods:

We conducted a prospective, cross-sectional study including 101 IHTs between theatre and Intensive care unit (ICU) . The most senior anaesthetist completed an adapted questionnaire. Frequency distributions were described, and Chi-square/Fisher's exact tests were performed to identify associations between IHT practices and categorical variables with anaesthetist seniority and first and subsequent IHTs.

Results:

The majority(72.97%) of the variables investigated were below the expected compliance rate(75%). Only 22.80% of anaesthetists were aware of IHT guidelines. Statistically significant differences were observed in IHT practices between senior and junior anaesthetists for: airway equipment taken(OR 0.25 95% CI 0.08-0.84), emergency drugs taken(inotropes(OR 1.49 95%CI 1.29-1.73), muscle relaxants(OR 5.81 95% CI 1.70-19.83), induction agents(OR 7.23 95% CI 1.86-28.13)) and awareness of IHT guidelines(OR 1.35 95% CI 1.20-1.53); and between first and subsequent transfers for: notification of the current condition(OR 0.23 95% CI 0.08-0.71), use of manual resuscitation bag(OR 3.64 95% CI 1.21-11.00), and awareness of IHT guidelines(OR 0.14 95% CI 0.04-0.49).

Conclusion:

The majority of IHT practices interrogated during this study were below the expected compliance level irrespective of the seniority of the anaesthetist or the number of IHTs. Resource limitations and education seem to be the major contributors.

Acknowledgements

I would like to thank the following people:

My supervisors, Sujai Mohanadasan and Zainub Jooma for their ongoing assistance, encouragement, and patience.

Mrs Zvifadzo Matsena Zingoni for her assistance and guidance during statistical analysis.

My wife and family for their constant encouragement during the study.

Table of Contents

Declaration.....	ii
Abstract.....	iii
Acknowledgements.....	iv
Table of Contents	v
List of Tables	viii
List of Figures.....	ix
Abbreviations	x
Statement.....	xi
Section 1: Draft Article	1
1.1. Abstract	3
1.2. Introduction	4
1.3. Method	18
1.4. Results	10
1.5. Discussion.....	17
1.6. Conclusion	22
1.7. Ethics approval and consent to participate.....	23
1.8. Consent for publication	23
1.9. Funding	23
1.10. Authors' contributions.....	23
1.11. Declaration of Competing Interest.....	23
1.12. Data availability	23
1.13. References:.....	24
1.14. Supplementary file 1: Questionnaire	28
Section 2: Authors guideline for Journal of Critical Care	38
Section 3: Proposal.....	53
3.1. Introduction.....	54

3.2.	Problem statement	58
3.3.	Aim	59
3.4.	Objectives.....	59
3.5.	Research assumptions	59
3.6.	Demarcation of study field	60
3.7.	Ethical considerations	61
3.8.	Research Methodology	62
3.8.1.	Research design.....	62
3.8.2.	Study population	63
3.8.3.	Study sample	63
3.8.4.	Data collection	64
3.8.5.	Data analysis	66
3.8.6.	Significance of the study	66
3.9.	Validity and reliability of the study.....	67
3.10.	Potential limitations of the study.....	67
3.11.	Project outline	69
3.11.1.	Time frame	69
3.11.2.	Financial plan	70
3.12.	References.....	71
3.13.	Appendices	75
3.13.1.	Appendix A: Permission from the Head of Department	75
3.13.2.	Appendix B: Information sheet	76
3.13.3.	Appendix C: Permission to use and adapt checklist	77
3.13.4.	Appendix D: Data Collection Sheet	78
3.13.5.	Appendix E: Permission from CEO of CMJAH	89
Section 4:	Annexures	90
4.1.	Annexure 1: Postgraduate Approval letter	90

4.2.	Annexure 2: Human research ethics committee clearance certificate	91
4.3.	Annexure 3: Plagiarism/Turnitin report	92

List of Tables

Section 1: Draft Article

Table 1-1: Overview of content of published IHT checklists and guidelines.....	6
Table 1-2: Capacity of various ICUs and number of admissions from theatre	8
Table 1-3: Patients' demographics.....	11
Table 1-4: Anaesthetists' demographics	12
Table 1-5: Comparison of IHT practices based on seniority of anaesthetist and number of IHTs done	14

Section 3: Proposal

Table 3-1: Overview of content of published intrahospital transportation (IHT) checklists and guidelines.....	56
Table 3-2: Capacity of various ICUs and number of admissions from theatre	61
Table 3-3: Time frame.....	69
Table 3-4: Financial plan.....	70

List of Figures

Section 1: Draft Article

Figure 1-1: Study flow diagram.....10

Figure 1-2: Potential barriers to practicing safe IHT guidelines. 13

Abbreviations

AA	Association of Anaesthetists
AAGBI	Association of Anaesthetists of Great Britain and Ireland
ANZCA	Australian and New Zealand College of Anaesthetists
CCSSA	Critical Care Society of South Africa
DA	Diploma in Anaesthesia
FCA	Fellowship of the College of Anaesthetists
HICs	High income countries
ICU	Intensive care unit
IHT	Intrahospital transportation
LMICs	Low-middle income countries
NICE	National Institute for Health and Care Excellence
SASA	South African Society of Anaesthesiologists

Statement

The Research Report consists of a draft article and appendices.

The article is prepared according to the Author Guidelines for Journal of Critical Care (Section 2) and may differ from the rest of the document.

The study proposal is included for background reference and is not for examination.

Section 1: Draft Article

Assessment of current practices of intrahospital transportation in critically ill patients at an academic hospital in South Africa

Bhavic Mawjee, BSc (Wits) MBBCH (Wits) DipPEC (SA) DA (SA) ^{a, *}

Sujai Mohanadasan, MBChB (UKZN) DA(SA) FCA(SA) MMED (Anaes) FANZCA ^b

Zainub Jooma MBBCH(Wits) DA(SA) FCA(SA) MMED (Anaes) Cert Crit Care (SA) ^a

^a Department of Anaesthesiology, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand

^b Department of Anaesthesiology, Kaitia Hospital, New Zealand

* Corresponding author at: B Mawjee

Department of Anaesthesiology

Charlotte Maxeke Johannesburg Academic Hospital

7 York Rd

Parktown

Johannesburg

2193

Bhavicmawjee@gmail.com

+2711 488 4344

Keywords: Intrahospital transportation, perioperative, current practices, critically ill, anaesthetist

Highlights:

- Poor awareness of intrahospital transportation guidelines in South African hospital.
- Low compliance level during intrahospital transportation in South African hospital.
- Experience of the anaesthetist did not change intrahospital transportation practice.
- The number of transfers done did not change intrahospital transportation practice.
- Resource limitations and education seem to be the major contributors to low compliance.

Abstract

Purpose:

There is a paucity of guidelines for safe intrahospital transportation(IHT) in a resource-limited setting. The aim of this study was to describe the practices of IHT of critically ill patients perioperatively by anaesthetists.

Methods:

We conducted a prospective, cross-sectional study including 101 IHTs between theatre and intensive care unit (ICU). The most senior anaesthetist completed an adapted questionnaire. Frequency distributions were described, and Chi-square/Fisher's exact tests were performed to identify associations between IHT practices and categorical variables with anaesthetist seniority and first and subsequent IHTs.

Results:

The majority(72.97%) of the variables investigated were below the expected compliance rate(75%). Only 22.80% of anaesthetists were aware of IHT guidelines. Statistically significant differences were observed in IHT practices between senior and junior anaesthetists for: airway equipment taken(OR 0.25 95% CI 0.08-0.84), emergency drugs taken(inotropes(OR 1.49 95%CI 1.29-1.73), muscle relaxants(OR 5.81 95% CI 1.70-19.83), induction agents(OR 7.23 95% CI 1.86-28.13)) and awareness of IHT guidelines(OR 1.35 95% CI 1.20-1.53); and between first and subsequent transfers for: notification of the current condition(OR 0.23 95% CI 0.08-0.71), use of manual resuscitation bag(OR 3.64 95% CI 1.21-11.00), and awareness of IHT guidelines(OR 0.14 95% CI 0.04-0.49).

Conclusion:

The majority of IHT practices interrogated during this study were below the expected compliance level irrespective of the seniority of the anaesthetist or the number of IHTs. Resource limitations and education seem to be the major contributors.

Introduction

Critically ill patients have life threatening injuries or illnesses that are often associated with reduced physiological reserves. These patients often require transfer from the intensive care unit (ICU) to other departments in the hospital for therapeutic or diagnostic purposes [1]. The risks associated with intrahospital transportation (IHT) need to be outweighed by the benefits from the procedure required [2,3]. Critically ill patients should be monitored and receive physiological support during patient transport identical to that which would be given while in the ICU [2]. Transportation of these patients to various departments require highly trained and skilled medical practitioners with appropriate essential equipment [2]. It has been accepted that system failures and errors can stem from a culture of non-compliance, perceptions of invulnerability, or inherently poor practices [3].

Some studies from high income countries (HICs) and some low- and middle-income countries (LMICs) have highlighted the risk of adverse events occurring during IHT [1,4,5]. These adverse events that occur during IHT may have a direct or indirect role in influencing the patient's risk of morbidity and mortality [4,6–8]. The incidence of adverse events occurring during IHT can vary from 1.7% to 67.9% [4,9] depending on the level of experience of the practitioner present. A South African study noted that in the absence of an adverse event, a medical officer transporting a patient to the ICU perioperatively, as opposed to a registrar, was associated a higher in-hospital mortality rate [5]. This might allude to different levels of experience or skill sets having an impact on the patient's condition during an IHT [4,5]. Although some adverse events are inevitable, many are preventable [4].

Several guideline documents and checklists have been produced to guide safe IHT and thereby decrease IHT related adverse events [1,4,5,10–17] . These are summarised in Table 1.

While many guidelines exist in HICs, there is a paucity of standardized protocols or guidelines for safe IHT of critically ill patients in a resource-limited setting, specifically in South Africa [4]. At the time this study was conducted, there were no guidelines published by the South African Society of Anaesthetists (SASA) or the Critical Care Society of South Africa (CCSSA) for IHT of critically ill patients. It is at the discretion of the medical practitioner transporting a critically ill patient to determine the

equipment and drugs required during the IHT. To the best of our knowledge, there are no known studies that describe IHT practices in any LMICs. The aim of this study was to describe the current practices of IHT of critically ill patients perioperatively by anaesthetists in a quaternary hospital in South Africa.

Table 1-1: Overview of content of published IHT checklists and guidelines

Author		Warren et al. [13]	Fanara et al. [11]	Jarden and Quirke [14]	Brunsveld-Reinders et al. [12]	ANZCA guideline[15]	Williams et al. [1]	Geldenhuys et al. [4]
Year		2004	2010	2010	2015	2015	2020	2020
Administrative guidelines						+		
Pre IHT	Patient assessed prior to transfer	+	+	+		+		+
	IHT to destination beneficial/ required	+					+	+
	Transport team required	+	+	+		+		
	Consent confirmed		+	+			+	
	Equipment and consumables present	+	+	+	+	+	+	+
	Sufficient oxygen level in cylinder	+	+		+	+	+	+
	Sufficient IV medication present	+	+	+	+	+	+	+
	Extra IV fluids and medication	+	+	+	+	+	+	+
	Enteral feeding and insulin stopped				+			
	Check tubes and lines		+	+	+	+	+	+
	Suctioned airway						+	
	Check and set monitor alarms	+	+	+	+	+	+	+
	Check and set transport ventilator alarms	+	+		+	+		+
	Preparation and equipment adapted to procedure. (MRI compatible)	+	+		+		+	
	Register baseline vital signs	+	+	+	+	+	+	+
	Notification of destination department	+			+	+	+	+
	Transport route clear		+			+		+

Author		Warren et al. [13]	Fanara et al. [11]	Jarden and Quirke [14]	Brunsveld-Reinders et al. [12]	ANZCA guideline [15]	Williams et al. [1]	Geldenhuys et al. [4]
Year of publication		2004	2010	2010	2015	2015	2020	2020
During IHT	Equipment checked	+		+	+	+		+
	Record IV fluids or medication given			+	+	+	+	+
	Vital signs recorded every 20min	+		+	+	+	+	+
Post IHT	Enteral feeds and insulin restarted				+			
	Humidifier switched on				+			
	HME filter changed				+			
	Suction bag changed if used			+	+			
	Transport bag restocked				+			+
	Equipment placed on charge							+
	Reported any adverse events on route			+	+	+	+	+
	Handed over to destination staff	+		+	+	+	+	+

Method

This was a prospective cross-sectional study that focused on IHTs between the ICU and operating theatre, conducted by an anaesthetist in a quaternary, academic hospital in Johannesburg, South Africa, from December 1st, 2021, to February 28th, 2022, during the second wave of the COVID-19 pandemic.

Approval was obtained from the Human Research Ethics Committee (M210708) and relevant authorities. The study included patients transported between theatre and the following ICUs: medical, surgical, trauma, neurosurgery, and cardiothoracic. Table 2 summarizes the capacity and average number of monthly admissions from the theatre for each ICU and highlights the differences in numbers during the COVID-19 pandemic.

A minimum sample size of 80 was determined using StatCalc® with the following assumptions: a population size of 100 transfers per month, an alpha error of 5%, a 95% confidence interval, and an expected minimum compliance of 75%.

The current IHT practices were not known in LMICs. Compliance to IHT guidelines is unlikely to be the same as HICs due to the limited resources, thus an expected compliance rate was adjusted to 75%.

Table 1-2: Capacity of various ICUs and number of admissions from theatre

ICU	Number of beds available	Discipline serves	Average number of admissions from theatre over a month	
			Pre/post covid peak	During covid peak
Main ICU	24	Medical, Surgical and covid patients	27	11
Cardio-thoracic ICU	6	Cardiothoracic	32	16
Neurosurgery ICU	8	Neurosurgery	16	17
Trauma ICU	12	Trauma	22	24

The study population consisted of any anaesthetist at the hospital who transported a critically ill patient between the theatre and ICU during the study period. In this study, a junior anaesthetist was defined as an anaesthetist with less than or equal to three years' anaesthetic experience and a senior anaesthetist was defined as an anaesthetist with more than three years' anaesthetic experience. A critically ill patient was defined as any patient requiring cardiovascular, respiratory, renal, or neurological organ support, needing ICU admission for mechanical ventilation, inotropic infusions, intensive monitoring, or other supportive measures.

The questionnaire used for the study was adapted from Williams et al's checklist (1). Adaptations were made to account for the use in a resource-limited environment, as the checklist was originally developed and designed for use in a HIC. The adapted questionnaire contained seven sections with 58 variables in total, namely: demographics of the patient and participant, patient destination, patient preparation, airway management, emergency drugs, adverse events, and potential barriers to adherence to guidelines (Supplementary file 1). Questionnaires were made available electronically and were completed by the most senior anaesthetist accompanying the patient after IHT. Participation in the study was voluntary and completion of the questionnaire implied consent.

The inclusion criteria for the study included patients aged >12 years, critically ill, undergoing IHT between the ICU and theatre, and IHT done during the perioperative period by an anaesthetist. Exclusion criteria included IHT of critically ill patients to/from a ward/high care setting, IHT to/from endoscopy and radiology suites, and incomplete questionnaires.

The data was analysed using IBM SPSS version 21 (SPSS version 21.0; IBM Corporation, Armonk, NY, USA) statistical software. Frequency distribution was expressed in numbers and percentages for categorical variables. Chi-square and Fisher's exact tests were performed and univariate odds ratio and 95% confidence interval was calculated where appropriate comparing junior with senior anaesthetists' practices and first IHT with subsequent IHT practices with categorical variables. Statistical significance was set at a p-value of < 0.05.

Results

A total of 104 questionnaires were completed, of which 101 met the inclusion criteria (Figure 1). Tables 3 and 4 provide information about the demographics of the patients and anaesthetists, respectively.




Figure 1-1: Study flow diagram

Table 1-3:Patients' demographics

Variables	n (%)
Age	
12-18	4 (3.96)
19-30	18 (17.82)
31-40	18 (17.82)
40-65	41 (40.59)
>65	20 (19.80)
ASA category	
ASA 1	1 (1.00)
ASA 2	10 (9.90)
ASA 3	61 (60.40)
ASA 4	26 (25.74)
ASA 5	3 (2.97)
ASA 6	0
Transfer from	
ICU	35 (34.65)
Theatre	66 (65.35)
IHT destination	
To ICU	78 (77.23)
To theatre	23 (22.77)
Reason for requiring ICU care	
Mechanical ventilation	60 (59.41)
Inotropic support	34 (33.66)
Intensive monitoring	90 (89.11)
Renal replacement therapy	12 (11.88)

Table 1-4: Anaesthetists' demographics

Variables	All n (%)	Seniority of anaesthetist	
		Junior n=13	Senior n= 88
Age			
25-30	19(18.81)	7(53.85)	12(13.64)
31-40	58(57.43)	5(38.46)	53(60.23)
41-60	15(14.85)	0	15(17.05)
>60	9(8.91)	1(7.69)	8(9.10)
Gender			
Female	58(57.43)	3(23.08)	55(62.50)
Male	43(42.57)	10(76.92)	33(37.50)
Level of Training			
Medical officer	5(4.95)	3(23.08)	2(2.27)
Registrar	91(90.10)	10(76.92)	81(92.05)
Consultant	5(4.95)	0	5 (5.68)
Number of times participated in study			
1	44(43.56)	8(61.54)	36 (40.90)
> 1	57(56.44)	5(38.46)	52 (59.10)

The study investigated 37 different variables related to IHT practices and found that 27 (72.97%) of these variables were below the expected compliance rate of 75%. Consent was checked by 91.10% of anaesthetists, and vitals were checked by 99% of anaesthetists prior to IHT.

Only 23 (22.80%) anaesthetists were aware of IHT guidelines, including guidelines from the Association of Anaesthetists (AA), National Institute for Health and Care Excellence (NICE), and the Association of Anaesthetists of Great Britain and Ireland (AAGBI). Among the 44 first-time participants, only 6.80% (n=3) were aware of IHT guidelines. Whereas, of those having subsequent transfers, 35.10% participants (n=20) were aware of IHT guidelines. This comparison was statistically significant (OR 0.14 95% CI 0.04-0.49, p-value = 0.001).

No anaesthetist reported a major adverse event occurring during IHT. These include accidental extubation, airway obstruction, hypotension requiring management, new onset arrhythmias, initiation of cardiopulmonary resuscitation, infusion pump failure,

monitor failure and ventilator failure. Most anaesthetists (67.30%) reported no barriers to adherence to IHT guidelines, while 32.70% reported potential barriers to practicing safe IHT guidelines (Figure 2).




Figure 1-2: Potential barriers to practicing safe IHT guidelines

Table 5 presents the current practices of anaesthetists during IHT of critically ill patients perioperatively between theatre and ICU, and the comparisons based on seniority and number of IHTs performed.

There were no statistically significant differences in IHT practices between junior and senior anaesthetists, except in five variables, namely: airway equipment taken during IHT (supraglottic devices), emergency drugs taken during IHT (inotropes, muscle relaxants, and induction agents) and awareness of IHT guidelines. There were statistically significant differences in IHT practices between first and subsequent transfers for the following variables: notification of the current condition, use of manual resuscitation bag, and awareness of IHT guidelines.

Table 1-5: Comparison of IHT practices based on seniority of anaesthetist and number of IHTs done

	All	Senior vs Junior anaesthetists		First vs subsequent IHT	
Practices done prior to IHT	n (%)	Odds Ratio (CI 95%)	p-value	Odds Ratio (CI 95%)	p-value
Notification of destination prior to IHT	99 (98.00) [#]	1.02 (0.99-1.06)	1.00	0.96 (0.90-1.02)	0.19
Expected time of arrival	67 (66.30)	1.16 (0.33-4.09)	1.00	0.47 (0.20-1.09)	0.08
Notification of current condition	83 (82.10) [#]	0.43 (0.11-1.58)	0.24	0.23 (0.08-0.71)	0.01*
Infusions required on arrival	49 (48.50)	1.83 (0.56-6.05)	0.31	1.57 (0.67-3.39)	0.29
Requirement of ventilator support	57 (56.40)	1.27 (0.39-4.20)	0.69	1.43 (0.64-3.18)	0.38
Route cleared of potential delays prior to IHT	66 (65.35)	1.22 (0.35-4.30)	1.00	0.51 (0.22-1.18)	0.11
IV line checked and flushed	98 (97.00) [#]	0.28 (0.02-3.32)	0.34	1.56 (0.14-17.82)	1.00
Monitors checked and used					
Pulse oximeter	99 (98.00) [#]	0.14 (0.01-2.35)	0.24	1.04 (0.99-1.09)	0.50
ECG	90 (89.10) [#]	-	0.35	3.93 (0.81-19.25)	0.11
NIBP or invasive BP monitoring	92 (91.00) [#]	-	0.60	7.02 (0.84-58.42)	0.07
Practices during IHT					
Airway equipment taken with during IHT					
Oropharyngeal/nasopharyngeal airway	23 (22.8)	1.61 (0.45-5.82)	0.49	1.00 (0.93-2.54)	0.99

Supplemental oxygen device (e.g.: Nasal prong oxygen)	55 (54.5)	0.97 (0.30-3.13)	0.96	0.86 (0.39-1.89)	0.70
Extra endotracheal tubes (appropriate sizes)	5 (5.00)	1.06 (1.01-1.12)	1.00	0.31 (0.03-2.86)	0.38
Emergency supraglottic device	74 (73.3)	0.25 (0.08-0.84)	0.02*	0.63 (0.26-1.54)	0.31
Laryngoscope with appropriate blade sizes	1 (1.00)	0.92 (0.79-1.08)	0.13	0.97 (0.93-1.02)	0.44
Airway management					
Airway was suctioned	49 (48.50)	0.67 (0.19-2.43)	0.54	0.74 (0.29-1.86)	0.52
ETT size and depth noted if patient was intubated	52 (51.50)	0.31 (0.06-1.54)	0.16	0.31 (0.07-1.93)	0.15
ETT tape/securing device checked and secured (if intubated)	59 (58.40)	0.18 (0.01-3.15)	0.30	0.74 (0.04-12.33)	1.00
IHT practices related to Transport ventilator					
Transport ventilator was used during IHT	45 (44.60)	6.00(0.71-50.47)	0.08	0.77 (0.29-2.09)	0.61
Battery level of ventilator checked	42 (41.60)	0.71 (0.07-7.71)	1.00	0.56 (0.07-4.36)	0.62
Ventilator checked for leaks	37 (36.60)	0.45 (0.01-2.26)	0.38	0.32 (0.08-1.35)	0.15
Ventilator settings and alarm limits set	43 (42.60)	0.46 (0.04-5.68)	0.49	1.19 (0.10-14.14)	1.00
IHT practices related to manual resuscitation bag					
Manual resuscitation bag was not taken	27 (26.70)	0.54 (0.10-3.04)	0.69	0.72 (0.29-2.00)	0.57
Manual resuscitation bag was used	19 (18.80)	0.34 (0.04-2.98)	0.43	3.64 (1.21-11.00)	0.02*
Appropriately sized masks were present	31 (30.70)	1.12 (0.29-4.29)	0.87	1.02 (0.36-2.77)	0.97
PEEP valve was attached to manual resuscitation bag	27 (26.70)	1.04 (0.23-4.67)	1.00	0.92 (0.31-2.68)	1.00
IHT practices related to oxygen cylinder					
Oxygen cylinder was taken	90 (89.10) #	- ^	0.35	2.23 (0.56-8.96)	0.34
Oxygen gauge was checked for leaks	92 (91.10) #	- ^	1.00	0.39 (0.03-4.39)	0.58
Oxygen cylinder was full	84 (83.20) #	- ^	0.35	1.73 (0.49-6.20)	0.54

IHT practices related to emergency drugs					
Adequate volumes of drugs were prepared	85 (84.20) #	1.03 (0.99-1.07)	1.00	1.04 (0.99-1.10)	0.52
Battery life of infusion pump was checked when infusion pumps taken during IHT	47 (46.50)	0.35 (0.05-2.29)	0.27	0.96 (0.18-5.24)	1.00
Drugs taken with during IHT					
Inotropes	72 (71.30)	1.49 (1.29-1.73)	0.02*	0.93 (0.39-2.22)	0.87
Muscle relaxant	27 (26.70)	5.81 (1.70-19.83)	0.01*	1.94 (0.80-4.73)	0.14
Induction agent	12 (11.90)	7.23 (1.86-28.13)	0.002*	0.39 (0.10-1.54)	0.17
Sedation	52 (51.50)	1.6 (0.49-5.28)	0.44	1.46 (0.66-3.23)	0.35
Analgesia	53 (52.50)	2.25 (0.65-7.85)	0.24	0.61 (0.27-1.34)	0.21
Awareness of any IHT guidelines	23 (22.80)	1.35 (1.20-1.53)	0.04*	0.14 (0.04-0.49)	0.001*

Above expected compliance level of 75%; * Significant association (p value < 0.05) ^ Calculation not done due to small numbers in each group

Discussion

Our study's primary finding was that only 27.03% of IHT practices that were investigated were above a compliance rate of 75%. This was unexpectedly low, even after having adjusted the compliance rate. This is below multiple HIC's IHT practices which range from 80.8% to 98.5% compliance [1,18–20]. Multiple studies have highlighted the risks associated with the IHT of critically ill patients and the need to improve patient safety [1,3,12,18,21–23]. There has been a recurrent theme of limited resources and/or the disconnect of knowledge and practice during IHT in this study.

Williams et al. [1] reported that 86.7% of IHT practices were in accordance with the ANZCA/CICM/ACEM intercollegiate guidelines. In this study, the authors conducted a prospective, pre-, and post-interventional study on the IHT of 76 critically ill patients in a tertiary academic hospital in Australia to compare compliance with locally recommended guidelines before and after the introduction of a transport checklist. A statistically significant difference in compliance was noted after the introduction of the transport checklist. This was not noticeable in our study if subsequent transfers were regarded as the post-interventional phase. In our study, there were statistically significant differences in only three IHT practices between first and subsequent transfers: notification of the current condition, use of manual resuscitation bag, and awareness of IHT guidelines.

While acknowledging the importance of all practices recommended by IHT guidelines, it is essential to recognize that some practices are more critical than others. In a prospective observational study by Winter et al. [20] compliance with recommended guidelines during IHT occurred 83.3% of the time. Although this study also highlighted compliance was not fully achieved, key practices were performed 100% of the time, namely the presence of manual resuscitation bag, infusion pumps with alarms and suction availability during IHT [20]. This is significantly higher than what was reported in our study; 73.3% of IHTs had the presence of a manual resuscitation bag and infusion pumps were checked in 46.5% of IHT. The presence of suction during IHT was not assessed because portable suction is not readily available at our institution. Each theatre has its own manual resuscitation bag allocated, and thus is available for use during IHT. This precludes resource limitations as being the only factor for the variance in IHT practices. Possible explanations

include a disconnect between knowledge and practice or a false sense of security due to no adverse events being reported in this study. The variances between other studies in HICs illustrated above could be a result of compliance rates being measured using different methods and guidelines in each publication.

Many of the IHT guidelines [1,11,13,15–17,23,24], often used in HICs, specify the presence of a specialist during IHT. Adherence to this recommendation occurred rarely in this study (5%), this is often due to a limited number of specialists in the department. A specialist is frequently responsible for multiple theatres with a registrar/medical officer present for the full duration of the surgery in each theatre. Due to this limitation, the use of a specialist to idealistically lead each IHT, can be difficult. The majority of participants in this study were registrars, a further explanation for the differences in current IHT practices.

Another explanation for the differences in compliance is the disconnect of knowledge and practice. A substantial proportion of participants, 67.30%, reported no barriers to safe IHT. However, overall compliance in this study was found to be 27.03%. The care for a critically ill patient and the IHT of these patients is included in the curriculum for the Diploma in Anaesthesia (DA), usually completed within the first year of anaesthesia practice, and the Fellowship of the College of Anaesthetists (FCA) final exit exam. Despite having received theoretical knowledge regarding safe transfer of patients, adherence to recommended practices is subpar. This precludes knowledge as being the only factor for the variance in IHT practices and increases the probability of participants transporting critically ill patients with a false sense of security or invulnerability.

Previous publications have found an association between the transferring doctor's level of experience and the occurrence of adverse events, alluding to a difference in practices based on seniority [5,25]. This was not found during our analysis of practices between junior and senior anaesthetists in most categories. When compared to senior anaesthetists, junior anaesthetists were more likely to take a rescue airway device (supraglottic device) during IHT. Senior anaesthetists were more likely to be aware of IHT guidelines and ensure adequate volumes of emergency drugs were available for IHT when compared to junior anaesthetists. The absence of standardized protocols or guidelines in the institution could infer that these practices were guided by previous adverse events. Senior anaesthetists may

also have better insight of the logistics involved and anticipate potential problems that they could encounter during IHT compared to junior anaesthetists. This could be attributed to experience, management of more complex patients, time spent in intensive care units and the amount of time senior anaesthetists have spent in the institution as part of their training.

The majority of the participants (77.20%) were not aware of IHT guidelines. This implies that participants' practices are rooted in tradition, departmental teachings, or long-standing habits, rather than being informed by evidence-based practices. This is in keeping with another South African study by Jordan et al, which showed that more than half of ICU nurses in the public and private sector were not aware of evidence-based practices [26].

Multiple studies have highlighted that the use of IHT checklists improve safety [1,3,12,21–23]. Fanara et al. had subdivided IHT-related adverse events into human-, equipment-, organisational- and patient-related factors [11]. A more significant change in practices was anticipated in our study as the questionnaires were perceived to function as a reminder or checklist for IHT. Despite being exposed to a checklist, practice of anaesthetists in this study did not change significantly except for differences in human- (notification of the current condition and organisational-related factors (awareness of IHT guidelines)). The questionnaires in our study were completed after IHT, while the majority of the aforementioned studies used checklists either during or before IHT and therefore found an improvement in compliance with recommended practices [1,12,16,23]. Although completion of a checklist after an IHT may not be an effective method to improve IHT practices, it is an effective method to assess differences in applied knowledge of IHT. Thus, the introduction of a standardized checklist prior to IHT might provide a more effective method to improve IHT practices in a resource-limited setting.

The recurrent theme of complacency during IHT is also highlighted by the decreased likelihood of a senior anaesthetists taking a supraglottic device with during IHT and anaesthetists during their subsequent IHT were less likely to take a manual bag resuscitator during transfer.

This alludes to anaesthetists, irrespective of their level of experience, not continuously applying what was learnt for safe IHT. This emphasizes the potential

benefit in introducing a standardized protocol or checklist for safe IHT in the LMIC setting. Thus, the introduction of a standardized protocol, checklist or guideline may improve IHT practices in a resource-limited setting. Simulation based training may aid in reducing the discrepancies in safe IHT practices between HICs and LMICs. Manggala et al.[27] have highlighted the benefits of high-fidelity simulations on transferring critically ill patients during the SARS COVID-19 pandemic. These benefits include improved global performance skills, communication, and teamwork during IHT of critically ill patients.

Although there is a lower-than-expected compliance to practices, no adverse events were reported in our study. We only enquired about major adverse events during IHT and not minor physiological changes. The possibility of self-reporting bias must be considered when conducting any questionnaire. The incidence of adverse events during IHT is reported to range from 1.70%-67.90% in other studies [4,9,21,22,28] . Veiga et al [8] found that the incidence of clinical and non-clinical events during IHT in another LMIC with a similar setting as our study was 7.50% and 8.00% respectively.

The majority of participants reported no barriers to adhering to safe IHT guidelines, while 10.9% highlighted that there were no standardized protocols present in the institution and 4% reported inadequate training for IHT. Due to the SARS COVID-19 pandemic, the institution's simulation-based training was suspended during the study period which could have contributed to some participants feeling there was inadequate training. Although competency based training and assessments for IHT are well established in many HICs, 95% of these trainees still reported the need for formal training in IHT [25]. The large disparity in trainees reporting the need for formal training in HIC and our institution highlights the apparent lack of significance placed on transfer care.

A strength of this study is that it focused on IHT practices of anaesthetists with varying levels of experience, transferring a heterogenous population of critically ill patients from different ICUs. The study was conducted prospectively, limiting recall bias. Standardized statistical tests were used, and the study is easily replicable. This allows this data to function as a baseline for other studies in LMICs.

Our study had some limitations. It was conducted in a single institution, investigating the IHT practice of only anaesthetists alone making it difficult to generalise these

findings to practice in other practitioners involved in the care of critically ill patients. Institution specific factors that affect IHT with specific reference to adverse events, namely human- (relating to transport teams), equipment-, organisational- and patient-related factors as described by Fanara et al., could not be delineated during data analysis [11].

The data was collected from self-reported questionnaires after IHT was completed. When comparing adherence rates between self-reported and objective measures of adherence, self-reported adherence rates have been shown to exceed the objective adherence rates [29]. While self-reported questionnaires may provide useful information with minimal impact on service delivery and are somewhat easier to implement in an institution, they are subject to bias and should not be used as the only measure of guideline adherence.

The data for comparisons between first vs subsequent IHT practices are at risk of bias as well. The Hawthorne effect could have been present as participants were aware their IHT practices were being observed. Cognitive bias is another consideration as participants were aware their transfer was the first IHT vs a subsequent IHT.

Another limitation of this study is that IHT of the paediatric population and IHT to remote sites were excluded; and these are known to have a higher incidence of adverse events and may require more specific equipment and expertise [30,31].

This study highlights areas which can improve IHT safety considerably in a relatively cost-effective manner. Further studies should focus on the perceived importance of IHT, what resources are available for safe IHT and the adverse events and/or physiological changes that occur during IHT in resource-limited setting. This will be required to afford the development of an IHT guideline suitable for a resource-limited setting. These studies should include other specialties and be conducted as a multicentre pre- and post-intervention observational study once an appropriate guideline has been proposed.

Bearing in mind the lack of previous studies, this study offers an important contribution to the literature and sets the path for future researchers. Research data can be used as a foundation for institutions in LMICs to develop guidelines tailored to

the resources available in their setting while ensuring adequate patient care is achieved.

Conclusion

This is one of the first studies to describe the current IHT practices in a LMIC and highlights the disparity in practice between HIC and LMIC. Although this is multifactorial, resource limitations, lack of significance placed on IHT practices and education seem to be the major contributors.

More than half of IHT practices interrogated during this study were below the expected compliance level irrespective of the seniority of the anaesthetist or the number of IHTs done during this study.

Practitioners performing IHT need appropriate training and the importance of safe IHT practice must be emphasised. There is a need for a clear and considered guideline to ensure standardized and safe IHT practice.

Ethics approval and consent to participate

Approval to conduct this study was obtained from Human Research Ethics Committee (M210708) and relevant departments.

Informed consent was implied from all participants on completion of the electronic questionnaire.

Consent for publication

Every author gave their approval on the final version of the manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

Bhavic Mawjee: Project administration, Conceptualization, Investigation, Methodology, formal analysis, Writing – original draft, Writing – review & editing.

Sujai Mohanadasan: Project administration, Conceptualization, Writing – review & editing, supervision.

Zainub Jooma: Project administration, Conceptualization, formal analysis, Writing – review & editing, supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. This research was completed as partial fulfilment of a Masters in Anaesthesiology degree at the University of the Witwatersrand.

Data availability

The de-identified datasets used and analysed during the current study are available from the corresponding author on reasonable request.

References:

- [1] Williams P, Karuppiyah S, Greentree K, Darvall J. A checklist for intrahospital transport of critically ill patients improves compliance with transportation safety guidelines. *Aust Crit Care* 2020;33:20–4. <https://doi.org/10.1016/j.aucc.2019.02.004>.
- [2] McLennon M. Use of a Specialized Transport Team for Intrahospital Transport of Critically Ill Patients: *Dimens Crit Care Nurs* 2004;23:225–9. <https://doi.org/10.1097/00003465-200409000-00008>.
- [3] Shirley PJ, Bion JF. Intra-hospital transport of critically ill patients: minimising risk. *Intensive Care Med* 2004;30:1508–10. <https://doi.org/10.1007/s00134-004-2293-6>.
- [4] Geldenhuys L, Wise R, Rodseth R. The impact of a bundled intrahospital transfer protocol on the safety of critically ill patients in a South African Metropolitan Hospital System. *South Afr J Anaesth Analg* 2020:139–48. <https://doi.org/10.36303/SAJAA.2020.26.3.2343>.
- [5] Seilbea L, de Vasconcellos K. Adverse events during the intrahospital transfer of critically ill perioperative patients in a South African tertiary hospital. *South Afr J Anaesth Analg* 2020:131–8. <https://doi.org/10.36303/SAJAA.2020.26.3.2307>.
- [6] Schwebel C, Clec'h C, Magne S, Minet C, Garrouste-Orgeas M, Bonadona A, et al. Safety of Intrahospital Transport in Ventilated Critically Ill Patients: A Multicenter Cohort Study. *Crit Care Med* 2013;41:1919–28. <https://doi.org/10.1097/CCM.0b013e31828a3bbd>.
- [7] Slim J. Safety of Intrahospital Transport in Ventilated Critically Ill Patients: A Multicenter Cohort Study. *J Emerg Med* 2013;45:978. <https://doi.org/10.1016/j.jemermed.2013.10.019>.
- [8] Veiga VC, Postalli NF, Alvarisa TK, Travassos PP, Vale RT da S, Oliveira CZ de, et al. Adverse events during intrahospital transport of critically ill patients in a large hospital. *Rev Bras Ter Intensiva* 2019;31. <https://doi.org/10.5935/0103-507X.20190003>.

- [9] Slave M, Scribante J, Perrie H, Lambat F. Carbon dioxide levels of ventilated adult critically ill post-operative patients on arrival to the intensive care unit. *South Afr J Crit Care* 2023;13–8. <https://doi.org/10.7196/SAJCC.2023.v39i1.655>.
- [10] Quenot J-P, Milési C, Cravoisy A, Capellier G, Mimoz O, Fourcade O, et al. Intrahospital transport of critically ill patients (excluding newborns) recommendations of the Société de Réanimation de Langue Française (SRLF), the Société Française d'Anesthésie et de Réanimation (SFAR), and the Société Française de Médecine d'Urgence (SFMU). *Ann Intensive Care* 2012;2:1. <https://doi.org/10.1186/2110-5820-2-1>.
- [11] Fanara B, Manzon C, Barbot O, Desmettre T, Capellier G. Recommendations for the intra-hospital transport of critically ill patients. *Crit Care* 2010;14:R87. <https://doi.org/10.1186/cc9018>.
- [12] Brunsveld-Reinders AH, Arbous MS, Kuiper SG, de Jonge E. A comprehensive method to develop a checklist to increase safety of intra-hospital transport of critically ill patients. *Crit Care* 2015;19:214. <https://doi.org/10.1186/s13054-015-0938-1>.
- [13] Warren J, Fromm RE, Orr RA, Rotello LC, Horst HM. Guidelines for the inter- and intrahospital transport of critically ill patients*: *Crit Care Med* 2004;32:256–62. <https://doi.org/10.1097/01.CCM.0000104917.39204.0A>.
- [14] Jarden RJ, Quirke S. Improving safety and documentation in intrahospital transport: Development of an intrahospital transport tool for critically ill patients. *Intensive Crit Care Nurs* 2010;26:101–7. <https://doi.org/10.1016/j.iccn.2009.12.007>.
- [15] Australasian College for Emergency Medicine (ACEM), Australian and New Zealand College of Anaesthetists (ANZCA), College of Intensive Care Medicine of Australia and New Zealand (CICM). Guideline for transport of critically ill patients 2015. <https://www.anzca.edu.au/getattachment/bd5938d2-d3ab-4546-a6b0-014b11b99b2f/PS52-Guideline-for-transport-of-critically-ill-patients#page=>.
- [16] Ash A, Whitehead C, Hughes B, Williams D, Nayyar V. Impact of a transport checklist on adverse events during intra-hospital transport of critically ill patients. *Aust Crit Care* 2015;28:49–50. <https://doi.org/10.1016/j.aucc.2014.10.033>.

- [17] Intensive Care Clinical Advisory Group. Intensive care services in New Zealand: a report to the Deputy Director-General, Clinical Services. Wellington, N.Z.: Ministry of Health; 2005.
- [18] Lozada-Martínez ID, Camargo-Martínez W, Agrawal A, Mishra R, Murlimanju BV, Shrivastava A, et al. Letter to the Editor. Intrahospital transport and SAH: possible impact on low- and middle-income countries. *J Neurosurg* 2021;135:1587–8. <https://doi.org/10.3171/2021.3.JNS21734>.
- [19] Lin S-J, Tsan C-Y, Su M-Y, Wu C-L, Chen L-C, Hsieh H-J, et al. Improving patient safety during intrahospital transportation of mechanically ventilated patients with critical illness. *BMJ Open Qual* 2020;9:e000698. <https://doi.org/10.1136/bmjopen-2019-000698>.
- [20] Winter MW. Intrahospital Transfer of Critically Ill Patients; a Prospective Audit within Flinders Medical Centre. *Anaesth Intensive Care* 2010;38:545–9. <https://doi.org/10.1177/0310057X1003800321>.
- [21] Beckmann U, Gillies DM, Berenholtz SM, Wu AW, Pronovost P. Incidents relating to the intra-hospital transfer of critically ill patients: An analysis of the reports submitted to the Australian Incident Monitoring Study in Intensive Care. *Intensive Care Med* 2004;30:1579–85. <https://doi.org/10.1007/s00134-004-2177-9>.
- [22] Farnoosh L, Hossein-nejad H, Beigmohammadi M-T, Seyed-Hosseini-Davarani S-H. Preparation and Implementation of Intrahospital Transfer Protocol for Emergency Department Patients to Decrease Unexpected Events. *Adv J Emerg Med* 2018. <https://doi.org/10.22114/ajem.v0i0.50>.
- [23] Canellas M, Palma I, Pontífice-Sousa P, Rabiais I. Checklist para o transporte intra-hospitalar seguro do doente crítico: A scoping review. *Enferm Glob* 2020;19:525–72. <https://doi.org/10.6018/eglobal.411831>.
- [24] Australasian College for Emergency Medicine, Australian and New Zealand College of Anaesthetists, Joint Faculty of Intensive Care Medicine. Minimum standards for intrahospital transport of critically ill patients. *Emerg Med Australas* 2003;15:202–4. <https://doi.org/10.1046/j.1442-2026.2003.00004.x>.

- [25] Spencer C, Watkinson P, McCluskey A. Training and assessment of competency of trainees in the transfer of critically ill patients. *Anaesthesia* 2004;59:1248–9. <https://doi.org/10.1111/j.1365-2044.2004.04017.x>.
- [26] Jordan PJ, Bowers CA, Morton D. Barriers to implementing evidence-based practice in a private intensive care unit in the Eastern Cape. *South Afr J Crit Care* 2016;32:50. <https://doi.org/10.7196/SAJCC.2016.v32i2.253>.
- [27] Manggala SK, Tantri AR, Sugiarto A, Sianipar IR, Prasetyono TOH. In situ simulation training for a better interprofessional team performance in transferring critically ill patients with COVID-19: a prospective randomised control trial. *Postgrad Med J* 2022;98:617–21. <https://doi.org/10.1136/postgradmedj-2021-141426>.
- [28] Knight P, Maheshwari N, Hussain J, Scholl M, Hughes M, Papadimos T, et al. Complications during intrahospital transport of critically ill patients: Focus on risk identification and prevention. *Int J Crit Illn Inj Sci* 2015;5:256. <https://doi.org/10.4103/2229-5151.170840>.
- [29] Adams A, Soumerai S, Lomas J, Ross-Degnan D. Evidence of self-report bias in assessing adherence to guidelines. *Int J Qual Health Care* 1999;11:187–92. <https://doi.org/10.1093/intqhc/11.3.187>.
- [30] Haydar B, Baetzel A, Elliott A, MacEachern M, Kamal A, Christensen R. Adverse Events During Intrahospital Transport of Critically Ill Children: A Systematic Review. *Anesth Analg* 2020;131:1135–45. <https://doi.org/10.1213/ANE.0000000000004585>.
- [31] Kulshrestha A, Singh J. Inter-hospital and intra-hospital patient transfer: Recent concepts. *Indian J Anaesth* 2016;60:451. <https://doi.org/10.4103/0019-5049.186012>.

Supplementary file 1: Questionnaire

Demographics of Patient

1. Age of patient *

Mark only one oval.

☐ ≤ 12

☐ 13-18

☐ 19-30

☐ 31-40

☐ 40-65

☐ ≥ 65

2. ASA category *

Mark only one oval.

☐ ASA 1

☐ ASA 2

☐ ASA 3

☐ ASA 4

☐ ASA 5

☐ ASA 6

3. Please select reasons for the patient requiring ICU care. More than one option can be selected *

Check all that apply.

- ☐ Mechanical ventilation
☐ Inotropic support
☐ Intensive monitoring
☐ Renal replacement therapy

Other: ☐ _____

Demographics of Participant

4. Age *

Mark only one oval.

- ☐ 25-30
☐ 31-40
☐ 41-60
☐ >60

5. Sex *

Mark only one oval.

- ☐ Female
☐ Male
☐ Prefer not to say
☐ Other: _____

6. Level of training (the most senior person involved in the IHT) *

Mark only one oval.

- ☐ Medical Officer
☐ Registrar
☐ Consultant

7. Years of experience in anaesthesia *

Mark only one oval.

- ☐ ≤ 3 years
☐ > 3 years

8. How many times have you participated in this study? *

Mark only one oval.

- ☐ One
☐ Two
☐ Three
☐ Other: _____

Patient Destination

9. Where was the patient being transferred FROM? *

Mark only one oval.

- ☐ Main ICU (576/567)
☐ Trauma ICU
☐ Cardio-thoracic ICU
☐ Neurosurgery ICU
☐ Theatre

10. Where was the IHT destination? *

Mark only one oval.

- ☐ Main ICU (576/567)
- ☐ Trauma ICU
- ☐ Cardio-thoracic ICU
- ☐ Neurosurgery ICU
- ☐ Theatre

Patient preparation prior to intrahospital transportation (IHT)

11. Was consent checked? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable

12. Were vitals recorded prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No

13. Was IHT destination notified? *

Mark only one oval.

- ☐ Yes
- ☐ No

14. If yes, which of the following actions were done? More than one option can be selected

Check all that apply.

- ☐ Expected time of arrival
- ☐ Notification of current condition of patient
- ☐ Infusions required on arrival (eg: preparation of inotropes)
- ☐ Requirement of ventilatory support

15. Was the route cleared if expected IHT had potential delays ? (eg: Elevators) *

Mark only one oval.

- ☐ Yes
- ☐ No

16. Was IV access confirmed and flushed prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No

17. Which of the following monitors were checked and taken with for IHT? More than one option can be selected *

Check all that apply.

- ☐ Pulse oximetry
- ☐ ECG
- ☐ NIBP or invasive BP monitoring

Other: ☐ _____

Airway management and Ventilatory support

18. Which of the following airway equipment was taken with for IHT? More than one option can be selected *

Check all that apply.

- ☐ Oropharyngeal/ nasopharyngeal airway
- ☐ Supplemental oxygen device (eg: Nasal prong o2)
- ☐ Extra endotracheal tube (appropriate sizes)
- ☐ Emergency supraglottic device
- ☐ Laryngoscope with appropriate blade sizes

Other: ☐ _____

19. Was the airway suctioned prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable

20. Was the ETT size and depth noted prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable, patient not intubated

21. Was the ETT tape/securing device checked and secured? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable, patient not intubated

22. Answer the following questions with regards to transport ventilation practice during IHT: *

Mark only one oval per row.

	Yes	No	Not applicable
Transport ventilator was used	☐	☐	☐
Battery level of ventilator checked	☐	☐	☐
Ventilator checked for leaks	☐	☐	☐
Ventilator settings and alarm limits set	☐	☐	☐

23. Answer the following questions with regards to manual ventilation during IHT: *

Mark only one oval per row.

	Yes	No	Not Available	Not applicable
Manual resuscitation bag was USED	☐	☐	☐	☐
Manual resuscitation bag was NOT taken	☐	☐	☐	☐
Appropriately sized masks were present	☐	☐	☐	☐
PEEP valve was attached to manual resuscitation bag	☐	☐	☐	☐

24. Answer the following questions with regards to the oxygen cylinder used during IHT: *

Mark only one oval per row.

	Yes	No	Not applicable
Oxygen cylinder was NOT taken	☐	☐	☐
Oxygen gauge was checked for leaks	☐	☐	☐
Oxygen cylinder was full	☐	☐	☐

Emergency drugs taken with during IHT

25. Were there adequate volume of drugs prepared prior to IHT? (eg: adrenaline infusion, sedation infusion) *

Mark only one oval.

- ☐ Yes
☐ No
☐ Not applicable

26. Was the battery life of the infusion pump checked? *

Mark only one oval.

- ☐ Yes
☐ No
☐ Not applicable

27. Which of the following drugs were taken with you during IHT? More than one option can be selected *

Check all that apply.

- ☐ Inotropes
☐ Muscle relaxant
☐ Induction agent
☐ Sedation
☐ Analgesia

Intrahospital transport guidelines and adverse events

28. Are you aware of any IHT guidelines? *

Mark only one oval.

☐ Yes

☐ No

29. If yes, for above question, please state which guideline you are aware of.

30. Were there any major adverse events from the IHT? If yes , please select appropriate box. More than one option can be selected. *

Check all that apply.

☐ No adverse event occurred

☐ Accidental extubation

☐ Airway obstruction

☐ Hypotension requiring management

☐ New onset arrhythmia

☐ Initiation of cardiopulmonary resuscitation

☐ Infusion pump failure

☐ Monitor failure

☐ Ventilator failure

Other: ☐ _____

31. Were there any barriers to adhering to guidelines for safe transportation of critically ill patients for current transfer, only select barriers that are applicable for current IHT. More than one option can be selected *

Check all that apply.

- ☐ No barriers
- ☐ Equipment availability
- ☐ Equipment inadequately functioning
- ☐ Time constraints (eg: time pressured from senior or surgical team)
- ☐ Limited consumables (eg: Limited number of syringes or drugs)
- ☐ Inadequate training
- ☐ Inadequate help with transporting patients
- ☐ No standardised protocols present in institution

This content is neither created nor endorsed by Google.

Google Forms

Section 2: Authors guideline for Journal of Critical Care



JOURNAL OF CRITICAL CARE

Improving Patient Care by Integrating Critical Care Systems Knowledge into Practice Behavior

AUTHOR INFORMATION PACK

TABLE OF CONTENTS

• Description	p.1
• Impact Factor	p.1
• Editorial Board	p.1
• Guide for Authors	p.4



ISSN: 0883-9441

DESCRIPTION

The Journal of Critical Care, the official publication of the World Federation of Intensive and Critical Care (WFICC) and the Society for Complex Acute Illness (SCAI), is a leading international, peer-reviewed journal that aims to improve patient care by publishing manuscripts that contain new or updated information that can be integrated into clinical practice, with the exception of experimental research.

The Editorial Board represents an international cross section of individuals actively involved in the disciplines of Critical Care Medicine.

The Journal welcomes the following paper types: Original Research Reviews Meta-analysis and Narrative Reviews: please follow our requirements article [here](#) before submitting your paper Research Letters: should not exceed 600 words, 5 references or 1 figure and/or table. No supplementary material permitted Clinical Challenges Medical Ethics in Intensive Care Medicine Education in Intensive Care Medicine Editorials Letters to Editor: should refer to an article published in *Journal of Critical Care* no more than three months after the articles "Available Online Date" Special Issues

Please see the journal's [Guide For Authors](#) for a more detailed explanation of submission requirements.

IMPACT FACTOR

2021: 4.298 © Clarivate Analytics Journal Citation Reports 2022

EDITORIAL BOARD

Editor-in-Chief

Jan Bakker, New York University Grossman School of Medicine, New York, New York, USA

Managing Editor

Camilla H. Davies, Grosse Pointe, Michigan, United States of America

Associate Editors

Eric Hoste, Ghent University, Ghent, Belgium

Sheila Nainan Myatra, Tata Memorial Hospital, Department of Anaesthesia, Mumbai, India

Jan de Waele, Ghent University, Ghent, Belgium

Section Editors

Ethics

Erwin Kompanje, Erasmus University Rotterdam, Rotterdam, Netherlands

Shahla Siddiqui, Harvard Medical School, Beth Israel Deaconess Medical Center, Department of Anesthesia Critical Care and Pain Medicine, Boston, Massachusetts, United States of America

Education

Glenn Hernandez, Pontifical Catholic University of Chile, Santiago, Chile

Iwan van der Horst, Maastricht University Medical Centre+, Department of Intensive Care Medicine, Maastricht, Netherlands

Cardiovascular

Olfa Hamzaoui, Paris-South University, Antoine Bécclère Hospital, Clamart, France

Jan Belohlavek, General University Hospital in Prague II, Department of Internal Cardiovascular Medicine, Praha, Czechia

Clinical Pharmacology

John Papadopoulos, NYU Langone Health, New York, New York, United States of America

Data Science in Critical Care

Elizabeth Luszczek, University of Minnesota Twin Cities, Minneapolis, Minnesota, United States of America

Joshua M. Tobin, The University of Texas Health Science Center at San Antonio, San Antonio, Texas, United States of America

Infection-Sepsis

Marc Leone, Aix-Marseille Université, Marseille, France

Jeffrey Lipman, University of Queensland, Brisbane, Queensland, Australia

Luciano Cesar Pontes Azevedo, University of Sao Paulo Medical School, SÃO PAULO, Brazil

Renal

Josee Bouchard, University of Montreal, Montréal, Quebec, Canada

Marlies Ostermann, Guy's and St Thomas' NHS Foundation Trust, London, United Kingdom

Neurology

Mathieu van der Jagt, Erasmus Medical Center, Rotterdam, Netherlands

Stephan Mayer, Henry Ford Hospital, Detroit, Michigan, United States of America

Metabolics-Nutrition

Jean Charles Preiser, Erasmus Hospital, Bruxelles, Belgium

Pulmonology

Frank M.P. van Haren, Australian National University Medical School, Canberra, Australia

Elizabeth Wilcox, Toronto Western Hospital, Toronto, Ontario, Canada

Sonography

Jose Luis Diaz-Gomez, Houston, Texas, United States of America

Sedation-Pain-Delirium

Christina Boncyk, Vanderbilt University Medical Center, Nashville, Tennessee, United States of America

Quality and Outcome of Care

Jorge Salluh, D'Or Institute for Research and Education, Rio de Janeiro, Brazil

Social Media

MARCELA LOPES, São Paulo, Brazil

Editorial Board

Sean Bagshaw, University of Alberta, Edmonton, Alberta, Canada

Satish Bhagwanjee, University of Washington Medical Center, Seattle, Washington, United States of America

Mark van den Boogaard, Radboud University Medical Center, Department of Intensive Care, Nijmegen, Netherlands

Hendrik Bracht, Ulm University Medical Center, Department of Anaesthesiology and Intensive Care Medicine, Ulm, Germany

Shari Brosnahan, NYU Langone Health, New York, New York, United States of America

Alexander Cavalcanti, Heart Hospital, SÃO PAULO, Brazil

Mauricio Cecconi, Milan, Italy

Davide Chiumello, Policlinico of Milan, Milano, Italy

Arnaldo Dubin, National University of La Plata, La Plata, Argentina
Martin Duenser, Kepler University Hospital, Linz, Austria
Richard Ferrer, University Hospital MutuaTerrassa, Terrassa, Spain
Seyed Mohammad Reza Hashemian, Shahid Beheshti University of Medical Sciences, Tehran, Iran
Judith Jacobi, Indiana University Health Inc, Indianapolis, Indiana, United States of America
Dimitrios Karakitsos, King Saud Medical City, Riyadh, Saudi Arabia
Ashish K. Khanna, Cleveland, Ohio, United States of America
Andrej Michalsen, Hospital Tettang, Tettang, Germany
Zsolt Molnar, University of Pécs, Pécs, Hungary
Gustavo Ospina Tascon, Icesi University, Valle del Lili Foundation, Cali, Colombia
Stephen M. Pastores, Weill Cornell Medicine, New York, New York, United States of America
Zhiyong Peng, Zhongnan Hospital of Wuhan University, Wuhan, China
Gregory Peterfreund, University of Southern California Keck School of Medicine, Los Angeles, California, United States of America
Joseph W. Rossano, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA
Ram Shanmugam, San Marcos, Texas, United States of America
Andrew F. Shorr, Medstar Washington Hospital Center, Pulmonary and Critical Care Medicine, Washington, District of Columbia, United States of America
Robert Stevens, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA
Yu-Chang Yeh, National Taiwan University Hospital, Department of Anesthesiology, Taipei City, Taiwan
Marieke Zegers, Radboudumc, Nijmegen, Netherlands

Senior Editorial Board
Rinaldo Bellomo, Austin Health, Heidelberg, Victoria, Australia
Philip D. Lumb, University of Southern California Keck School of Medicine, Los Angeles, California, United States of America
Didier Payen, University of Paris Societies and Humanities Faculty, Paris, France
Michael Pinsky, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America
Charles L. Sprung, Jerusalem Bio Park, Jerusalem, Israel
Jean Louis Vincent, Brussels, Belgium

Statistical Board
Ricardo Castro, Pontifical Catholic University of Chile, Santiago, Chile
Irina Chis Ster, St George's University of London Institute for Infection and Immunity, London, United Kingdom
Wan-Jie Gu, Medical College of Nanjing University, Nanjing Drum Tower Hospital, Nanjing, China
Lucas Petri Damiani, HCor Research Institute, São Paulo, Brazil
Fernando Zampieri, São Paulo, Brazil

GUIDE FOR AUTHORS

Your Paper Your Way

We now differentiate between the requirements for new and revised submissions. You may choose to submit your manuscript as a single Word or PDF file to be used in the refereeing process. Only when your paper is at the revision stage, will you be requested to put your paper in to a 'correct format' for acceptance and provide the items required for the publication of your article.

To find out more, please visit the Preparation section below.

Article Types

The Journal welcomes the following paper types:

Original Research

Reviews

Meta-analysis and Narrative Reviews: please follow our requirements article [here](#) before submitting your paper.

Research Letters: should not exceed 600 words, 5 references or 1 figure and/or Table. No supplementary material permitted.

Clinical Challenges

Medical Ethics in Intensive Care Medicine

Education in Intensive Care Medicine

Editorials

Letter to the Editor: should refer to an article published in Journal of Critical Care no more than three months after the articles "Available Online Date".

Special Issues

The Journal of Critical Care provides a forum for the publication of original peer-reviewed articles with the goal of improving patient care by integrating critical care systems knowledge into practice behavior. The journal represents the World Federation of Societies of Intensive and Critical Care Medicine (WFSICCM), an organization of 42 national intensive/critical care societies representing some 32,000 physicians and allied health professionals. With this responsibility to the WFSICCM comes an international focus in systems research in constrained resource environments.

We accept research articles and review articles as well as those in a seminar or tutorial format. Topics covered are all aspects of Health Services Research, the interface of critical care, anesthesiology, and pain, as well as tutorials for residency education core competencies.

For the seminar format, the articles should be directed to the resident or practicing healthcare professional. We are particularly interested in your up-to-date evaluation of the topic. Dealing with subjects of current and sometimes controversial educational and research themes is acceptable. The seminar format lends itself to a more informal presentation. Express your own viewpoints, but feel free to discuss other viewpoints as well. If you are uncertain, please indicate the degree of your uncertainty. We are looking for an absolutely honest evaluation of the topic.

Manuscripts are accepted for consideration on the condition that they are contributed solely to the Journal of Critical Care. No substantial part of a paper may have been or may be published elsewhere, except for an abstract of 200 words or less.

Manuscripts will be critically reviewed by the Editor with appropriate independent referees drawn from the Editorial Board and other experts. Acknowledgments to other investigators for advice or data must be substantiated by written authorization specifically granting permission to authors. Upon submission, authors will be required to fill out an ethics in publishing form. All submissions must also include a conflict of interest statement.

Submission checklist

You can use this list to carry out a final check of your submission before you send it to the journal for review. Please check the relevant section in this Guide for Authors for more details.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded:

Manuscript:

- Include keywords
 - All figures (include relevant captions)
 - All tables (including titles, description, footnotes)
 - Ensure all figure and table citations in the text match the files provided
 - Indicate clearly if color should be used for any figures in print
- Graphical Abstracts / Highlights files (where applicable)*
Supplemental files (where applicable)

Further considerations

- Manuscript has been 'spell checked' and 'grammar checked'
- All references mentioned in the Reference List are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)
- A competing interests statement is provided, even if the authors have no competing interests to declare
- Journal policies detailed in this guide have been reviewed
- Referee suggestions and contact details provided, based on journal requirements

For further information, visit our [Support Center](#).

BEFORE YOU BEGIN

Plagiarism

Plagiarism and duplicate submission are serious acts of misconduct. Plagiarism is defined as unreferenced use of published or unauthorized use of unpublished ideas, and may occur at any stage of planning, researching, writing, or publication. Plagiarism takes many forms, from 'passing off' another's paper as the author's own paper, to copying or paraphrasing substantial parts of another's paper (without attribution), to claiming results from research conducted by others. Plagiarism in all its forms constitutes unethical publishing behavior and is unacceptable. Duplicate submission/publication occurs when two or more papers, without full cross-reference, share the same hypothesis, data, discussion points, or conclusions.

Manuscript Suitability

To understand the scope of publishing in our journal, please refer to our Aims and Scopes in our journal home page. If you would like to check suitability of your paper, please send the abstract and cover letter along with your complete contact information to jcritcare@gmail.com and the editor will get back to you.

Ethics in publishing

Please see our information on [Ethics in publishing](#).

Declaration of interest

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential competing interests include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. Authors must disclose any interests in two places: 1. A summary declaration of interest statement in the title page file (if double anonymized) or the manuscript file (if single anonymized). If there are no interests to declare then please state this: 'Declarations of interest: none'. 2. Detailed disclosures as part of a separate Declaration of Interest form, which forms part of the journal's official records. It is important for potential interests to be declared in both places and that the information matches. [More information](#).

Declaration of generative AI in scientific writing

The below guidance only refers to the writing process, and not to the use of AI tools to analyse and draw insights from data as part of the research process.

Where authors use generative artificial intelligence (AI) and AI-assisted technologies in the writing process, authors should only use these technologies to improve readability and language. Applying the technology should be done with human oversight and control, and authors should carefully review and edit the result, as AI can generate authoritative-sounding output that can be incorrect, incomplete or biased. AI and AI-assisted technologies should not be listed as an author or co-author, or be cited as an author. Authorship implies responsibilities and tasks that can only be attributed to and performed by humans, as outlined in Elsevier's [AI policy for authors](#).

Authors should disclose in their manuscript the use of AI and AI-assisted technologies in the writing process by following the instructions below. A statement will appear in the published work. Please note that authors are ultimately responsible and accountable for the contents of the work.

Disclosure instructions

Authors must disclose the use of generative AI and AI-assisted technologies in the writing process by adding a statement at the end of their manuscript in the core manuscript file, before the References list. The statement should be placed in a new section entitled 'Declaration of Generative AI and AI-assisted technologies in the writing process'.

Statement: During the preparation of this work the author(s) used [NAME TOOL / SERVICE] in order to [REASON]. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

This declaration does not apply to the use of basic tools for checking grammar, spelling, references etc. If there is nothing to disclose, there is no need to add a statement.

Preprints

Please note that [preprints](#) can be shared anywhere at any time, in line with Elsevier's [sharing policy](#). Sharing your preprints e.g. on a preprint server will not count as prior publication (see '[Multiple, redundant or concurrent publication](#)' for more information).

Submission declaration and verification

Submission of an article implies that the work described has not been published previously (except in the form of an abstract, a published lecture or academic thesis, see '[Multiple, redundant or concurrent publication](#)' for more information), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service [Crossref Similarity Check](#). All authors MUST disclose details of preprint posting, including DOI and licensing terms.

Preprint posting on SSRN

In support of [Open Science](#), this journal offers its authors a free preprint posting service. Preprints provide early registration and dissemination of your research, which facilitates early citations and collaboration.

During submission to Editorial Manager, you can choose to release your manuscript publicly as a preprint on the preprint server [SSRN](#) once it enters peer-review with the journal. Your choice will have no effect on the editorial process or outcome with the journal. Please note that the corresponding author is expected to seek approval from all co-authors before agreeing to release the manuscript publicly on SSRN.

You will be notified via email when your preprint is posted online and a Digital Object Identifier (DOI) is assigned. Your preprint will remain globally available free to read whether the journal accepts or rejects your manuscript.

For more information about posting to [SSRN](#), please consult the [SSRN Terms of Use](#) and [FAQs](#).

Use of inclusive language

Inclusive language acknowledges diversity, conveys respect to all people, is sensitive to differences, and promotes equal opportunities. Content should make no assumptions about the beliefs or commitments of any reader; contain nothing which might imply that one individual is superior to another on the grounds of age, gender, race, ethnicity, culture, sexual orientation, disability or health condition; and use inclusive language throughout. Authors should ensure that writing is free from bias, stereotypes, slang, reference to dominant culture and/or cultural assumptions. We advise to seek gender neutrality by using plural nouns ("clinicians, patients/clients") as default/wherever possible to avoid using "he, she," or "he/she." We recommend avoiding the use of descriptors that refer to personal attributes such as age, gender, race, ethnicity, culture, sexual orientation, disability or health condition unless they are relevant and valid. When coding terminology is used, we recommend

to avoid offensive or exclusionary terms such as "master", "slave", "blacklist" and "whitelist". We suggest using alternatives that are more appropriate and (self-) explanatory such as "primary", "secondary", "blocklist" and "allowlist". These guidelines are meant as a point of reference to help identify appropriate language but are by no means exhaustive or definitive.

Reporting sex- and gender-based analyses

Reporting guidance

For research involving or pertaining to humans, animals or eukaryotic cells, investigators should integrate sex and gender-based analyses (SGBA) into their research design according to funder/ sponsor requirements and best practices within a field. Authors should address the sex and/or gender dimensions of their research in their article. In cases where they cannot, they should discuss this as a limitation to their research's generalizability. Importantly, authors should explicitly state what definitions of sex and/or gender they are applying to enhance the precision, rigor and reproducibility of their research and to avoid ambiguity or conflation of terms and the constructs to which they refer (see Definitions section below). Authors can refer to the [Sex and Gender Equity in Research \(SAGER\) guidelines](#) and the [SAGER guidelines checklist](#). These offer systematic approaches to the use and editorial review of sex and gender information in study design, data analysis, outcome reporting and research interpretation - however, please note there is no single, universally agreed-upon set of guidelines for defining sex and gender.

Definitions

Sex generally refers to a set of biological attributes that are associated with physical and physiological features (e.g., chromosomal genotype, hormonal levels, internal and external anatomy). A binary sex categorization (male/female) is usually designated at birth ("sex assigned at birth"), most often based solely on the visible external anatomy of a newborn. Gender generally refers to socially constructed roles, behaviors, and identities of women, men and gender-diverse people that occur in a historical and cultural context and may vary across societies and over time. Gender influences how people view themselves and each other, how they behave and interact and how power is distributed in society. Sex and gender are often incorrectly portrayed as binary (female/male or woman/man) and unchanging whereas these constructs actually exist along a spectrum and include additional sex categorizations and gender identities such as people who are intersex/have differences of sex development (DSD) or identify as non-binary. Moreover, the terms "sex" and "gender" can be ambiguous—thus it is important for authors to define the manner in which they are used. In addition to this definition guidance and the SAGER guidelines, the [resources on this page](#) offer further insight around sex and gender in research studies.

Author contributions

For transparency, we encourage authors to submit an author statement file outlining their individual contributions to the paper using the relevant CRediT roles: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Resources; Software; Supervision; Validation; Visualization; Roles/Writing - original draft; Writing - review & editing. Authorship statements should be formatted with the names of authors first and CRediT role(s) following. [More details and an example](#).

Article transfer service

This journal uses the Elsevier Article Transfer Service to find the best home for your manuscript. This means that if an editor feels your manuscript is more suitable for an alternative journal, you might be asked to consider transferring the manuscript to such a journal. The recommendation might be provided by a Journal Editor, a dedicated [Scientific Managing Editor](#), a tool assisted recommendation, or a combination. If you agree, your manuscript will be transferred, though you will have the opportunity to make changes to the manuscript before the submission is complete. Please note that your manuscript will be independently reviewed by the new journal. [More information](#).

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (see [more information](#) on this). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. [Permission](#) of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations. If

excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has [preprinted forms](#) for use by authors in these cases.

For gold open access articles: Upon acceptance of an article, authors will be asked to complete a 'License Agreement' ([more information](#)). Permitted third party reuse of gold open access articles is determined by the author's choice of [user license](#).

Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. [More information](#).

Elsevier supports responsible sharing

Find out how you can [share your research](#) published in Elsevier journals.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement, it is recommended to state this.

Open access

Please visit our [Open Access page](#) for more information.

Elsevier Researcher Academy

[Researcher Academy](#) is a free e-learning platform designed to support early and mid-career researchers throughout their research journey. The "Learn" environment at Researcher Academy offers several interactive modules, webinars, downloadable guides and resources to guide you through the process of writing for research and going through peer review. Feel free to use these free resources to improve your submission and navigate the publication process with ease.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the [English Language Editing service](#) available from Elsevier's Author Services.

Submission

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

Submit your article

Please submit your article via <https://www.editorialmanager.com/jcrc/default.aspx>.

NIH Public Access Policy Compliance

To comply with the NIH Public Access Policy, Elsevier will deposit to PubMed Central (PMC) author manuscripts on behalf of authors reporting NIH funded research. The NIH policy requires that NIH-funded authors submit to PubMed Central (PMC), or have submitted on their behalf, their peer-reviewed author manuscripts, to appear on PMC no later than 12 months after final publication. Elsevier will send to PMC the final peer-reviewed manuscript, which was accepted for publication and sent to Elsevier's production department, and that reflects any author-agreed changes made in response to peer-review comments. Elsevier will authorize the author manuscript's public access posting 12 months after final publication. Following the deposit by Elsevier, authors will receive further communications from the NIH with respect to the submission. Note: Authors must declare their NIH funding (or the other funding bodies listed below) when completing the copyright transfer form.

PREPARATION

Queries

For questions about the editorial process (including the status of manuscripts under review) or for technical support on submissions, please visit our [Support Center](#).

Peer review

This journal operates a single anonymized review process. All contributions will be initially assessed by the editor for suitability for the journal. Papers deemed suitable are then typically sent to a minimum of two independent expert reviewers to assess the scientific quality of the paper. The Editor is responsible for the final decision regarding acceptance or rejection of articles. The Editor's decision is final. Editors are not involved in decisions about papers which they have written themselves or have been written by family members or colleagues or which relate to products or services in which the editor has an interest. Any such submission is subject to all of the journal's usual procedures, with peer review handled independently of the relevant editor and their research groups. [More information on types of peer review.](#)

Use of word processing software

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the [Guide to Publishing with Elsevier](#)). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

Article structure

Subdivision - unnumbered sections

Divide your article into clearly defined sections. Each subsection is given a brief heading. Each heading should appear on its own separate line. Subsections should be used as much as possible when cross-referencing text: refer to the subsection by heading as opposed to simply 'the text'.

Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

Material and methods

Provide sufficient details to allow the work to be reproduced by an independent researcher. Methods that are already published should be summarized, and indicated by a reference. If quoting directly from a previously published method, use quotation marks and also cite the source. Any modifications to existing methods should also be described.

Results

Results should be clear and concise.

Discussion

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Please clearly indicate the given name(s) and family name(s) of each author and check that all names are accurately spelled. You can add your name between parentheses in your own script behind the English transliteration. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.

• **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. This responsibility includes answering any future queries about Methodology and Materials. **Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.**

• **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Highlights

Highlights are optional yet highly encouraged for this journal, as they increase the discoverability of your article via search engines. They consist of a short collection of bullet points that capture the novel results of your research as well as new methods that were used during the study (if any). Please have a look at the examples here: [example Highlights](#).

Highlights should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point).

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Abstract

Note: Research papers should contain a structured abstract of no more than 200 words which contain the following:

1. Purpose: Why was this study done?
2. Materials and Methods: What was the source of the data generated? How was it obtained?
3. Results: Findings should be objectively reported and statistical significance indicated (if appropriate).
4. Conclusions: Do not include a summary at the end of the paper.

Graphical abstract

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. You can view [Example Graphical Abstracts](#) on our information site.

Keywords

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Formatting of funding sources

List funding sources in this standard way to facilitate compliance to funder's requirements:

Funding: This work was supported by the National Institutes of Health [grant numbers xxxx, yyyy]; the Bill & Melinda Gates Foundation, Seattle, WA [grant number zzzz]; and the United States Institutes of Peace [grant number aaaa].

It is not necessary to include detailed descriptions on the program or type of grants and awards. When funding is from a block grant or other resources available to a university, college, or other research institution, submit the name of the institute or organization that provided the funding.

If no funding has been provided for the research, it is recommended to include the following sentence:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Artwork

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the published version.
- Submit each illustration as a separate file.
- Ensure that color images are accessible to all, including those with impaired color vision.

A detailed [guide on electronic artwork](#) is available.

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. [Further information on the preparation of electronic artwork.](#)

Figure captions

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules and shading in table cells.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Reference links

Increased discoverability of research and high quality peer review are ensured by online links to the sources cited. In order to allow us to create links to abstracting and indexing services, such as Scopus, Crossref and PubMed, please ensure that data provided in the references are correct. Please note that incorrect surnames, journal/book titles, publication year and pagination may prevent link creation. When copying references, please be careful as they may already contain errors. Use of the DOI is highly encouraged.

A DOI is guaranteed never to change, so you can use it as a permanent link to any electronic article. An example of a citation using DOI for an article not yet in an issue is: VanDecar J.C., Russo R.M., James D.E., Ambueh W.B., Franke M. (2003). Aseismic continuation of the Lesser Antilles slab beneath northeastern Venezuela. *Journal of Geophysical Research*, <https://doi.org/10.1029/2001JB000884>. Please note the format of such citations should be in the same style as all other references in the paper.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

Data references

This journal encourages you to cite underlying or relevant datasets in your manuscript by citing them in your text and including a data reference in your Reference List. Data references should include the following elements: author name(s), dataset title, data repository, version (where available), year, and global persistent identifier. Add [dataset] immediately before the reference so we can properly identify it as a data reference. The [dataset] identifier will not appear in your published article.

Preprint references

Where a preprint has subsequently become available as a peer-reviewed publication, the formal publication should be used as the reference. If there are preprints that are central to your work or that cover crucial developments in the topic, but are not yet formally published, these may be referenced. Preprints should be clearly marked as such, for example by including the word preprint, or the name of the preprint server, as part of the reference. The preprint DOI should also be provided.

References in a special issue

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

Reference management software

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support [Citation Style Language styles](#), such as [Mendeley](#). Using citation plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide. If you use

reference management software, please ensure that you remove all field codes before submitting the electronic manuscript. [More information on how to remove field codes from different reference management software.](#)

Reference style

Text: Indicate references by number(s) in square brackets in line with the text. The actual authors can be referred to, but the reference number(s) must always be given.

List: Number the references (numbers in square brackets) in the list in the order in which they appear in the text.

Examples:

Reference to a journal publication:

[1] Van der Geer J, Hanraads JAJ, Lupton RA. The art of writing a scientific article. *J Sci Commun* 2010;163:51–9. <https://doi.org/10.1016/j.Sc.2010.00372>.

Reference to a journal publication with an article number:

[2] Van der Geer J, Hanraads JAJ, Lupton RA. The art of writing a scientific article. *Heliyon*. 2018;19:e00205. <https://doi.org/10.1016/j.heliyon.2018.e00205>

Reference to a book:

[3] Strunk Jr W, White EB. *The elements of style*. 4th ed. New York: Longman; 2000.

Reference to a chapter in an edited book:

[4] Mettam GR, Adams LB. How to prepare an electronic version of your article. In: Jones BS, Smith RZ, editors. *Introduction to the electronic age*, New York: E-Publishing Inc; 2009, p. 281–304.

Reference to a website:

[5] Cancer Research UK. Cancer statistics reports for the UK, <http://www.cancerresearchuk.org/aboutcancer/statistics/cancerstatsreport/>; 2003 [accessed 13 March 2003].

Reference to a dataset:

[dataset] [6] Oguro M, Imahiro S, Saito S, Nakashizuka T. Mortality data for Japanese oak wilt disease and surrounding forest compositions, Mendeley Data, v1; 2015. <https://doi.org/10.17632/xwj98nb39r.1>.

Note shortened form for last page number. e.g., 51–9, and that for more than 6 authors the first 6 should be listed followed by 'et al.' For further details you are referred to 'Uniform Requirements for Manuscripts submitted to Biomedical Journals' (*J Am Med Assoc* 1997;277:927–34) (see also [Samples of Formatted References](#)).

Journal abbreviations source

Journal names should be abbreviated according to the [List of Title Word Abbreviations](#).

Video

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the file in one of our recommended file formats with a preferred maximum size of 150 MB per file, 1 GB in total. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including [ScienceDirect](#). Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our [video instruction pages](#). Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

Data visualization

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions [here](#) to find out about available data visualization options and how to include them with your article.

Supplementary material

Supplementary material such as applications, images and sound clips, can be published with your article to enhance it. Submitted supplementary items are published exactly as they are received (Excel or PowerPoint files will appear as such online). Please submit your material together with the article and supply a concise, descriptive caption for each supplementary file. If you wish to make changes to

supplementary material during any stage of the process, please make sure to provide an updated file. Do not annotate any corrections on a previous version. Please switch off the 'Track Changes' option in Microsoft Office files as these will appear in the published version.

Research data

This journal encourages and enables you to share data that supports your research publication where appropriate, and enables you to interlink the data with your published articles. Research data refers to the results of observations or experimentation that validate research findings. To facilitate reproducibility and data reuse, this journal also encourages you to share your software, code, models, algorithms, protocols, methods and other useful materials related to the project.

Below are a number of ways in which you can associate data with your article or make a statement about the availability of your data when submitting your manuscript. If you are sharing data in one of these ways, you are encouraged to cite the data in your manuscript and reference list. Please refer to the "References" section for more information about data citation. For more information on depositing, sharing and using research data and other relevant research materials, visit the [research data](#) page.

Data linking

If you have made your research data available in a data repository, you can link your article directly to the dataset. Elsevier collaborates with a number of repositories to link articles on ScienceDirect with relevant repositories, giving readers access to underlying data that gives them a better understanding of the research described.

There are different ways to link your datasets to your article. When available, you can directly link your dataset to your article by providing the relevant information in the submission system. For more information, visit the [database linking](#) page.

For [supported data repositories](#) a repository banner will automatically appear next to your published article on ScienceDirect.

In addition, you can link to relevant data or entities through identifiers within the text of your manuscript, using the following format: Database: xxxx (e.g., TAIR: AT1G01020; CCDC: 734053; PDB: 1XFN).

Research Elements

This journal enables you to publish research objects related to your original research – such as data, methods, protocols, software and hardware – as an additional paper in a [Research Elements journal](#).

Research Elements is a suite of peer-reviewed, open access journals which make your research objects findable, accessible and reusable. Articles place research objects into context by providing detailed descriptions of objects and their application, and linking to the associated original research articles. Research Elements articles can be prepared by you, or by one of your collaborators.

During submission, you will be alerted to the opportunity to prepare and submit a manuscript to one of the Research Elements journals.

More information can be found on the [Research Elements](#) page.

Data statement

To foster transparency, we encourage you to state the availability of your data in your submission. This may be a requirement of your funding body or institution. If your data is unavailable to access or unsuitable to post, you will have the opportunity to indicate why during the submission process, for example by stating that the research data is confidential. The statement will appear with your published article on ScienceDirect. For more information, visit the [Data Statement](#) page.

AFTER ACCEPTANCE

Online proof correction

To ensure a fast publication process of the article, we kindly ask authors to provide us with their proof corrections within two days. Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to

MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors. If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

Other Funding Body Policies

Elsevier has also worked with the following funding bodies to ensure that our authors can comply with their policies:

- Arthritis Research Campaign (UK)
- British Heart Foundation (UK)
- Cancer Research (UK)
- Chief Scientist Office
- Department of Health (UK)
- Howard Hughes Medical Institute (US)
- Medical Research Council (UK)
- Wellcome Trust (UK)

For full details on how these policies are implemented, please see complete information at: <https://www.elsevier.com/about/open-science/open-access/agreements>.

AUTHOR INQUIRIES

Visit the [Elsevier Support Center](#) to find the answers you need. Here you will find everything from Frequently Asked Questions to ways to get in touch.

You can also [check the status of your submitted article](#) or find out [when your accepted article will be published](#).

© Copyright 2018 Elsevier | <https://www.elsevier.com>

Section 3: Proposal

Proposal: Assessment of current practices of intrahospital transportation in critically ill patients at an academic hospital in South Africa

Name: Bhavic Mawjee

Student number:361961

Supervisor:	Dr S Mohandas Department of Anaesthesia
Co-supervisor:	Dr Z Jooma Department of Anaesthesia

1.1. Introduction

Critically ill patients have life threatening injuries or illnesses that are often associated with reduced physiological reserves. Critically ill patients can be defined as any patient presenting with dysfunction or failure of one or more vital organs [1,2]. These patients are often required to be transferred from the intensive care unit (ICU) to other departments in the hospital for therapeutic or diagnostic purposes [3]. The risks associated with intrahospital transportation (IHT) need to be outweighed by the benefits from the procedure required [4,5]. Transportation of these patients to various departments require highly trained and skilled medical practitioners with appropriate essential equipment [4].

Multiple studies from high income countries (HICs) and some low- and middle-income countries (LMICs) have highlighted the risk of adverse events occurring during IHT [3,6,7]. The adverse events during IHT can have a direct or indirect role in influencing the patient's risk of morbidity and mortality [7–10]. These adverse events may also lead to an increased incidence of long-term complications such pneumothoraxes, atelectasis, deep vein thrombosis, prolonged length of intensive care unit stay, ventilator-associated pneumonia [11], raised intracranial pressure and cardiac arrest [7,12,13].

Although some adverse events are inevitable, many are preventable [7]. Incidence of adverse events occurring during IHT can vary from 1.7% to 67.9% [7,14] depending on the level of experience of the practitioner present. Physiologic changes that occur during IHT of critically ill patients can have a significant impact on the patient's mortality and morbidity [3,6,13,15]. A South African study noted that in the absence of an adverse event, a medical officer transporting a patient to the ICU perioperatively, as opposed to a registrar, was associated a higher mortality rate [6]. This might allude to different levels of experience or skill sets having an impact on the patient's condition during an IHT [6,7]. Critically ill patients being transported should be monitored and receive physiological support identical to that which would be given while in the ICU [15].

Fanara et al. [16] have subdivided IHT related adverse events into human-, equipment-, organisational- and patient-related factors [16]. This study highlighted patient-related factors such as physiologic changes in response to transportation as

not modifiable [16]. Modifiable factors such as human-related, equipment-related, and organisational-related factors need to be minimised to mitigate adverse events during IHT [8,16]. These factors are often related to poor planning [17,18].

Slave et al. [14] studied IHT of 47 critically ill patients ventilated with a manual resuscitator bag (as opposed to a transport ventilator) from theatre to ICU in a tertiary hospital in Johannesburg, South Africa [14]. They demonstrated an adverse event occurred in 26 patients; of which 41.7% were patient-related, 8.3% were equipment-related and 50% were organisational-related [14]. This finding is in keeping with another South African study conducted at a tertiary hospital in Durban where an adverse events incidence of 23.4% was found [6]. [14]. Meneguín et al. [19] reported an adverse rate of 11.3% in 459 IHT of critically ill patients in a tertiary hospital in Sao Paulo, Brazil. During these IHT, 82.7% were due to patient-related factors, while institutional-related (9.6%) and equipment-related factors (7.7%) followed [19].

Data from the Australian Incident Monitoring study was analysed by Beckmann et al [13]; 191 incidents that occurred during IHT were reviewed. It was found that 39% of adverse events were equipment-related while the other 61% was patient- or staff-related. The cause of most adverse events were multifactorial however human-related factors were highlighted to be the most common [13]. The human-related factors included poor preparation prior to transportation, haste, failure to follow protocols, errors of judgement and errors of problem recognition [13,17]. Geldenhuys et al. [7] devised a bundled intervention attempting to minimize the human-, equipment- and organisational-related factors contributing to adverse events [7]. The interventional bundle included: IHT protocol, a transport bag with emergency drugs based on the protocol and IHT simulation training [7].

Several guideline documents and checklists have been produced in order to decreased IHT related adverse events [1,3,6,7,16,18,20–24]. These are summarised in Table 1 and are recommended practises for safe IHT.

Table 3-1: Overview of content of published intrahospital transportation (IHT) checklists and guidelines

Author		Warren et al. [20]	Fanara et al. [16]	Jarden and Quirke [21]	Brunsveld-Reinders et al. [18]	ANZCA guideline[22]	Williams et al. [3]	Geldenhuys et al. [7]
Year		2004	2010	2010	2015	2015	2020	2020
Administrative guidelines						+		
Pre IHT	Patient assessed prior to transfer	+	+	+		+		+
	IHT to destination beneficial/ required	+					+	+
	Transport team required	+	+	+		+		
	Consent confirmed		+	+			+	
	Equipment and consumables present	+	+	+	+	+	+	+
	Sufficient oxygen level in cylinder	+	+		+	+	+	+
	Sufficient IV medication present	+	+	+	+	+	+	+
	Extra IV fluids and medication	+	+	+	+	+	+	+
	Enteral feeding and insulin stopped				+			
	Check tubes and lines		+	+	+	+	+	+
	Suctioned airway						+	
	Check and set monitor alarms	+	+	+	+	+	+	+
	Check and set transport ventilator alarms	+	+		+	+		+
	Preparation and equipment adapted to procedure. (MRI compatible)	+	+		+		+	
	Register baseline vital signs	+	+	+	+	+	+	+
	Notification of destination department	+			+	+	+	+
	Transport route clear		+			+		+

Author		Warren et al. [20]	Fanara et al. [16]	Jarden and Quirke [21]	Brunsveld-Reinders et al. [18]	ANZCA guideline [22]	Williams et al. [3]	Geldenhuys et al. [7]
Year of publication		2004	2010	2010	2015	2015	2020	2020
During IHT	Equipment checked	+		+	+	+		+
	Record IV fluids or medication given			+	+	+	+	+
	Vital signs recorded every 20min	+		+	+	+	+	+
Post IHT	Enteral feeds and insulin restarted				+			
	Humidifier switched on				+			
	HME filter changed				+			
	Suction bag changed if used			+	+			
	Transport bag restocked				+			+
	Equipment placed on charge							+
	Reported any adverse events on route			+	+	+	+	+
	Handed over to destination staff	+		+	+	+	+	+

Many of these guidelines recommend the IHT is done by ICU trained staff and are led by an intensivist [4,7,16,23,24]. Safely transporting critically ill patients has already become a workplace training objective with formal training provided in many HICs [25]. In a resource limited environment, the use of an intensivist as a team leader during transfers may not be feasible due the scarcity of specialists in the critical care environment [7].

These guidelines and checklists have a common goal of improving the quality of transport and thus minimizing adverse events and physiological changes that occur during IHT, however, they are of limited significance unless the medical practitioner at the bedside translates them into effective action [5].

Williams et al. [3] conducted a prospective pre- and post- interventional study at a tertiary hospital in Melbourne [3]. The primary objective of their study was to assess compliance to the guidelines recommended by the Australian and New Zealand College of Anaesthetists (ANZCA). Williams et al. [3] collected all the data points prior to IHT being commenced with the exception of variables which are normally completed post-IHT such as handover and documentation of observations [3]. Compliance to guidelines was reported as 86.7% prior to the introduction of the checklist [3]. After the introduction of checklist and education given to nursing staff and junior staff, compliance was improved from 86.7% to 90% [3]. Although Williams et al. [3] found no statistically significant improvement in compliance to guidelines, the use of a pre-transport checklist has shown to improve safety and quality in transporting critically ill patients [3,7,18,24]. Ash et al. [24] and Geldenhuys et al. [7] conducted prospective interventional studies focusing on adverse events during IHT after the introduction of checklists. Both these studies highlighted a significant decrease in adverse events during IHT [7,24]. The development of these checklists, by the respective authors, were guided by current guidelines, recommendations and previous checklists published [7,24]. This may allude to the use of checklists improving compliance to IHT guidelines.

1.2. Problem statement

Adverse events often occur during IHT of critically ill patients [6,7,10,24]. While many guidelines exist in HICs, there is a paucity of standardized protocols or guidelines for safe IHT of critically ill patients in a resource limited setting [7]. At the time of this

study, there are no guidelines published by the South African Society of Anaesthesiologists (SASA) or the Critical Care Society of South Africa (CCSSA) for transporting critically ill patients. After discussions with the ICU consultants at the academic hospital where the study will be conducted, no institutional policies regarding IHT of critically ill patients are present. It is the discretion of the medical practitioner transporting a critically patient to determine the equipment and drugs required during the IHT. A review of the current literature did not identify studies conducted on current practices of IHT of critically ill patients perioperatively by anaesthesiologists in South Africa. Conducting a study, focusing on current practises and potential barriers to safe IHT, may highlight patterns unique to resource limited countries. This data has the potential to advocate for the development of standardized protocols and guidelines, adequate resource allocation, improved patient care and a standardized level of training regarding safely transporting critically ill patients perioperatively.

1.3. Aim

1. To assess current practice of IHT of critically ill patients perioperatively by anaesthesiologists
2. To identify possible barriers to practicing recommended guidelines.

1.4. Objectives

1. Describe current practice during IHT of critically ill patients perioperatively by anaesthesiologists
2. Compare practice between junior and senior anaesthesiologists
3. Describe barriers to following recommended IHT guidelines
4. Compare IHT practice of first transfer to practice of IHT of subsequent transfer

1.5. Research assumptions

The following definitions will be used in the study:

Critically ill patient: Any patient requiring cardiovascular, respiratory, renal or neurological organ support, needing ICU admission for mechanical ventilation, inotropic infusions, intensive monitoring or other supportive measures [7].

Intensive Care Unit (ICU): A specialized area in the hospital; providing intensive and specialized medical and nursing care, enhanced capacity for monitoring and multiple

modalities of physiologic organ support to sustain life during a period of life threatening organ dysfunction [2]. The ICUs in this study will include main ICU, trauma ICU, neurosurgery ICU and cardiothoracic ICU.

Anaesthesiologist: HPCSA registered medical practitioner part of the anaesthesia department at Charlotte Maxeke Johannesburg Academic Hospital (CMJAH), which includes medical officers, registrars, and consultants.

Medical officer: Is a qualified doctor practising in the Department of Anaesthesiology under specialist supervision.

Registrar: Is a qualified doctor who is registered with the Health Professional Council of South Africa as a trainee anaesthesiologist.

Consultant: A qualified anaesthesiologist who has completed advanced training in anaesthesia in accordance with the requirements of the Fellowship of the College of Anaesthesiologists and is registered with the Health Professions Council of South Africa, as an anaesthesiologist. A career medical officer, who is a medical officer with more than 10 years of experience, will be considered a consultant in this study.

Junior anaesthesiologists: An anaesthesiologist with experience as an anaesthesiologist for less than or equal to three years.

Senior anaesthesiologists: An anaesthesiologist with experience as an anaesthesiologist for more than three years.

Perioperatively: The period surrounding a surgical procedure, including the preoperative, intraoperative, and postoperative periods.

Intrahospital transport (IHT): Transport of critically ill patients from one area of a hospital to another area within the hospital. IHT in this study will only be limited to transfers between ICU and theatre complex at any time.

Subsequent transfer: the mean of number of IHT done by participants. Used for statistical analysis to compare practices of first IHT to subsequent IHT.

1.6. Demarcation of study field

Study will be conducted CMJAH, Gauteng, South Africa. Areas of focus include main ICU, trauma ICU, neurosurgery ICU, cardiothoracic ICU, and theatre complex.

CMJAH is a quaternary hospital in Johannesburg and is part of the University of Witwatersrand circuit. There are approximately 21 theatres operational daily. The table below summarises the capacity and average number of admissions per month from theatre for each ICU mentioned above.

Table 3-2: Capacity of various ICUs and number of admissions from theatre

ICU	Number of beds available	Discipline ICU serves	Average number of admissions from theatre over a month	
			Pre/post covid peak	During covid peak
Main ICU	24	Medical, Surgical and covid patients	27	11
Cardio-thoracic ICU	6	Cardiothoracic	32	16
Neurosurgery ICU	8	Neurosurgery	16	17
Trauma ICU	12	Trauma	22	24

1.7. Ethical considerations

Approval to conduct this study will be obtained from Human Research Ethics Committee (HREC) (medical), Graduate Studies Committee of the University of the Witwatersrand, the head of department of Anaesthesiology (Appendix A) along with the various Head of Departments for the respective ICUs. Permission to conduct study in various ICUs in CMAJH has been requested (Appendix A).

The study will be conducted in accordance with the Declaration of Helsinki [12] and the South African Guidelines for Good Clinical Practice [13]. The principal investigator will complete the required ethics courses as per the HREC of the university.

The study will not impact service delivery as the online form will not take longer than 10 minutes to complete. This will be achieved by devising questions in a succinct manner and using an easy-to-read checkbox or multiple-choice format. Participants will be able to access the questionnaire via a QR code or link provided via email.

Minimising the risk of spreading the SARS-CoV2 virus will be achieved by requesting participants to complete questionnaire on their own mobile device.

Participants will be given information on study and ethics approval prior to starting the questionnaire (Appendix B). Participation in study will be completely voluntarily and consent will be implied on completion of questionnaire. Once the questionnaire is fully completed, participants will receive voucher to redeem a cup of coffee from theatre complex cafeteria, Olives & Plates. This will be done to encourage participation. Participants will not be allowed to submit the questionnaire if any questions are left blank. This will deter any participants who wish to skew the data in study to receive the voucher code.

Anonymity of the participants will be maintained as no personal data, login details, email address or IP tracking will be necessary to access the questionnaire.

Permission has been obtained for use and adaptation of checklist from Williams et al (Appendix C).

Permission to conduct the study has been provisionally obtained by the CEO of CMJAH, Ms. Bogoshi. (Appendix E). The condition for approval includes HREC approval and post-graduate studies committee approval.

Raw data of study will be stored in a Microsoft excel file for six years. This will be stored on the Google forms database and backed up on a stand-alone pc. The raw data will only be accessible to the authors of the study, and data safety will be ensured by encrypting the file and use of a unique login details to the survey database.

If the results of the survey show that there is underutilisation of equipment available for transportation or poor practice of IHT of critically ill patients, the Head of the Department of Anaesthesiology as well as the Heads of Department of the various ICUs at our institution will be informed and suggestions will be made to improve awareness and education regarding the available equipment and standards of practise.

1.8. Research Methodology

1.8.1. Research design

This study will be a prospective, descriptive study design. Questionnaires will be completed by the most senior anaesthesiologist accompanying patient after IHT.

The study will be a prospective study as data will be collected for the duration of the study by anaesthesiologists after transporting critically ill patients between ICU and theatre complex. The study will describe current practices of IHT of critically ill patients by anaesthesiologists at CMJAH.

1.8.2. Study population

The study's population will be any anaesthesiologist transferring any critically ill patient (as per the definition in this study) between any of the mentioned ICUs and theatre at CMJAH during the study period.

1.8.3. Study sample

1.8.3.1. Sample size

After consultation with a biostatistician, using an average of 100 transfers of critically ill patients between the theatre complex and respective ICUs per month (based on the average number of transfers per month in the last month), a sample size of 75 transfers was determined to be adequate. This calculation was done using StatCalc® with the following assumptions: random sampling for survey/descriptive study with confidence interval of 95% and expected frequency of 75%.

Williams et al. [3] reported 86.7% of IHT prior to intervention were in compliance with recommended guidelines. This study was conducted in a HIC where resource limitations are minimal. It would be unrealistic to assume compliance to IHT guidelines are the same as a HIC due to the limited resources available in the current setting. An expected frequency of 75% is used as the current practices are not known in resource limited countries, thus an assumption of the adherence to standard practise occurring is set as 75%.

The sample size will be determined over the duration of three months or until the sample size is achieved.

1.8.3.2. Sampling method

In this study, non-probability sampling will be used. Individuals in the study population that are readily accessible will be selected to participate, thus a convenience sampling method will be used. Questionnaires will be made available electronically to anaesthesiologists transporting critically ill patients between ICU and theatre complex.

1.8.3.3. Inclusion criteria

- Age of patient > 12
- Critically ill patient
- IHT between ICU and theatre at CMJAH
 - Main ICU (576/567)
 - Trauma ICU
 - Neurosurgical ICU
 - Cardiothoracic ICU
- IHT done during perioperative period
- IHT done by anaesthesiologist

1.8.3.4. Exclusion criteria

- IHT of critically ill patient to ward/ high care setting
 - Obstetric high care
 - Folate wards
 - Trauma high care (377)
 - Neonatal ICU
- Patients placed on cardio-pulmonary bypass or assist devices
- IHT to/from MRI suite or other imaging intervention
- IHT to/from endoscopy suites
- Incomplete questionnaires
- Patients admitted to the ICU not meeting the definition for critically ill as defined in this study

1.8.4. Data collection

Questionnaires will be completed by the most senior anaesthesiologist accompanying patient after IHT .The questionnaire used for study was adapted from Williams et al. [3]. The goal of that study was to assess if compliance to their guidelines would change after the introduction of a checklist. The checklist was developed using the guideline entitled: Transport of critically ill patients. This is an intercollegiate guideline from the Australian and New Zealand College of Anaesthetists (ANZCA) and College of Intensive Care Medicine of Australia and New Zealand (CICM) and Australasian College for Emergency Medicine (ACEM) [22]. The checklist was designed for use in a tertiary academic hospital in a HIC. The variables

mentioned in the checklist were selected by the authors, Williams et al. [3] after discussions with the consultant Anaesthesiologist and Intensivists in a tertiary hospital in Melbourne [3]. These variables covered the major sections of the intercollegiate guidelines.

Adaptations to this checklist have been done to account for use in resource limited environment [7]. The adapted questionnaire contains seven sections namely: demographics of patient, demographics of participant, patient destination, patient preparation, airway management, emergency drugs, adverse events, and potential barriers to adherence to guidelines. The adapted questionnaire can be reviewed in Appendix D.

The general section of the original checklist has been removed; this was done to maintain anonymity of patient data. A demographics section was added to allow for different levels of experience as anaesthesiologist to be compared. Patient destination information was included to attempt to identify if any behaviours changed based on destination.

The questionnaire elaborates on equipment taken during IHT as the department currently does not have a standardized airway/transport bag. End-tidal carbon dioxide monitoring and metaraminol were excluded from the questionnaire as they are not available in our setting.

The subsection, adverse events, was included to identify if any potential major adverse events occurred during IHT. Although this is not the main theme of the study, this information will be used to correlate current practices with adverse events. The major adverse events chosen for the questionnaire was based on the study by Seilbea and de Vasconcellos [2].

The questionnaire will be hosted by Google forms. The questionnaire will not require any personal details or login details. This will be accessible via a link provided or QR code.

Participants will receive an email containing a synopsis of the study and link to the questionnaire. Flyers with the QR code for the questionnaire will be placed in common areas to remind participants of current study; these will include notice boards in ICU and in the theatre complex and the anaesthesia tearoom. The

questionnaire will be completed by the participant using their own mobile phone, minimising the risk of spreading the SARS-CoV2 virus. Participants will be encouraged to complete the questionnaire by offering them a unique voucher code after completion of study. This voucher code will allow participants to redeem a cup of coffee from the theatre complex cafeteria, Olives & Plates. Should they wish to participate in the study, a QR code will be made available to the participant after IHT is completed. Each IHT of a patient, IHT to theatre or IHT to ICU, will be counted as a single event. This allows for multiple transfers of the same patient on the same day or different days to be included. The practices of a patient being transported to theatre from ICU may differ to the IHT of the same patient returning to ICU from theatre, thus each IHT is counted as a single event. These differences may be due to equipment availability, different participants, or changes in clinical condition of patient.

1.8.5. Data analysis

Data collected will be captured onto spread sheets using Microsoft Excel 365. Data collected from the questionnaire will be summarised with the help of a biostatistician by using various descriptive statistics.

Categorical data will be expressed as percentages and frequencies.

Depending on the distribution of the data, variables will be described using means and standard deviations or median and interquartile ranges.

Comparison between junior and senior anaesthesiologists' IHT practices will be done by performing a chi-square test or a fisher exact test if the chi-square test assumptions are violated. A p-value of 0.05 will determine if a statistically significant difference is present. An odds ratio with a 95% confidence interval will be done, comparing junior to senior anaesthetist IHT practices and first to subsequent IHT for each variable.

1.8.6. Significance of the study

A South African study has shown an adverse event rate of 63.3% [7] during IHT of critically ill patients. Safe transporting practices have been shown to reduce adverse events during IHT [9,17,18,26], thus improving patient safety and care.

There is currently a paucity of data of current practises of IHT of critically ill patients in South Africa. There are no known or widely accepted IHT safety guidelines for

critically ill patients available at CMJAH or the hospitals associated with the University of Witwatersrand circuit. This study's results will assist in establishing the baseline practice of anaesthesiologists at CMJAH and may highlight any patterns or potential barriers to practising according to recommended guidelines. These results may also be utilised to develop a standardized protocol and assist in adequate resource allocation. The results may assist in identifying the need for more simulations/teaching about safe practises of transporting critically ill patients perioperatively to be incorporated into registrar training.

Identifying potential barriers to safe practises is another objective of the study. This can have a potential of improving patient care and safety as well as improving the burden of care on fellow anaesthesiologists in the department.

1.9. Validity and reliability of the study

The validity and reliability of the study will be maintained by the following:

- Use of an appropriate study design
- A clearly stated definition of critically ill patients
- Use of an adapted questionnaire which was validated and used in another published study [3]
- Calculation of the sample size in consultation with a biostatistician
- Analysis of data with the assistance of a biostatistician

1.10. Potential limitations of the study

- Limited to current practises done by anaesthesiologists between theatre and ICU
 - Cannot be generalized to transfers to other parts of the hospital
 - Cannot be generalized to other specialties transporting critically ill patients
- Self-reported survey
 - Can lead to bias
 - Over reporting
 - Recall bias
- Survey can act as prompt for checks to be done prior to IHT
 - Risk of skewing data
- IHT done during perioperative period

- Excludes IHT for non-surgical interventions, thus cannot be generalised to all patients who are critically ill. Remote sites have different levels of expertise as opposed to theatre; the latter has a set-up is more appropriate for management and resuscitation of a critically ill patient

1.11. Project outline

1.11.1. Time frame

Table 3-3: Time frame

Month	Feb 21	Mar 21	April 21	May 21	June 21	July 21	Aug 21	Sept 21	Nov 21	Dec 21	Jan 22	Feb 22
Proposal	X	X										
Ethics approval			X	X								
Postgraduate approval			X	X								
Data collection					X	X	X					
Data analysis								X	X			
Submission										X	X	X

1.11.2. Financial plan

Table 3-4: Financial plan

Description	Price per item	Number of pages/items	Copies	Total
Printing proposal	R2/page	33 pages	1	R66
Printing ethics	R2/page	10 pages	2	R40
Coffee	R25	100 cups		R2 500
Final Document	R2/Page	100 pages	3	R600
Binding	R300/copy		1	R300
Total				R3 506

The above-mentioned costs of the study will be incurred by the principal investigator.

1.12. References

- [1] Quenot J-P, Milési C, Cravoisy A, Capellier G, Mimoz O, Fourcade O, et al. Intrahospital transport of critically ill patients (excluding newborns) recommendations of the Société de Réanimation de Langue Française (SRLF), the Société Française d'Anesthésie et de Réanimation (SFAR), and the Société Française de Médecine d'Urgence (SFMU). *Ann Intensive Care* 2012;2:1. <https://doi.org/10.1186/2110-5820-2-1>.
- [2] Marshall JC, Bosco L, Adhikari NK, Connolly B, Diaz JV, Dorman T, et al. What is an intensive care unit? A report of the task force of the World Federation of Societies of Intensive and Critical Care Medicine. *J Crit Care* 2017;37:270–6. <https://doi.org/10.1016/j.jcrc.2016.07.015>.
- [3] Williams P, Karuppiyah S, Greentree K, Darvall J. A checklist for intrahospital transport of critically ill patients improves compliance with transportation safety guidelines. *Aust Crit Care* 2020;33:20–4. <https://doi.org/10.1016/j.aucc.2019.02.004>.
- [4] McLenon M. Use of a Specialized Transport Team for Intrahospital Transport of Critically Ill Patients: *Dimens Crit Care Nurs* 2004;23:225–9. <https://doi.org/10.1097/00003465-200409000-00008>.
- [5] Shirley PJ, Bion JF. Intra-hospital transport of critically ill patients: minimising risk. *Intensive Care Med* 2004;30:1508–10. <https://doi.org/10.1007/s00134-004-2293-6>.
- [6] Seilbea L, de Vasconcellos K. Adverse events during the intrahospital transfer of critically ill perioperative patients in a South African tertiary hospital. *South Afr J Anaesth Analg* 2020:131–8. <https://doi.org/10.36303/SAJAA.2020.26.3.2307>.
- [7] Geldenhuys L, Wise R, Rodseth R. The impact of a bundled intrahospital transfer protocol on the safety of critically ill patients in a South African Metropolitan Hospital System. *South Afr J Anaesth Analg* 2020:139–48. <https://doi.org/10.36303/SAJAA.2020.26.3.2343>.
- [8] Schwebel C, Clec'h C, Magne S, Minet C, Garrouste-Orgeas M, Bonadona A, et al. Safety of Intrahospital Transport in Ventilated Critically Ill Patients: A Multicenter Cohort Study*. *Crit Care Med* 2013;41:1919–28. <https://doi.org/10.1097/CCM.0b013e31828a3bbd>.

- [9] Slim J. Safety of Intrahospital Transport in Ventilated Critically Ill Patients: A Multicenter Cohort Study. *J Emerg Med* 2013;45:978.
<https://doi.org/10.1016/j.jemermed.2013.10.019>.
- [10] Veiga VC, Postalli NF, Alvarisa TK, Travassos PP, Vale RT da S, Oliveira CZ de, et al. Adverse events during intrahospital transport of critically ill patients in a large hospital. *Rev Bras Ter Intensiva* 2019;31. <https://doi.org/10.5935/0103-507X.20190003>.
- [11] Bercault N, Wolf M, Runge I, Fleury J-C, Boulain T. Intrahospital transport of critically ill ventilated patients: A risk factor for ventilator-associated pneumonia—A matched cohort study*. *Crit Care Med* 2005;33:2471–8.
<https://doi.org/10.1097/01.CCM.0000185644.54646.65>.
- [12] Knight P, Maheshwari N, Hussain J, Scholl M, Hughes M, Papadimos T, et al. Complications during intrahospital transport of critically ill patients: Focus on risk identification and prevention. *Int J Crit Illn Inj Sci* 2015;5:256.
<https://doi.org/10.4103/2229-5151.170840>.
- [13] Beckmann U, Gillies DM, Berenholtz SM, Wu AW, Pronovost P. Incidents relating to the intra-hospital transfer of critically ill patients: An analysis of the reports submitted to the Australian Incident Monitoring Study in Intensive Care. *Intensive Care Med* 2004;30:1579–85. <https://doi.org/10.1007/s00134-004-2177-9>.
- [14] Slave M, Scribante J, Perrie H, Lambat F. Carbon dioxide levels of ventilated adult critically ill post-operative patients on arrival to the intensive care unit. *Masters of Medicine in the branch of Anaesthesiology. University of Witwatersrand, 2020.*
Unpublished data
- [15] Esmail R, Banack D, Cummings C, Duffett-Martin J, Shultz J, Thurber T, et al. Is Your Patient Ready for Transport? Developing an ICU Patient Transport Decision Scorecard. *Healthc Q* 2006;9:80–6. <https://doi.org/10.12927/hcq.2013.18376>.
- [16] Fanara B, Manzon C, Barbot O, Desmettre T, Capellier G. Recommendations for the intra-hospital transport of critically ill patients. *Crit Care* 2010;14:R87.
<https://doi.org/10.1186/cc9018>.
- [17] Bergman L, Pettersson M, Chaboyer W, Carlström E, Ringdal M. Improving quality and safety during intrahospital transport of critically ill patients: A critical

incident study. *Aust Crit Care* 2020;33:12–9.

<https://doi.org/10.1016/j.aucc.2018.12.003>.

[18] Brunsveld-Reinders AH, Arbous MS, Kuiper SG, de Jonge E. A comprehensive method to develop a checklist to increase safety of intra-hospital transport of critically ill patients. *Crit Care* 2015;19:214.

<https://doi.org/10.1186/s13054-015-0938-1>.

[19] Meneguín S, Alegre PHC, Luppi CHB. Characterization of the intrahospital transport of critically ill patients. *Acta Paul Enferm* 2014;27:115–9.

<https://doi.org/10.1590/1982-0194201400021>.

[20] Warren J, Fromm RE, Orr RA, Rotello LC, Horst HM. Guidelines for the inter- and intrahospital transport of critically ill patients*: *Crit Care Med* 2004;32:256–62.

<https://doi.org/10.1097/01.CCM.0000104917.39204.0A>.

[21] Jarden RJ, Quirke S. Improving safety and documentation in intrahospital transport: Development of an intrahospital transport tool for critically ill patients.

Intensive Crit Care Nurs 2010;26:101–7. <https://doi.org/10.1016/j.iccn.2009.12.007>.

[22] Australasian College for Emergency Medicine (ACEM), Australian and New Zealand College of Anaesthetists (ANZCA), College of Intensive Care Medicine of Australia and New Zealand (CICM). Guideline for transport of critically ill patients 2015. <https://www.anzca.edu.au/getattachment/bd5938d2-d3ab-4546-a6b0-014b11b99b2f/PS52-Guideline-for-transport-of-critically-ill-patients#page=>. Date accessed: 02 Apr 2021.

[23] Intensive Care Clinical Advisory Group. Intensive care services in New Zealand: a report to the Deputy Director-General, Clinical Services. Wellington, N.Z.: Ministry of Health; 2005. Available from:

<https://www.health.govt.nz/system/files/documents/publications/intensivecare.pdf>.

Date accessed: 28 Mar 2021.

[24] Ash A, Whitehead C, Hughes B, Williams D, Nayyar V. Impact of a transport checklist on adverse events during intra-hospital transport of critically ill patients. *Aust Crit Care* 2015;28:49–50. <https://doi.org/10.1016/j.aucc.2014.10.033>.

- [25] Spencer C, Watkinson P, McCluskey A. Training and assessment of competency of trainees in the transfer of critically ill patients. *Anaesthesia* 2004;59:1248–9. <https://doi.org/10.1111/j.1365-2044.2004.04017.x>.
- [26] Bergman LM, Pettersson ME, Chaboyer WP, Carlström ED, Ringdal ML. Safety Hazards During Intrahospital Transport: A Prospective Observational Study. *Crit Care Med* 2017;45:e1043–9. <https://doi.org/10.1097/CCM.0000000000002653>.

1.13. Appendices

1.13.1. Appendix A: Permission from the Head of Department

Dr B Mawjee

Department of Anaesthesiology

University of the Witwatersrand

Bhavicmawjee@gmail.com

March 2021

Dear Prof (Department)

I am currently conducting research for the completion of my M Med in Anaesthesiology titled: Assessment of current practices of intrahospital transportation (IHT) in critically ill patients at an academic hospital in South Africa

I hereby request permission to conduct my research by means of a self-administered electronic questionnaire over a three-month period. This will be completed by the most senior anaesthesiologist accompanying the patient after IHT. The protocol for my proposed study is attached to this email. The aims of this study are to assess current practice of IHT of critically ill patients perioperatively by anaesthesiologists at CMJAH and to identify possible barriers to practicing recommended guidelines.

A South African study has shown an adverse event rate of 63.3% during IHT of critically ill patients. Safe transporting practices have been shown to reduce adverse events during IHT, thus improving patient safety and care.

There are no known or widely accepted IHT safety guidelines for critically ill patients available at CMJAH or the hospitals associated with the University of Witwatersrand circuit. This study's results will assist in establishing the baseline practice of anaesthesiologists at CMJAH and may highlight any patterns or potential barriers to practising according to recommended guidelines. These results may also be utilised to develop a standardized protocol and assist in adequate resource allocation.

Should you require more information, please do not hesitate to contact me on 0711917585 or Bhavicmawjee@gmail.com.

Permission to conduct the study will be obtained from the Human Research Ethics Committee (Medical) and the Graduate Studies Committee of the University of the Witwatersrand.

Thanking you in advance. Sincerely

Dr Bhavic Mawjee

Registrar: Department of Anaesthesiology

I hereby give permission for this study

Name

Signature

Date

1.13.2. Appendix B: Information sheet

Dear Colleague,

My name is Bhavic Mawjee. I am a registrar in the Department of Anaesthesiology at the University of the Witwatersrand, and I am currently busy with my MMed degree. I would like to invite you to participate in my research study titled: Assessment of current practices of intrahospital transportation (IHT) in critically ill patients at an academic hospital in South Africa. Identifying the current practice of transporting critically ill patients between ICU and theatre complex within the department will assist in the continued professional development in the department, improve patient outcomes and reduce hospital costs.

Participation is completely voluntary; consent will be implied on completion of this self-administered questionnaire. The questionnaire is anonymous. Confidentiality will be maintained as only the researcher and supervisor will have access to the raw data. Should you not wish to participate you can withdraw from the study at any time prior to the submission of the questionnaire.

The questionnaire should not take longer than 10 minutes to complete and participants are encouraged not to share the information provided.

Before completion of this questionnaire, please ensure that you understand the above information. Any questions regarding this study or want access of the final report, you can contact the following:

Bhavic Mawjee – 0711917585, bhavicmawjee@gmail.com

My supervisors:

Dr Sujai Mohanadasana at iajus24@gmail.com

Dr Zainub Jooma at zainub.jooma@gmail.com

The final report can be accessed at the Wits Institutional Repository Environment on DSpace (WIReDSpace) at www.wiredspace.wits.ac.za.

Permission to conduct the study has been obtained from the Human Research Ethics Committee (Medical) of the University of the Witwatersrand, Johannesburg. A principal function of this Committee is to safeguard the rights and dignity of all human subjects who agree to participate in a research project and the integrity of the research.

If you have any concern regarding the way the study is being conducted, please contact the Chairperson of this Committee, Professor Clement Penny, who may be contacted on telephone number 011 717 2301, or by e-mail on Clement.Penny@wits.ac.za. The telephone numbers for the Committee secretary are 011 717 2700/1234 and the e-mail addresses are Zanele.Ndlovu@wits.ac.za and Rhulani.Mukansi@wits.ac.za

Thank you for taking time to read this letter.

March 2021

1.13.3. Appendix C: Permission to use and adapt checklist

From: Peter Williams
Sent: Monday, 14 September 2020 10:42
To: Bhavic Mawjee
Subject: Re: Request access to questionnaire assessing compliance with transportation safety guidelines

Hi Bhavic,

No problem. Please find it attached. I hope it helps.

Cheers,

Pete

On Tue, 8 Sep 2020 at 15:28, Bhavic Mawjee <bhavicmawjee@gmail.com> wrote:

Good day Dr Williams

My name is Bhavic Mawjee, I'm a medical officer in anaesthesia at Charlotte Maxeke Johannesburg Academic Hospital, Johannesburg, South Africa. I'd like to request permission from you and your colleagues to gain access and adapt the questionnaire used in your study "A checklist for intrahospital transport of critically ill patients improves compliance with transportation safety guidelines".

I intend to conduct a study for my Mmed, "Assessment of adherence of intrahospital transportation safety guidelines in critically ill patients perioperatively at Wits circuit".

We often transport critically ill patients between the Intensive care unit and theatre, however there are varying levels of compliance to transportation safety guidelines.

The aim of the study would be: to assess adherence to recommended intrahospital transfer guidelines and to identify possible barriers to practicing these recommended guidelines.

This study intends to highlight the deficiencies that occur during the transportation of critically ill patients and thus motivate for more education and attention regarding transportation safety guidelines. The ultimate goal is to ensure patient safety and minimize adverse events during transporting critically ill patients perioperatively.

Should you require more information, please do not hesitate to contact me

Take care and hope you have a good day.

1.13.4. Appendix D: Data Collection Sheet

Transferring a critically patient between theatre and ICU? Please assist by participating in my MMed Study, Assessment of current practices of intrahospital transportation (IHT) in critically ill patients at an academic hospital in South Africa.

INCLUSION CRITERIA	EXCLUSION CRITERIA
Critically ill patient*	Patients placed on cardio-pulmonary bypass
IHT between Theatre at CMJAH and Main ICU (576/567), Trauma ICU, Neurosurgical ICU or Cardiothoracic ICU	IHT of critically ill patient to ward/ high care setting including Obstetric high care, Folateng wards, Trauma high care (377) and Neonatal ICU
IHT done during perioperative period	IHT to/from MRI suite or other imaging intervention
IHT done by Anaesthesiologist	IHT to/from endoscopy suites
Age > 12 years old	Patient not meeting the definition of critically ill patient

*Critically ill patient: any patient requiring EITHER cardiovascular or respiratory or renal or neurological organ support, needing ICU admission for mechanical ventilation, inotropic infusions, intensive monitoring, or other supportive measures.

Please complete questionnaire using this link or QR code.

Google forms Link and QR code: <https://forms.gle/g61uwHc7uFzoW4Sh9>



Demographics of Patient

1. Age of patient *

Mark only one oval.

- ☐ ≤ 12
- ☐ 13-18
- ☐ 19-30
- ☐ 31-40
- ☐ 40-65
- ☐ ≥ 65

2. ASA category *

Mark only one oval.

- ☐ ASA 1
- ☐ ASA 2
- ☐ ASA 3
- ☐ ASA 4
- ☐ ASA 5
- ☐ ASA 6

3. Please select reasons for the patient requiring ICU care. More than one option can be selected *

Check all that apply.

- ☐ Mechanical ventilation
☐ Inotropic support
☐ Intensive monitoring
☐ Renal replacement therapy

Other: ☐ _____

Demographics of Participant

4. Age *

Mark only one oval.

- ☐ 25-30
☐ 31-40
☐ 41-60
☐ >60

5. Sex *

Mark only one oval.

- ☐ Female
☐ Male
☐ Prefer not to say
☐ Other: _____

6. Level of training (the most senior person involved in the IHT) *

Mark only one oval.

- ☐ Medical Officer
☐ Registrar
☐ Consultant

7. Years of experience in anaesthesia *

Mark only one oval.

- ☐ ≤ 3 years
☐ > 3 years

8. How many times have you participated in this study? *

Mark only one oval.

- ☐ One
☐ Two
☐ Three
☐ Other: _____

Patient Destination

9. Where was the patient being transferred FROM? *

Mark only one oval.

- ☐ Main ICU (576/567)
☐ Trauma ICU
☐ Cardio-thoracic ICU
☐ Neurosurgery ICU
☐ Theatre

10. Where was the IHT destination? *

Mark only one oval.

- ☐ Main ICU (576/567)
- ☐ Trauma ICU
- ☐ Cardio-thoracic ICU
- ☐ Neurosurgery ICU
- ☐ Theatre

Patient preparation prior to intrahospital transportation (IHT)

11. Was consent checked? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable

12. Were vitals recorded prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No

13. Was IHT destination notified? *

Mark only one oval.

- ☐ Yes
- ☐ No

14. If yes, which of the following actions were done? More than one option can be selected

Check all that apply.

- ☐ Expected time of arrival
- ☐ Notification of of current condition of patient
- ☐ Infusions required on arrival (eg: preparation of inotropes)
- ☐ Requirement of ventilatory support

15. Was the route cleared if expected IHT had potential delays ? (eg: Elevators) *

Mark only one oval.

- ☐ Yes
- ☐ No

16. Was IV access confirmed and flushed prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No

17. Which of the following monitors were checked and taken with for IHT? More than one option can be selected *

Check all that apply.

- ☐ Pulse oximetry
- ☐ ECG
- ☐ NIBP or invasive BP monitoring

Other: ☐ _____

Airway management and Ventilatory support

18. Which of the following airway equipment was taken with for IHT? More than one option can be selected *

Check all that apply.

- ☐ Oropharyngeal/ nasopharyngeal airway
- ☐ Supplemental oxygen device (eg: Nasal prong o2)
- ☐ Extra endotracheal tube (appropriate sizes)
- ☐ Emergency supraglottic device
- ☐ Laryngoscope with appropriate blade sizes

Other: ☐ _____

19. Was the airway suctioned prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable

20. Was the ETT size and depth noted prior to IHT? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable, patient not intubated

21. Was the ETT tape/securing device checked and secured? *

Mark only one oval.

- ☐ Yes
- ☐ No
- ☐ Not applicable, patient not intubated

22. Answer the following questions with regards to transport ventilation practice during IHT: *

Mark only one oval per row.

	Yes	No	Not applicable
Transport ventilator was used	☐	☐	☐
Battery level of ventilator checked	☐	☐	☐
Ventilator checked for leaks	☐	☐	☐
Ventilator settings and alarm limits set	☐	☐	☐

23. Answer the following questions with regards to manual ventilation during IHT: *

Mark only one oval per row.

	Yes	No	Not Available	Not applicable
Manual resuscitation bag was USED	☐	☐	☐	☐
Manual resuscitation bag was NOT taken	☐	☐	☐	☐
Appropriately sized masks were present	☐	☐	☐	☐
PEEP valve was attached to manual resuscitation bag	☐	☐	☐	☐

24. Answer the following questions with regards to the oxygen cylinder used during IHT: *

Mark only one oval per row.

	Yes	No	Not applicable
Oxygen cylinder was NOT taken	☐	☐	☐
Oxygen gauge was checked for leaks	☐	☐	☐
Oxygen cylinder was full	☐	☐	☐

Emergency drugs taken with during IHT

25. Were there adequate volume of drugs prepared prior to IHT? (eg: adrenaline infusion, sedation infusion) *

Mark only one oval.

- ☐ Yes
☐ No
☐ Not applicable

26. Was the battery life of the infusion pump checked? *

Mark only one oval.

- ☐ Yes
☐ No
☐ Not applicable

27. Which of the following drugs were taken with you during IHT? More than one option can be selected *

Check all that apply.

- ☐ Inotropes
☐ Muscle relaxant
☐ Induction agent
☐ Sedation
☐ Analgesia

Intrahospital transport guidelines and adverse events

28. Are you aware of any IHT guidelines? *

Mark only one oval.

☐ Yes

☐ No

29. If yes, for above question, please state which guideline you are aware of.

30. Were there any major adverse events from the IHT? If yes , please select appropriate box. More than one option can be selected. *

Check all that apply.

- ☐ No adverse event occurred
- ☐ Accidental extubation
- ☐ Airway obstruction
- ☐ Hypotension requiring management
- ☐ New onset arrhythmia
- ☐ Initiation of cardiopulmonary resuscitation
- ☐ Infusion pump failure
- ☐ Monitor failure
- ☐ Ventilator failure

Other: ☐ _____

31. Were there any barriers to adhering to guidelines for safe transportation of critically ill patients for current transfer, only select barriers that are applicable for current IHT. More than one option can be selected *

Check all that apply.

- ☐ No barriers
- ☐ Equipment availability
- ☐ Equipment inadequately functioning
- ☐ Time constraints (eg: time pressured from senior or surgical team)
- ☐ Limited consumables (eg: Limited number of syringes or drugs)
- ☐ Inadequate training
- ☐ Inadequate help with transporting patients
- ☐ No standardised protocols present in institution

This content is neither created nor endorsed by Google.

Google Forms

1.13.5. Appendix E: Permission from CEO of CMJAH



Enquiries: Ms. TT Mahlangu

Email: Thandi.Mahlangu4@gauteng.gov.za

Tel: 011 488 3365

Ref: 1/7/2

Date: 17 August 2021

GP_202106_040

To: Dr Bhavic Mawjee

RE: PROVISIONAL APPROVAL

TITLE: ASSESSMENT OF CURRENT PRACTICES OF INTRAHOSPITAL TRANSPORTATION IN CRITICALLY ILL PATIENTS AT AN ACADEMIC HOSPITAL IN SOUTH AFRICA

Permission to conduct the above-mentioned study is provisionally granted subject to:

1. Submitting an Ethics Clearance Certificate from the University of Witwatersrand Human Research and Ethics Committee (HREC).
2. Submitting a proof of post graduation approval.

Please submit the above documents at your earliest convenience for final approval to be granted. Please note that your study cannot commence until a final approval letter is issued.

Supported / ~~Not Supported~~

Signed by: Jayshina Punwasi
Signed at: 2021-08-19 16:07:24 +02:00
Reason: Witnessing Jayshina Punwasi

Lawtrust

Dr J. Punwasi
Clinical Director

Approved / ~~Not Approved~~

Signed by: Gladys Magugudi Bogoshi
Signed at: 2021-08-20 06:21:30 +02:00
Reason: Witnessing Gladys Magugudi Bo

GAUTENG PROVINCE

Ms. Bogoshi

CEO: CMJAH

Section 4: Annexures

1.14. Annexure 1: Postgraduate Approval letter



Private Bag 3 Wits, 2050
Fax: 027117172119
Tel: 02711 7172076

Reference: Mrs Sandra Benn
E-mail: sandra.benn@wits.ac.za

06 January 2023
Person No: 361961
PAG


Dr B Mawjee
Unit 19,06 Bompas Road
Dunkeld West
2196
South Africa

Dear Dr Bhavic Mawjee

Master of Medicine in Anaesthesia: Approval of Title

We have pleasure in advising that your proposal entitled *Assessment of current practices of intrahospital transportation in critically ill patients at an academic hospital in South Africa* has been approved. Please note that any amendments to this title have to be endorsed by the Faculty's higher degrees committee and formally approved.

Yours sincerely

A handwritten signature in black ink, appearing to read 'Sandra Benn'.

Mrs Sandra Benn
Faculty Registrar
Faculty of Health Sciences

1.15. Annexure 2: Human research ethics committee clearance certificate



R49 Dr B Mawjee

HUMAN RESEARCH ETHICS COMMITTEE (MEDICAL) CLEARANCE CERTIFICATE NO. M210708

NAME:
(Principal Investigator)

Dr B Mawjee

DEPARTMENT:

School of Clinical Medicine
Department of Anaesthesiology
Medical School
University

PROJECT TITLE:

*Assessment of current practices of intrahospital
transportation in critically ill patients at an academic
hospital in South Africa*

DATE CONSIDERED:

2021/07/30

DECISION:

Approved unconditionally

CONDITIONS:


NOTE:

If contact information regarding student study participants is required,
please contact the Registrar's office - <Nicoleen.Potgieter@wits.ac.za>

SUPERVISOR:

Mr S Mohanadasan and Dr Z Jooma

APPROVED BY:


Dr CB Penny, Chairperson, HREC (Medical)

DATE OF APPROVAL:

2021/10/08

This Clearance Certificate is valid for 5 years from the date of approval. An extension may be applied for.

1.16. Annexure 3: Plagiarism/Turnitin report

Write up final draft bhavic
23.docx
by Bhavic Mawjee

Submission date: 05-Jun-2023 08:17PM (UTC+0200)
Submission ID: 2109684536
File name: Write_up_final_draft_bhavic_23.docx (874.76K)
Word count: 6643
Character count: 37208

Write up final draft bhavic 23.docx

ORIGINALITY REPORT

11 %	10 %	9 %	0 %
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.ajol.info Internet Source	2 %
2	hdl.handle.net Internet Source	1 %
3	deepblue.lib.umich.edu Internet Source	1 %
4	www.ncbi.nlm.nih.gov Internet Source	1 %
5	www.researchgate.net Internet Source	1 %
6	coek.info Internet Source	1 %
7	www.scielo.br Internet Source	<1 %
8	docksci.com Internet Source	<1 %
9	journals.lww.com Internet Source	<1 %

10	www.discoveryjournals.org Internet Source	<1 %
11	api.cqc.org.uk Internet Source	<1 %
12	link.springer.com Internet Source	<1 %
13	samajournals.co.za Internet Source	<1 %
14	docplayer.net Internet Source	<1 %
15	"Supplement 2 September 2011", Intensive Care Medicine, 2011 Publication	<1 %
16	C. Spencer. "Training and assessment of competency of trainees in the transfer of critically ill patients", Anaesthesia, 12/2004 Publication	<1 %
17	Peter Williams, Sathappan Karuppiah, Kate Greentree, Jai Darvall. "A checklist for intrahospital transport of critically ill patients improves compliance with transportation safety guidelines", Australian Critical Care, 2020 Publication	<1 %
18	josr-online.biomedcentral.com Internet Source	<1 %

19	admin.concern.net Internet Source	<1 %
20	gs.amegroups.com Internet Source	<1 %
21	Carmen Voigt-Graf. "Migration and Transnational Families in Fiji: Comparing Two Ethnic Groups", International Migration, 2008 Publication	<1 %
22	ccn.aacnjournals.org Internet Source	<1 %
23	mdpi-res.com Internet Source	<1 %
24	rbti.org.br Internet Source	<1 %
25	www.karger.com Internet Source	<1 %
26	Tanakorn Apornpong, Beatriz Grinsztejn, Michael Hughes, Justin Ritz et al. "Antiretroviral hair levels, self-reported adherence, and virologic failure in second-line regimen patients in resource-limited settings", AIDS, 2021 Publication	<1 %
27	core.ac.uk Internet Source	<1 %

28 edoc.site <1 %
Internet Source

29 q8c7r2f7.stackpathcdn.com <1 %
Internet Source

Exclude quotes On

Exclude matches < 8 words

Exclude bibliography On