

REFERENCES

- Abdel-Fattah, Y.R., Saeed, H.M., Gohar, Y.M. and El-Baz, M.A. (2005). Improved production of *Pseudomonas aeruginosa* uricase by optimization of process parameters through statistical experimental designs. *Proc. Biochem.*; **40**: 1707-1714.
- Ahlin, P., Kristi, J., Kristi, A. and Vrečer, F. (2002). Investigation of polymeric nanoparticles as carriers of enalaprilat for oral administration. *Int. J. Pharm.*; **239**: 113-120.
- Anderson, N.H., Bauer, M., Boussac, N., Khan-Malek, R., Munden, P. and Sardaro, M. (1998). An evaluation of fit factors and dissolution efficiency for the comparison of in vitro dissolution profiles. *J. Pharm. Biomed. Anal.*; **17**: 811-822.
- Aquil, M. and Ali, A. (2002). Monolithic matrix type transdermal drug delivery systems of pinacidil monohydrate: *in vitro* characterization. *Eur. J. Pharm. Biopharm.*; **54**: 161-164.
- Batycky, R.P., Hanes, J., Langer, R. and Edwards, D.A. (1997). A theoretical approach of erosion and macromolecular drug release from biodegradable microspheres. *J. Pharm. Sci.*; **86**: 1464-1477.
- Bloomfield, N.S. and Butler, W.C. (2000). Robustness testing using experimental design of a flow-through dissolution method for a product where the actives have markedly differing solubility properties. *Int. J. Pharm.*; **209**: 55-61.
- Brannon-Peppas, L. (1997). Polymers in controlled drug delivery. *Medical Plastics and Biomaterials Magazine*. MPB Article Index, pgs 1-15.
- Chafi, N., Kolli, M., Vergnaud, J.M. and Montheard, J.P. (1994). Amines release from schiff bases polymers and diffusion from dosage forms with eudragit RL in acidic medium. *J. Appl. Polym. Sci.*; **43**: 1837-1847.

- Chafi, N., Montheard, J.P. and Vergnaud, J.M. (1992). Release of 2-aminothiazole from polymeric carriers. *Int. J. Pharm.*; **67**: 265-274.
- Chattaraj, S.C., Swarbrick, J. and Kanfer, I. (1995). A simple diffusion cell to monitor drug release from semi-solid dosage forms. *Int. J. Pharm.*; **120**: 119-124.
- Chen, S., Wu, Y., Mi, F., Lin, Y., Yu, L. and Sung, H. (2004). A novel pH-sensitive hydrogel composed of N, O-carboxylmethyl chitosan and alginate cross-linked by genipin for protein drug delivery. *J. Control Release.*; **96**: 285-300.
- Chourasia, M.K. and Jain, S.K. (2003). Pharmaceutical approaches to colon targeted drug delivery systems. *J. Pharm. Pharmaceut. Sci.*; **6**: 33-66.
- Chu, L., Liang, Y., Chen, W., Ju, X. and Wang, H. (2004). Preparation of glucose-sensitive microcapsules with a porous membrane and functional gates. *Coll. Surf. B: Biointerf.*; **37**:9-14.
- Cole, E.T., Scott, R.A., Conner, A.L., Wilding, I.R., Petereit, H.U., Schminke, C., Beckert, T. and Cade, D. (2002). Enteric coated HPMC capsules designed to achieve intestinal targeting. *Int. J. Pharm.*; **231**: 83-95.
- Colombo, P., Bettini, R., Santi, P. and Peppas, N.A. (2000). Swellable matrices for controlled drug delivery: gel-layer behaviour, mechanisms and optimal performance. *PSST.*; **3**: 198-204.
- Correia, M.A. (2000). In "Drug Biotransformation", ed. Katzung, B.G. in *Basic and Clinical Pharmacology*. 8th edition, pgs 51-63.
- Costa, F.O., Sousa, J.J.S., Pais, A.A.C.C. and Formosinho, S.J. (2003). Comparison of dissolution profiles of ibuprofen pellets. *J. Control Release.*; **89**: 199-212.
- Cui, X. and Yan, D. (2005). Preparation, characterization and crystalline transitions of odd-even polyamides 11,12 and 11,10. *Eur. Polym. J.*; **41**: 863-870.

- Cui, X., Liu, Z. and Yan, D. (2004). Synthesis and characterization of novel even-odd nylons based on undecanoic acid. *Eur. Poly. J.*; **40**: 1111-1118.
- Culotta, E. and Koshland jr., D.E. (1991). Buckyballs: a wide open playing field for chemists. *Sci.*, **254**: 1706-1709.
- Davis, S.S. and Illum, L. (1988). Polymeric microspheres as drug carriers. *Biomat.*; **9**: 111-115.
- Dolorico, A.M.T., Ramin, T., Hung, V.O. and Ronald, N.G. (2003). Short-term and long-term visual and astigmatic results of an opposing 10-0 nylon double running suture technique for penetrating keratoplasty. *J. Americ. Coll. Surg.*; **197**: 991-999.
- Doria-Serrano, M.C., Ruis-Trevino, F.A., Rios-Arciga, C., Hernandez-Esparza, M. and Santiago, P. (2001). Physical characteristics of poly (vinylalcohol) and calcium alginate hydrogels for the immobilization of activated sludge. *Biomacromolecules.*; **2**: 568-574.
- Douglas, S.J., Davis, S.S. and Illum, L. (1987). Nanoparticles in drug delivery. *Ther. Drug. Carr. Syst.*; **3**: 233-261.
- Dürig, T. and Fassihi, R. (2000). Evaluation of floating and sticking extended release delivery systems: an unconventional dissolution test. *J. Control Release.*; **67**: 37-44.
- El-Sherbiny, I.M., Lins, R.J., Abdel-Bary, E.M. and Harding, D.R.K. (2005). Preparation, characterization, swelling and *in-vitro* drug release behaviour of poly [N-acryloylglycine-chitosan] inter-polymeric pH and thermally-responsive hydrogels. *Eur. Polym. J.*; **41**: 2584-2591.
- Fornes, T.D., Yoon, P.J. and Paul, D.R. (2003). Polymer matrix degradation and color formation in melt processed nylon6/clay nanocomposites. *Polym.*; **44**: 7545-7556.

- Franco, L., Navarro, E., Subirana, J.A. and Puiggali, J. (1998a). Incorporation of glycine in even-even nylons disrupts their characteristics all-trans conformation. *Polym.*; **39**: 5553-5560.
- Franco, L., Subirana, J.A. and Puiggali, J. (1998b). Incorporation of glycine in even-even nylons 6,10 and 12,10. *Polym.*; **40**: 2429-2438.
- Frohoff-Hulsmann, M.A., Lippold, B.C. and McGinity, J.W. (1999). Aqueous ethyl cellulose dispersion containing plasticizers of different water solubility and hydroxypropyl methyl-cellulose as coating material for diffusion pellets II: properties of sprayed films. *Eur. J. Pharm. Biom.*; **48**: 67-75.
- García-Encina, G., Torres, D., Seijo, B. and Vila Jato, J.L. (1993). In vivo evaluation of nylon-coated diclofenac-resin complexes. *J. Control Release.*; **23**: 201-207.
- Gaymans, R.J. and Sikkema, D.J. (1999). In "Aliphatic polyamides", ed. Allen, S.G., Bevington, C.B. and Price, C. in *Comprehensive Polymer Science*. Vol. **5**, pgs 357-373.
- Ghosh, S. (2004). Recent research and development in synthetic polymer-based drug delivery systems. *J. Chem. Res.*; **4**: 241-246.
- Gohel, M.C. and Amin, A.F. (1999). Formulation, design and optimization of modified release microspheres of diclofenac sodium. *Drug Dev. Ind. Pharm.*; **25**: 247-251.
- Goupy, J. (2005). What kind of experimental design for finding and checking the robustness of analytical methods. *Analyt. Chim. Acta.*; **544**: 184-190.
- Govender, S., Pillay, V., Chetty, D.J., Essack, S.Y., Dangor, C.M. and Govender, T. (2005). Optimization and characterization of bioadhesive controlled release tetracycline microspheres. *Int. J. Pharm.*; **306**: 24-40.
- Hite, M., Federici, S., Turner, S. and Fassihi, R. (2005). Novel design of a self-correcting monolithic controlled release system for tramadol. <MMS\Drug%20Delivery%

- 20Technology%20-%20Article%20 Index.htm>. *Drug Deliv. Ttech.*; Article Index, pgs 1-13 [Accessed: 20.03.2006].
- Hopf, G. (2001). Nylon 12- Huels in comparison to other nylons. *Chemische Werke Huels AG*, West Germany, pgs 1-22.
- Huang, S.J. (1999). In "Biodegradation", ed. Allen, S.G., Bevington, C.B. and Price, C. in *Comprehensive Polymer Science*. Vol. **6**, pgs 597-606.
- Huang, X. and Brazel, C.S. (2001). On the importance and mechanisms of burst release in matrix-controlled drug delivery systems. *J. Control Release.*; **73**: 121-136.
- Huang, Y., Li, W. and Yan, D. (2003). Crystalline transition of nylons 12,20 and 10,20. *Eur. Polym. J.*; **39**: 1113-1140.
- Ives, H.E. (2000). In "Diuretic Agents", ed. Katzung, B.G. in *Basic and Clinical Pharmacology*. 8th edition, pgs 245-264.
- Iza, M., Stoianovia, G., Viora, L., Grossiord, L. and Courraze, G. (1998). Hydrogels of poly (ethylene glycol): mechanical characterization and release of a model drug. *J. Control Release.*; **52**: 41-51.
- Jamzad, S., Tutunji, L. and Fassihi, R. (2005). Analysis of macromolecular changes and drug release from hydrophilic matrix systems. *Int. J. Pharm.*; **292**: 75-85.
- Jones, D.S., Woolfson, D.A., Djokic, J. and Coulter, W.A. (1996). Development and mechanical characterization of bioadhesive semi-solid, polymeric systems containing tetracycline for the treatment of periodontal diseases. *Pharm. Res.*; **13**: 1734-1738.
- Jones, N.A., Atkins, E.D.T., Hill, M.J., Cooper, S.J. and Franco, L. (1997). Chain-folded lamellar crystals of aliphatic polyamides. Investigation of nylons 4 8, 4 10, 4 12, 6 10, 6 12, 6 18, and 8 12. *Polym.*; **38**: 2689-2699.

- Kang, M., Myung, S. and Jin, H. (2006). Nylon 6,10 and carbon nanotube composite by in-situ interfacial polymerization. *Polym.*; **47**: 3961-3966.
- Karanchi, A.A. and Khan, M.A. (1996). Box-Behnken design for the optimization of formulation variables of indomethacin co-precipitates with polymer mixtures. *Int. J. Pharm.*; **131**: 9-17.
- Karatas, A. and Baykara, T. (2001). Studies on indomethacin inserts prepared by water-soluble polymers- II. The relation between dissolution rate and swelling behaviour. *Il. Farmaco.*; **56**: 97-202.
- Katz, J. (2001). Developments in medical polymers for biomaterial applications, MDDI Article Index, pgs 1-2.
- Kiely, D.E., Chen, L. and Lin, T.H. (1994). Hydroxylated nylons based on unprotected esterified D-glucaric acid by simple condensation reactions. *J. Amer. Chem. Soc.*; **116**: 571-578.
- Kim, H., Knowlesand, J.C., Kim, H. (2004). Hydroxyapatite/poly (ϵ -caprolactone) composite coatings on hydroxyapatite porous bone scaffold for drug delivery. *Biomat.*; **25**: 1279-1287.
- Kincl, M., Turk, S. and Vrečer, T.F. (2005). Application of experimental design methodology in development and optimization of drug release method. *Int. J. Pharm.*; **291**: 39-49.
- Konishi, R. and Ito, M. (2004). Relation between ductility and segmental mobility on nylon 6. *Polym.*; **45**: 5191-5198.
- Koulouri, E.G., Georgaki, A.X. and Kallitsis, J.K. (1997). Reactive compatibilization of aliphatic polyamides with functionalized polyethylenes. *Polym.*; **38**: 4185-4192.
- Kumar, R. and Münstedt, H. (2004). Silver ion release from antimicrobial polyamide/silver composites. *Biomat.*; **26**: 2081-2088.

- Lai, S.Y. and Becker, D.G. (2004). Sutures and Needles. <<http://www.emedicine.com/ent/topic38.htm>>. WebMD, pgs 1-13 [Accessed: 26.06.2006].
- LaVan, D.A., McGuire, T. and Langer, R. (2003). Review on small scale systems for *in vivo* drug delivery. *Nat. Biotech.*; **21**:1184-1191.
- Lee, M.J., Patricia, A.D. and Laurie, A.C. (2003). Aesthetic and predictable correction of the inverted nipple. *Aesthet. Surg. J.*; **23**: 353-356.
- Li, W., Nadig, D., Rasmussen, H.T., Patel, K. and Shah, T. (2005). Sample preparation optimization for assay of active pharmaceutical ingredients in a transdermal drug delivery system using experimental designs. *J. Pharm. Biomed. Anal.*; **37**: 493-498.
- Li, Y., Yan, D. and Zhu, X. (2001). Crystal forms of nylon 10,12 crystallized from melt and after solution casting. *Eur. Polym. J.*; **37**: 1849-1853.
- Liu, L., Khang, G., Rhee, J.M. and Lee, H.B. (2000). Monolithic osmotic tablet system for nifedipine delivery. *J. Control Release.*; **67**: 309-322.
- Lu, E., Jiang, Z., Zhang, Q. and Jiang, X. (2003). A water-insoluble drug monolithic osmotic tablet system utilizing gum Arabic as an osmotic, suspending and expanding agent. *J. Control Release.*; **92**: 375-382.
- Lundborg, G., Dahlin, L., Dohi, D., Kanje, M. and Terada, N. (1997). A new type of "bioartificial" nerve graft for bridging extended defects in nerves. *J. Han. Surg. (British and European Volume)* **22B**: 299-303.
- Madan, P.L. and Chareonboonsit, P. (1989). Nylon microcapsules. II. Effects of selected variables on theophylline release. *Pharm. Res.*; **6**: 714-718.
- Makino, K., Mack, E.J., Okano, T. and Wan Kim, S. (1990). A microcapsule self-regulating delivery system for insulin. *J. Control Release.*; **12**: 235-239.
- Martin, L., Wilson, C.G., Koosha, F., Tetteiy, L., Gray, A.J., Senel, S. and Uchegbu, J.F. (2002). The release of model macromolecules may be controlled by the

- hydrophobicity of palmitoyl glycol chitosan hydrogels. *J. Control Release.*; **80**: 87-100.
- McGinity, J.W. and Co, G.W. (1985). Nylon encapsulated pharmaceuticals. *Meth. Enzymol.*; **112**: 87-101.
- Mehuys, E., Vervaet, C. and Remon, J.P. (2004). Hot-melt extruded ethylcellulose cylinders containing a HPMC-Gelucire® core for sustained drug delivery. *J. Control Release.*; **94**: 273-280.
- Middleton, J.C. and Tipton, A.J. (1998). Synthetic biodegradable polymers as medical devices. *Medical Plastics and Biomaterials Magazine*. MPB Article Index, pgs 1-10.
- Miyajima, M., Koshika, A., Okada, J., Kusia, A. and Ikeda, M. (1998). Factors influencing the diffusion-controlled release of papaverine from poly (L-lactic acid) matrix. *J. Control Release.*; **56**: 85-94.
- Mona, M. (2000). Multiple modes of drug delivery: Technologies such as microchips and microspheres are enabling the therapeutic use of proteins. *Mod. Dru. Discov*; **3**: 30-34.
- Moore, J.W. and Flanner, H.H. (1996). Mathematical comparison of dissolution profiles. *Pharm. Tech.*; **20**: 64-74.
- Muhammad, N.A., Biosvert, W., Harns, M.R. and Weiss, J. (1991). Modifying the release properties of eudragit L30D. *Drug Dev. Ind. Pharm.*; **17**: 2497-2509.
- Murdan, S. (2003). Electro-responsive drug delivery from hydrogels. *J. Control Release.*; **92**: 1-17.
- Murthy, N.S. (1999). Interactions between crystalline and amorphous domains in semi-crystalline polymers: small-angle X-ray scattering studies of the Brill transition in nylon 6,6. *Macromol.*; **32**: 5594.

- Nadig, D.E. (2002). In "Preparation of drug samples for analysis", in *Hand book of Pharmaceutical Analysis*. Mercel Decker.
- Narasimhan, B. (2001). Mathematical models describing polymer dissolution: consequences for drug delivery. *Adv. Drug Deliver. Rev.*; **48**:195-210.
- Nazzal, S., Nutan, M., Palamakula, A., Shah, R., Zaghloul, A.A. and Khan, M.A. (2002). Optimization of a self-nanemulsified tablet dosage form of ubiquinone using response surface methodology: effect of formulation ingredients. *Int. J. Pharm.*; **240**: 103-114.
- Nutan, M.T.H., Soliman, M.S., Taha, E.I. and Khan, M.A. (2005). Optimization and characterization of controlled release multi-particulate beads with starch acetate. *Int. J. Pharm.*; **294**: 89-101.
- Ostad, S.N. and Gard, P.R. (2000). Cytotoxicity and teratogenicity of chlorhexidine diacetate released from hollow nylon fibres. *J. Pharm. Pharmacol.*; **52**: 779-784.
- Ostad, S.N., Farhadkhani, M., Minaiee, B., Malihi, G. and Abdollahi, M. (2002). The blood pressure and dermal sensitivity effects of nylon hollow fiber releasing glycerin trinitrate *in vivo*. *Daru.*; **10**: 125-129.
- Ostad, S.N., Malhi, J.S. and Gard, P.R. (1998). *In vitro* cytotoxicity and teratogenicity of norethisterone and levonorgestrel released from hollow nylon monofilaments. *J. Control Release.*; **50**: 179-186.
- Pedley, D.G., Skelly, P.J. and Tighe, B.J. (1980). Hydrogels in biomedical applications. *Br. Polym. J.*; **12**: 99-110.
- Peeters, R. and Kinget, R. (1993). Film-forming polymers for colonic drug delivery, I. Synthesis, physical and chemical properties of methyl derivatives of eudragit S. *Int. J. Pharm.*; **94**: 125-134.

- Peppas, N.A. and Langer, R. (1994). New challenges in biomaterials. *Sci.*; **273**: 1715-1720.
- Peschka, R., Dennehy, C. and Szoka, F.C. (1998). A simple *in vitro* model to study the release kinetics of liposome encapsulated material. *J. Control Release.*; **56**: 41-51.
- Phares, K., Cho, M., Johnson, K. and Swarbrick, J. (1995). Drug transport across nylon 6,10 films: Influence of synthesis variables. *Pharm. Res.*; **12**: 248-256.
- Pharmaceutical Codex. (1994). Principles and Practice of Pharmaceutics. Ed. Walter Lund, the Pharmaceutical press, London.
- Pillay, V. and Danckwerts, M.P. (2002). Textural profiling and statistical optimization of crosslinked calcium-alginate-pectinate-cellulose acetophthalate gelisphere matrices. *J. Pharm. Sci.*; **91**: 2559-2570.
- Pillay, V. and Fassihi, R. (1998). Evaluation and comparison of dissolution data derived from different modified release dosage forms: an alternative method. *J. Control Release.*; **55**: 45-55.
- Pillay, V. and Fassihi, R. (1999a). *In vitro* release modulation of cross-linked pellets for site-specific drug delivery to the gastrointestinal tract, I. Comparison of pH-responsive drug release and associated kinetics. *J. Control Release.*; **59**: 229-242.
- Pillay, V. and Fassihi, R. (1999b). *In vitro* release modulation of cross-linked pellets for site-specific drug delivery to the gastrointestinal tract, II. Physicochemical characterization of calcium-alginate, calcium-pectinate and calcium-alginate-pectinate pellets. *J. Control Release.*; **59**: 243-256.
- Pillay, V. and Fassihi, R. (1999c). Electrolyte-induced compositional heterogeneity: a novel approach to rate-controlled oral drug delivery. *J. Pharm. Sci.*; **88**: 1140-1148.
- Pillay, V. and Fassihi, R. (2000a). In situ electrolyte interactions in a disc-compressed system for up-curving and constant drug delivery. *J. Control Release.*; **67**: 55-65.

- Pillay, V. and Fassihi, R. (2000b). A novel approach for constant rate delivery of highly soluble bioactives from a simple monolithic system. *J. Control Release.*; **67**: 67-78.
- Pillay, V., Danckwerts, M.P., Muhidinov, Z. and Fassihi, R. (2005a). Novel modulation of drug delivery using binary zinc-alginate-pectinate polyspheres for zero-order kinetics over several days: Experimental design strategy to elucidate the crosslinking mechanism. *Drug Dev. Ind. Pharm.*; **31**: 191-207.
- Pillay, V., Sibanda, W. and Danckwerts, M.P. (2005b). Sequential design of a novel PVA based crosslinked ethylenic homopolymer for extended drug delivery. *Int. J. Pharm.*; **301**: 89-101.
- Pouton, C.W. (2001). Polymeric materials for advanced drug delivery. *Adv. Drug Deliver. Rev.*; **53**: 1-3.
- Ramchandani, M. and Robinson, D. (1998). *In vivo* and *in vitro* release of ciprofloxacin from PLGA 50:50 implants. *J. Control Release.*; **54**: 167-175.
- Ramraj, R., Farrell, S. and Loney, N.W. (1999). Mathematical modeling of controlled release from a hollow fiber. *J. Memb. Sci.*; **162**: 73-81.
- Sánchez-Lafuente, C., Rabasco, A.M., Álvarez-Fuentes, J. and Fernández-Arévalo, M. (2002). Eudragit® RS-PM and Ethocel® 100 Premium: influence over the behaviour of didanosine inert matrix system. *II Farmaco.*; **57**: 649-656.
- Sastry, S.V. and Khan, M.A. (1998). Aqueous based polymeric dispersion: Plackett-Burman design for screening of formulation variables of atenolol gastrointestinal therapeutic system. *Pharm. Acta Helv.*; **73**: 105-112.
- Sastry, S.V., Reddy, I.K. and Khan, M.A. (1996). Optimization and characterization of controlled release pellets coated with an experimental latex: II. Cationic drug. *Int. J. Pharm.*; **196**: 179-195.

- Schliecker, G., Schmidt, C., Fuchs, S., Ehinger, A., Sandow, J. and Kissel, T. (2004). *In vitro* and *in vivo* correlation of buserelin release from biodegradable implants using statistical moment analysis. *J. Control Release.*; **94**: 25-37.
- Scholsky, K.M. and Fitch, R.M. (1986). Controlled release of pendant bioactive materials from acrylic polymer colloids. *J. Control Release.*; **3**: 87-108.
- Seitz, B., Langenbucher, A., Kühle, M. and Naumann, G.O.H. (2003). Impact of graft diameter on corneal power and the regularity of postkeratoplasty astigmatism before and after suture removal. *Ophthalmol.*; **110**: 2162-2167.
- Shah, N., Zhang, G., Apelian, V., Zeng, F., Infeld, M.H. and Malick, A.W. (1993). Prediction of drug release from hydroxypropylmethylcellulose (HPMC) matrices: effect of polymer concentration. *Pharm. Res.*; **10**: 1693-1695.
- Shenoy, D.B. and Amiji, M.M. (2005). Poly (ethyleneoxide)-modified poly (ϵ -caprolactone) nanoparticles for targeted delivery of tamoxifen in breast cancer. *Int. J.Pharm.*; **293**: 261-270.
- Sibanda, W., Pillay, S., Danckwerts, M.P., Viljoen, A.M., Vuuren, S. and Khan, R.A. (2004). Experimental design for the formulation and optimization of novel cross-linked oilispheres developed for *in vitro* site-specific release of *Mentha piperita* oil. *AAPS Pharm. Sci. Tech.*; **5**: 1-14.
- Siepmann, J. and Gopferich, A. (2001). Mathematical modeling of bioerodible, polymeric drug delivery systems. *Adv. Drug Deliver. Rev.*; **48**: 229-247.
- Siepmann, J. and Peppas, N.A. (2001a). Mathematical modeling of controlled drug delivery. *Adv. Drug Deliver. Rev.*; **48**: 137-138.
- Siepmann, J. and Peppas, N.A. (2001b). Modeling of drug release from delivery systems based on hydroxypropyl methylcellulose (HPMC). *Adv. Drug Deliver. Rev.*; **48**: 139-157.

- Sikorski, P., Jones, N.A., Atkins, E.D.T. and Hills, M.J. (2001). Measurement of the intersheet shear along the chain axis in nylon 6. *Macromol.*; **34**: 1673.
- Singh, S.K., Dodge, J., Durrani, M.J. and Khan, M.A. (1995). Optimization and characterization of controlled release pellets coated with an experimental latex: I. Anionic drug. *Int. J. Pharm.*; **125**:243-255.
- Sinha, V.R. and Khosla, L. (1998). Bioabsorbable polymers for implantable therapeutic systems. *Drug Dev. Ind. Pharm.*; **24**: 38.
- Sohier, J., Vlugt, T.J.H., Cabrol, N., Van Blitterswijk, C., de Groot, K. and Bezemer, J.M. (2006). Dual release of proteins from porous polymeric scaffolds. . *J. Control Release.*; **111**: 95-106.
- Song, Y., Sun, P., Henry, L.L. and Sun, B. (2005). Mechanisms of structure and performance controlled thin film composite membrane formation via interfacial polymerization process. *J. Memb. Sci.*; **251**: 67-79.
- Steward, P.A. (1995). A literature review of pharmaceutical controlled release methods and devices, <http://www.initium.demon.co.uk/rel_nf.htm>. [Accessed: 14.05.2005].
- Symn, C., Kala, W., Van den Mooter, G. and Kinget, R. (1995). Synthesis and *in vitro* biodegradation of poly (ether-ester) azo polymers designed for colon targeting. *Int. J. Pharm.*; **121**: 211-216.
- Torres, D., Seijo, B., García-Encina, G., Alonso, M. and Vilajato, J.L. (1990). Microencapsulation of ion-exchange resins by interfacial nylon polymerization. *Int. J. Pharm.*; **59**: 9-17.
- Van M.W. (1996). The efficacy of a single continuous nylon suture for control of post keratoplasty astigmatism. *Trans. Am. Ophthalmol. Soc.*; **94**: 1157–1180.
- Velasco, M.V., Ford, J.L., Rowe, P., Rajabi-Siahboomi, A.R. (1999). Influence of drug:hydroxypropylmethylcellulose ratio, drug and polymer particle size and

- compression force on the release of diclofenac sodium from HPMC tablets. *J. Control Release.*; **57**: 75-85.
- Vogelson, C.T. (2001). Advances in drug delivery systems. *Nanotech.*; **4**: 49-52.
- Vyas, S.P., Sood, A., Venugopalan, P. and Venkatesan, N. (2000). Preparation and characterization of microencapsulated gelspheres for controlled oral theophylline delivery. *J. Microencap.*; **17**:767-775.
- Wehrle, P., Karner, D. and Benita, S. (1996). Sequential statistical optimization of a positively charged submicron emulsion of miconazole. *Pharm. Dev. Technol.*; **1**: 97-111.
- Williams III., R.O., Reynolds, T.D., Cabelka, T.D., Sykora, M.A. and Mahaguna, V. (2002). Investigation of excipients type and level on drug release from controlled release tablets containing HPMC. *Pharm. Dev. Technol.*; **7**: 181-193.
- Yang, L. and Fassihi, R. (1997a). Examination of drug solubility, polymer types, hydrodynamics and loading dose on drug release behaviour from a triple-layer asymmetric configuration delivery system. *Int. J. Pharm.*; **155**: 219-229.
- Yang, L. and Fassihi, R. (1997b). Modulation of diclofenac release from a totally soluble controlled release drug delivery system. *J. Control Release.*; **44**: 135-140.
- Yuk, S.H., Cho, S.H., Lee, H.B. (1992). Electric current-sensitive drug delivery systems using sodium alginate/poly-acrylic acid composites. *Pharm. Res.*; **9**: 955-957.
- Zajc, N., Obreza, A., Bele, M. and Srcic, S. (2005). Physical properties and dissolution behaviour of nifedipine/mannitol solid dispersions prepared by hot melt method. *Int. J. Pharm.*; **291**: 51-58.