

The present study also addressed the American Heart Association's call to overcome the paucity of research concerning the effects of exercise on CHD patients' psychological health (Fletcher et al., 1992). An attempt was made to determine whether the frequency and duration of CHD patients' exercise would effect their anxiety and depression. The present sample did experience a reduction in anxiety and depression after six months of exercise. However, contrary to expectations, the frequency of patients' exercise was not associated with this reduction in anxiety and depression. While the duration of the present sample's exercise was not associated with their reduction in depression, it limited their reduction in anxiety. Though surprising, this result is consistent with research that has revealed an increase in anxiety following participation in an exercise programme (e.g., De Geus et al., 1993; Stern & Cleary, 1981; Van Dixhoorn et al., 1983). The physical symptoms of increased duration of exercise are similar to those of cardiovascular distress (Ledwidge, 1980; Weg, 1985). It is possible that patients associate the symptoms induced by increased duration with the symptoms of cardiovascular distress and, consequently, experience anxiety when exercising. In complying with the duration component of their exercise prescriptions, patients may also believe that they are being stretched beyond their physical abilities and are placing their lives at risk. The risk patients associate with the increased duration exercise is likely to cause anxiety. Given the present finding that exercise duration acts as an impediment to psychological recovery, the relationship between exercise and CHD patients' psychological health will require close scrutiny in future research.

Practical Implications of the Present Study

Myocardial infarction survivors constitute the largest group of people who are at risk of suffering a fatal cardiac event (Kamarck & Jennings, 1991). Indeed, some studies have shown that over 70% of sudden cardiac death victims had a prior history of myocardial infarctions (Kamarck & Jennings, 1991). It is thus imperative that every

effort is made to enhance the effectiveness of rehabilitation. A primary goal of rehabilitation is to help CHD patients regain their ability to pursue a full and productive life and reduce their risk of reinfarction. This is achieved by increasing CHD patients' cardiovascular fitness through exercise training. The present study demonstrates that the duration of exercise needs to be emphasised in exercise based rehabilitation programmes. Supervised exercise based cardiac rehabilitation programmes remain the most effective means of reducing the risk of fatal and non-fatal cardiac events and also, of reducing the behaviours that initiate these (Kamarck & Jennings, 1991). While many of the benefits of cardiac rehabilitation are intangible (Froelicher et al., 1993), the present study shows that the duration of exercise provides a primary line of defence against reinfarction (Curfman, 1993).

An additional goal of existing rehabilitation programmes is to inform patients of the relationship between risk factors (e.g., smoking, diet, sedentary lifestyle) and CHD and encourage them to change their lifestyles to eradicate these risk factors (Wright, Carbonari, & Voyles, 1992). The results of the present study suggest that this should be extended to the provision of interventions designed to modify the toxic components of Type A behaviour. While Type A components were unassociated with physiological strain in the present study, the anger-in and anger-out components predicted reduced recovery from anxiety and depression. The psychophysiological effects of anxiety and depression pose a substantial threat to an already compromised cardiovascular system (Brown & Munford, 1984; Garrity & Klein, 1975; Rejeski et al., 1985). The progression of CHD is slow with the result that the magnitude of this threat may not be apparent in the short term (Orha et al., 1985). As the focus of the present study was on the first six months of recovery, it was not possible to determine whether the negative effect of anger on anxiety and depression would have resulted in an increased risk of reinfarction in the long term. However, given the relationship between anxiety, depression and increased risk of reinfarction, future rehabilitation programmes may be enhanced by designing

interventions which alter the toxic components of Type A behaviour which impede recovery from anxiety and depression.

In identifying which of the Type A behaviours are benign, and which are toxic, the present study makes it possible to identify sub-groups within the Type A population who are at greater risk of suffering subsequent psychological strain. The present study also provides future research with a valid and reliable questionnaire that can be used to identify the individual components of Type A behaviour which are manifested by CHD patients. With this knowledge it will be possible to design intervention strategies which reduce those components of Type A behaviour which exert a deleterious effect and cultivate those which exert a positive effect (Bluen et al., 1990). Differentiation of the toxic and benign effects of the various Type A components, and the treatment thereof, has important implications for the design of rehabilitation programmes. Active, vigorous individuals are often labelled Type A's despite the fact that they do not exhibit the potentially dangerous hostility, impatience irritability and anger, characteristic of Type A behaviour (Friedman & Booth-Kewley, 1987b). It would be unfortunate if these individuals were made to alter their seemingly healthy behaviour because of the incorrect definition and assessment of Type A behaviour. Further, as achievement striving behaviour enhances self-esteem and social, economical and personal stature, modification of this aspect of Type A behaviour may have negative social and personal consequences for the individual (Cohen, 1980). Thus, from both the literature and the present findings it is evident that intervention strategies should focus on the modification of CHD patients' anger and the cultivation of their achievement striving behaviour.

Limitations of the Present Study

The results of the present study support the notion that Type A behaviour is a

multidimensional construct wherein the different dimensions exert differential effects. Also supported, is the notion that it is insufficient to view exercise compliance exclusively in terms of attendance and that it is in fact, the duration of exercise which predicts changes in cardiorespiratory condition. It is possible, however, that the value of this research is limited by several flaws in the conceptualisation of the model, the composition of the sample, the study design and the methodology employed. The various limitations, together with suggestions for future research, are outlined below.

The present research addresses a fundamental weakness of past Type A research by examining the multidimensional, and not the global, nature of the construct. Hostility has also been erroneously conceptualised and treated as a global construct (Thoreson & Powell, 1992) where, in fact, it is multidimensional (Suarez & Williams, 1990). While the present research does not make the mistake of examining hostility globally, it includes only one dimension of the construct, namely, acting-out hostility. The measure of acting-out hostility is consistent with the original definition of Type A behaviour (Friedman & Rosenman, 1974) and the type of hostility associated with CHD (Carmody et al., 1989; Dembroski & Costa, 1987). However, it is possible that other dimensions of hostility (e.g., potential for hostility, cynical hostility; neurotic hostility) exert differential effects on other, equally salient outcomes of CHD diagnosis (Suarez & Williams, 1990). For example, past research has demonstrated that neurotic hostility predicts CHD patients' anxiety (e.g., Siegman et al., 1987). Given the likelihood of differential effects, future research should both specify and consider the predictive role of all dimensions of hostility on the various different psychological and physiological outcomes seen in samples of CHD patients.

A further limitation concerns the possible incorrect composition of the sample. For example, Carney et al. (1987) have found that depression is related strongly to CHD.

Other studies (e.g., Carney, Rich & Freeland, 1988) have shown that depression is an independent predictor of secondary cardiac events, while the results of Powell et al.'s (1993) study suggest that depression is associated with CHD mortality. Research also suggests that anxiety is a precursor of CHD (Matthews, 1982; Nielson & Dobson, 1980). Friedman and Booth-Kewley's (1988) meta-analysis shows that both anxiety and depression are reliably associated with CHD. Thus, future models should also consider anxiety and depression as potential predictors of CHD.

It has been argued in the present research (see Chapter 4) and elsewhere (e.g., Ganster et al., 1991), that it is the heightened cardiovascular reactivity associated with the toxic components of Type A behaviour that predisposes individuals to CHD risk. It has also been argued that these behavioural components occur in response to excessively challenging environmental situations (Price, 1982). Should this be the case, then any attempt to determine a conclusive relationship between Type A components and CHD risk, or Type A's response thereto, should include an assessment of the quality, frequency and intensity of excessively challenging environmental stimuli experienced by CHD patients (Suarez & Williams, 1990). While such measurement was beyond the scope of the present study, future research should identify and quantify challenging environmental stimuli and assess these in relation to CHD. Such research might provide evidence for the belief that acute stressors precipitate sudden cardiac death in humans in the same way as they have been found to do in animals (Kamarck & Jennings, 1991).

Several limitations of the present study are imposed by the demographic composition of the sample. By only including a small number of women, the present study can be criticised for contributing to the paucity of information concerning all aspects of cardiovascular disease and rehabilitation in women (Chesney, 1993; Wenger et al., 1993). Women of childbearing age have been excluded from past research as have elderly women who frequently have coexisting diseases (Wenger,

1992). Further, the less frequent use of intensive or invasive diagnostic and corrective treatments for CHD in women than men (despite an equal severity of CHD) (Ayanian & Epstein, 1991) has contributed to the fact that fewer women find themselves included in studies of secondary prevention. Similarly, less women than men are referred to exercise based rehabilitation programmes by medical practitioners despite the fact that the therapeutic benefit is equal for both sexes (Oldridge, LaSalle, & Jones, 1980). Those females who are referred for rehabilitation, tend to comply with the programme to a lesser extent than their male counterparts because of a higher frequency of coexisting illnesses, familial responsibilities and various other psychosocial factors (Powell et al., 1993; Wenger et al., 1993).

The rate of CHD morbidity and mortality among women is substantial and increasing (Johnson & Morse, 1990; Wenger, Speroff, & Packard, 1993). The prognosis following either medical and surgical therapy for CHD is poorer for women than for men (Wenger et al., 1993). Further, the incidence of early death following myocardial infarction (Greenland, Reicher-Reiss, Goldbourt, & Behar, 1991; Lerner & Kannel, 1986) and the rate of in-hospital deaths after cardiovascular surgery is higher among women than men (Loop et al., 1983; Weintraub, Wenger, Jones, Craver, & Guyton, 1992). The magnitude of the association between Type A behaviour and CHD is also as strong for women as it is for men (Miller et al., 1991). However, recent research (e.g., Norris, deGuzman, Sobel, Brooks, & Haywood, 1993; Powell et al., 1993) suggests that the psychosocial predictors of secondary CHD mortality among women differ considerably from the standard predictors of mortality among men. In the present study, t tests did not yield any significant differences between men and women on the Type A and exercise predictor variables. Further attempts to reveal any differences between male and female CHD patients' changes in anxiety, depression and maximal oxygen uptake by means of t tests, also failed to yield significant results. From the results of the t tests conducted in the present study, it is evident that there are no differences in the male and female

psychological and behavioural predictor variables and in their psychological and physiological outcome variables. However, given that past research has revealed different relationships between psychosocial predictors and secondary CHD mortality among men and women, it is advisable that the relationship between personality, behaviour and CHD as it applies to women be examined in greater detail (Wenger et al., 1993). What is needed is sex specific research concerning the psychosocial factors associated with the cause, prevention, treatment and prognosis of CHD in women (Chesney, 1993; Wenger et al., 1993).

A second limitation imposed by the sample concerns the under-representation of black CHD patients. Cross-sectional research shows that the distribution of Type A behaviour among white samples is equal to that found in black samples and that the association between behaviour, CHD risk factors and CHD outcomes is the same for both races (Anderson et al., 1986; Thoreson & Powell, 1992). Research conducted in the United States (Becker et al., 1993) has shown that the incidence of CHD is significantly greater among blacks than among whites and that blacks have a significantly lower rate of subsequent survival. In comparison to whites, blacks undergo less invasive cardiac procedures such as coronary artery bypass grafts, percutaneous transluminal angioplasty and cardiac catheterisation (Whittle et al., 1993). Research into the incidence of risk factors and mortality among, black, white and Latina women has established a differential distribution of clinical outcomes according to race (Norris et al., 1993). There is thus a racial difference in the contraction and survival of CHD wherein blacks receive less of the treatment necessary to relieve symptoms and enhance prognosis (Ayanian, 1993). The lower survival and treatment rates is paralleled by the under representation of blacks in research (Becker et al., 1993; Keil et al., 1993). Indeed, there is a relative paucity of information regarding racial differences in the risk of contracting CHD and in subsequent survival rates (Becker et al., 1993). In view of this deficiency, t tests were conducted in the present study to compare blacks and whites on the predictor

and outcome variables. No significant differences were found between blacks and whites on the psychological and behavioural predictor variables. The t tests also failed to reveal any significant differences between the two racial groups' changes in anxiety, depression and maximal oxygen uptake. However, given the potentially higher risk indicated in past research (e.g., Becker et al., 1993), it is imperative that future research identify those factors which render different race groups particularly prone to CHD.

The medical heterogeneity of the sample constitutes both a strength and a weakness of the present study. The present sample is characterised by a wide variety of types and combinations of cardiac events and surgical procedures. While this increases the generalisability of the study, it decreases the specificity of the results. There is evidence to suggest that the various cardiac events and interventions are associated with different risk factors and outcomes (Wright, 1992). For example, CHD patients who are subjected to surgery (e.g., coronary artery bypass graft and percutaneous transluminal angioplasty) have a better psychological and physiological prognosis than patients whose treatment is limited to medication (Soloff, 1978a). The psychological and risk factor profile, and the consequent outcome, may then vary according to the underlying type and treatment of disease. In view of this, the medical heterogeneity of the present sample may have confounded the results. To address this issue, two subgroups of patients were created and compared on the outcome variables by means of t tests. The first group comprised patients who had only experienced a myocardial infarction while the second comprised those patients who had undergone a coronary artery bypass graft, percutaneous transluminal angioplasty or coronary artery angiography but had not experienced a myocardial infarction. Results of the t tests showed no significant difference between the two groups' changes in anxiety, depression and maximal oxygen uptake. This suggests that the results of the present study were not influenced by the type and treatment of CHD. However, in view of the potential difference between groups (Soloff, 1978a;

Wright, 1992), future research should examine the relationship between Type A and exercise components and psychological and physiological outcomes within specific sub-groups of CHD patients and compare these to other groups (e.g., myocardial infarction vs. coronary artery bypass graft patients). In taking this step, future research would be able to separate the effects of the various Type A and exercise components on the different disease endpoints.

A related weakness is the examination of the Type A construct within the exclusive context of CHD. While the association between Type A behaviour, the components thereof, and CHD is well documented, it is possible that the construct is equally predictive of other diseases (Wright, 1992). Future research should compare the relationships between Type A components and psychological and physiological morbidity found in CHD samples with samples comprising both healthy and other disease groups. Examples of diseases, which might be associated with Type A behaviour and therefore, should be considered in future research, include asthma (Everaerd, Vroman, & van der Elst, 1990), hypertension (Godaert (1990), peptic ulcers, thyroid disorders and rheumatoid arthritis (Rimé et al., 1989). Only by comparing the relationships between Type A behaviour and various different diseases will it be possible to establish whether the Type A components are specifically associated with CHD, or are, in fact, risk factors for disease in general.

The design of the present study constitutes a further limitation. Given that replication remains the most robust form of validation (Grossarth-Maticek & Eysenck, 1991), use of a control group would have enhanced the scientific validity of the present study. It has been argued in the present research and elsewhere (e.g., Evans, 1990; Ragland & Brand, 1988b) that the Type A behaviour demonstrated by CHD patients is inherently different to that which is displayed by healthy individuals. Further, the association between Type A behaviour and CHD is far greater among healthy samples than it is among CHD samples (Miller et al., 1991).

Demographic factors such as sex, age or culture may alter the strength of the relationship between Type A behaviour and CHD, and consequently, bias the results according to the demographic characteristics of the sample (Miller et al., 1991). A matched controlled study would reveal whether there are differences in the behaviour demonstrated by healthy individuals and CHD patients and whether this has a differential effect on health. The incorporation of a control group would also help determine the efficacy of the rehabilitation programme (Oldridge, 1988). While ideal, the inclusion of a matched sample of healthy subjects in the present study was beyond the scope of the present research.

A further number of limitations are associated with the present study's reliance on self-report measures of Type A components. Same-source effects may have inflated the correlations between psychological variables (Terry, 1992). The use of self-reports may have generated further sources of bias. For example, in seeking both an explanation and repentance for their condition, CHD patients are suspected of exaggerating their risk related behaviours such as anger and impatience irritability (Evans, 1990; Wright, 1992). This syndrome may result in an inaccurate inflation of scores derived from self-report measures of Type A components. In turn, this could result in positive and significant correlations between the Type A components and disease which are a product of patients' exaggerated perceptions and not their behaviour *per se* (Wright, 1992).

A second source of bias concerns the negative connotations associated with Type A behaviour which may result in socially desirable responses (Tett et al., 1992). As a means of maintaining high self-esteem, Type A's typically describe themselves in a self-appreciatory manner (Furnham, 1990b). Type A's also tend to recall more positive than negative traits about themselves (Furnham, Borovoy & Henley, 1986). As a consequence, desirable traits such as ambition and achievement striving are over-reported (Emmons & McAdams, 1991) and negative traits such as time-urgency

and aggressive-hostility are under-reported (Furnham, 1990b). Thus, there is a strong possibility that Type A's responses to self-report Type A measures are tainted by their need to maintain self-esteem, and this influence undermines the accuracy of measurement (Furnham, 1990b). The influence of social desirability may be one of the reasons why self-report measures of Type A behaviour are poorly correlated with CHD (Miller et al., 1991).

A related issue is that many of the studies (e.g., Anderson et al., 1986; Svebak et al., 1992) which have revealed an association between Type A components and indices of CHD have been based on the Structured Interview (Rosenman et al., 1964a, 1964b). The Structured Interview is considered to be the most accurate measure of Type A behaviour and to possess the strongest predictive validity for CHD (Ganster et al., 1991; Palmer et al., 1992). The Structured Interview is far better able to predict physiological strain than the self-report measures of Type A behaviour (Palmer et al., 1992). Given the relative power of the Structured Interview, it is possible that, had it been used in the present study, relationships between the individual Structured Interview factors, hostility, anger, and impatience and maximal oxygen uptake would have occurred. However, the Structured Interview could not be used in the present study for the following reasons. First, the Structured Interview is impractical for use in large sample studies (Ganster et al., 1991; Newton & Keenan, 1990) such as the present one. The primary reason for this is that the Structured Interview requires the use of audio-visual equipment, 15 to 20 minutes for administration and approximately the same amount of time for scoring (Abbott & Peters, 1988). Second, the Structured Interview can only be administered by specially trained, certified interviewers (Abbott & Peters, 1988). As this training is currently unavailable in South Africa it was not possible to use the Structured Interview in the present study. It is primarily for these reasons that a self-report questionnaire presented the only option. Though there are a number of limitations

associated with the use of questionnaires, they have been found to predict health outcomes in general, and CHD in particular (Powell, 1987).

The number of assumptions tests show that four of the measures of the Type A components are particularly robust. The same tests indicate that the fifth, acting-out hostility, while statistically acceptable, is less robust than the others. In particular, the data was not as close to linear as would have been ideal. Use of the acting-out subscale of the HDHQ may have limited the possibility of identifying relationships between hostility and changes in anxiety, depression and maximal oxygen uptake in the present study. The standardisation of the HDHQ for normal adults, and consequently its validity, is considered by some (e.g., Henderson, Davidson, Lewis, Clark, & Baiki, 1977) to be deficient. The psychometric weaknesses of the HDHQ highlighted by past research may have accounted for the relatively lower reliability coefficient (see Table 8.2) and coefficient of congruence and higher root mean square (see Table 8.7) of the hostility subscale in the present study. Despite the fact that the inherent psychometric weaknesses of the HDHQ may have compromised the reliability coefficient and the comparison statistics, the acting-out hostility subscale still demonstrated acceptable reliability and factor stability across samples, and consequently an absence of measurement error.

The relative inconsistency of the Type A Component Questionnaire across samples constitutes a potential limitation of the present study. In the second study of the present research, the data was subjected to confirmatory factor analysis. Five methods of factor extraction were used, namely, Kaiser's eigenvalue greater than one criterion, Kaiser's MSA criterion, the communality estimate, component saturation, and Cattell's scree test. The analysis yielded a five factor solution comprising 26 items. In the third study, the 26 items were subjected to the same five methods of factor extraction. One of these items generated a communality estimate below the .20 cut off, while a second failed to load on any of the five factors. These two items

were eliminated from the analysis conducted in study three. It can be argued that the inconsistency of the questionnaire across samples undermined its construct validity.

An aim of factor analysis is to reduce the number of variables to the extent that the information present in the original set of variables is retained (Guadagnoli & Velicer, 1988). With confirmatory factor analysis it is possible to determine whether the selected items tap the constructs defined prior to analysis (Welch et al., 1990). Therefore, confirmatory factor analyses serves to test the conceptual hypotheses concerning the proposed factor structure of a construct (Briggs & Cheek, 1986). In the present research, the nature of the five factors remained consistent over samples (see tables 7.2 and 8.6). As the items tapped the *a priori* operationally defined constructs in both samples, the conceptual hypothesis was supported. The similarity of the factors across samples further suggests that essentially all of the information of the original set of variables was preserved in the second set of variables. Thus, it can be argued that the elimination of the two items which failed to meet the extraction criteria refined the questionnaire rather than undermined its construct validity.

The six month time period used in the present study may have limited the potential for finding significant changes in psychological strain. Several studies (e.g., Palmer et al., 1992; Ragland & Brand, 1988b) have shown that both the global Type A measure and specific Structured Interview derived components such as hostility, anger and impatience do not predict recurrent CHD in the year following a coronary event. Some studies (e.g., Shephard et al., 1985) have shown that the aerobic power gains in the first year of exercise are relatively small. The results of these studies may explain why other studies (e.g., De Geus et al., 1993) have failed to reveal a relationship between exercise and enhanced psychological condition. The average maximal oxygen uptake gain of 15% seen in the present study was modest (De Geus

et al., 1993). Thus, with the focus of the present study being on the first six months of exercise compliance it is possible that the gains in aerobic power were of too low a magnitude to exert an effect on psychological health. While the American Heart Association (Erb et al., 1979) maintains that six months of exercise will result in the physical reconditioning of most CHD patients, it is possible that more time is required for the improvements in physical condition to effect psychological health.

Studies (e.g., Barefoot et al., 1983; Haynes et al., 1980) which have used a longer period of follow-up have shown that Type A components predict physiological strain. The duration of the present study may have been too short to detect any such chronic relationship between the toxic components of Type A behaviour and an increasingly compromised cardiovascular system. It is possible then, that a longer follow-up period would have revealed an association between the components of Type A behaviour and recurrent CHD. However, cognisance must be taken of the fact that the longer the time period, the greater will be the effect (Kerlinger, 1986). With longitudinal studies, time constitutes an additional variable and it is difficult to determine whether any changes that occur in the outcome variables are the result of the effect of the predictor variables or of time itself (Kerlinger, 1986). Therefore, it is suggested that future research examine the long term effects of Type A components on physiological health while simultaneously controlling for the effects of time.

The analysis component of the present research is also characterised by several limitations. A primary limitation of multiple regression, and indeed, all regression techniques, is that while it is able to establish relationships, it is unable to confirm the causal mechanisms underlying these (Montgomery & Peck, 1982). Moreover, the success of stepwise multiple regression is contingent on distributional assumptions (Montgomery & Peck, 1982). This dependency renders multiple regression particularly sensitive to sampling bias (Carmody et al., 1989). To ensure that the

independent variables examined in the present study comprise the best set of predictors of the outcome variables, and consequently, that the regression model is robust, this variable set must be applied to additional samples (Carmody et al., 1989). Application of the present model to additional samples was beyond the scope of the present study.

The present study can be criticised for using a .60 cut off point for reliability tests. While Nunnally (1967) maintains that a cut-off point of .60 is sufficient, others (e.g., James & James, 1989) state that a minimal cut-off point of .70 is required to confirm reliability. In the present study, the achievement striving, impatience irritability and acting-out hostility measures fall short of the more stringent .70 cut-off point. However, it must be remembered that, because they comprise items of different response lengths, the reliability of the measures of Type A components is assessed not by the customary Cronbach's Alpha, but by Kristof's formula. Unlike Cronbach's alpha, which reflects the most conservative estimate (Cortina, 1993), Kristof's formula provides a more exact, measure of internal reliability (Gilmer & Feldt, 1983). Given the precision with which Kristof's formula estimates reliability, it is believed that the use of a .60 cut-off point will not have disguised the existence of any measurement error in the present study.

A further limitation concerns the argument of statistical significance versus statistical meaning. While the relevant correlations are significant, the meaning of this significance is limited by their small sizes. Similarly, the relatively small amount of variance explained suggests that while the predictor variables account for a significant amount of the variance in the dependent variable, a relatively small amount of variance is explained by the independent variables included in the model.

The inclusion of additional key variables (e.g., perceived control, competition) in future models may increase the amount of explained variance.

Conclusion

The coverage on Type A behaviour in the popular press has been so extensive that the construct has become an almost endearing excuse for excessive, often commercially desirable, but potentially destructive behaviour (Gordon & Gibbons, 1991). It is the potential for destruction which dictates that this perception should not be allowed to persist. Nor too, should the construct be discarded because of past inconsistency and controversy. Clearly, the questionable credibility of the construct in the past can be attributed to the inaccurate conceptualisation and measurement of the Type A construct (Thoreson & Powell, 1992).

The results of the present study concerning the Type A components demonstrate that Type A behaviour has negative concomitants (Steinkamp, 1990) which are associated with similarly negative affective disorders (Lee, 1992). By redefining the construct as multidimensional and using a psychometric measure which adequately reflects these in the present study it was possible to demonstrate that the components of Type A behaviour do in fact, impose a significant and differential effect on anxiety and depression. Due to a number of limitations, the present study was unable to determine whether the components of Type A behaviour exert similar effects on physiological health. By overcoming these, future research may be able to establish whether Type A components also exert differential effects on physiological condition.

The relationship between the aerobic fitness generated by regular exercise, be it during work or leisure, and reduced risk of CHD has been demonstrated by past research (De Geus et al., 1990). However, while the scientific and popular literature extol the virtues of exercise, it has been unable to determine precisely how exercise affects physiological condition (Desharnais et al., 1993). The present study reveals that it is the duration, and not the frequency, of exercise which is the key determinant of cardiovascular fitness. The present study also shows that change in anxiety is influenced negatively by the duration of exercise compliance and that neither the frequency nor duration of exercise predicts a change in depression. While numerous physiological, biochemical and psychological theories have been proposed to justify the relationship between exercise and psychological health, none has received conclusive empirical support and therefore, the possibility that exercise offers little more than a placebo effect on psychological condition cannot be denied (Desharnais et al., 1993). This possibility will need to be investigated in future research.

In conclusion, stress is an inherent part of life requiring ongoing adaptation (Cohen, 1981). It also remains a primary risk factor for CHD (Grossarth-Maticcek & Eysenck, 1991). It follows then, that any disposition liable to enhance the experience and consequences of stress requires attention. So too, does any factor able to mediate the negative effects of stress. Despite the existence of a considerable volume of research, knowledge of the process of adjustment to CHD remains fragmented (Johnson & Morse, 1990). Therefore, it is imperative that research continues to explore the relationships between Type A components, exercise and the psychological and physiological health of CHD patients. To this end, a refined model of the predictors of CHD patients' psychological and physiological health is

developed in Chapter 9. Incorporating the present findings with additional variables likely to influence prognosis, the model proposed in Chapter 9 outlines the implications for future research concerning CHD patients.

CHAPTER NINE

FUTURE RESEARCH IMPLICATIONS: TOWARD A REFINED MODEL OF CHD PATIENTS' BEHAVIOUR

The aim of the present research was to identify key factors involved in the process of recovery from CHD and establish the extent to which these predict prognosis. Two areas were examined, namely, Type A behaviour and exercise. The present research arose out of past research's failure to assess adequately these areas within the context of cardiac rehabilitation. On the one hand, Type A research has been limited by the inaccurate conceptualisation and measurement of the construct (Edwards et al, 1990a). Research which has conceptualised and measured Type A behaviour as a multidimensional construct has not examined the differential effects of the components on CHD patients' prognosis for recovery. On the other hand, extant research into the influence of exercise on CHD patients' psychological and physiological health is inconclusive and marred by methodological weaknesses (e.g., Kavanagh et al., 1977; NEHDP, 1981; Shephard et al., 1981; Valliant & Asu, 1985). In the present thesis, three studies were conducted to address these conceptual and empirical limitations of past research.

The first study compared the changes in anxiety and depression and the number of secondary cardiac events and associated surgical procedures of 20 cardiac patients who had completed a six month exercise programme with 20 who had been admitted to, but elected to withdraw from, the same programme for reasons other than ill health. No significant differences between the experimental and control groups' change in anxiety and depression and number of cardiac and surgical events were found. In the second study, the Type A Component Questionnaire was developed and validated to measure five of the key Type A components manifested by CHD patients, namely, achievement striving, impatience irritability, acting-out hostility, anger-in and anger-out. Thus, the second study generated a reliable and


valid self-report scale comprising independent measures of five of the primary components of the multidimensional Type A construct. The third study developed and tested a model of exercise compliance and Type A components as predictors of the psychological and physiological strain manifested by cardiac rehabilitation patients. This study supported the hypothesis that the predictors were differentially and independently related to psychological and physiological outcomes. However, the value of the model depicted in Figures 8.1 and 8.2 is limited by its exclusion of variables which are potentially central to the process of adjustment and to the prediction of prognosis. Therefore, the aim of Chapter 9 is to extend the present thesis to include additional factors likely to influence prognosis, and consequently, identify areas for future research. As can be seen from Figure 9.1, the model synthesises the variables examined in Study 1, 2 and 3 with antecedents, covariates, moderator and outcome variables not considered in the earlier model. A discussion of these and some of the key findings of the present research and the corresponding implications for future research follows.

Antecedents

The present research establishes that Type A behaviour comprises both toxic and benign components. With this knowledge, future rehabilitation programmes will be able to design interventions which simultaneously modify those components which exert a negative effect, and cultivate those which exert a benign effect. However, given the tenacity of Type A behaviour, it will only be possible to design interventions which generate enduring change if the primary personal and environmental factors which motivate and maintain these components are known (Price, 1982). Knowledge of these components will also prove crucial to understanding Type A's potential increased risk of CHD (Strube, 1988). Hence, the revised model makes provision for antecedent variables. It is in the theoretical formulations of Type A behaviour provided by Glass (1977a), Price (1982) and Strube (1987) that the primary antecedents can be found.

Figure 9.1

Revised Model of the Causes and Consequences of CHD Patients' Behaviour



As the need for control is a factor underlying much of Type A behaviour (Lee et al., 1990), it is included in the revised model as an antecedent. Glass' (1977a) model of the need for control has been discussed in Chapter 4 and will not be repeated here. Suffice to say that CHD undermines Type A's sense of control (Johnson & Morse, 1990). Consequently, the process of adjustment to CHD is dominated by the struggle to re-establish a sense of control (Johnson & Morse, 1990). It is in the struggle to achieve this control that achievement striving, impatient, irritable, hostile and angry behaviours manifest themselves (Matthews, 1982). The need to control the process of adjustment to CHD also increases compliance with an exercise based rehabilitation programme (Soloff, 1978a; 1978b). However, the act of struggling to achieve control may influence metabolic functions to the extent that it increases the risk of CHD (Friedman & Rosenman, 1959). Therefore, the response to control can be seen as both an antecedent of Type A and compliance behaviour and as a potential source of physiological strain.

A related factor likely to generate and maintain achievement striving, impatience, irritability, hostility, anger-in and anger-out is Type A's style of self-evaluation, which has at its core the need to prove oneself (Matthews, 1982; Price, 1982). In one of the most comprehensive theoretical models of Type A behaviour, Price (1982) asserts that much of Type A's behaviour is motivated by dysfunctional personal beliefs, arising out of socio-cultural values. Specifically, Type A's believe that vital resources are in scarce supply, the environment in which they find themselves is essentially hostile and that there exists no universal moral principle. These beliefs dictate that Type A's constantly try to prove and evaluate themselves in terms of excessively high standards of performance. They are also associated with Type A's fear that they will be judged unworthy if they fail to achieve sufficient observable accomplishments. Together, these beliefs motivate Type A's to achieve increasingly higher goals and to direct impatience, irritability, hostility, and anger at anyone or anything which threatens to hinder their ability to do so. The need to prove oneself

is also likely to influence Type A CHD patients' exercise performance through over compliance, though research has yet to examine this relationship. While Price's (1982) model is revered and referred to extensively in the Type A literature, little attempt has been made to test whether Type A CHD patients' style of self-evaluation stems from these beliefs and whether the need to prove oneself is in fact an antecedent of the various Type A and exercise compliance components (Yuen & Kuiper, 1992). Hence, the inclusion of the need to prove oneself in the revised model.

Strube's (1987) model of self-appraisal is also based on the premise that a dysfunctional style of self-evaluation spawns many of the Type A behaviours (Yuen & Kuiper, 1992). This model asserts that Type A's consider the achievement of goals to be contingent on acquiring accurate appraisals of their abilities (Strube, 1987). Consequently, Type A behaviour is motivated by the need to minimise uncertainty regarding these abilities (Yuen & Kuiper, 1992). To acquire the diagnostic information necessary for accurate appraisals, Type A's resort to exaggerated achievement striving, and aggressive and time urgent behaviour (Strube, 1988; Strube et al., 1986). As Type A's may fail to realise that they simply lack certain abilities, they may strive unnecessarily and consequently, subject themselves to considerable stress (Strube, 1988). Given that the period of recovery from CHD is characterised by substantial uncertainty, Type A CHD patients are particularly prone to these stress inducing activities. However, rehabilitation programmes do provide an arena wherein Type A's can acquire accurate information concerning their physical abilities. Rehabilitation programmes offer Type A CHD patients a regular opportunity to test their exercise performance and consequently, acquire information regarding ability (Naughton & Hellerstein, 1971). If exercise performance is seen as a means of achieving an accurate appraisal of ability, then Strube's concept of self-appraisal may act as an antecedent of exercise compliance among Type A CHD patients. While Strube's (1987) model fails to elucidate Type A's

underlying need to minimise uncertainty regarding ability, it does suggest that an aberrant style of self-appraisal is an antecedent of Type A behaviour. Therefore, self-appraisal is included as an antecedent in the revised model.

Within the context of cardiac rehabilitation, behaviour is further anteceded by the numerous stressors generated by CHD. An initial premise of the present research was, that while CHD is a manifestation of stress induced strain, it is also the source of subsequent stress and strain. This argument was supported by Study 3 which demonstrated that the anger-in and anger-out components of cardiac patients' behaviour, thought to be generated by stress, induced anxiety and depression. However, the present research failed to measure the number and intensity of stressors experienced. Therefore, it was unable to establish whether the stressors which characterise the recovery period are in fact both antecedents of Type A behaviour and directly related to subsequent strain. The model depicted in Figure 9.1 lists the stressors discussed in Chapter 2 which are likely to act as antecedents of achievement striving, impatience irritability, hostility, anger-in and anger-out and the frequency and duration of exercise compliance. It is likely that these stressors will also be directly related to strain. To determine whether a direct relationship exists between these stressors and both the Type A and compliance components and subsequent strain, future research should measure the frequency and intensity of CHD patients' exposure to specific stressors.

The Type A Components: Achievement Striving, Impatience Irritability, Hostility,
Anger-in and Anger-out

Study 2 of the present research demonstrated that the Type A behaviour manifested by cardiac patients comprises achievement striving, impatience irritability, hostility, anger-in and anger-out. Study 3 demonstrated that each of these act independently and differentially on measures of psychological health and thus, can be divided into

benign and toxic categories. By reconciling all five dimensions in a single model of Type A behaviour, Study 1 overcame two of the primary limitations of past research, namely, the failure to include the highly toxic components of anger and hostility and the failure to examine the incidence and effects of the five components in a high risk sample.

With the use of the Type A Component Questionnaire developed and validated in the present research it is possible to measure the components independently and identify specific relationships between the components and strain. With the use of this questionnaire it will also be possible to establish whether each component gains from its relationship with the other components (Carver, 1989). In other words, the Type A Component Questionnaire makes it possible to determine whether synergies exist between components. The present findings demonstrate that measuring the components independently not only enhances understanding of the Type A construct, but also provides an opportunity for determining which of the components are necessary and important (Carver, 1989) in the prediction of prognosis. However, to continue to focus on these Type A behaviours to the exclusion of other potential psychological and behavioural risk factors is scientifically impermissible (Grossarth-Maticek & Eysenck, 1991). Research by Cramer (1991) suggests that the Type A correlates, trait neuroticism and psychological distress (i.e., neurotic symptoms) may predict CHD. So too may other correlates of Type A behaviour such as hyperalertness (Price, 1982), authoritarianism (Byrne et al., 1989), and extraversion (Eysenck & Fulker, 1983). Therefore, future research should explore the relevance and prevalence of other components associated with Type A behaviour and increased risk of a negative prognosis.

Exercise Compliance

Study 1 of the present research failed to yield significant differences between the levels of strain of the experimental and control groups of CHD patients. However, it is in the possible causes of the non significant results that Study 1 makes a contribution to extant knowledge. First, these results emphasise the inadequacy of measuring compliance exclusively in terms of attendance (Rejeski et al., 1984; Sharkey, 1984). Second, they demonstrate that, in the short term, hard disease end points are an insufficiently sensitive measure of physiological strain and that future research needs to utilise a measure able to detect more subtle changes in physiological condition (Bethell & Mullee, 1990; Pisa et al., 1985). These issues were addressed in Study 3 by measuring the frequency and duration of exercise compliance in relation to a more sensitive index of cardiovascular condition, namely, maximal oxygen uptake. As this study showed that it is the duration measure of exercise compliance which is significantly related to maximal oxygen uptake, it is clear that research can no longer rely on attendance figures alone to examine the relationship between exercise and physiological condition.

Covariates

To limit the possibility of achieving spurious results, the present research controlled potentially confounding variables by including them in the model as covariates. The variables considered as possible covariates were age, sex, race, marital status, occupation and education. Of these, age and occupation were significantly related to the predictor variables and were included as covariates in the model tested in Study 3. Two additional variables which may impose a confounding effect, are pharmacological treatment and the genetic component of maximal oxygen uptake. Inclusion of these two variables as covariates was beyond the scope of the present research. Therefore, in the revised model, the list of covariates is extended to include

pharmacological treatment and the genetic component of maximal oxygen uptake.

Research indicates that a number of the drugs used in the treatment of CHD diminish the intensity of Type A behaviour (cf. Powell, 1987; Schneider et al., 1982). For example, the beta-blocking drugs prescribed for many CHD patients not only reduce the incidence of reinfarction by 25%, but also, reduce Type A behaviour (Abbott & Peters, 1988). As has been outlined in previous chapters, a heightened cardiovascular response to stress is believed to be the mechanism linking Type A behaviour to CHD. There is some evidence to suggest that beta-blockers might inhibit Type A's exaggerated cardiovascular response to stress (Haynes & Matthews, 1988). The influence of beta-blockers on Type A assessment may also render Type A classifications invalid (Matthews, 1988). Thus, given its relationship to Type A behaviour, pharmacological treatment constitutes a potential covariate (Hartely et al., 1987). A variable likely to be related to exercise compliance, and consequently maximal oxygen uptake, is genetic predisposition. Sibling and twin studies show that as much as 38% of maximal oxygen uptake is genetically determined (Bouchard, Boulay, Simoneau, & Pérusse, 1988). This explains why CHD patients' responses to training differ so greatly. By including approximately 40% of the variance in maximal oxygen uptake as a covariate, future research would be better able to identify the cause of the remaining variance in maximal oxygen uptake (Sandvik et al., 1993). The list of covariates included in the revised model is by no means exhaustive and should be extended or condensed according to the specific characteristics of the samples examined by future research.

Moderator Variables

Friedman and Booth-Kewley (1988) suggest that the modest size of the Type A-CHD relationship revealed in their meta-analysis may be the result of past research's failure to examine relevant moderating variables. Critical to the development of a

model for use in future research on CHD patients is an understanding of the factors which moderate the relationship between Type A and compliance behaviour and CHD (Powell, 1987). The present research indicates that, despite experiencing considerable stress, cardiac patients do not demonstrate significant levels of strain. As it is probable that situational and personality factors serve to develop resilience to stress, and consequently, modify strain (Ganellen & Blaney, 1984; Kobasa, Maddi & Courington, 1981), provision is made in the revised model for moderating variables. In interactive models, such as the one proposed here, the inclusion of moderating variables serves to increase the percentage of explained variance (Moosebrugger & Schermelleh-Engel, 1991).

Social Support

A variable likely to moderate the relationship between cardiac patients' stress and strain is social support (Orth-Gomér & Undén, 1990). The constant tension generated by Type A's impatience and hostility damages personal relationships to the extent that Type A's are prone to relatively unsupportive social networks (Abbott & Peters, 1988). The resulting social isolation is believed to contribute to physical morbidity and early mortality (Christensen & Smith, 1993). Indeed, several studies have revealed a significant relationship between the quantity and quality of social support and CHD morbidity and mortality (Abbott & Peters, 1988; Friedman & Booth-Kewley, 1987a; Lepore et al., 1993; Orth-Gomér & Undén, 1990). It is thought that, by protecting the individual from the pathological effects of extreme cardiovascular reactivity, social support reduces the impact of stress on the cardiovascular system (Lepore et al., 1993). A number of studies have demonstrated that social support reduces cardiovascular reactivity (e.g., Gerin, Pieper, Levy, & Pickering, 1992; Kamarck, Manuck & Jennings, 1990; Lepore et al., 1993). Therefore, it is possible that social support modifies CHD risk.

The individual's level of comfort with social support may also exert a moderating effect on exercise compliance, and consequently, physiological condition. Cardiac rehabilitation programmes are typically based on group activities, characterised by considerable social interaction and support (Blumenthal et al., 1982). For the socially extroverted individual comfortable with this milieu, support is likely to act as both a buffer against stress and as an inspiration to persist with healthy life-style changes (Rodin & Salovey, 1989). Comfort with social situations has been found to relate to compliance with a cardiac rehabilitation programme (Blumenthal et al., 1982). However, for the socially introverted patient, the supportive and interactive rehabilitation environment is likely to be so stressful as to cause these patients to withdraw from the programme (Blumenthal et al., 1982). Encouraging social introverts to remain in the programme may prove futile, as stress alters the physiological responses associated with exercise to the extent that it undermines the cardiovascular benefit of exercise (Roth et al., 1990).

Social support is also believed to moderate the relationship between stress and psychological strain (Bolger & Eckenrode, 1991). Rook (1987) asserts that social relationships not only render support with problems, but also provide a valuable source of companionship in the form of shared activities. Rook's (1987) research into this phenomenon reveals that the companionship derived from shared activities enhances psychological well-being and buffers the psychological impact of stress.

It may be that social support also exerts a negative moderating effect. Underlying social support is the norm of reciprocity wherein those rendering support will expect to receive a return for their efforts (Gouldner, 1960). According to Johnson and Morse (1990), CHD patients are often unable to reciprocate in a manner they would consider appropriate. The resulting sense of imbalance and indebtedness hinders patients' ability to participate fully in social relationships (Johnson & Morse, 1990). In this context, social support is likely to exacerbate the experience of stress and

strain. Clearly, social support has the potential to moderate the relationship between CHD patients' stress and strain either positively or negatively.

Locus of Control

A second moderator variable included in the revised model is locus of control. Type A's are believed to possess an internal locus of control (Ganster et al., 1991; Perloff, Yarnold, & Feltzer, 1988). An internal locus of control reflects a belief that the achievement of satisfaction is under the individual's personal control (Newton & Keenan, 1990) and protects the individual from the effects of stress (Kobasa, Maddi, & Zola, 1983). Individuals who are internally oriented tend to utilise adaptive health responses in terms of preventative and rehabilitative measures (Taylor & Cooper, 1988). Thus, internally oriented Type A's are likely to approach rehabilitation with the intention of controlling and hence facilitating their own process of recovery. Research shows that CHD patients who are internally oriented cope better with the process of adaptation than externally oriented patients (Terry, 1992). However, an internal locus of control is also positively associated with cardiovascular reactivity (Houston, 1986). It is possible that, while an internal locus of control increases resistance to stress, the act of imposing that control generates physiological responses which may culminate in strain. Thus, it is probable that locus of control will moderate the relationship between Type A behaviours and strain.

Perception of Health

A third moderator included in the model is the perception of health, be it optimistic or pessimistic. CHD patients' perception of their disease affects their reactions to symptoms and feelings of vulnerability (Martin & Lee, 1992). Patients' perception of their illness plays an important role in the process of adaptation to CHD (Desharnais et al., 1990). Maeland and Havick (1987) report a better rate of recovery

among patients who demonstrated a positive perception of their health and future functioning and few inaccurate beliefs concerning the limitations imposed by CHD. In comparison to pessimists, optimists have been found to enjoy a faster rate of recovery from a coronary artery bypass graft (Scheier et al., 1989), be more resistant to depression (Carver & Gaines, 1987) and to adopt more problem focused methods of coping (Scheier, Weintraub, & Carver, 1989). Therefore, it is likely that patients' optimistic or pessimistic perceptions of their health will moderate the process of recovery.

Outcome Variables

Anxiety, Depression and Mixed Anxiety-Depressive Disorder

The present research examined the impact of the psychological and behavioural predictors on anxiety, depression and maximal oxygen uptake. In the third study of this research, anger-out predicted less reduction in anxiety while both anger-in and anger-out predicted less reduction in depression. Contrary to expectations, impatience irritability and hostility were not associated with anxiety or depression. The duration of exercise also predicted less reduction in anxiety but was not associated positively with depression. Further, the frequency of exercise was related to neither anxiety nor depression. The results of past research suggest that a significant relationship between exercise and a reduction in these disorders largely occurs among patients diagnosed as clinically anxious or depressed (De Geus et al., 1993). Thus, the non significant relationship can be attributed to the fact that the present sample were not severely anxious or depressed. Given that the experience of anxiety and depression is pervasive among CHD patients (Brown & Munford, 1984; Hackett & Cassem, 1975), they are retained as measures of strain in the revised model. However, it is asserted that the conceptualisation and measurement of

anxiety and depression should be adapted to suit the psychological status of the sample.

Studies utilising samples characterised by low anxiety and depression at the inception of the research should accommodate the new diagnostic category included in the DSM-IV, namely, mixed anxiety-depressive disorder (American Psychiatric Association, 1994). The conceptual overlap between anxiety and depression translates to an average correlation of .61 between scales measuring the two disorders and a shared variance ranging between 37% and 48% (Endler & Parker, 1990; Gershon et al., 1991; Gotlib, 1984). Anxiety and depression are now viewed as coexisting parts of a general psychological distress state or response to stress (Endler & Parker, 1990). Symptoms of anxiety and depression frequently coexist in CHD populations (Gershon et al., 1992). Further, the two disorders only become distinct from one another as the severity and intensity thereof increases (Gershon et al., 1992). As the present research shows, the anxiety and depression manifested by CHD patients is not severe. A scale designed to measure the less severe category of mixed anxiety-depressive disorder could have important implications for assessing the strain experienced by CHD patients. With the application of such a scale it would be possible to determine whether the less severe form of mixed anxiety-depressive disorder is a pervasive disorder and whether, given sufficiently stressful conditions, can develop into more severe anxiety and depression (Gershon et al., 1991). By including anxiety and depression as both independent and combined measures of strain, the revised model makes provision for samples demonstrating varying degrees of these disorders. It also provides future research with the opportunity of comparing the utility of the two approaches to the measurement of anxiety and depression.

Maximal Oxygen Uptake and Coronary Artery Angiography

The relevance of maximal oxygen uptake as a measure of physiological outcome was supported by the present research which revealed a significant predictive relationship between the duration of exercise compliance and increased maximal oxygen uptake. However, contrary to expectations, the third study of the present research failed to reveal any relationships between the Type A components, exercise frequency and maximal oxygen uptake. One possible explanation for this is that while maximal oxygen uptake is a more sensitive measure of cardiovascular condition than hard disease end points, it may still be relatively insensitive to subtle changes in the further development of atherosclerosis. A coronary artery angiography provides the most precise measure of the extent of atherosclerotic build-up (Miller et al., 1991) and a reliable means of predicting subsequent CHD (Loscalzo, 1990). Unlike maximal oxygen uptake it is not a functional assessment and therefore does not provide information concerning changes in cardiorespiratory fitness (Schneiderman, Weiss, & Kaufman, 1989). Therefore, in the interests of greater precision, future research should measure outcome in terms of both angiographic reports and maximal oxygen uptake. By employing the measures in tandem, future research will be able to determine changes in both the extent of atherosclerosis and in levels of cardiorespiratory fitness with considerable accuracy.

Achievement of Control

The revised model includes achievement of control as a measure of recovery. To date the majority of cardiac rehabilitation studies have considered recovery in terms of return to work (e.g., Greenland & Briody, 1984; Smith & O'Rourke, 1988; World

Health Organization, 1981). These goals may not be consistent with the goals set by the patient themselves. Research implies that CHD patients' primary goal in the adjustment process is to regain control of their bodies and their lives (Johnson & Morse, 1990). The belief that control has been achieved enhances the ability to manage stress and cope with difficult and distressing circumstances (Ben-Sira, 1983). Conversely, the belief that control is evasive results in the abandonment of the battle to adjust and the acceptance of a permanent role as an invalid (Johnson & Morse, 1990). Therefore, future research should include a measure of recovery which is relevant to CHD patients, namely, the achievement of control.

Other Diseases

The interface between personality and disease has been studied widely and it is generally accepted that numerous illnesses have substantial psychological components (Carmody, Crossen, & Wiens, 1989). However, it is the relationship between Type A behaviour and CHD which has drawn researchers' attention (Rimé et al., 1989). As a consequence little attempt has been made to determine whether Type A behaviour predicts other diseases. As Type A behaviour generates a chronic state of stress, it is likely to predict other stress related diseases (Matteson & Ivancevich, 1980). For example, behavioural and personality components are implicated in the pathogenesis of cancer (Rodin & Salovey, 1989). Psychological factors, which include Type A behaviour among them have also been found to influence the development and progression of somatic illnesses such as skin disorders (Kaptein, 1990), asthma (Everaerd et al., 1990; Rimé et al., 1989), hypertension (Godaert, 1990), peptic ulcers, thyroid problems and rheumatoid arthritis (Rimé et al., 1989). While, the present research focused on the relationship between Type A behaviour and secondary CHD, it is possible that Type A behaviour

may be a general disease prone condition rather than a specific risk factor for CHD (Rimé et al., 1989). Therefore, it is important that future research consider the relationship between the construct and other diseases.

In sum, the revised model retains the original outcome variables of anxiety, depression and maximal oxygen uptake. To accommodate samples characterised by non-clinical anxiety and depression, the revised model suggests the measurement of the less severe outcome, Mixed anxiety-depressive disorder. In order for future research to determine the relationship between the predictor variables and subtle changes in atherosclerotic build-up, the revised model advocates the additional assessment of cardiovascular condition by means of coronary angiography. Also included is an outcome measure consistent with the goals set by CHD patients, namely the achievement of control. The revised model further suggests that Type A research should be extended beyond CHD, to examine the impact of the various components on other diseases.

Feedback Loops

In their conceptualisation of Type A behaviour, Friedman and Rosenman (1974) described the construct as a mode of response to environmental challenges to self-appraisal. While Type A behaviour was originally defined in terms of person-environment interactions, it has typically been studied as a personality trait (Thoreson & Powell, 1992). To restore Type A behaviour to its status as an interactional construct, it is necessary to examine the interdependent development and function of thoughts, perceptions, emotions, behaviours and physiological processes which occur in response to the environment over time (Thoreson & Powell,

1992). Thus, it is important that future research examines the reciprocal, as opposed to the linear, relationships between the various factors in the model. To accommodate this need, the revised model advocates the inclusion of a feedback loop between the antecedents and the outcome variables (Smith & Anderson, 1986). The inclusion of a feedback loop is motivated by the assertion that the way in which Type A's evaluate themselves and the extent to which they feel that their need for control is being fulfilled is influenced by how they perceive their physiological response to the cardiac event and their ability to cope with it (Bandura, 1982). Thus, Type A's positive or negative perception of their ability to adjust during the process of recovery will influence their future style of self-evaluation.

Conclusion

An aim of the present research was to reconcile components of Type A behaviour and exercise in a model of predictors of recovery from CHD. Since its identification in the 1950's the Type A construct has enjoyed erratic popularity. Following the significant results of the WCGS and Framingham study the construct was met with considerable enthusiasm. With the inconclusive and often negative results generated by subsequent studies, enthusiasm waned to the point of extreme scepticism (Thoreson & Powell, 1992). This led the proponents of Type A behaviour to investigate the multidimensional nature of the construct. These studies showed that the construct comprised both benign and toxic components each of which exerted a differential effect. The reconceptualisation and study of Type A behaviour as a multidimensional versus a global construct has rekindled enthusiasm to the extent that the concept of an independent relationship between psychological and behavioural constructs and health has achieved renewed credibility. Certainly, the

results of the present research support this revival. The present findings that achievement striving, impatience irritability, hostility anger-in and anger-out are differentially and independently related or unrelated to measures of health demonstrates that the term Type A behaviour may be, as suggested by Carver (1989), little more than a convenient summary of the various components which contribute to it. Despite these encouraging findings, it is recognised that the conceptualisation and measurement of Type A behaviour in the present research may not be sufficiently comprehensive or interactional to reduce the confusion surrounding the Type A construct.

Research into the prognostic role of aerobic exercise has been more consistent and numerous studies have demonstrated a relationship between exercise and reduced psychological strain and CHD risk (De Geus et al., 1990). While this relationship has been firmly established, little attempt has been made to look beyond the generic concept of exercise to determine the precise means by which exercise exerts such effects (Desharnais et al., 1993; Sharkey, 1984). By examining the components of exercise, and establishing that it is the duration of exercise which determines cardiovascular fitness, the present research contributes to an understanding of the mechanisms underlying the relationship between exercise and reduced CHD risk. It remains a task of future research to design studies able to identify the means by which exercise influences psychological strain.

To increase understanding of the predictive relationship between exercise and Type A components and the psychological and physiological strain of CHD patients, the present chapter proposes a transactional model of CHD patients' behaviour. The factors which trigger the model are the various CHD events and interventions. These

generate a number of stressors and cognitive antecedents, which in turn, motivate the achievement striving, impatience irritability, hostility, and anger expression brought to predict prognosis. Also included as predictors of prognosis are the frequency and duration of exercise compliance. Physiological prognosis is measured in terms of VO_2 max and angiographic readings. Provision is also made in the model for the relationship between Type A behaviour and other diseases. Psychological outcomes in the revised model are anxiety and depression, mixed anxiety-depression disorder and achievement of control. The relationship between the predictors and physiological and psychological outcomes is moderated by social support, locus of control and health perceptions. The revised model acknowledges the interdependent relationship between factors and provides for the impact of changes over time (Smith & Anderson, 1986). By testing this model it will be possible to establish which combinations of Type A and exercise compliance components increase the risk of persistent disease and which predict recovery. With this knowledge, it will be possible to design rehabilitation interventions which enhance health and reduce CHD risk through the modification of toxic, and cultivation of salutary, behaviours.

Summary

In conclusion, the present research has attempted to advance knowledge of the factors involved in the prediction of CHD patients' prognosis for psychological and physiological recovery. With respect to the role of exercise, the present research reveals that it is the duration of performance which influences cardiovascular condition. In so doing, it clarifies the means by which exercise exerts an influence on health. Given that the majority of extant studies have used the more convenient measure of exercise frequency, this finding may explain the equivocal results of past exercise research. The present research also answers the call for the recognition and study of Type A behaviour as a multidimensional construct (Thoreson & Powell, 1992). By defining CHD patients' Type A behaviour as comprising five, independent components, the present research reduces some of the ambiguity surrounding the construct. It also advances Type A component research by examining the differential impact of these components on CHD patients' psychological and cardiovascular health. The present research makes a further contribution to the theory and practice of Type A research through its development of a reliable and valid measure of CHD patients' Type A behaviour. While the present thesis may have contributed to existing knowledge, the work on the relationship between psychological and behavioural predictors of secondary CHD is far from complete. For, even when all classical and known psychosocial risk factors are considered, there remains a substantial amount of unaccounted for variance in the incidence of CHD (Chesney, 1993). Hence, there is a critical need to continue the search for variables which can be declared as indisputable risk factors of primary and secondary CHD.

REFERENCES

- Abbott, A.V., & Peters, R.K. (1988). Type A behavior and coronary heart disease: An update. American Family Physician, 38(5), 105-110.
- Akaike, H. (1974). A new look at the statistical model identification. IEEE Transactions on Automatic Control, AC-19, 716-723.
- Allen, J.K., Becker, D.M., & Swank, R.T. (1990). Factors related to functional status after coronary artery bypass surgery. Heart and Lung, 19(4), 337-343.
- Aitchler, L., & Motta, R. (1994). Effects of aerobic and nonaerobic exercise on anxiety, absenteeism, and job satisfaction. Journal of Clinical Psychology, 50(6), 829-840.
- American Psychiatric Association (1980). Diagnostic and statistical manual of mental disorders (3rd ed.). Washington DC.
- Anastasi, A. (1968). Psychological Testing (3rd ed.). New York: MacMillan.
- Anderson, N.B., Williams, R.B., Lane, J.D., Hancy, T., Simpson, S., & Houseworth, S.J. (1986). Type A behavior, family history of hypertension, and cardiovascular responsivity among black women. Health Psychology, 5(4), 393-406.
- Antonovsky, H., & Sagy, S. (1985). The development of a sense of coherence and its impact on responses to stress situations. Journal of Social Psychology, 126(2), 213-225.
- Astrand, P.O., & Rodahl, K. (1977). Textbook of Work Physiology. New York: McGraw-Hill.
- Atlas, J.A., & DiScipio, W.J. (1992). Correlations of Beck Depression Inventory and Reynolds Adolescent Depression Scale. Psychological Reports, 70, 621-622.
- Auld, F. (1985). IPAT Anxiety Scale. In D.J. Keyser, & R.C. Sweetland (Eds.), Test critiques (Vol. 2) (pp. 357-362). Kansas City: Test Corporation of America.
- Ayanian, J.Z. (1993). Heart disease in black and white. New England Journal of Medicine, 329(9), 656-657.
- Ayanian, J.Z., & Epstein, A.M. (1991). Differences in the use of procedures between women and men hospitalized for coronary heart disease. New England Journal of Medicine, 325, 221-225.
- Baer, P.E., Collins, F.H., Bourenhoff, G.C., & Ketchel, M.F. (1979). Assessing personality factors in essential hypertension with a brief self-report instrument. Psychosomatic Medicine, 16, 721-730.
- Bandura, A. (1982). Self-efficacy mechanisms in human agency. American Psychologist, 37, 122-147.
- Bandura, A. (1991). Social cognitive theory of self-regulation. Organizational Behaviour and Human Decision Processes, 50, 248-287.
- Barefoot, J.C., Dahlstrom, W.C., & Williams, R.B. (1983). Hostility, CHD incidence and total mortality: A 25 year follow-up study of 255 physicians. Psychometric Medicine, 45, 59-63.

Barefoot, J.C., Peterson, B.L., Harrell, F.E., Hlatky, M.A., Pryor, D.B., Haney, T.L., Blumenthal, J.A., Siegler, I.C., & Williams, R.B. Jr. (1989). Type A behavior and survival: A follow-up study of 1,467 patients with coronary artery disease. American Journal of Cardiology, 64(8), 427-432.

Barling, J., Bluen, S.D., & Moss, V. (1990). Dimensions of Type A behaviour and marital dissatisfaction - Disentangling the effects of Achievement Striving and Impatience-Irritability. Journal of Psychology, 124(3), 311-319.

Barling, J., & Charbonneau, D. (1992). Disentangling the relationship between Type A behaviour, performance and health. Journal of Organizational Behavior, 13(4), 369-377.

Barling, J., & Milligan, J. (1987). Some psychological consequences of striking: A six month longitudinal study. Journal of Occupational Psychology, 8, 127-138.

Barnett, P.A., & Gotlib, I.H. (1988). Psychosocial functioning and depression: Distinguishing among antecedents, concomitants, and consequences. Psychological Bulletin, 104, 97-126.

Barnett, V., & Lewis, T. (1978). Outliers in statistical data. Chichester: John Wiley & Sons.

Bass, C. (1984). Psychosocial outcome after coronary bypass surgery. Circulation, 66(5), III 24-28.

Baum, A., Grunberg, N.E., & Singer, J.E. (1982). The use of psychological and neuroendocrine measurements in the study of stress. Health Psychology, 1, 217-236.

Beck, A.T. (1967). Depression: Clinical, experimental, and theoretical aspects. New York: Harper & Row.

Beck, A.T. (1970). Depression: Causes and treatment. Philadelphia: University of Pennsylvania Press.

Beck, A.T. (1987). Cognitive models of depression. Journal of Cognitive Psychotherapy, an International Quarterly, 1, 5-37.

Beck, A.T., & Steer, R.A. (1984). Internal consistencies of the original and revised Beck Depression Inventory. Journal of Clinical Psychology, 40, 1365-1367.

Beck, A.T., Ward, C.H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for assessing depression. Archives of General Psychiatry, 4, 53-63.

Becker, L.B., Han, B.H., Meyer, P.M., Wright, F.A., Rhodes, K.V., Smith, D.W., Barrett, J., & the CPR Chicago Project. (1993). Racial differences in the incidence of cardiac arrest and subsequent survival. New England Journal of Medicine, 329(9), 600-606.

Bechr, T.A., & Newman, J.E. (1978). Job stress, employee health, and organizational effectiveness: A facet analysis, model, and literature review. Personnel Psychology, 31, 665-699.

Belgian-French Pooling Project. (1984). Assessment of Type A behaviour by the Bortner scale and ischaemic heart disease. European Heart Journal, 5, 440-446.

Belsey, D.A. (1991). Conditioning diagnostics: Collinearity and weak data in regression. New York: John Wiley & Sons.

Belsey, D.A., Kuh, E., & Welsh, R.E. (1980). Regression diagnostics - identifying influential data and sources of collinearity. New York: John Wiley & Sons.

- Belkin, L. (1990). The love-your-heart guide for the 1990s. Chicago: Contemporary Books.
- Bengtsson, K. (1983). Rehabilitation after myocardial infarction. Scandinavian Journal of Rehabilitative Medicine, 15, 1-9.
- Ben-Sira, Z. (1983). Personal control over aversive stimuli and its relationship to stress. Psychological Bulletin, 80, 286-303.
- Bergman, L.R., & Magnusson, D. (1986). Type A behaviour: A longitudinal study from childhood to adulthood. Psychosomatic Medicine, 48(1/2), 134-142.
- Berkowitz, L. (1983). The experience of anger as a parallel process in the display of impulsive 'angry' aggression. In R.G. Geen & E.I. Donnerstein (Eds.), Aggression: theoretical and empirical reviews. Vol 1. (pp. 103-133). New York: Academic Press.
- Bethell, H.J.N., & Mullee, M.A. (1990). A controlled trial of community based coronary rehabilitation. British Heart Journal, 64, 370-375.
- Biaggio, M.K., & Godwin, W.H. (1987). Relation of depression to anger and hostility constructs. Psychological Reports, 61, 87-90.
- Biddles, B.J., Slavings, R.L., & Anderson, D.S. (1985). Panel studies and causal inference. Journal of Applied Behavioural Science, 21(10), 79-93.
- Bluen, S.D. (1991). Predicting successful career choice. Unpublished masters dissertation, University of the Witwatersrand, South Africa.
- Bluen, S.D. (1986). Consequences and moderators of industrial relations stressors. Unpublished doctoral dissertation, University of the Witwatersrand, South Africa.
- Bluen, S.D., & Barling, J. (1987). Stress and the industrial relations process: Development of the Industrial Relations Event Scale. South African Journal of Psychology, 17(4), 150-159.
- Bluen, S.D., Barling, J., & Burns, W. (1990). Predicting sales performance, job satisfaction, and depression by using the Achievement Strivings and Impatience-Irritability dimensions of Type A behavior. Journal of Applied Psychology, 75(2), 212-216.
- Bluen, S.D., & Donald, C. (1991). The nature and measurement of in-company industrial relations climate. South African Journal of Psychology, 21(1), 12-20.
- Bluen, S.D., & Jubiler-Lurie, V. (1990). Some consequences of labour-management negotiations: Laboratory and field studies. Journal of Organizational Behavior, 11, 105-118.
- Bluen, S.D., & Odesnik, J. (1988). Township unrest: Development of the Township Life Events Scale. South African Journal of Psychology, 18(2), 50-57.
- Blumenthal, J.A., Barefoot, J.C., Burg, M.M., & Williams, R.B. (1987). Psychological correlates of hostility among patients undergoing coronary angiography. British Journal of Medical Psychology, 60, 349-355.
- Blumenthal, J.A., O'Toole, L.C., & Haney, T. (1984). Behavioral Assessment of the Type A Behavior Pattern. Psychosomatic Medicine, 46, 415-423.
- Blumenthal, J.A., Williams, R.B., Kong, Y., Schanberg, S.M., & Thompson, L.W. (1978). Type A Behavior Pattern and Coronary Atherosclerosis. Circulation, 58(4) 634-639.

- Blumenthal, J.A., Williams, S., Wallace, A.G., Williams, R.B., & Needles, T.L. (1982). Physiological and psychological variables predict compliance to prescribed exercise therapy in patients recovering from myocardial infarction. Psychosomatic Medicine, *44*, 141-147.
- Blumenthal, J.A., Williams, S., Williams, R.B., & Wallace, A.G. (1980). Effects of exercise on the Type A (coronary prone) behavior pattern. Psychosomatic Medicine, *42*(2), 289-296.
- Bolger, N., & Eckenrode, J. (1991). Social relationships, personality, and anxiety during a major stressful event. Journal of Personality and Social Psychology, *61*(3), 440-449.
- Boogaard, M.A.K. (1984). Rehabilitation of the female patient after myocardial infarction. Nursing Clinics of North America, *19*(3), 433-439.
- Booth-Kewley, S., & Friedman, H.S. (1987). Psychological Predictors of Heart Disease: A Quantitative Review. Psychological Bulletin, *101*(3), 343-362.
- Bortner, R.W. (1969). A short rating scale as a potential measure of pattern A behavior. Journal of Chronic Diseases, *22*, 87-91.
- Bouchard, C., Boulay, M.R., Simoneau, J., Lortie, G., & Pérusse, L. (1988). Hereditary and trainability of aerobic and anaerobic performances. An update. Sports Medicine, *5*, 69-73.
- Bozdogan, H. (1987). Model selection and Akaike's Information Criterion (AIC): The general theory and its analytical extensions. Psychometrika, *52*(3), 345-370.
- Bradley, C. (1988). Stress and diabetes. In S. Fisher & J. Reason (Eds.), Handbook of life stress, cognition and health. Chichester: John Wiley & Sons.
- Braestrup, C. (1982). Neurotransmitters and CNS disease: Anxiety. Lancet, *2*, 1030-1034.
- Brief, A.P., Rude, D.E., & Rabinowitz, S. (1983). The impact of Type A behavior pattern on subjective work load and depression. Journal of Occupational Behavior, *4*, 157-164.
- Briggs, S.R., & Cheek, J.M. (1986). The role of factor analysis in the development and evaluation of personality scales. Journal of Personality, *54*(1), 106-148.
- Brink, T.L., Yesavage, J., Lum, D., Hiersema, P., Adey, N., & Rose, T. (1982). Screening tests for geriatric depression. Clinical Gerontologist, *1*, 37-44.
- Brown, G.H & Harris, T. (1978). The social origins of depression: A study of psychiatric disorder in women. London: Tavistock.
- Brown, J.D. (1988). Environmental and Nuclear Threats. In S. Fisher & J. Reason (Eds.), Handbook of life stress, cognition and health. Chichester: John Wiley & Sons.
- Brown, J.D. (1991). Staying fit and staying well: Physical fitness as a moderator of life stress. Journal of Personality and Social Psychology, *60*(4), 555-561.
- Brown, M.A., & Munford, A. (1984). Rehabilitation of post MI depression and psychological invalidism: A pilot study. International Journal of Psychiatry in Medicine, *13*(4), 281-298.
- Bruce, R.A. (1971). Exercise testing of patients with coronary heart disease: Principles and normal standards for evaluation. Annals of Clinical Research, *3*, 323-332.
- Bruhn, J., Chandler, B., & Wolf, S. (1969). A psychological study of survivors and nonsurvivors of myocardial infarction. Psychosomatic Medicine, *31*, 8-19.

Brunson, B.I., & Matthews, K.A. (1981). The Type A coronary-prone behavior pattern and reactions to uncontrollable stress. An analysis of performance strategies, affect, and attributions during failure. Journal of Personality and Social Psychology, 40, 90-918.

Budnick, H.N. (1991). Heart to heart. A guide to the psychological aspects of heart disease. Santa Fe: Health Press

Bumberry, W., Oliver, J.M., McClure, J.N. (1978). Validation of the Beck Depression Inventory in a university population using psychiatric estimates as the criterion. Journal of Consulting and Clinical Psychology, 46, 150-155.

Burke, R.J. (1984). Beliefs and fears underlying Type A behaviour. Psychological Reports, 54, 655-662.

Burke, R.J., & Deszca, E. (1984). What makes Sammy run - so fast and aggressively? Beliefs and fears underlying Type A behavior. Journal of Occupational Behaviour, 5, 219-227.

Burns, W. (1992). Correlates of the multidimensional components of Type A behaviour. Unpublished doctoral thesis, University of the Witwatersrand.

Burns, W., & Bluen, S.D. (October, 1990). A multidimensional model of the components of Type A behaviour. Paper presented at the Eighth National Congress of the Psychological Association of South Africa, Port Elizabeth.

Burns, W., & Bluen, S. (1992). Assessing a multidimensional Type A behaviour scale. Personality and Individual Differences, 13(9), 977-986.

Byrne, D.G., Reinhart, M.I., & Heaven, P.C.L. (1989). Type A behaviour and the authoritarian personality. British Journal of Medical Psychology, 62, 163-172.

Byrne, D.G., & Rosenman, R.H. (1986). Type A Behaviour and the experience of affective discomfort. Journal of Psychosomatic Research, 30(6), 663-672.

Caine, T.M., Foulds, G.A., & Hope, K. (1967). Manual of the Hostility and Direction of Hostility Questionnaire. London: University of London Press.

Campbell, I.A., Burgess, P.M., & Finch, S.J. (1984). A factorial analysis of Beck Depression Inventory Scores. Journal of Clinical Psychology, 40, 992-996.

Cantor, J.R., Zillman, D., & Day, K.D. (1978). Relationships between cardiorespiratory fitness and physiological responses to films. Perceptual and Motor Skills, 46, 1123-1130.

Caplan, R.D. (1971). Organizational stress and individual strain: A social psychological study. Unpublished doctoral dissertation, University of Michigan. Ann Arbor.

Carey, M.P., Kelley, M.L., Buss, R.R., & Scott, W.O. (1986). Relationship of activity to depression in adolescents: Development of the adolescent activities checklist. Journal of Consulting and Clinical Psychology, 54, 320-322.

Carlson, N.R. (1981). Physiology of behavior. (3rd ed.). Boston: Allyn & Bacon.

Carmelli, D., Rosenman, R., Chesney, M., Fabsitz, R., Lee, M., & Borhani, N. (1988). Genetic heritability and shared environmental influences of Type A measures in the NHLBI twin study. American Journal of Epidemiology, 127(5), 1041-1052.

Carmody, T.P., Crossen, J.R., & Wiens, A.N. (1989). Hostility as a health risk factor: Relationship with neuroticism, Type A behavior, attentional focus and interpersonal style. Journal of Clinical Psychology, 45(5), 754-762.

Carney, R.M., Rich, M.W., & Freeland, K.E. (1988). Major depressive disorder predicts cardiac events in patients with coronary artery disease. Psychosomatic Medicine, 50, 627-633.

Carney, R.M., Rich, M.W., Tevelde, A., Saini, J., Clark, K., & Jaffe, A. (1987). Major depressive disorder in coronary artery disease. American Journal of Cardiology, 60, 1273-1275.

Carver, C.S. (1989). How should multifaceted personality constructs be tested? Issues illustrated by self-monitoring, attributional style, and hardiness. Journal of Personality and Social Psychology, 57, 577-585.

Carver, C.S., Coleman, E., Glass, D.C. (1976). The Coronary-prone behavior pattern and the suppression of fatigue on a treadmill test. Journal of Personality and Social Psychology, 33(4), 460-466.

Carver, C.S., & Humphries, C. (1982). Social psychology and the Type A coronary-prone behavior pattern. In G.S. Sanders, & J. Suls (Eds.), Social Psychology of Health and Illness. (pp. 33-64). Hillsdale, New Jersey: Erlbaum.

Case, R.B. (1988). Type A behavior. American Heart Journal, 116(6 part 1), 1653-1653.

Case, R.B., Heller, S.S., Case, N.B., & Moss, A.J. (1985). Type A Behavior and survival after acute myocardial infarction. New England Journal of Medicine, 312, 737-741.

Cassem, N.H., & Hackett, T.P. (1977). Psychological aspects of myocardial infarction. Medical Clinics of North America, 61, 711-721.

Cattell, R.B. (1966). Handbook of multivariate experimental psychology. Chicago: Rand McNally.

Cattell, R.B., & Scheier, I.H. (1976). The IPAT anxiety scale. Champaign: Institute for Personality and Ability Testing.

Cattell, R.B., Scheier, I.H., & Madge, E.M. (1968). Handbook for the IPAT anxiety scale questionnaire. Pretoria: Human Sciences Research Council.

Cattell, R.B., & Vogelmann, S. (1977). A comprehensive trial of the scree and KG criteria for determining the number of factors. Journal of Multivariate Behavioral Research, 12, 289-325.

Cay, E., Vetter, N., Philips, A.E., & Dugard, P. (1972). Psychological status during recovery from an acute heart attack. Journal of Psychosomatic Research, 16, 425-435.

Cay, E., Zaitsev, V.P., & Rudnicki, S. (1985). Psychological factors in patients after myocardial infarction. In V. Kallio & E. Cay (Eds.), Rehabilitation after myocardial infarction. The European experience. (pp. 33-53). Copenhagen: World Health Organization.

Chatterjee, S., & Price, B. (1977). Regression analysis by example. New York: John Wiley & Sons.

Chesney, M.A. (1993). Social isolation, depression and heart disease: Research on women broadens the agenda. Psychosomatic Medicine, 55, 426-433.

Chesney, M.A., Black, J.W., Chadwick, J.H., & Rosenman, R.H. (1981). Psychological correlates of the Type A behavior pattern. Journal of Behavioral Medicine, 4, 217-229.

- Chesney, M.A., Ekman, P., Friesen, W.V., Black, G.W., & Hecker, H.L. (1990). Type A Behavior Pattern: Facial behavior and speech components. Psychosomatic Medicine, 53, 307-319.
- Christensen, A.J., & Smith, T.W. (1993). Cynical hostility and cardiovascular reactivity during self-disclosure. Psychosomatic Medicine, 55, 193-202.
- Christensen, L.B. (1985). Experimental methodology (3rd ed.). Newton: Allyn & Bacon.
- Chung, E.K. (1983). Exercise Electrocardiography. Practical Approach (2nd ed.). Baltimore: Williams & Wilkins.
- Cinciripini, P.M. (1986). Cognitive stress and cardiovascular reactivity. II. Relationship to atherosclerosis, arrhythmias, and cognitive control. American Heart Journal, 112(5), 1051-1065.
- Cleveland, S.E., & Johnson, D.L. (1962). Personality patterns in young males with coronary disease. Psychosomatic Medicine, 24, 600-611.
- Cochrane, R. (1973). Hostility and neuroticism among unselected essential hypertensives. Journal of Psychosomatic Research, 17, 215-218.
- Cohen, F. (1981). Stress and bodily illness. Psychiatric Clinics of North America, 4(2), 269-286.
- Cohen, F. (1987). Measurement of Coping. In S.V. Kasl & C.L. Cooper (Eds.), Stress and health: Issues in research methodology (pp. 283-318). New York: John Wiley & Sons.
- Cohen, F., & Lazarus, R.S. (1979). Coping with the stresses of illness. In G.C. Stone, F. Cohen, & N.E. Adler (Eds.), Health psychology (pp. 217-254). San Francisco: Jossey-Bass.
- Cohen, J., & Cohen, P. (1983). Applied multiple regression / correlation analysis of the behavioral sciences. Hillsdale, New Jersey: Erlbaum.
- Cohen, J.B. & Reed, D. (1985). Type A behavior and coronary heart disease among Japanese men in Hawaii. Journal of Behavioral Medicine, 8, 343-352.
- Cohen, J.B., Syme, S.L., Jenkins, C.D., Kagan, A., & Zyzanski, S.J. (1979). Cultural context of Type A behavior and risk for CHD: A study of Japanese-American males. Journal of Behavioral Medicine, 2, 375-384.
- Cohen, S., & Wills, T.A. (1985). Stress, social support, and the buffering hypothesis. Psychological Bulletin, 98, 310-357.
- Cohn, K., Kamm, B., Feteih, N., Brand, R., & Goldschlager, N. (1979). Use of treadmill score to score and quantify ischaemic response and predict extent of coronary disease. Circulation, 59(2), 286-296.
- Concise Oxford Dictionary (7th ed.). (1982). Oxford: Clarendon Press.
- Constant, J. (1986). Prognostic information from early post-infarction exercise testing. American Journal of Medicine, 81(4), 655-660.
- Contrada, R.J., & Krantz, D.S. (1987). Measurement bias in health psychology research designs. In S.V. Kasl & C.L. Cooper (Eds.), Stress and health: Issues in research methodology (pp. 57-78). New York: John Wiley.

- Cook, R.D., & Weisberg, S. (1982). Residuals and influence in regression. London: Chapman & Hall.
- Cook, W.W., & Medley, D.M. (1954). Proposed hostility and pharisaic-virtue scales for the MMPI. Journal of Applied Psychology, 38, 414-418.
- Cooper, C.L., & Marshall, J. (1976). Occupational sources of stress. Journal of Occupational Psychology, 49, 11-28.
- Cooper, K.H., Meyer, B.U., Blide, R., Pollock, M., & Gibbons, L. (1991). The important role of fitness determination and stress testing in predicting coronary incidence. Annals of the New York Academy of Sciences, 301, 642-654.
- Cornett, S.J., & Watson, J.E. (1984). Cardiac rehabilitation: An interdisciplinary team approach. New York: John Wiley.
- Coronary Drug Project Research Group (1974). Factors influencing long term prognosis after recovery from myocardial infarction: three year findings of the Coronary Drug Project. Journal of Chronic Diseases, 27(6), 267-285.
- Cortina, J.M. (1993). What is coefficient alpha? An examination of theory and applications. Journal of Applied Psychology, 78(1), 98-104.
- Costa, P.T., & McCrae, R.R. (1987). Neuroticism, somatic complaints and disease: Is the bark worse than the bite? Journal of Personality, 55, 299-316.
- Cox, T. (1985). The nature and measurement of stress. Ergonomics, 28(8), 1155-1163.
- Cox, T. (1988). Psychobiological factors in stress and health. In S. Fisher & J. Reason (Eds.), Handbook of life stress, cognition and health. Chichester: John Wiley.
- Cramer, D. (1991). Type A behaviour pattern, extraversion, neuroticism and psychological distress. British Journal of Medical Psychology, 64, 73-83.
- Crean, P. C. & Fox, K.M. (1987). Exercise electrocardiography in coronary artery disease. Quarterly Journal of Medicine, New Series, 62(237), 7-13.
- Crews, D.J., & Landers, D.M. (1987). A meta-analytic review of aerobic fitness and reactivity to psychosocial stressors. Medicine and Science in Sports and Exercise, 19, 114-120.
- Crider, A.B., Goethals, G.R., Kavanagh, R.D., & Solomon, P.R. (1983). Psychology. Glenview: Scott, Foresman and Company.
- Cronbach, L.J. (1951). Coefficient alpha and the internal structure of tests. Psychometrika, 16, 297-334.
- Cronbach, L.J. (1970). Essentials of Psychological Testing (3rd ed.) New York: Harper & Row.
- Cudeck, R., & Brown, M.W. (1983). Cross-validation of covariance structures. Multivariate Behavioral Research, 1, 245-276.
- Cunningham, D.A. (1981). The relationship of exercise to the recurrence rate of myocardial infarction in men - Ontario Exercise-Heart Collaborative Study. Medicine and Science in Sports and Exercise, 13, 99.

Cupelli, V., Brettoni, M., Attina, D.A., Laverone, E., Arcangeli, G., Cupelli, G., Bucchino, G., Bini, G., & Giuliano, G. (1984). Cardiovascular response to maximal exercise in elderly healthy people. Journal of Sports Medicine and Physical Fitness, 24, 273-279.

Cureton, E.E., & D'Agostino, R.B. (1983). Factor analysis: An applied approach. Hillsdale: Lawrence Erlbaum.

Curfman, G.D. (1993). Is exercise beneficial - or hazardous - to your heart? New England Journal of Medicine, 329(23), 1730-1731.

Danilewitz, L. (1987). The family functioning patterns of coronary heart diseases patients. Unpublished masters dissertation, University of Pretoria, South Africa.

De Geus, E.J.C., Van Doornen, L.J.P., & Orlebeke, J.F. (1993). Regular exercise and aerobic fitness in relation to psychological make-up and physiological stress reactivity. Psychosomatic Medicine, 55, 347-363.

De Geus, E.J.C., Van Doornen, L.J.P., de Visser, D.C., & Orlebeke, J.F. (1990). Existing and training induced differences in aerobic fitness: Their relationship to physiological response patterns during different types of stress. Psychophysiology, 27(4), 457-478.

Delamater, R.J., & McNamara, J.R. (1987). Expression of anger: Its relationship to assertion and social desirability among college women. Psychological Reports, 61, 131-134.

Dembroski, T.M., & Costa, P.T. (1987). Coronary prone behavior: Components of the Type A pattern and hostility. Journal of Personality, 55(2), 211-233.

Dembrowski, T.M., MacDougall, J.M., Costa, P.T., & Grandits, G.A. (1989). Components of hostility as predictors of sudden death and myocardial infarction in the Multiple Risk Factor Intervention Trial. Psychosomatic Medicine, 51, 514-522.

Dembrowski, T.M., MacDougall, J.M., Shields, J.L., Pettito, J., & Lushene, R. (1978). Components of the Type A coronary-prone behavior pattern and cardio-vascular responses to psychomotor performance challenge. Journal of Behavioral Medicine, 1, 159-176.

Dembrowski, T.M., MacDougall, J.M., Williams, R.B., Haney, T.L., & Blumenthal, J.A. (1985). Components of Type A, hostility, and anger-in: Relationship to angiographic findings. Psychosomatic Medicine, 47(3), 219-233.

Denolin, H. (1985). Rehabilitation as part of comprehensive care. In V. Kallio & E. Cay (Eds.), Rehabilitation after myocardial infarction, the European experience. Copenhagen: WHO. p. 24.

Desharnais, R., Godin, G., Jobin, J., & Valois, P. (1990). Optimism and health-relevant cognitions after a myocardial infarction. Psychological Reports, 67, 1131-1135.

Desharnais, R., Jobin, J., Côté, C., Lévesque, L., & Godin, G. (1993). Aerobic exercise and the placebo effect: A controlled study. Psychosomatic Medicine, 55, 149-154.

Detre, K., Holubkov, R., Kelsey, S., Cowley, M., Kent, K., Williams, D., Myler, R., Faxon, D., Holmes, D., Bourassa, M., Block, P., Gosselin, A., Bentivoglio, L., Leatherman, L., Dorros, G., King, S., Galichia, J., Al-Bassam, M., Leon, M., Robertson, T., & Passamani, E. (1988). Percutaneous transluminal coronary angioplasty in 1985-1986 and 1977-1981. New England Journal of Medicine, 318(5), 265-270.

(DSM-III) Diagnostic and Statistical Manual of Mental Disorders (3rd ed.). (1980). Washington, D.C.: American Psychiatric Association.

(DSM-IV) Diagnostic and Statistical Manual of Mental Disorders (4th ed.). (1994). Washington, D.C.: American Psychiatric Association

Diamond, E.L. (1982). The role of anger and hostility in essential hypertension and coronary heart disease. Psychological Bulletin, *92*, 410-433.

Digenio, A.G., Sim, J.G.M., Krige, K., Stewart, A., Morris, R., Dowdeswell, R.J., & Padayachee, G.N. (1991). The Johannesburg cardiac rehabilitation programme. South African Medical Journal, *79*, 183-187.

Dillon, W.R., & Goldstein, M. (1984). Multivariate analysis: Methods and applications. New York: John Wiley & Sons.

Dimsdale, J.E. (1977). Emotional causes of sudden death. American Journal of Psychiatry, *134*, 1361-1366.

Dimsdale, J.E., Gilbert, J., Hutter, A.M., Hackett, T.P., & Block, P.C., (1981). Predicting cardiac morbidity based on risk factors and angiographic findings. American Journal of Cardiology, *47*, 73-76.

Dimsdale, J.E., Hackett, T.P., Block, P.C., & Hutter, A.M.. (1978). Emotional correlates of the Type A behavior pattern. Psychosomatic Medicine, *40*, 580-583.

Dimsdale, J.E., Hackett, T.P., Hutter, A.M., Block, P.C., Catanzano, D.M., & White, P.J. (1979). Type A behavior and angiographic findings. Journal of Psychosomatic Research, *23*, 273-276.

Dishman, R.K. (1986). Exercise compliance: A new view for public health. Physician and Sports Medicine, *14*, 127-145.

Dobson, C.B. (1982). Stress. The hidden adversary. Lancaster: MTP Press.

Dongier, M. (1974). Psychosomatic aspects in myocardial infarction in comparison with angina pectoris. Psychotherapeutics and Psychosomatics, *23*, 123-131.

Doyle Gentry, W., & Stewart, M.A. (1985). Psychological effects of exercise training in coronary-prone individuals and in patients with symptomatic coronary heart disease. In N.K. Wenger (Ed.), Exercise and the heart (2nd ed.). (pp. 255-260). Philadelphia: F.A. Davis.

Drever, J. (1958). A dictionary of psychology. Harmondsworth: Penguin.

Duckworth, D.H. (1985). Is the 'organizational stress' construct a red herring? A reply to Glowinkowski and Cooper. Bulletin of the British Psychological Society, *38*, 401-404.

Dunbar, F. (1943). Psychosomatic Diagnosis. New York: Hoeber Press.

Dyck, D.G., Moser, C.G., & Janisse, M.P. (1987). Type A behavior and situation specific perceptions of control. Psychological Reports, *60*, 991-999.

Eaker, E.D., Abbott, R.D., & Kannel, W.B. (1989). Frequency of uncomplicated angina pectoris in Type A Compared with Type B persons. American Journal of Cardiology, *63*, 1042-1045.

Eaker, E.D., & Castelli, W.P. (1988). Type A behavior and mortality from coronary disease in the Framingham study. New England Journal of Medicine, *319*, 1480-1481.

Edwards, J.R., & Baglioni, A.J., Jr. (1991). Relationship between Type A behavior pattern and mental and physical symptoms: A comparison of global and component measures. Journal of Applied Psychology, 76(2), 276-290.

Edwards, J.R., Baglioni, A.J.Jr., & Cooper, C.L. (1990a). Examining the relationships among self-report measures of the Type A behavior pattern: The effects of dimensionality, measurement error, and differences in underlying constructs. Journal of Applied Psychology, 75(4), 440-454.

Edwards, J.R., Baglioni, A.J.Jr., & Cooper, C.L. (1990b). The psychometric properties of the Bortner Type A Scale. British Journal of Psychology, 81, 315-333.

Edwards, J.R. & Cooper C. L. (1990). The person-environment fit approach to stress: Recurring problems and some suggested solutions. Journal of Organizational Behavior, 11, 293-307.

Eliot, R.A. (1979). Stress and the major cardiovascular disorders. Mount Kisko: Futura.

Endler, N.S., Cox, B.J., Parker, J.D.A., & Bagby, R.M. (1992). Self-reports of depression and state-trait anxiety: Evidence for differential assessment. Journal of Personality and Social Psychology, 63(5), 832-838.

Endler, N.S., & Parker, J.D.A. (1990). State and trait anxiety, depression and coping styles. Australian Journal of Psychology, 42(2), 207-220.

Engel, G.L. (1970). Sudden death and the "medical model" in psychiatry. Canadian Psychiatric Association Journal, 15, 527-538.

Engel, G.L. (1971). Sudden and rapid death during psychological stress. Folklore or folk wisdom? Annals of Internal Medicine, 129, 725-731.

Engelbreton, T. O., Matthews, K.A. (1992). Dimensions of hostility in men, women and boys: Relationships to personality and cardiovascular responses to stress. Psychosomatic Medicine, 54, 311-323.

Engelbreton, T. O., Matthews, K.A., & Scheier, M.F. (1989). Relations between anger expression and cardiovascular reactivity: Reconciling inconsistent findings through matching hypothesis. Journal of Personality and Social Psychology, 57(3), 513-521.

Erb, B.D., Fletcher, G.F., & Sheffield, T.L. (1979). Standards for cardiovascular exercise treatment programs. American Heart Association Subcommittee on Rehabilitation. Target Activity Group. Circulation, 59, 1084A-1090A.

Erdman, R.A.M. (1990). Myocardial infarction and cardiac rehabilitation. In A.A., Kaptein, H.M. van der Ploeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. Psychological treatment of somatic disorders (pp. 127-145). Chichester: John Wiley.

Erdman, R.A.M., Duivenvoorden, H.J., Verhage, F., Kazemier, M., & Hugenholtz, P.G. (1986). Predictability of beneficial effects in cardiac rehabilitation: A randomized clinical trial of psychosocial variables. Journal of Cardiopulmonary Rehabilitation, 6, 206-213.

Evans, P.D. (1990). Type A behaviour and coronary heart disease: When will the jury return? British Journal of Psychology, 81, 147-157.

Everaerd, W., Vromans, I.S.Y., & van der Elst, A.M.C. (1990). Asthma. In A.A., Kaptein, H.M. van der Ploeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. psychological treatment of somatic disorders (pp. 147-157). Chichester: John Wiley.

Eysenck, H.J., Fulker, D. (1983). The components of Type A behaviour and its genetic determinants. Personality and Individual Differences, 4, 499-505.

Falkner, B., Onesti, G., & Hamstra, B. (1981). Stress response characteristics of adolescents with high genetic risk for essential hypertension: A five year follow up. Clinical and Experimental Hypertension, 3, 583-591.

Fava, M., Littman, A., Lamoreaux, S., Milani, R., Shera, D., MacLaughlin, R., Cassem, E., Leaf, A., Marchio, B., Bolognesi, E., & Guaraldi, G. (1992). Psychological, behavioral and biochemical risk factors for coronary artery disease among American and Italian male corporate managers. American Journal of Cardiology, 70, 1412-1416.

Fazio, R.H., Cooper, M., Dayson, K., & Johnson, M. (1981). Control and the coronary-prone behavior pattern A: Responses to multiple situation demands. Personality and Social Psychology Bulletin, 7, 97-102.

Feather, N.T., & Volkmer, R.E. (1991). Task preference in relation to achievement striving and impatience-irritability components of Type A behavior. Australian Journal of Psychology, 43, 23-27.

Feather, N.T., & Volkmer, R.E. (1988). Preference for situations involving effort, time pressure, and feedback in relation to Type A behavior, locus of control, and test anxiety. Journal of Personality and Social Psychology, 55, 266-271.

Feldt, L.S., & Brennan, R.L. (1989). Reliability. In R.L. Linn (Ed.). Educational measurement (3rd ed.). (pp. 105-146). London: Collier MacMillan.

Finman, R., & Berkowitz, L. (1989). Some factors influencing the effect of depressed mood on anger and overt hostility toward another. Journal of Research on Personality, 23, 70-84.

Finnegan, D.L., & Suler, J.R. (1985). Psychological factors associated with maintenance of improved health behaviors in postcoronary patients. Journal of Psychology, 119(1), 87-94.

Fishman-Alschuler, C., & Alschuler, A.S. (1984). Developing healthy responses to anger: The counselors role. Journal of Counseling and Development, 63, 26-29.

Fletcher, G.F., Blair, S.N., Blumenthal, J., Caspersen, C., Chaitman, B., Epstein, S., Falls, H., Sivarajan Froelicher, E.S., Froelicher, V.F., & Pina, I.L. (1992). Statement on exercise. Benefits and recommendations for physical activity programs for all Americans. A statement for health professionals by the committee on exercise and cardiac rehabilitation of the council on clinical cardiology, American Heart Association. Circulation, 86(1), 340-344.

Folkins, C.H. (1976). Effects of physical training on mood. Journal of Clinical Psychology, 32(2), 385-388.

Folkins, C.H., & Amsterdam, E.A. (1977). Control and modification of stress emotions through chronic exercise. In F.A. Amsterdam, J.H. Wilmore, & A.N. DeMaria (Eds.), Exercise and cardiovascular health and disease. New York: Yorke.

Folkins, C.H., & Sime, W.E. (1981). Physical fitness training and mental health. American Psychologist, 36(4), 373-389.

Folkman, S., & Lazrus, R.S. (1986). Stress processes and depressive symptomatology. Journal of Abnormal Psychology, 95, 107-113.

- Follick, M.J., Ahern, D.K., Gorkin, L., Niaura, R., Herd, J.A., Ewart, C., Schron, E.B., Kornfeld, D.S., & Capone, R.J. (1990). The relationship of psychosocial and stress reactivity variables to ventricular arrhythmias in the cardiac arrhythmia pilot study (CAPS). American Journal of Cardiology, 66, 63-67.
- Fontana, A.F., Kerns, R.D., Rosenberg, R.L., Marcus, J.L., & Colonese, K.L. (1986). Exercise training for cardiac patients: adherence, fitness, and benefits. Journal of Cardiopulmonary Rehabilitation, 6, 4-15.
- Forgays, D.K. (1992). Type A behavior and parenting stress in mothers with young children. Current Psychology: Research & Reviews, 11(1), 3-19
- Foulds, G.A. (1959). The relative stability of personality measures compared with diagnostic measures. Journal of Mental Science, 105, 783-787.
- Foulds, G.A. (1965). Personality and personal illness. London: Tavistock.
- Foulds, G.A., Caine, T.M., & Creasy, M.A. (1960). Aspects of extra and intro-punitive expression in mental illness. Journal of Mental Sciences, 106, 599-609.
- Francis, K.T. (1981). Perceptions of anxiety, hostility, and depression in subjects exhibiting the coronary-prone behavior pattern. Psychiatric Research, 16(3), 183-190.
- French, J.R.P., & Caplan, R.D. (1972). Organizational stress and individual strain. In A.G. Marrow (Ed.), The failure of success (pp. 31-66). New York: AMACOM.
- Friedberg, C.K. (1966). Diseases of the heart (3rd ed.). Philadelphia: W.B. Saunders.
- Friedman, H.S. (1989). The role of emotional expression in coronary heart disease. In A.W. Siegman & T.M. Dembroski (Eds.), In search of coronary-prone behavior: Beyond Type A. (pp. 149-168). Hillsdale: Erlbaum.
- Friedman, H.S., & Booth-Kewley, S. (1987a). Personality, Type A behavior, and coronary heart disease: The role of emotional expression. Journal of Personality and Social Psychology, 53(4), 783-792.
- Friedman, H.S., & Booth-Kewley, S. (1987b). The disease prone personality. American Psychologist, 42(6), 539-555.
- Friedman, H.S., & Booth-Kewley, S. (1988). Validity of the Type A construct: A reprise. Psychological Bulletin, 104(2), 381-384.
- Friedman, M., & Rosenman, R.H. (1959). Association of Specific Overt Behaviour Pattern with Blood and Cardiovascular Findings. Journal of the American Medical Association, 169, 1286-1296.
- Friedman, M., & Rosenman, R.H. (1974). Type A behavior and your heart. New York: Knopf.
- Friedman, M., Rosenman, R.H., & Carrol, V. (1958). Changes in the serum cholesterol and blood clotting time in men subjected to cyclic variation of occupational stress. Circulation, 17, 852-861.
- Friedman, M., St. George, S., Byers, S.O., & Rosenman, R.H. (1960). Excretion of catecholamines, 17-ketosteroids, 17-hydroxycorticoids, and 5-hydroxyindole in men exhibiting a particular behavior pattern (A) associated with high incidence of clinical coronary artery disease. Journal of Clinical Investigation, 39, 758.

Friedman, M., Thoreson, C., Gill, J., Ulmer, D., Powell, L., Price, V., Brown, B., Thompson, L., Rabin, D., Bressi, W., Bousg, E., Levy, R., & Dixon, T. (1986). Alteration of Type A behavior and its effects on cardiac recurrences in post myocardial infarction patients: Summary results of the recurrent coronary prevention project. American Heart Journal, 112, 653-665.

Froelicher, V.F., Myers, J., Follansbee, W.P., & Labovitz, A.J. (1993). Exercise and the heart (3rd ed.). St. Louis: Mosby.

Gullager, C. (1996). A factor analytic study on the validity of a union commitment scale. Journal of Applied Psychology, 17(1), 129-136.

Funkhstein, D., King, S.H., & Drolette, M. (1954). The direction of anger during a laboratory stress-inducing situation. Psychosomatic Medicine, 16, 404-413.

Furnham, A. (1990a). The Protestant work ethic and Type A behaviour: A pilot study. Psychological Reports, 66, 323-328.

Furnham, A. (1990b). The Type A behavior pattern and the perception of self. Personality and Individual Differences, 11(8), 841-851.

Furnham, A., Burvey, A., & Henley, S. (1986). Type A behavior pattern, the recall of personality information, and self-evaluation. British Journal of Medical Psychology, 59, 353-374.

Furnham, A., & Linfoot, J. (1987). The Type A behavior pattern and the need to prove oneself: A conditional study. Current Psychological Research and Reviews, 6(2), 125-135.

Fuster, V. (1993). Coronary thrombolysis - A perspective for the practicing physician. New England Journal of Medicine, 329(10), 723-725.

Gallagher, D., Nies, G., & Thompson, L.W. (1982). Reliability of the BDI with older adults. Journal of Consulting and Clinical Psychology, 50, 152-153.

Gallant, L.D., & Chrisjohn, R.D. (1985). The Jenkins Activity Survey: A valid measure of Type A behavior? University of Western Ontario. London: Ontario.

Gamble, G.O., & Matteson, M.T. (1992). Type A behavior, job satisfaction, and stress among black professionals. Psychological Reports, 70, 43-50.

Ganellen, R.J., & Blaney, P.H. (1984). Hardiness and social support as moderators of the effects of life stress. Journal of Personality and Social Psychology, 71(1), 102-110.

Ganster, D.C., Mayes, B.T., Sime, W.E., & Tharp, G.D. (1982). Managing organizational stress: A field experiment. Journal of Applied Psychology, 67(5), 533-542.

Ganster, D.C., Schaubroeck, J., Sime, W.E., & Mayes, B.T. (1991). The nomological validity of the Type A personality among employed adults. Journal of Applied Psychology, 76(1), 143-168.

Garrity, T., & Klein, R. (1975). Emotional response and clinical severity as early determinants of six-month mortality after myocardial infarction. Heart and Lung, 4(5), 730-737.

Gastorf, J.W., Suls, J., Sanders, G.S. (1980). Type A coronary-prone behavior pattern and social facilitation. Journal of Personality and Social Psychology, 38(5), 773-780.

- Gatewood-Colwell, G., Kaczmarek, M., & Ames, M.H. (1989). Reliability and validity of the Beck Depression Inventory for a white and Mexican-American gerontic population. Psychological Reports, *65*, 1163-1166.
- Gentry, W.D. (1985). Relationship of anger-coping styles and blood pressure among black Americans. In M.A. Chesney & R.H. Rosenman (Eds.), Anger and hostility in cardiovascular and behavioral disorders. New York: McGraw-Hill.
- Gentry, W.D., Balder, L., Oude-Weme, J.D., Musch, F., & Gary, H.E. (1983). Type A/B differences in coping with acute myocardial infarction: Further considerations. Heart and Lung, *12*, 212-214.
- Gentry, W.D., Chesney, A.P., Gary, H.G., Hall, R.P., & Harburg, E. (1982). Habitual anger-coping styles. I. Effect on mean blood pressure and risk for essential hypertension. Psychosomatic Medicine, *44*, 273-281.
- Gerbing, D.W., & Anderson, J.C. (1988). An updated paradigm for scale development incorporating unidimensionality and its assessment. Journal of Marketing Research, *25*, 186-192.
- Gerin, W., Pieper, C., Levy, R., & Pickering, T.G. (1992). Social support in social interaction: A moderator of cardiovascular reactivity. Psychosomatic Medicine, *54*, 324-336.
- Gerahon, S., Rosenbaum, J.F., Reynolds, C.F., Perel, J.M., Barlow, D.H., Kupfer, D.J., & Mann, J.J. (1992). Mixed anxiety-depression - a new diagnostic category for DSM-IV? Treatment implications. Journal of Clinical Psychiatry, *52*(10), 441-442.
- Giese, H., & Schomer, H.H. (1986). Life-style changes and mood profile of cardiac patients after an exercise rehabilitation program. Journal of Cardiopulmonary Rehabilitation, *6*, 30-37.
- Gilbert, D.G., & McArthur, S.F. (1988). Heart rate responses to quantified doses of nicotine as a function of anger expression, Type A behavior, and nicotine tolerance. Psychophysiology, *25*(4), 450.
- Gilbert, P. (1991). Essay review. Depression: New books and conceptualizations of an old problem. British Journal of Psychology, *82*, 119-126.
- Gilmer, J.S., & Feldt, L.S. (1983). Reliability estimation for a test with parts of unknown lengths. Psychometrika, *48*(1), 99-111.
- Giubbini, R., Galli, M., Campini, R., Bosimini, E., Becivelli, W., & Tavazzi, L. (1991). Effects of mental stress on myocardial perfusion in patients with ischaemic heart disease. Circulation, *83*(4), II-100-II-107.
- Glass, D.C. (1977a). Behavior patterns, stress, and coronary disease. Hillsdale: Erlbaum.
- Glass, D.C. (1977b). Stress, behavior patterns, and coronary disease. American Scientist, *65*, 177-187.
- Glass, D.C., & Carver, C.S. (1980). Helplessness and the coronary-prone personality. In J. Garber & M.E. Seligman (Eds.), Human helplessness: theory and applications (p. 232) New York: Academic.
- Godaert, G. (1990). Hypertension. In A.A., Kaptein, H.M. van der Floeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. psychological treatment of somatic disorders (pp. 109-126). Chichester: John Wiley.

Goldberg, B., & Folkins, C.H. (1974). Relationship of body-image to negative emotional attitudes. Perceptual and Motor Skills, 39, 1053-1054.

Goldberg, P. (1978). Executive health. New York: McGraw-Hill.

Gordon, N. & Gibbons, L. (1991). The complete heart recovery guide. Cape Town: Oxford University Press.

Gordon, T. & Kannel, W.B. (1971). Premature mortality from coronary heart disease: The Framingham study. Journal of the American Medical Association, 215, 1617-1625.

Gorsuch, R.L. (1983). Factor analysis. Hillsdale: Lawrence Erlbaum.

Gothb, I.H. (1984). Depression and general psychopathology in university students. Journal of Abnormal Psychology, 93, 19-30.

Gouldner, A.W. (1960). The norm of reciprocity: A preliminary statement. American Sociological Review, 25, 161-178.

Greenglass, E.R., & Julkunen, J. (1991). Cook-Medley Hostility, anger, and the Type A behavior pattern in Finland. Psychological Reports, 68, 1059-1066.

Greenland, P., & Briody, M.E. (1984). Rehabilitation of the MI survivor. Cardiac Rehabilitation, 75(1), 79-96.

Greenland, P., Reicher-Reiss, H., Goldbourt, U., Behar, S. (1991). In-hospital and 1-year mortality in 1,524 women after myocardial infarction: comparison with 4,315 men. Circulation, 83, 484-491.

Greer, S., Ramsay, I., Bagley, C. (1973). Neurotic and thyrotoxic anxiety: Clinical, psychological and physiological measurements. British Journal of Psychiatry, 122, 549-554.

Grodzinski, E., Jetté, M., Blümchen, G., & Borer, J.S. (1987). Effects of a four-week training program on left ventricular function as assessed by radionuclide ventriculography. Journal of Cardiopulmonary Rehabilitation, 7, 517-524.

Groebner, D.F., & Shannon, P.W. (1985). Business Statistics. A decision-making approach (2nd Ed.). Columbus, Ohio: A Bell & Howell.

Groenman, N.H., Vlaeyen, J.W.S., van Eek, & Schuerman, J.A. (1990). Chronic pain. In A.A., Kaptein, H.M. van der Ploeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. psychological treatment of somatic disorders (pp. 51-66). Chichester: John Wiley.

Grossarth-Maticek, R. & Eysenck, J. (1990). Personality, stress and disease: Description and validation of a new inventory. Psychological Reports, 66, 355-373.

Grossarth-Maticek, R. & Eysenck, J. (1991). Personality, stress and motivational factors in drinking as determinants of risk for cancer and coronary heart disease. Psychological Reports, 59, 1027-1043.

Guadagnoli, E., & Velicer, W.F. (1988). Relation of sample size to the stability of component patterns. Psychological Bulletin, 103(2), 265-275.

Guadagnoli, E., & Velicer, W.F. (1991). A comparison of pattern matching indices. Multivariate Behavioral Research, 26, 323-343.

- Guiry, E., Conroy, R.M., Hickey, N., & Mulcahy, R. (1987). Psychological response to an acute coronary event and its effect on subsequent rehabilitation and lifestyle change. Clinical Cardiology, 10, 256-260.
- Gundle, M., Bozman, R., Tate, S., Raft, D., McLaurin, L. (1980). Psychosocial outcome after coronary artery surgery. American Journal of Psychiatry, 137, 1591-1594.
- Gut, E. (1989). Productive and unproductive depression: Success and failure of a vital process. London: Routledge.
- Guyatt, G., Walter, S., & Norman, G. (1987). Measuring change over time: assessing the usefulness of evaluative instruments. Journal of Chronic Diseases, 40(2), 171-178.
- Hackett, T.P. (1985). Depression following myocardial infarction. Psychosomatics, 26, 23-30.
- Hackett, T.P., & Cassem, N.H. (1975). Coronary care: patient psychology. Dallas: American Heart Association.
- Hackett, T.P., & Cassem, N.H. (1978). Psychologic aspects of rehabilitation after myocardial infarction. In N.K. Wenger, & H.K. Hellerstein (Eds.), Rehabilitation of the coronary patient. New York: John Wiley.
- Hackett, T.P., & Cassem, N.H. (1982). Coping with cardiac disease. Advanced Cardiology, 31, 212-217.
- Hafner, R.J. (1974). Relationships between personality and autonomic nervous reactions to stress. Journal of Psychosomatic Research, 18, 181-185.
- Haft, J.I. (1974). Cardiovascular injury induced by sympathetic catecholamines. Progress in Cardiovascular Diseases, 17, 73-86.
- Hakstian, A.R., Rogers, W.T., & Cattell, R.B. (1982). The behavior of number-of-factors rules with simulated data. Journal of Multivariate Behavioral Research, 17, 193-219.
- Hammon, H.K. (1985). Exercise for coronary heart disease patients: Is it worth the effort? Journal of Cardiopulmonary Rehabilitation, 5, 531-539.
- Hansson R.O., Hogan, R., Johnson, J.A., & Schroeder, D. (1983). Disentangling Type A behavior: The role of ambition, insensitivity, and anxiety. Journal of Research in Personality, 17, 186-197.
- Harburg, E., Gleiberman, L., Russell, M., & Cooper, M.L. (1991). Anger-coping styles and blood pressure in black and white males: Buffalo, New York. Psychosomatic Medicine, 53, 153-164.
- Hardy, J.H., & Smith, T.W. (1988). Cynical hostility and vulnerability to disease: Social support, life stress, and physiological response to conflict. Health Psychology, 7, 447-459.
- Horn, H.H. (1976). Modern factor analysis (2nd Ed.). Chicago: University of Chicago Press.
- Harris, C.W. (1967). On factors and factor scores. Psychometrika, 32, 363-379.
- Hart, K.E. (1984). Anxiety management training and anger control for Type A individuals. Journal of Behavioral Therapy and Experimental Psychiatry, 15(2), 133-139.

Hartley, L.H., Foreyt, J.P., Alderman, M.H., Chesney, M.A., Friedman, M., Hendrix, G.H., Herd, J.A., Levenson, R.M., Ruberman, W., & Thoreson, C.E. (1987). Task Force 6: Secondary prevention of coronary artery disease. Circulation, 76(1), 168-173.

Haskell, W.L. (1978). Cardiovascular complications during exercise training of cardiac patients. Circulation, 57(5), 920-924.

Hassmén, P., Stahl, R., & Borg, G. (1993). Psychophysiological responses to exercise in Type A/B men. Psychosomatic Medicine, 55, 178-184.

Hatzenbuehler, L.C., Parpai, M., & Matthews, L. (1983). Classifying college students as depressed or nondepressed using the Beck Depression Inventory: An empirical analysis. Journal of Consulting and Clinical Psychology, 51, 360-366.

Haynes, S.G., Feinleib, M., & Kannel, W.B. (1980). The relationship of psychosocial factors to coronary heart disease in the Framingham Study: III. Eight-year incidence of coronary heart disease. American Journal of Epidemiology, 111, 37-58.

Haynes, S.G., Feinleib, M., Levine, S., Scotch, N., & Kannel, W.B. (1978). The relationship of psychosocial factors to coronary heart disease in the Framingham Study. II. Prevalence of coronary heart disease. American Journal of Epidemiology, 107(5), 384-402.

Haynes, S.G., Levine, S., Scotch, N., Feinleib, M., & Kannel, W.B. (1978). The relationship of psychosocial factors to coronary heart disease in the Framingham Study. I. Methods and risk factors. American Journal of Epidemiology, 107(5), 362-383.

Haynes, S.G. & Matthews, K.A. (1988). Coronary-prone behavior: Continuing evolution of the concept. Annals of Behavioral Medicine, 10(2), 47-59.

Heilbrun, A.B., Palchanis, N., & Friedberg, E. (1986). Self-report measurement of Type A behavior: Toward refinement and improved prediction. Journal of Personality Assessment, 50(4), 525-539.

Heller, S., Frank, K., & Kornfeld, D. (1974). Psychological outcome following open-heart surgery. Archives of Internal Medicine, 134, 108-114.

Hellerstein, H.K., & Franklin, B.A. (1984). Exercise testing and prescription. In N.K. Wenger & H.K. Hellerstein (Eds.), Rehabilitation of the coronary patient (2nd ed.). (pp. 197-284). New York: Wiley.

Helmreich, R.L., Spence, J.T., & Pred, R.S. (1988). Making It without losing it: Type A, achievement motivation, and scientific attainment revisited. Personality and Social Psychology Bulletin, 14(3), 495-504.

Henderson, S., Davidson, J.A., Lewis, I.C., Gillard, H.N., & Baikie, A.G. (1977). An assessment of hostility in a population of adolescents. Archives of General Psychiatry, 34, 706-711.

Henley, S., & Furnham, A. (1989). The Type A behavior pattern and self-evaluation. British Journal of Medical Psychology, 62, 51-59.

Hickson, R.C., Hagberg, J.M., Ehsani, A.A., & Holloszy, J.O. (1981). Time course of the adaptive responses of aerobic power and heart rate to training. Medical Science, Sports and Exercise, 13, 17-20.

Hildebrand-Saints, L., & Weary, G. (1989). Depression and social information gathering. Personality and Social Psychology Bulletin, 15(2), 150-160.

- Hinkin, T.R., & Schriesheim, C.A. (1989). Development and application of new scales to measure the French and Raven (1959) bases of social power. Journal of Applied Psychology, 74(4), 561-567.
- Hinton, J.W., Rotheiler, E., & Howard, A. (1991). Confusion between stress and state anxiety in a much used self-report 'stress' inventory. Personality and Individual Differences, 12(1), 91-94.
- Hlatky, M.A., Haney, T.L., Barefoot, J.C., Califf, R.M., Mark, D.B., Pryor, D.B., & Williams, R.B. (1980). Medical, psychological and social correlates of work disability among men with coronary artery disease. American Journal of Cardiology, 58(10), 911-915.
- Hogan, R., & Nicholson, R.A. (1988). The meaning of personality test scores. American Psychologist, 43(8), 621-626.
- Holden, R.R., & Hickman, D. (1987). Computerized versus standard administration of the JAS (Form T). Journal of Human Stress, 13, 175-179.
- Holloszy, J.O. (1983). Exercise, health, and aging: A need for more information. Medicine and Science in Sports and Exercise, 15(1), 1-5.
- Holmes D.S., & Roth, D.L. (1985). Association of aerobic fitness with pulse rate and subjective responses to psychological stress. Psychophysiology, 22(5), 525-529.
- Holmes T.H., & Masuda, M. (1974). Life change and illness susceptibility. In B.S. Dohrenwend & B.P. Dohrenwend (Eds.), Stressful life events: Their nature and effects (pp. 45-72). New York: John Wiley & Sons.
- Hopkins, P.N. & Williams, R.R. (1981). A survey of 246 suggested coronary risk factors. Atherosclerosis, 40, 1-52.
- Hossack K.F., & Hartwig, R. (1982). Cardiac arrest associated with supervised cardiac rehabilitation. Journal of Cardiac Rehabilitation, 2, 402-408.
- Hotelling, H. (1933). Analysis of a complex of statistical variables into principle components. Journal of Educational Psychology, 24, 417-441, 498-520.
- House, J.S. (1974). Occupational stress and coronary heart disease: A review and theoretical integration. Journal of health and Social Behavior, 15, 12-27.
- Houston, B.K. (1983). Physiological responsivity and the Type A behavior pattern. Journal of Research in Personality, 17, 22-39.
- Houston, B.K. (1986). Psychological variables and cardiovascular and neuroendocrine reactivity. In K.A. Matthews, S.M. Weiss, T. Detre, T.M. Dembroski, B. Falkner, S.B. Manuck, & R.B. Williams (Eds.), Handbook of stress, reactivity, and cardiovascular disease (pp. 220-230). New York: Wiley.
- Houston, B.K., Smith, T.W., & Zurawski, R.M. (1986). Principle dimensions of the Framingham Type A Scale: Differential relationships to cardiovascular reactivity and anxiety. Journal of Human Stress, 12, 105-112.
- Houston, B.K., & Vavak, C.R. (1991). Cynical hostility: Developmental factors, psychosocial correlates, and health behaviors. Health Psychology, 10(1), 9-17.
- Howard, J.H., Cunningham, D.A., & Rechnitzer, P.A. (1976). Health Patterns associated with Type A behavior: A managerial population. Journal of Human Stress, 2, 24-31.

- Hubbard, S.G., & Ferguson, G. (1992). Recovering from coronary bypass surgery. New York: Harper.
- Huck, S.W., & McClean, R.A. (1975). Using a repeated measures ANOVA to analyze data from a pretest-posttest design: A potentially confusing task. Psychological Bulletin, 82, 511-518.
- Hughes, J.R. (1984). Psychological effects of habitual aerobic exercise: A critical review. Preventative Medicine, 13, 66-78.
- Ismail, A.H., & Young, R.J. (1977). Effect of chronic exercise on the personality of adults. Annals of the New York Academy of Sciences, 301, 958-969.
- Jaeger, E. (1990). EOS Sprint: Instruction Manual (Version 4.0). Würzburg, Germany: Erich Jaeger GmbH & CoKG.
- Jamal, M. (1990). Relationship of job stress and Type - A behavior to employees' job satisfaction, organizational commitment, psychosomatic health problems, and turnover motivation. Human Relations, 43(8), 727-738.
- Jamal, M., & Baba, V.V. (1991). Type A behavior, its prevalence and consequences among women nurses: an empirical examination. Human Relations, 44(11), 1213-1228.
- James, L.R., & James, L.P. (1989). Causal modelling in organizational research. In C.L. Cooper, & I. Robertson (Eds.) International Review of Industrial and Organizational Psychology. London: John Wiley & Sons.
- Jamieson, J.L. & Lavoie, N.F. (1987). Type A behavior, aerobic power, and cardiovascular recovery from a psychosocial stressor. Health Psychology, 6(4), 361-371.
- Janisse, M.P., & Dyck, D.G. (1989). The Type A behavior pattern and coronary heart disease: physiological and psychological dimensions. In M.P. Janisse (Ed.), Individual differences, stress, and health psychology (pp. 57-71). New York: Springer-Verlag.
- Janisse, M.P., Edguer, N., & Dyck, D.G. (1986). Type A behavior, anger expression and reactions to anger imagery. Motivation and Emotion, 8, 373-388.
- Järvikoski, A., & Härkäpää, K. (1988). Type A behaviour and life-events. British Journal of Medical Psychology, 61, 353-363.
- Jenkins, C.D. (1976). Recent evidence supporting Psychological and social risk factors for coronary disease I-II. New England Journal of Medicine, 294, 987-994, & 1033-1038.
- Jenkins, C.D. (1988). Epidemiology of cardiovascular disease. Journal of Consulting and Clinical Psychology, 56(3), 324-332.
- Jenkins, C.D., Rosenman, R.H., & Friedman, M. (1967). Development of an objective psychological test for the determination of the coronary-prone behavior pattern in employed men. Journal of Chronic Diseases, 20, 371-379.
- Jenkins, C.D., & Zyzanski, S.J. (1975). Short Type A scale from the Activity Survey, Form N. (Unpublished manuscript).
- Jenkins, C.D., Zyzanski, S.J., & Rosenman, R.H. (1971). Progress toward validation of a computer-scored test for the Type A coronary prone behavior pattern. Psychosomatic Medicine, 33, 193-202.
- Jennings, J.R. (1984). Cardiovascular reactions and impatience in Type A and B college students. Psychosomatic Medicine, 46(5), 424-439.

- Johnson, E.H. (1984). Anger and anxiety as determinants of elevated blood pressure in adolescents. Unpublished doctoral dissertation, University of South Florida, Tampa.
- Johnson, E.H., & Broman, C.L. (1987). The relationship of anger expression to health problems among black Americans in a national survey. Journal of Behavioral Medicine, 10(2), 103-116.
- Johnson, H.J., & Sarason, I.G. (1978). Life stress, depression and anxiety: Internal-external control as a moderator variable. Journal of Psychosomatic Research, 22, 205-208.
- Johnson, J.L., & Morse, J.M. (1990). Regaining control: The process of adjustment after myocardial infarction. Heart and Lung, 19(2), 126-135.
- Johnston, M. (1988). Impending surgery. In S. Fisher & J. Reason (Eds.), Handbook of life stress, cognition and health. Chichester: John Wiley & Sons.
- Johnston, M., Weinman, J., & Marteau, T.M. (1990). Health psychology in hospital settings. In A.A., Kaptein, H.M. van der Ploeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. psychological treatment of somatic disorders (pp. 15-32). Chichester: John Wiley.
- Jolliffe, I.T. (1986). Principal component analysis. New York: Springer-Verlag.
- Jones, N.L., & Campbell, E.J.M. (1982). Clinical exercise testing (2nd ed.). Philadelphia: W.B. Saunders Company.
- Joreskog, K.G., & Sorbom, D. (1986). Lisrel VI. Chicago: National Education Resources.
- Julius, M., Harburg, E., Cottingham, E.M., & Johnson, E.H. (1986). Anger-coping types, blood pressure, and all cause mortality: A follow-up in Tecumseh, Michigan (1971-1983). American Journal of Epidemiology, 124(2), 220-233.
- Kahn, S. (1986). The role of anger in managerial effectiveness. Unpublished masters thesis, University of Cape Town.
- Kaiser, H.F. (1960). Relating factors between studies based on individuals. Unpublished manuscript. University of Illinois.
- Kaiser, H.F. (1970). A second-generation Little Jiffy. Psychometrika, 35(4), 401-415.
- Kaiser, H.F., & Rice J. (1974). Little Jiffy, Mark IV. Educational and Psychological Measurement, 34, 111-117.
- Kallio, V., Hämäläinen, H., Hakila, J., & Luurila, O.J. (1979). Reduction in sudden deaths by multifactorial intervention programme after acute myocardial infarction. Lancet, 2, 1092-1094.
- Kamarck, T.W., & Jennings, J.R. (1991). Biobehavioral factors in sudden cardiac death. Psychological Bulletin, 109(1), 42-75.
- Kamarck, T.W., Manuck, S.B., & Jennings, J.R. (1990). Social support reduces cardiovascular reactivity to psychological challenge: A laboratory model. Psychosomatic Medicine, 52, 42-58.
- Kannel, W.B. (1982). Exercise and sudden death. Journal of the American Medical Association, 248, 3143-3144.

- Kannel, W.B., & Gordon, T. (1974). The Framingham study. Washington D.C.: U.S. Government Printing Office.
- Kaptein, A.A. (1990). Skin disorders. In A.A., Kaptein, H.M. van der Ploeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. psychological treatment of somatic disorders (pp. 217-230). Chichester: John Wiley.
- Kasl, S.V. (1983). Pursuing the link between stressful life experiences and disease: A time for reappraisal. In C.L. Cooper, (Ed.) Stress research (pp. 79-102). New York: John Wiley.
- Kass, G.V. (1975). Significance testing in, and some extensions of, automatic interaction deletion. Unpublished master's thesis, University of the Witwatersrand, South Africa.
- Kauth, M.R. & Zettle, R.D. (1990). Validation of depression measures in adolescent populations. Journal of Clinical Psychology, 46(3), 291-295.
- Kavanagh, T., Shephard, R.J., Doney, H., & Pandit, V. (1973). Intensive exercise in coronary rehabilitation. Medical Science and Sports, 5, 34.
- Kavanagh, T., Shephard, R.J., & Kennedy, J. (1977). Characteristics of post-coronary marathon runners. Annals of the New York Academy of Science, 301(656), 455-465.
- Kavanagh, T., Shephard, R.J., Pandit, V., & Doney, H. (1970). Exercise and hypnotherapy in the rehabilitation of the coronary patient. Archives of Physical Medicine and Rehabilitation, 51, 578-587.
- Kavanagh, T., Shephard, R.J., & Tuck, J.A. (1975). Depression after myocardial infarction. Canadian Medical Association Journal, 113, 23-27.
- Kavanagh, T., Shephard, R.J., Tuck, J.A., & Qureshi, S. (1977). Depression following myocardial infarction: The effects of distance running. Annals of the New York Academy of Sciences, 301, 1029-1038.
- Keil, J.E., Sutherland, S.E., Knapp, R.G., Lackland, D.T., Gazes, P.C., & Tyroler, H.A. (1993). Mortality rates and risk factors for coronary disease in black as compared with white men and women. New England Journal of Medicine, 329(2), 73-78.
- Kellerman, J.J. (1973). Physical conditioning in patients after myocardial infarction. Results of a comparative study and 9 years follow-up. Schweizerische Medizinische Wochenschrift, 103, 79-85.
- Kellerman, J.J. (1981). Cardiac rehabilitation: Reminiscences, international variations, and experiences. Journal of Cardiac Rehabilitation, 1, 43-50.
- Keltikangas-Järvinen, L. (1990). Alexithymia and Type A behavior compared in psychodynamic terms of personality. British Journal of Medical Psychology, 63, 131-135.
- Kempe, C. (1945). Rorschach method and psychosomatic diagnosis: Personality traits of patients with rheumatic disease, hypertensive cardiovascular disease, coronary occlusion, and fracture. Psychosomatic Medicine, 7, 85-88.
- Kennedy, H.G. (1992). Anger and irritability. British Journal of Psychiatry, 161, 145-153.
- Kent, K.M. (1987). Coronary Angioplasty. A Decade of experience. New England Journal of Medicine, 316(18), 1148-1149.
- Kerlinger, F.N. (1964). Foundations of behavioural research. New York: Holt, Rinehart and Winston.

- Kerlinger, F.N. (1986). Foundations of behavioural research (3rd. ed.). Orlando: Holt, Rinehart & Winston.
- Kerlinger, F.N., & Pedhazur, E. (1973). Multiple regression in behavioral research. New York: Holt Rinehart & Winston.
- Kernis, M.H., Grannemann, & Barclay, L.C. (1989). Stability and level of self-esteem as predictors of anger arousal and hostility. Journal of Personality and Social Psychology, 56(6), 1013-1022.
- Kim, J.O. & Mueller, C.W. (1978). Factor analysis. statistical methods and practical issues. Beverly Hills: Sage Publications.
- Kinnaird, L.S., Yoham, M.A.S., & Kieval, Y.M. (1982). Patient compliance in rehabilitation programs. Nursing Clinics of North America, 17, 523-532.
- Klein, D.F. (1981). Anxiety reconceptualized. In D.F. Klein & J.G. Rabkin (Eds.), Anxiety: New research and changing concepts (pp. 235-263). New York: Raven Press.
- Kleinbaum, D.G., & Kupper, L.L. (1978). Applied regression analysis and other multivariable methods. North Scituate, Massachusetts: Duxbury Press.
- Kliewer, W. (1991). Coping in middle childhood: Relations to competence, Type A behavior, monitoring, blunting, and locus of control. Developmental Psychology, 27(4), 689-697.
- Kliewer, W., & Weidner, G. (1987). Type A behavior and aspirations: A study of parents' and children's goal setting. Developmental Psychology, 23(2), 204-209.
- Kligfield, P. & Okin, P.M. (1994). Evolution of exercise electrocardiogram. American Journal of Cardiology, 73, 1209-1210.
- Kobasa, S., Maddi, S.R., & Courington, S. (1981). Personality and constitution as mediators in the stress-illness relationship. Journal of Health and Social Behavior, 22, 368-378.
- Kobasa, S., Maddi, S.R., & Zola, M.L. (1983). Type A and hardiness. Journal of Behavioral Medicine, 6, 41-51.
- Kopac, C.A. (1990). The environment and its relationship to the Type A behavior pattern in middle and old age, a pilot study. Psychological Reports, 66, 768-770.
- Koskenvuo, M., Kaprio, J., Rose, R.J., Kesäniemi, A., Sarna, S., Heikkilä, K., & Langinvainio, H. (1988). Hostility as a risk factor for mortality and ischaemic heart disease in men. Psychosomatic Medicine, 50, 330-340.
- Krantz, D.S., Arabian, J.M., Davis, J.E., & Parker, J.S. (1982). Type A behavior and coronary artery bypass surgery: Intraoperative blood pressure and preoperative complication. Psychosomatic Medicine, 44, 273-294.
- Krantz, D.S., Grunberg, N.E., & Baum, A. (1985). Health psychology. Annual Review of Psychology, 36, 349-383.
- Krantz, D.S., & Manuck, S.B. (1984). Acute psychophysiological reactivity and risk of cardiovascular disease: A review and methodologic critique. Psychological Bulletin, 96, 435-464.
- Kristof, W. (1974). Estimation of reliability and true score variance from a split of a test into three arbitrary parts. Psychometrika, 39, 491-499.

- Kuller, L.H. (1976). Epidemiology of cardiovascular disease: current perspectives. American Journal of Epidemiology, 104, 425-456.
- Lakka, T.A., Venäläinen, J.M., Rauramaa, R., Salonen, R., Tuomilehto, J., & Salonen, J. (1994). Relation of leisure-time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction in men. New England Journal of Medicine, 330(22), 1549-1554.
- Landau, C., Lange, R.A., & Hillis, L.D. (1994). Percutaneous Transluminal Angioplasty. New England Journal of Medicine, 330(14), 981-993.
- Landy, F.J. (1986). Stamp collecting versus science. American Psychologist, 41(11), 1183-1192.
- LaRue, A., Bank, L., Jarvik, L., & Hetland, M. (1979). Health in old age: How do physicians' ratings and self-ratings compare? Journal of Gerontology, 34, 687-691.
- Lassner, J.B., Matthews, K.A., & Stoney, C.M. (1994). Are cardiovascular reactors to asocial stress also reactors to social stress? Journal of Personality and Social Psychology, 66(1), 69-77.
- Layman, E.M. (1974). Psychological effects of physical activity. In J.H. Wilmore (Ed.), Exercise and sports sciences reviews. New York: Academic Press.
- Lazarus, R.S. (1966). Psychological stress and the coping process. New York: McGraw-Hill.
- Leaf, A., & Ryan, T.J. (1990). Prevention of coronary artery disease. A medical imperative. New England Journal of Medicine, 323(20), 1416-1419.
- Leak, G., & Flotte, K. (1987). Factor structure and factorial replication of a new measure of the Type A behavior pattern. Psychological Reports, 60, 35-38.
- Ledwidge, B. (1980). Run for your mind: Aerobic exercise as a means of alleviating anxiety and depression. Canadian Journal of Behavioral Science Review, 12(2), 126-140.
- Lee, C. (1992). The relations of personality and cognitive style on job and class performance. Journal of Organizational Behavior, 13, 175-185.
- Lee, C., Ashford, S.J., & Bobko, P. (1990). Interactive effects of "Type A" behavior and perceived control on worker performance, job satisfaction, and somatic complaints. Academy of Management Journal, 33, 870-881.
- Lee, R., & Foley, P.P. (1986). Is the validity of a test constant throughout the test score range? Journal of Applied Psychology, 71(4), 641-644.
- Lees-Haley, P.R. (1989). Malingering traumatic mental disorder on the Beck Depression Inventory: Cancer phobia and toxic exposure. Psychological Reports, 65, 623-626.
- Leikin, L.J. (1990). Type A behaviour and health locus of control: Another view on Perloff et al. (1988). British Journal of Medical Psychology, 63, 81-84.
- Leon, G.R., Finn, S.E., Murray, D., & Bailey, J.M. (1988). Inability to predict cardiovascular disease from hostility scores or MMPI items related to Type A behavior. Journal of Consulting and Clinical Psychology, 56(4), 597-600.
- Lepore, S.J., Allen, K.A.M., & Evans, G.W. (1993). Social support lowers cardiovascular reactivity to an acute stressor. Psychosomatic Medicine, 55, 518-524.

- Lerner, D.J., & Kannel, W.B. (1986). Patterns of coronary heart disease morbidity and mortality in the sexes: A 26-year follow-up of the Framingham population. American Heart Journal, 111, 383-390.
- Leventhal, H., & Tomarken, A. (1987). Stress and illness: Perspectives from health psychology. In S.V. Kasl & C.L. Cooper (Eds.), Stress and health: Issues in research methodology (pp. 27-55). New York: John Wiley & Sons.
- Levine, M.S. (1977). Canonical analysis and factor comparison. Beverly Hills: Sage.
- Lewis-Beck, M.S. (1980). Applied regression: An introduction. London: Sage.
- Linn, B.S., & Linn, M.W. (1980). Objective and self-assessed health in the old and very old. Social Science Medicine, 14, 311-315.
- Lipschitz, D.S. (1987). The influence of exercise and Type A behaviour on the stress-strain relationship. Unpublished honours dissertation, University of the Witwatersrand, South Africa.
- Lloyd, G.G., & Cawley, R.H. (1983). Distress or illness? A study of psychological symptoms after myocardial infarction. British Journal of Psychiatry, 142, 120.
- Loop, F.D., Golding, L.R., MacMillan, J.P., Cosgrove, D.M., Lytle, B.W., & Sheidon, W.C. (1983). Coronary artery surgery in women compared with men: analyses of risks and long-term results. Journal of the American College of Cardiology, 1, 383-390.
- Lopez, F.G., & Thurman, C.W. (1986). A cognitive behavioral investigation of anger among college students. Cognitive Therapy and Research, 10(2), 245-256.
- Loscalzo, J. (1990). Regression of coronary atherosclerosis. New England Journal of Medicine, 323(19), 1337-1339.
- Lovallo, W.R., Sausen, K.P., Altmiller, D., Everson, M.S., & Wilson, M.F. (1991). Type A behavior differentially influences blood pressure during periods of low vs. high stress in medical students. Psychosomatic Medicine, 53, 211-212.
- Löwel, H., Dobson, A., Keil, U., Herman, B., Hobbs, M., Mustaniemi, H., & Tuomilehto, J. (1993). Coronary heart disease case fatality in four countries. Circulation, 88(6), 2524-2531.
- Lundberg, U., & Forsman, L. (1975). Adrenal-medullary and adrenal-cortical responses to understimulation and overstimulation: Comparison between Type A and Type B persons. Biological Psychology, 9, 79-89.
- Lundberg, U., Hedman, M., Melin, M.S.G., & Frankenhaeuser, M. (1989). Type A behavior in healthy males and females as related to physiological reactivity and blood lipids. Psychosomatic Medicine, 51, 113-122.
- MacDougall, J.M., Dembrowski, T.M., Dimsdale, J.E., & Hackett, T.P. (1985). Components of Type A, hostility, and anger-in: Further relationships to angiographic findings. Health Psychology, 4(2), 137-152.
- Maeland, J.G., & Havick, O.E. (1987). Psychological predictors for return to work after a myocardial infarction. Journal of Psychosomatic Research, 31, 471-481.

Maes, S., & Van Veldhoven, A. (1990). From health behaviour to health behaviour change. In A.A., Kaptein, H.M. van der Ploeg, Garssen, B., P.J.G. Schreurs, & R. Beunderman, (Eds.), Behavioural medicine. Psychological treatment of somatic disorders (pp.33-47). Chichester: John Wiley.

Maes, S., Vingerhoets, A., & Van Heck, G. (1987). The study of stress and disease: some developments and requirements. Social Science Medicine, 25(6), 567-578.

Malan, L. (1989). The joint moderating effect of the AHA syndrome and physical exercise on the work stress-strain relationship. Unpublished honours dissertation, University of the Witwatersrand, South Africa.

Malcolm, A.T., Janisse, M.P., & Dyck, D.G. (1984). Type A behavior, heart rate and pupillary response: Effects of cold pressor and ego threat. Journal of Psychosomatic Research, 28, 27-34.

Manuck, S.B., & Krantz, D. (1986). Psychophysiological reactivity in coronary heart disease and essential hypertension. In K.A. Matthews, S. Weiss, T. Detre, T. Dembroski, B. Falkner, S. Manuck, & R. Williams (Eds.), Handbook of stress, reactivity, and cardiovascular disease. New York: John Wiley.

Manuck, S.B., Morrison, R.L., Bellack, A.S., & Polefrone, J.M. (1985). Behavioral factors in hypertension: Cardiovascular responsivity, anger, and social competence. In M.A. Chesney & R.H. Rosenman, R.H. (Eds.), Anger and hostility in cardiovascular and behavioral disorders (pp. 149-172). New York: McGraw-Hill.

Marmot, M.G., & Madge, N. (1987). An epidemiological perspective on stress and health. In S.V. Kasl & C.L. Cooper (Eds.), Stress and health: Issues in research methodology (pp. 3-26). New York: John Wiley & Sons.

Martin, P., & Lee, H. (1992). Indicators of active and passive coping in myocardial infarction victims. Journal of Gerontology: Psychological Sciences, 47(4), 238-241.

Martin, R.A., Kuiper, N.A., & Westra, H.A. (1989). Cognitive and affective components of the Type A behavior pattern: Preliminary evidence of a self-worth contingency model. Personality and Individual Differences, 10, 771-784.

Martinsen, E.W. (1987). Interaction of exercise and medication in the psychiatric patient. In W.P. Morgan & S.E. Goldston (Eds.), Exercise and mental health (pp. 85-95). Washington: Hemisphere.

Martocchio, J.J. (1994). Effects of conceptions of ability on anxiety, self-efficacy, and learning in training. Journal of Applied Psychology, 79(6), 819-825.

Mason, C.H. & Perreault, W.D. (1991). Collinearity, power and the interpretation of multiple regression analysis. Journal of Marketing Research, 28, 268-280.

Mathews, A. (1990). Why worry? The cognitive function of anxiety. Behavioral Research and Therapy, 28(6), 455-468.

Matteson, M.T., & Ivancevich, J.M. (1980). The coronary-prone behaviour pattern: A review and appraisal. Social Science and Medicine, 14, 337-351.

Matthews, K.A. (1982). Psychological perspectives on the Type A behavior pattern. Psychological Bulletin, 91, 293-323.

- Matthews, K.A. (1988). Coronary heart disease and Type A behaviors: Update on and alternative to the Booth-Kewley and Friedman (1987) Quantitative Review. Psychological Bulletin, 104(3), 373-380.
- Matthews, K.A., Beane, W.E., Helmreich, R.L., & Lucker, G.W. (1980). Pattern A, achievement striving, and scientific merit: Does Pattern A help or hinder? Journal of Personality and Social Psychology, 5, 962-967.
- Matthews, K.A., Glass, D., Rosenman, R., & Bortner, R.W. (1977). Competitive drive, pattern A and coronary heart disease: A further analysis of some data from the Western Collaborative Group Study. Journal of Chronic Disorders, 30, 489-498.
- Matthews, K.A., & Haynes, S.G. (1986). Type A behavior pattern and coronary disease risk. American Journal of Epidemiology, 123, 923-960.
- Matthews, K.A., Rakaczky, J., Stoney, C.M., & Manuck, S.B. (1987). Are cardiovascular responses to behavioral stressors a stable individual difference variable in childhood? Psychophysiology, 24(4), 464-473.
- Mayes, B.T., Sime, W.E., & Ganster, D.C. (1984). Convergent validity of Type A Behavior pattern scales and their ability to predict physiological responsiveness in a sample of female public employees. Journal of Behavioral Medicine, 7(1), 83-107.
- Mayou, R. (1983). A controlled trial of early rehabilitation after myocardial infarction. Journal of Cardiac Rehabilitation, 3, 397-402.
- Mayou, R., Foster, A., & Williamson, B. (1978). The psychological and social effects of myocardial infarction on wives. British Medical Journal, 1, 699-701.
- McClelland, D.C., & Burnham, D. (1976, March-April). Power is the great motivator. Harvard Business Review, pp. 100-111.
- McGlynn, G.H., Franklin, B., Lauro, G., & McGlynn, I.K. (1983). The effect of aerobic conditioning and induced stress on state-trait anxiety, blood pressure, and muscle tension. Journal of Sports Medicine, 23, 241-351.
- Meakin, C.J. (1992). Screening for depression in the medically ill. The future of paper and pencil tests. British Journal of Psychiatry, 160, 212-216.
- Mechanic, D. (1962). Students under stress. New York: Free Press.
- Meininger, J.C., Stashenko, E.E., & Hayman, L.L. (1991). Type A behavior in children: Psychometric properties of the Matthews Youth Test for Health. Nursing Research, 40(4), 221-227.
- Melamed, S., Harari, G., & Green, M.S. (1993). Type A behavior, tension, and ambulatory cardiovascular reactivity in workers exposed to noise stress. Psychosomatic Medicine, 55, 185-192.
- Meltzer, L.E. (1973). Anxiety in cardiovascular disease. International Congress Series, Number 305. Proceedings of the Symposium held at St. Lucas Hospital, Amsterdam, The Netherlands.
- Meninger, K.A., & Meninger, W.C. (1936). Psychoanalytic observations in cardiac disorders. American Heart Journal, 11, 10.
- Milesis, C.A. (1987). Prediction of treadmill performance from clinical characteristics in healthy persons. Journal of Cardiopulmonary Rehabilitation, 7, 365-373.

- Millar-Craig, M.W., Bishop, C.N., & Raftery, E.B. (1978). Circadian variation of blood pressure. Lancet, 1, 795-797.
- Miller, T.Q., Turner, C.W., Tindale, R.S., Posavac, E.J., & Dugoni, B.L. (1991). Reasons for the trend toward null findings in research on Type A behavior. Psychological Bulletin, 110(3), 469-485.
- Mitchell, T.R. (1985). An evaluation of the validity of correlational research conducted in organizations. Academy of Management Review, 10, 192-205.
- Mittleman, M.A., Maclure, M., Tofler, G.H., Sherwood, J.B., Goldberg, R.J., & Muller, J.E. (1993). Triggering of acute myocardial infarction by heavy physical exertion. New England Journal of Medicine, 329(23), 1677-1683.
- Montgomery, D.C., & Peck, E.A. (1982). Introduction to linear regression analysis. New York: John Wiley & Sons.
- Moosbrugger, H., & Schermelleh-Engel, K. (1991). Determinants of pain experience: Perceived competence, trait anxiety, trait depression and moderator effects. Personality and Individual Differences, 12(12), 1261-1266.
- Morris, J.N., Clayton, D.G., Everitt, M.G., Semmence, A.M., & Burgess, E.H. (1990). Exercise in leisure time: Coronary attack and death rates. British Heart Journal, 63, 325-324.
- Moschides, V.M. (1988). The influence of medical checkups and Type A behaviour on physiological and psychological strain. Unpublished honours dissertation, University of the Witwatersrand, South Africa.
- Musante, L., MacDougall, J.M., Dembroski, T.M., & Van Horn, A.E. (1983). Component analyses of the Type A coronary-prone behavior pattern in male and female college students. Journal of Personality and Social Psychology, 45, 1104-1117.
- Myers, J.L. (1979). Fundamentals of experimental design (3rd. ed.). Boston: Allyn & Bacon.
- Myrtek, M., & Greenlee, M.W. (1984). Psychophysiology of Type A behavior pattern: A critical analysis. Journal of Psychosomatic Research, 28, 455-466.
- Nakano, K. (1991). Operant self-control procedure in modifying Type A behavior. Journal of Behavior Therapy and Experimental Psychiatry, 21(4), 249-255.
- Napper, N. (1983). Stress - The silent scream. Training and Development Forum, 10-15.
- Naughton, J.P., & Hellerstein, H.K. (1971). Exercise testing and exercise training in coronary heart disease. New York: Academic Press.
- Neale, J.M., & Liebert, R.M. (1980). Science and behaviour (2nd Ed.). Englewood Cliffs, New Jersey.: Prentice-Hall.
- Newton, T.J., & Keenan, A. (1990). The moderating effect of the Type A behavior pattern and locus of control upon the relationship between change in job demands and change in psychological strain. Human Relations, 43(12), 1229-1255.
- Nielson, W.R., & Dobson, K.S. (1980). The coronary-prone behavior pattern and trait anxiety: Evidence for discriminant validity. Journal of Consulting and Clinical Psychology, 48(4), 546-547.

- Norris, R.M., Caughey, D.E., Deeming, L.W., Mercer, C.J., & Scott, P.J. (1970). Coronary Prognostic Index for predicting survival after recovery from acute myocardial infarction. Lancet, 2, 485-487.
- Norris, S.L., deGuzman, M., Sobel, E., Brooks, S., & Haywood, L.J. (1993). Risk factors and mortality among black, Caucasian, and Latina women with acute myocardial infarction. American Heart Journal, 126(6), 1312-1319.
- Novaco, R.W. (1976). The functions and regulation of the arousal of anger. American Journal of Psychiatry, 133(10), 1124-1128.
- Novaco, R.W. (1985). Anger and its therapeutic regulation. In M.A. Chesney & R.H. Rosenman (Eds.), Anger and hostility in cardiovascular and behavioral disorders (pp. 203-226). New York: McGraw-Hill.
- Novick, M.R., & Lewis, C. (1967). Coefficient alpha and the reliability of composite measurements. Psychometrika, 32, 1-13.
- Nunnally, J.C. (1967). Psychometric theory. New York: McGraw-Hill.
- Nunnally, J.C. (1978). Psychometric theory. New York: McGraw-Hill.
- O'Brien, R.G., & Kaiser, M.K. (1985). MANOVA method for analyzing repeated measures designs: An extensive primer. Psychological Bulletin, 97(2), 316-333.
- O'Connor, G.T., Buring, J.E., Yusuf, S., Goldhaber, S.Z., Olnstead, E.M., Paffenbarger, R.S., & Hennekens, C.H. (1989). An overview of randomized trials of rehabilitation with exercise after myocardial infarction. Circulation, 80, 234-244.
- Oberman, A. (1983, March). Does exercise prevent heart disease? Modern Medicine, pp. 37-49.
- Obrist, P.A. (1981). Cardiovascular psychophysiology. A perspective. New York: Plenum Press.
- Öhman, A., Nerdby, H., & Svebak, S. (1989). Components of Type A behavior and task-induced cardiovascular activation. Psychophysiology, 26(1), 81-88.
- Olke, S. (1990, April 30). Mortality: Heart disease is world's worst scourge. Washington Post, p. A2.
- Oldridge, N.B. (1988). Cardiac rehabilitation exercise programme. Compliance and compliance-enhancing strategies. Sports Medicine, 6, 42-55.
- Oldridge, N.B., & Jones, N.L. (1983). Improving patient compliance in cardiac exercise rehabilitation: Effects of written agreement and self-monitoring. Journal of Cardiac Rehabilitation, 3, 257-262.
- Oldridge, N.B., LaSalle, D., & Jones, N.L. (1980). Exercise rehabilitation of female patients with coronary heart disease. American Heart Journal, 100, 755-757.
- O'Leary, K.D., & Turkewitz, H. (1978). Methodological errors in marital and child treatment research. Journal of Consulting and Clinical Psychology, 46, 747-758.
- Olson, C.L. (1976). On choosing a test statistic in multivariate analysis of variance. Psychological Bulletin, 83(4), 579-586.

Orha, I., Nikolaeva, L.F., & Velasco, J. (1985). Secondary prevention. In V. Kallio & E. Cay (Eds.), Rehabilitation after myocardial infarction. The European experience. (pp. 73-88). Copenhagen: World Health Organization.

Orth-Gomér, K., & Undén, A.L. (1990). Type A behavior, social support, and coronary risk: Interaction and significance for mortality in cardiac patients. Psychosomatic Medicine, 52, 59-72.

Osler, W. (1892). Lectures on angina pectoris and allied states. New York: D Appelton and Company.

Oxford reference concise medical dictionary (3rd ed.). (1990). Oxford: Oxford University Press.

Paffenbarger, R.S., Hyde, R.T., Wing, A.L., & Steinmetz, C.H. (1984). A natural history of athleticism and cardiovascular health. Journal of the American Medical Association, 252(4), 491-495.

Palmer, K.J., Leppinmäke, P.M., Jones, M., & Tennant, C. (1992). The relation of the Type A behaviour pattern, factors of the Structured Interview, and anger to survival after myocardial infarction. Australian Journal of Psychology, 44(1), 13-19.

Pauly, J.T., Palmer, J.A., Wright, C.C., & Pfeiffer, G.J. (1982). The effects of a 14-week employee fitness program on selected physiological and psychological parameters. Journal of Occupational Medicine, 24, 457-463.

Payne, R. (1965). Individual differences in the study of occupational stress. In C.L. Cooper & R. Payne (Eds.), Causes, coping, and consequences of stress at work (pp. 209-232). New York: John Wiley.

Pearson, K. (1901). On lines and planes of closest fit to systems of points in space. Philosophy Magazine, 2(6), 559-572.

Pedhazur, E.J. (1980). Multiple regression in behavioral research: Explanation and prediction. New York: Holt, Rinehart & Winston.

Perloff, L., Yarnold, P.R., & Feltzer, B. (1988). Control theory and Type A behavior. British Journal of Medical Psychology, 61, 365-368.

Perry, A.R., Kane, K.M., Bernesser, K.J., & Spicker, P.T. (1990). Type A behavior, competitive achievement-striving, and cheating among college students. Psychological Reports, 66, 459-465.

Philip, A.E. (1968). The constancy of structure of a hostility questionnaire. British Journal of Social and Clinical Psychology, 7, 16-18.

Pierce, E.F., Weltman, A., Seip, R.L., & Snead, D. (1990). Effects of training specificity on the lactate threshold and VO_2 max. International Journal of Sports Medicine, 11, 267-272.

Pisa, Z., Denolin, H., & Lamm, G. (1985). Rehabilitation after myocardial infarction: Where does current knowledge lead? In V. Kallio & E. Cay (Eds.), Rehabilitation after myocardial infarction. The European experience (pp. 131-135). Copenhagen: World Health Organization.

Pollock, M.L., Miller, H.S., Linnerud, A.C., & Cooper, K.H. (1975). Frequency of training as a determinant for improvement in cardiovascular function and body composition of middle-aged men. Archives of Physical and Medical Rehabilitation, 56, 126-130.

- Posner, J.D., Gorman, K.M., Klein, H.S., & Woldow, A. (1986). Exercise capacity in the elderly. American Journal of Cardiology, *57*, 52C-58C.
- Powell, L.H. (1987). Issues in the measurement of the Type A behaviour pattern. In S.V. Kasl & C.L. Cooper (Eds.). Stress and health: Issues in research methodology (pp. 231-282). New York: John Wiley.
- Powell, L.H., Shaker, L.A., Jones, B.A., Vaccarino, L.V., Thoreson, C.E., & Pattillo, J.R. (1993). Psychosocial predictors of mortality in 83 women with premature acute myocardial infarction. Psychosomatic Medicine, *55*, 426-433.
- Powell, L.H. & Thoreson, C.E. (1985). Behavioral and psychological determinants of long-term prognosis after myocardial infarction. Chronic Diseases, *38*(3), 253-263.
- Powell, L.H. & Thoreson, C.E. (1988). Effects of Type A behavioral counseling and severity of prior acute myocardial infarction on survival. American Journal of Cardiology, *62*, 1159-1163.
- Price, V.A. (1982). What is Type A? A cognitive social learning model. Journal of Occupational Behaviour, *3*, 109-129.
- Quick, J.C., & Quick, J.D. (1984). Preventative stress management at the organisational level. Personnel, *61*(5), 24-34.
- Ragland, D.R. & Brand, R.J. (1988a). Coronary heart disease mortality in the western collaborative group study. Follow-up of 22 years. American Journal of Epidemiology, *127*(3), 462-475.
- Ragland, D.R. & Brand, R.J. (1988b). Type A Behavior and mortality from coronary heart disease. New England Journal of Medicine, *318*, 65-69.
- Rahe, R.H., Captain, M.C., Ward, H.W., Commander, M.C., & Hayes, V. (1979). Brief group therapy in myocardial infarction rehabilitation: Three- to four-year follow-up of a controlled trial. Psychosomatic Medicine, *41*(3), 229-242.
- Räikkönen, K. & Keltinkangas-Järvinen, L. (1992). Childhood hyperactivity and the mother-child relationship as predictors of risk Type A behaviour in adolescence: A six year follow-up. Personality and Individual Differences, *13*(3), 321-327.
- Rawlings, J.O. (1988). Applied regression analysis: A research tool. Pacific Grove: Wadsworth & Brooks.
- Rawson, H.E., Bloomer, K., & Kendall, A. (1994). Stress, anxiety depression, and physical illness in college students. Journal of Genetic Psychology, *155*(3), 321-330.
- Ray, J.J., & Simons, L.A. (1982). Is authoritarianism the main element of the coronary-prone personality? British Journal of Medical Psychology, *55*, 215-218.
- Rejeski, W.J., Morley, D., & Miller, H.S. (1984). The Jenkins Activity Survey: Exploring its relationship with compliance to exercise prescription and MET gain within a cardiac rehabilitation setting. Journal of Cardiac Rehabilitation, *4*, 90-94.
- Rejeski, W.J., Morley, D., Sotile, W. (1985). Cardiac rehabilitation: A conceptual framework for psychological assessment. Journal of Cardiac Rehabilitation, *5*, 172-180.
- Review panel on coronary-prone behavior and coronary heart disease (1981). Coronary-prone behavior and coronary heart disease: A critical review. Circulation, *63*, 1199-1215.

- Rhodewalt, F., & Fairfield, M. (1991). An alternative approach to Type A behavior and health: Psychological reactance and medical noncompliance. In M.J. Strube (Ed.), Type A behavior (pp. 292-312). Newbury Park: Sage.
- Rimé, B., Ucross, C.G., & Bestgen, Y. (1989). Type A behaviour pattern: Specific coronary risk factor or general disease-prone condition? British Journal of Medical Psychology, 62, 229-240.
- Rodin, J., & Salovey, P. (1989). Health Psychology. Annual Review of Psychology, 40, 533-579.
- Roman, O. (1985). Do randomised trials support the use of cardiac rehabilitation? Journal of Cardiac Rehabilitation, 5, 93-96.
- Rook, K.S. (1987). Social support vs. companionship: Effects on life stress, loneliness, and evaluations by others. Journal of Personality and Social Psychology, 52, 1132-1147.
- Rosenberg, S.J., Peterson, R.A., & Hayes, J.R. (1987). Coping behaviors among depressed and nondepressed medical inpatients. Journal of Psychosomatic Research, 31, 653-658.
- Rosenman, R.H. (1978). The interview method of assessment of the coronary-prone behavior pattern. In T.M. Dembroski, S.M. Weiss, J.L. Shields, S.G. Haynes, & M. Feinleib (Eds.), Coronary-prone behavior (pp. 55-69). New York: Springer-Verlag.
- Rosenman, R.H. (1986). Health consequences of anger and implications for treatment. Activas Nervosa Superior, 28(1), 1-23.
- Rosenman, R.H., Brand, R.J., Jenkins, C.D., Friedman, M., Strauss, R., & Wurm, M. (1975). Coronary heart disease in the Western Collaborative Group Study: Final follow-up experience of 8½ years. Journal of the American Medical Association, 233, 872-877.
- Rosenman, R.H., & Chesney, M.A. (1982). Stress, Type A behavior, and coronary disease. In L. Goldberger, and S. Breznitz (Eds.), Handbook of stress (pp.547-565). New York: Free Press.
- Rosenman, R.H., & Friedman, M. (1974). Neurogenic factors in pathogenesis of coronary heart disease. Medical Clinics of North America, 58, 269-279.
- Rosenman, R.H., Friedman, M., Straus, R., Wurm, M., Kositchek, R., Hahn, W., & Werthessen, N.T. (1964a). A predictive study of coronary heart disease. Journal of the American Medical Association, 189(1), 113-124.
- Rosenman, R.H., Friedman, M., Straus, R., Wurm, M., Kositchek, R., Hahn, W., & Werthessen, N.T. (1964b). The predictive study of coronary heart disease. The Western Collaborative Group Study. Journal of the American Medical Association, 189(1), 15-22.
- Ross, R., & Glomset, J.A. (1973). Atherosclerosis and the arterial smooth cell. Science, 180, 1332-1339.
- Roth, D.L., Bachtler, S.D., & Fillingim, R.B. (1990). Acute emotional and cardiovascular effects of stressful mental work during aerobic exercise. Psychophysiology, 27(6), 694-701.
- Roth, D.L. & Holmes, D.S. (1987). Influence of aerobic exercise training and relaxation training on physical and psychological health following stressful life events. Psychosomatic Medicine, 49, 355-365.

Rovario, S., Holmes, D.S., & Holmsten, R.D. (1984). Influence of a cardiac rehabilitation program and the cardiovascular, psychological and social functioning of cardiac patients. Journal of Behavioural Medicine, 7, 61-81.

Rozanski, A., Bairey, C.N., Krantz, D.S., Friedman, J., Resser, K.J., Moreli, M., Hilton-Chalfen, S., Hestrin, L., Bietendorf, J., & Berman, D.S. (1988). Mental Stress and the induction of silent myocardial ischaemia in patients with coronary artery disease. New England Journal of Medicine, 318(16), 1005-1012.

Rozanski, A., Krantz, D.S., & Bairey, C.N. (1991). Ventricular responses to mental stress testing in patients with coronary artery disease. Circulation, 83(4), II-137-II-144.

Ruberman, W., Weinblatt, E., Goldberg, J.D., Chaudhary, B.S. (1984). Psychosocial influences on mortality after myocardial infarction. New England Journal of Medicine, 311, 552-559.

Runyon, R.P., & Haber, A. (1980). Fundamentals of behavioral statistics. Reading: Addison-Wesley.

Russek, L.G., King, S.H., Russek, S.J., & Russek, H.I. (1990). The Harvard mastery of stress study 35-year follow-up: Prognostic significance of patterns of psychophysiological arousal and adaptation. Psychosomatic Medicine, 52, 271-285.

Sackett, D.L., & Haynes, R.B. (1976). Compliance with therapeutic regimes. Baltimore: John Hopkins University Press.

Sami, M., Kraemer, H., DeBusk, R.F. (1979). The prognostic significance of serial exercise testing after myocardial infarction. Circulation, 60, 1238-1246.

Sandvik, L., Erikssen, J., Thaulow, E., Erikssen, G., Mundal, R., & Rodahl, K. (1993). Physical fitness as a predictor of mortality among healthy, middle-aged Norwegian men. New England Journal of Medicine, 328(8), 533-537.

Schaefer, A., Brown, J., Watson, C.G., Piemel, D., DeMotts, J., Howard, M.T., Petrik, N.M., & Balleweg, B.J. (1985). Comparison of the validities of the Beck, Zung, and MMPI depression scales. Journal of Consulting and Clinical Psychology, 53(3), 415-418.

Scheier, M.F., Matthews, K.A., Owens, J.F., Magovern, G.J., Lefebvre, R.C., Abbott, R.A., & Carver, C.S. (1989). Dispositional optimism and recovery from artery bypass surgery: The beneficial effects on physical and psychological well-being. Journal of Personality and Social Psychology, 57, 1024-1040.

Scheier, M.F., Weintraub, J.K., & Carver, C.S. (1986). Coping with stress: Divergent strategies of optimists and pessimists. Journal of Personality and Social Psychology, 51, 1257-1264.

Schill, T., & Wang, S. (1990). Correlates of the MMPI-2 Anger Content Scale. Psychological Reports, 67, 800-802.

Schmitt, N.V., & Colligan, M.J. (1984). Lack of stress-health relationships: A base rate problem. Journal of Community Psychology, 12, 245-252

Schneider, E.L., & Reed, J.D. (1985). Life extension. New England Journal of Medicine, 312(18), 1159-1168.

Schneider, R.H., Friedrich, G., Neus, J., & Ruddel, J. (1982). Effect of B-blockers on Type A coronary-prone behavior. Psychosomatic Medicine, 44, 129-130.

Schneider, R.H., Julius, S., & Karunas, R. (1989). Ambulatory blood pressure monitoring and laboratory reactivity in Type A behavior and components. Psychosomatic Medicine, 51, 290-305.

Schneiderman, N., Weiss, S.M., & Kaufman, P.G. (Eds.) (1989). Handbook of research methods in cardiovascular behavioural medicine. New York: Plenum.

Schomer, H.H. (1985). Exertion therapy for the highly anxious individual. South African Journal for Research in Sport, Physical Education and Recreation, 8(1), 63-77.

Schomer, H.H. & Noakes, T.D. (1983). The Psychological effect of exercise training on patients after a myocardial infarction. South African Medical Journal, 64, 473-475.

Schonwetter, D.J., & Janisse, M.P. (1991). Alcohol consumption, anger and hostility: A link to coronary heart disease. Personality and Individual Differences, 12(10), 1049-1055.

Schwartz, D.P., Burish, T.G., O'Rourke, D.F., & Holmes, D.S. (1986). Influence of personal and universal failure on the subsequent performance of persons with Type A and Type B behavior patterns. Journal of Personality and Social Psychology, 51, 459-462.

Seals, D.R., Hagberg, J.M., Hurley, B.F., Ehsani, A.A., & Holloszy, J.O. (1984). Endurance training in older men and women I. Cardiovascular responses to exercise. Journal of Applied Physiology. Respiratory, Environmental and Exercise Physiology, 37, 1024-1029.

Segers, M.J., & Mertens, C. (1977). Personality aspects of coronary heart disease related behavior. Journal of Psychosomatic Research, 21, 79-85.

Seligman, M.E.P. (1975). Helplessness: On depression, development, and death. San Francisco: W.H. Freeman.

Selye, H. (1974). Stress without distress. Philadelphia: J.B. Lippincott & Co.

Selye, H. (1976). The Stress of life (rev. ed.). New York: McGraw-Hill.

Sharkey, B.J. (1984). Physiology of fitness (2nd ed.). New York: Human Kinetics.

Shaw, L.W. (1981). Effects of a prescribed supervised exercise program on mortality and cardiovascular morbidity in patients after a myocardial infarction. The National Exercise and Heart Disease Project. American Journal of Cardiology, 48, 39-46.

Shekelle, R.B., Gale, M., & Norusis, M. (1985). Type A score (Jenkins Activity Survey) and risk of recurrent coronary heart disease in the Aspirin Myocardial Infarction Study. American Journal of Cardiology, 56, 221-225.

Shekelle, R.B., Gale, M., Ostfeld, A.M., & Paul, O. (1983). Hostility, risk of coronary heart disease, and mortality. Psychosomatic Medicine, 45, 109-114.

Shekelle, R.B., Hulley, S.B., Neaton, J.D., Billings, J.H., Borhani, N.O., Gerace, T.A., Jacobs, D.R., Lasser, N.L., Mittlemark, M.B., & Stamler, J. (1985). The MRFIT Behavior Pattern Study. Type A behavior and incidence of coronary heart disease. American Journal of Epidemiology, 122(4), 559-570.

Shephard, R.J. (1979). Cardiac rehabilitation in perspective. In M.L. Pollock & D.H. Schmidt (Eds.), Heart disease and rehabilitation (pp. 521-547). Boston: Houghton Mifflin.

Shephard, R.J. (1985). Exercise regimens after myocardial infarction: Rationale and results. In N.K. Wenger (Ed.), Exercise and the heart (2nd ed.) (pp. 145-157). Philadelphia: F.A. Davis.

- Stephward, R.J., Corey, P., & Kavanagh, T. (1981). Compliance and the prevention of myocardial infarction. Medical Science of Sports and Exercise, 13(1), 1-5.
- Stephward, R.J., Kavanagh, T. & Klavara, P. (1985). Mood state during postcoronary cardiac rehabilitation. Journal of Cardiopulmonary Rehabilitation, 5, 480-484.
- Siegel, S.I. & Alloy, L.B. (1990). Interpersonal perceptions and consequences of depressive-significant other relationships. Journal of Abnormal Psychology, 99, 361-373.
- Siezman, A.W., Dembrowski, T.M., & Ringel, N. (1987). Components of hostility and the severity of coronary artery disease. Psychosomatic Medicine, 49(2), 127-133.
- Simons, A.D., McGowan, C.R., Epstein, L.H., Kupfer, D.J., & Robertson, R.J. (1985). Exercise as a treatment for depression: An update. Clinical Psychology Review, 5, 553-568.
- Singh, J., Singh, S., Singh, S., Singh, A., & Malhotra, R.P. (1970). Sex life and psychiatric problems after myocardial infarction. Journal of the Association of Physicians of India, 18, 503-507.
- Skinner, J.E. (1985). Psychosocial stress and sudden cardiac death: Brain mechanisms. In R.E. Beamish, P.K. Singal, N.A. Dhalla (Eds.), Stress and heart disease (pp. 44-59). Boston: Martinus Nijhoff.
- Sloan, R.P., & Bigger, J.T. (1991). Biobehavioral factors in cardiac arrhythmia pilot study (CASS). Circulation, 83(4), II-52-II-57.
- Smith, G.R., & O'Rourke, D.F. (1988). Return to work after myocardial infarction. Journal of the American Medical Association, 259(11), 1673-1677.
- Smith, L.D. (1985). What do existing questionnaire measures of Type A behaviour measure? Unpublished Honours Dissertation. University of Cape Town, South Africa.
- Smith, T.W., & Anderson, N.B. (1986). Models of personality and disease: An interactional approach to Type A behavior and cardiovascular risk. Journal of Personality and Social Psychology, 50, 1163-1173.
- Smith, T.W., & Frohm, K.D. (1985). What's so unhealthy about hostility? Construct validity and psychosocial correlates of the Cook and Medley Ho Scale. Health Psychology, 4, 503-520.
- Smith, T.W., & O'Keeffe, J.L. (1988). Cross-situational consistency of cardiovascular reactivity. Biological Psychology, 27, 237-243.
- Soloff, P.H. (1978a). Medically and surgically treated coronary patients in cardiovascular rehabilitation: A comparative study. International Journal of Psychiatry in Medicine, 9(1), 93-106.
- Soloff, P.H. (1978b). The liaison psychiatrist in cardiovascular rehabilitation: An overview. International Journal in Psychiatry in Medicine, 8(4), 393-402.
- South African Department of Education, Arts, & Sciences. National Bureau of Educational and Social Research. (1967). Manual for the Jr.-Sr. High School Personality Questionnaire. Pretoria: Bureau.
- Specchia, G., Falcone, C., Traversi, E., La Rovere, M., Guasti, L., De Micheli, G., Ardisssino, D., & De Servi, S. (1991). Mental stress as a provocative test in patients with various clinical syndromes of coronary heart disease. Circulation, 83(4), II-108-II-114.

Spector, P.E. (1977). What to do with significant multivariate effects in the multivariate analysis of variance. Journal of Applied Psychology, 62, 158-163.

Spence, J.T., Helmreich, R.L., & Pred, R.S. (1987). Impatience versus achievement striving in the Type A pattern: Differential effects on students' health and academic achievement. Journal of Applied Psychology, 72, 522-528.

Spence, J.T., Pred, R.S., & Helmreich, R.L. (1989). Achievement strivings, scholastic aptitude, and academic performance: A follow-up to "Impatience versus achievement striving in the Type A pattern". Journal of Applied Psychology, 74(1), 176-178.

Spielberger, C.D., Jacobs, G.A., Russell, S.F., & Crane, R. (1983). Assessment of anger: The State-Trait Anger Scale. In J.N. Butcher & C.D. Spielberger (Eds.), Advances in personality assessment (Vol. 2) (pp. 159-187). Hillsdale: Erlbaum.

Spielberger, C.D., Johnson, E.H., Russell, S.F., Crane, R., Jacobs, G.A., & Worden, T.J. (1985). The experience and expression of anger: Construction and validation of an anger expression scale. In M.A. Chesney & R.H. Rosenman (Eds.), Anger and hostility in cardiovascular and behavioral disorders (pp. 5-30). New York: McGraw-Hill.

Spielberger, C.D., Krasner, S.S., & Solomon, E.P. (1988). The experience, expression, and control of anger. In M.P. Janisse (Ed.), Health psychology: Individual differences and stress (pp. 89-108). New York: Springer-Verlag.

Stedman's Medical Dictionary (1961). Baltimore: Williams & Wilkins.

Steinberg, A., & Ritzman, R.F. (1990). A living systems approach to understanding the concept of stress. Behavioral Science, 35(20), 138-146.

Steinkamp, M.W. (1990). The social concomitants of competitive and impatient / aggressive components of the Type A behavior pattern in preschool children: Peer responses and teacher utterances in a naturalistic setting. Journal of Personality and Social Psychology, 59(6), 1287-1295.

Stern, M.J., & Cleary, P. (1981). National Exercise and Heart Disease Project. Psychosocial changes observed during a low-level exercise program. Archives of Internal Medicine, 141, 1463-1467.

Stern, M.J., Gorman, P.A., & Kaslow, P. (1983). The group counsel vs exercise therapy study: A controlled intervention with subjects following myocardial infarction. Archives of Internal Medicine, 143, 1719-1725.

Stern, M.J., Pascale, L., & McLoone, J. (1976). Psychosocial adaptation following an acute myocardial infarction. Journal of Chronic Diseases, 29, 513-526.

Strasser, H. (1981). Physiological measures of mental load. In E.N. Corlett & J. Richardson (Eds.), Stress, work designs, and productivity (pp. 77-89). New York: John Wiley.

Strube, M.J. (1987). A self-appraisal model of the Type A behavior pattern. In R. Hogan & W.H. Jones (Eds.), Perspectives on personality: Theory, measurement, and interpersonal dynamics (Vol 2.), (pp. 201-250). Greenwich, Connecticut: JAI Press.

Strube, M.J. (1988). Performance attributions and the Type A behavior pattern: Causal sources versus causal dimensions. Personality and Social Psychology Bulletin, 14(4), 709-721.

Strube, M.J. (1989). Evidence of the type in Type A behaviour: A taxometric analysis. Journal of Personality and Social Psychology, 56(6), 972-987.

Strube, M.J., Berry, J.M., Lott, L., Fogelman, R., Steinhart, G., Moergen, S., & Davison, L. (1986). Self-schematic representation of the Type A and B behavior patterns. Journal of Personality and Social Psychology, 51, 170-180.

Strube, M.J., & Werner, C. (1985). Relinquishment of control and Type A behaviour pattern. Journal of Personality and Social Psychology, 48(3), 688-701.

Suarez, E.C., & Williams, R.B. (1989). Situational determinants of cardiovascular and emotional reactivity in high and low hostile men. Psychosomatic Medicine, 51, 404-418.

Suarez, E.C., & Williams, R.B. (1990). The relationship between dimensions of hostility and cardiovascular reactivity as a function of task characteristics. Psychosomatic Medicine, 52, 558-570.

Suarez, E.C., Williams, R.B., Kuhn, C.M., Zimmerman, E.H., Schanberg, S.M. (1991). Diabehavioral basis of coronary-prone behavior in middle-aged men. Part II: Serum cholesterol, the Type A behavior pattern, and hostility as interactive modulators of physiological reactivity. Psychosomatic Medicine, 53, 528-537.

Suls, J., & Marco, C.A. (1990). Relationship between JAS - and FTAS - Type A behavior and man - CHD illness: A prospective study controlling for negative affectivity. Health Psychology, 9(4), 479-492.

Suls, J., & Sanders, G.S. (1988). Type A behavior as a general risk factor. Journal of Behavioral Medicine, 11, 201-226.

Suls, J., & Wan, C.K. (1989). The relation between Type A behavior and chronic emotional distress: A meta-analysis. Journal of Personality and Social Psychology, 57(3), 503-512.

Svebak, S., Knardahl, S., Nordby, H., Aakvaag, A. (1992). Components of Type A behavior pattern as predictors of neuroendocrine and cardiovascular reactivity in challenging tasks. Personality and Individual Differences, 13(6), 733-744.

Sweetland, R.C., & Keyser, D.J. (1984). Tests. Kansas City: Test Corporation of America.

Tabachnick, B.G., & Fidell, L.S. (1983). Using multivariate statistics. New York: Harper and Row.

Tang, C.S., & Critelli, J.W. (1990). Depression and judgment of control: Impact of a contingency on accuracy. Journal of Personality, 54(4), 717-727.

Taylor, H., & Cooper, C.L. (1988). Organisational change - threat or challenge? The role of individual differences in the management of stress. Journal of Organizational Change Management, 1(1), 68-80.

Taylor, H., & Cooper, C.L. (1989). The stress-prone personality: A review of the research in the context of occupational stress. Stress Medicine, 5, 17-27.

Taylor, J.A. (1953). A personality scale of manifest anxiety. Journal of Abnormal and Social Psychology, 48, 285-290.

Tennant, C.C., Langeluddecke, P.M., Fulcher, G., & Wilby, J. (1987). Anger and other psychological factors in coronary atherosclerosis. Psychological Medicine, 17, 425-431.

Terry, D.J. (1992). Stress, coping and coping resources as correlates of adaptation in myocardial infarction patients. British Journal of Clinical Psychology, 31, 215-225.

- Tett, R.P., Bobocel, R., Hafer, C., Lees, M.C., Smith, C.A., & Jackson, D.N. (1992). The dimensionality of Type A behavior within a stressful work simulation. Journal of Personality, 60(3), 533-551.
- Thiel, H.G., Parker, D., & Bruce, T.A. (1973). Stress factors and the risk of myocardial infarction. Journal of Psychosomatic Research, 17, 43.
- Thomas, S.A., & Friedman, E. (1990). Type A behavior and cardiovascular responses during verbalization in cardiac patients. Nursing Research, 39(1), 48-53.
- Thoreson, C.E., & Powell, L.H. (1992). Type A behavior pattern: New perspectives on theory, assessment, and intervention. Journal of Consulting and Clinical Psychology, 60(4), 595-604.
- Townner, E.A., & Contrada, R.J. (1988). Effects of behavioral stress and Type A behavior on cardiovascular reactivity in mildly withdrawn smokers and non-smokers. Psychophysiology, 25(4), 486.
- Trelawny-Ross, C., & Russel, O. (1987). Social and psychological responses to myocardial infarction: Multiple determinants of outcome at six months. Journal of Psychosomatic Research, 31, 125-130.
- Tucker, L.R., & Lewis, C. (1973). The reliability coefficient for maximum likelihood factor analysis. Psychometrika, 38, 1-10.
- Tunstall Pedoe, D.S. (1990). Exercise and heart disease: Is there still a controversy? British Heart Journal, 64(5), 293-294.
- Uhl, G.S., Hopkik, A.C., Hickman, J.R., Fischer, & Medina, A. (1984). Predictive implications of clinical and exercise variables in detecting significant coronary artery disease in asymptomatic men. Journal of Cardiac Rehabilitation, 4, 245-252.
- Valliant, P.M., & Asu, M.E. (1985). Exercise and its effects on cognition and physiology in older adults. Perceptual and Motor Skills, 63, 955-961.
- Van Dixhoorn, J., der Loos, J., & Duivenvoorden, H.J. (1983). Contribution of relaxation technique to the rehabilitation of myocardial infarction patients. Psychotherapy and Psychosomatics, 40, 137-147.
- Van Egeren, L.F., & Sparrow, A.W. (1990). Ambulatory monitoring to assess real-life cardiovascular reactivity in Type A and Type B Subjects. Psychosomatic Medicine, 52, 297-306.
- Van Pragg, H.N. (1982). Neurotransmitters and CNS diseases: Depression. Lancet, 2, 1259-1263.
- Verma, R.M. (1974). A study of diagnostic differences of some functional psychoses by means of self-report technique and behaviour rating with an examination of punitiveness as a variable in the psychotic disorders. Acta Psychiatrica Scandinavica, 50, 569-576.
- Verrier, R.L. (1987). Mechanisms of behaviorally induced arrhythmias. Circulation, 76(1), 48-56.
- Verrier, R.L., DeSilva, R.A., & Lown, B. (1983). Psychological factors in cardiac arrhythmias and sudden death. In D.S. Krantz, A. Baum, & J.E. Singer (Eds.), Handbook of psychology and health (Vol. 3). Cardiovascular disorders and behavior. Hillsdale: Erlbaum.

- Vinoda, K.S. (1966). Personality characteristics of attempted suicides. British Journal of Psychiatry, 112, 1143.
- Vitaliano, P.P., Russo, J., Bailey, S.L., Young, H.M., & McCann, B.S. (1993). Psychosocial factors associated with cardiovascular reactivity in older adults. Psychosomatic Medicine, 55, 164-177.
- Wagner, E., & Williams, R.S. (1987). Rehabilitation after myocardial infarction. In K.G. Andreoli, D.P. Zipes, A.G. Wallace, M.R. Kinney, & V.K. Fowkes (Eds.), Comprehensive cardiac care (6th ed.). St.Louis: C.V. Mosby.
- Walding, M.F. (1991). Pain, anxiety and powerlessness. Journal of Advanced Nursing, 16, 388-397.
- Ward, C.H., & Eisler, R.M. (1987). Type A behavior, achievement striving, and a dysfunctional self-evaluation system. Journal of Personality and Social Psychology, 53(2), 318-326.
- Ward, M.M., Chesney, M.A., Swan, G.E., Black, G.W., Parker, S.D., & Rosenman, R.H. (1986). Cardiovascular Responses in Type A and Type B men to a series of stressors. Journal of Behavioral Medicine, 9(1), 43-48.
- Warren, J.V., & Subak-Sharpe, G.J. (1984). Surviving your heart attack. The Duke University complete heart treatment program. Garden City: Doubleday.
- Weg, J.G. (1985). Therapeutic exercise in patients with chronic obstructive pulmonary disease. In N.K. Wenger (Ed.), Exercise and the Heart (2nd ed.). (pp. 261-275), Philadelphia: F.A. Davis.
- Weidner, G., Friend, R., Ficarroto, T.J., & Mendell, N.R. (1989). Hostility and cardiovascular reactivity to stress in women and men. Psychosomatic Medicine, 51(1), 36-45.
- Weidner, G., Sexton, G., McLellan, R., Conner, S., & Matarazzo, J.D. (1987). The role of Type A behavior and hostility in an elevation of plasma lipids in adult women and men. Psychosomatic Medicine, 49(22), 136-145.
- Weinblatt, E., Goldberg, J.D., Ruberman, W., Frank, C.W., Monk, M.A., & Chaudhary, B.S. (1982). Mortality after first myocardial infarction: Search for a secular trend. Journal of the American Medical Association, 247, 1576.
- Weintraub, W.S., Wenger, N.K., Jones, E.L., Craver, J.M., & Guyton, R.A. (1992). Changing demography of coronary surgery patients: Differences between men and women. Circulation, 86, Supplement I, 1772.
- Weisse, C.S. (1992). Depression and immunocompetence: A review of the literature. Psychological Bulletin, 111, 475-489.
- Weissman, A., & Beck, A.T. (1978). Development and validation of the Dysfunctional Attitude Scale: A preliminary investigation. Paper presented at the annual meeting of the American Educational and Research Association, Toronto, Canada.
- Welch, G., Hall, A., & Walkey, F. (1990). The replicable dimensions of the Beck Depression Inventory. Journal of Clinical Psychology, 46(6), 817-827.

Weltman, A., Snead, D., Stein, R., Seip, R., Schurrer, R., Rutt, R., & Weltman, J. (1990). Reliability and validity of a continuous incremental treadmill protocol for the determination of lactate threshold, fixed blood lactate concentrations, and VO_2 max. International Journal of Sports Medicine, 11, 26-32.

Wenger, N.K. (1992). Exclusion of the elderly and women from coronary trials: Is their quality of compromised? Journal of the American Medical Association, 268, 1417-1422.

Wenger, N.K., Speroff, L., & Packard, B. (1993). Cardiovascular health and disease in women. New England Journal of Medicine, 329(4), 247-256.

Whittle, J., Conigliaro, J., Good, C.B., & Lofgren, R.P. (1993). Racial differences in the use of invasive cardiovascular procedures in the Department of Veterans Affairs Medical System. New England Journal of Medicine, 329(9), 621-627.

Widimsky, J., & Broustet, J.P. (1985). Cardiopulmonary function after myocardial infarction: Effects of bed-rest, disease and training. In V. Kallio & E. Cay (Eds.), Rehabilitation after myocardial infarction. The European experience. (pp. 9-21). Copenhagen: World Health Organization.

Wildt, A.R., & Ahtola, O.T. (1980). Analysis of covariance. California: Sage.

Williams, R.B., Barefoot, J.C., Haney, T.L., Harrell, Jr., F.E., Blumenthal, J.A., Pryor, D.B., & Peterson, B. (1988). Type A behavior and angiographically documented coronary atherosclerosis in a sample of 2,289 patients. Psychosomatic Medicine, 50, 139-152.

Williams, R.B., Barefoot, J.C., & Shekelle, R.B. (1985). The health consequences of hostility. In M.A. Chesney & R.H. Rosenman (Eds.), Anger and hostility in cardiovascular and behavioral disorders (173-185). Washington: Hemisphere.

Williams, R.B., Haney, T.L., Lee, K.L., Kong, Y.H., Blumenthal, J.A., & Whalen, R.F. (1980). Type A behavior, hostility, and coronary atherosclerosis. Psychosomatic Medicine, 42, 539-549.

Willich, S.N., Lewis, M., Löwel, H., Arntz, H., Schubert, F., & Schröder, R. (1993). Physical exertion as a trigger of acute myocardial infarction. New England Journal of Medicine, 329(23), 1684-1690.

Wilson-Barnett, J. (1979). A review of research into the experience of patients suffering from coronary thrombosis. International Journal of Nursing Studies, 16(2), 183-198.

Wise, T.N., & Rosenthal, J.B. (1982). Depression, illness beliefs and severity of illness. Journal of Psychosomatic Research, 26(2), 247-253.

Wood, C. (1981, November 5). The psyche and the heart. New Scientist, pp. 366-369.

Wood, D.T. (1977). The relationship between state anxiety and acute physical activity. American Corrective Therapy Journal, 31, 67-69.

Woods, P., & Eurns, J. (1984). Type A behavior and illness in general. Journal of Behavioral Medicine, 7, 411-415.

World Health Organization. (1981). Prevention of coronary heart disease. Copenhagen: World Health Organization.

World Health Organization Standard Programme. (1982). Dynavit Meditronic 40 operating instructions. Kaiserslautern, Germany: Keifer Dynavit.

- Wright, L. (1988). The Type A Behavior Pattern and Coronary Heart Disease. American Psychologist, 43(1), 2-14.
- Wright, L. (1992). Are the physical and TABP risk factors for heart disease unique to CHD? Journal of Clinical Psychology, 48(6), 705-710.
- Wright, L., Carbonari, J., & Voyles, W. (1992). A factor analytic study of risk variables for CHD. Journal of Clinical Psychology, 48(2), 165-170.
- Wrigley, C.S., & Niehaus, J.O. (1955). The matching of two sets of factors. American Psychologist, 10, 418-419.
- Wyndham, C.H. (1979). Mortality from cardiovascular disease in the various populations in the RSA. South African Medical Journal, 56, 1023-1030.
- Yuen, S.A., & Kuiper, N.A. (1992). Type A and self-evaluations: A social comparison perspective. Personality and Individual Differences, 13(5), 549-562.
- Zillman, D. (1983). Transfer of excitation in emotional behavior. In J.T. Cacioppe & R.E. Petty (Eds.), Social psychophysiology (pp. 335-368). Hillsdale: Lawrence Erlbaum.
- Zwick, W.R. & Velicer, W.F. (1982). Factors influencing four rules for determining the number of components to retain. Multivariate Behavioral Research, 17, 253-269.
- Zyzanski, S.J., & Jenkins, G.D. (1970). Basic dimensions within the coronary-prone behavior pattern. Journal of Chronic Diseases, 23, 781-795.

APPENDICES

APPENDIX A

Measuring Instruments used in the Present Research

The Illinois Personality Assessment Test Anxiety Scale

Inside this booklet you will find forty questions, dealing with difficulties that most people experience at one time or another. It will help a lot in self-understanding if you mark yes or, not etc., To each, frankly and truthfully, to describe any problems you may have. Never pass over an item but give an answer to every single one. Your answers will be entirely confidential. Do not spend time puzzling over them. Answer each one immediately, the way you want to at this moment (not last week, or usually). You may have answered questions like this before, but answer them as you feel now.

- | | | | |
|--|---|--|------------------------------------|
| 1 I find that my interests in people and amusements, tend to change fairly rapidly. | True
<input type="checkbox"/> | In between
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 2 If people think poorly of me I can still go on quite happily and without worrying too much. | True
<input type="checkbox"/> | In between
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 3 I like to wait till I am sure that what I am saying is correct, before I put forward an argument. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 4 I am inclined to let my actions get influenced by feelings of jealousy. | Sometimes
<input type="checkbox"/> | Seldom
<input type="checkbox"/> | Never
<input type="checkbox"/> |
| 5 If I had my life to live over again I would:
(A) plan very differently, (B) want it the same. | A
<input type="checkbox"/> | In between
<input type="checkbox"/> | B
<input type="checkbox"/> |
| 6 In general, I admire my parents. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 7 I find it hard to "take 'no' for an answer", even when I know what I ask is impossible. | True
<input type="checkbox"/> | In between
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 8 I doubt the honesty of people who are more friendly than I would naturally expect them to be. | True
<input type="checkbox"/> | In between
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 9 In demanding and enforcing obedience my parents (or guardians) were:
(A) always very reasonable,
(B) often unreasonable. | A
<input type="checkbox"/> | In between
<input type="checkbox"/> | B
<input type="checkbox"/> |
| 10 I need my friends more than they seem to need me. | Rarely
<input type="checkbox"/> | Sometimes
<input type="checkbox"/> | Often
<input type="checkbox"/> |
| 11 I feel sure that I could "pull myself together" in an emergency. | Always
<input type="checkbox"/> | Often
<input type="checkbox"/> | Seldom
<input type="checkbox"/> |
| 12 As a child I was afraid of the dark. | Often
<input type="checkbox"/> | Sometimes
<input type="checkbox"/> | Never
<input type="checkbox"/> |
| 13 People sometimes tell me that I show my excitement in voice and manner too obviously. | Yes
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 14 If people take advantage of my friendliness I:
(A) soon forget and forgive,
(B) resent it and hold it against them. | A
<input type="checkbox"/> | In between
<input type="checkbox"/> | B
<input type="checkbox"/> |
| 15 I find myself upset rather than helped by the kind of personal criticism that many people make. | Often
<input type="checkbox"/> | Occasionally
<input type="checkbox"/> | Never
<input type="checkbox"/> |
| 16 Often I get angry with people too quickly. | True
<input type="checkbox"/> | In between
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 17 I feel restless as if I want something but do not know what. | Very Rarely
<input type="checkbox"/> | Sometimes
<input type="checkbox"/> | Often
<input type="checkbox"/> |
| 18 I sometimes doubt whether people I am talking to are really interested in what I am saying. | True
<input type="checkbox"/> | In between
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 19 I have always been free from any vague feelings of ill-health, such as funny pains in my head, stomach or heart. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |

- | | | | |
|---|---------------------------------------|---|---------------------------------------|
| 20 In discussion with some people, I get so annoyed that I can hardly trust myself to speak. | Sometimes
<input type="checkbox"/> | Rarely
<input type="checkbox"/> | Never
<input type="checkbox"/> |
| Through getting "worked-up" I use up more energy than most people in getting things done. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |
| I make a point of not being absent-minded or forgetful. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 23 I never find it very difficult and unpleasant to deal with people. In fact, I always stick to my original ideas. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 24 I often get over-excited and "rattled" in stressful situations. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 25 I occasionally have vivid dreams that disturb my sleep. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 26 I always have enough energy when faced with difficulties. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 27 I sometimes find myself counting things for no apparent reason. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 28 Most people are a little queer mentally, though they do not like to admit it. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 29 If I make an awkward social mistake I soon forget it. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 30 I feel grumpy and just do not want to see people:
(A) occasionally, (B) rather often. | A
<input type="checkbox"/> | In between
<input type="checkbox"/> | B
<input type="checkbox"/> |
| 31 I am brought almost to tears by having things go wrong. | Never
<input type="checkbox"/> | Very Rarely
<input type="checkbox"/> | Sometimes
<input type="checkbox"/> |
| 32 In the midst of social groups I am nevertheless sometimes overcome by feelings of loneliness and worthlessness. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 33 I wake in the night and, through worry, have some difficulty in sleeping again. | Often
<input type="checkbox"/> | Sometimes
<input type="checkbox"/> | Never
<input type="checkbox"/> |
| 34 My spirits generally stay high no matter how many troubles I meet. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 35 I sometimes feel guilty or very sorry over quite small matters. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 36 My nerves get on edge so that certain sounds, e.g. a screechy hinge, are unbearable and give me the shivers. | Often
<input type="checkbox"/> | Sometimes
<input type="checkbox"/> | Never
<input type="checkbox"/> |
| 37 If someone badly upsets me I generally calm down again quite quickly. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |
| 38 I tend to tremble or perspire when I think of a difficult task ahead. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 39 I usually fall asleep quickly, in a few minutes, when I go to bed. | Yes
<input type="checkbox"/> | In between
<input type="checkbox"/> | No
<input type="checkbox"/> |
| 40 I sometimes get very excited or "worked-up" as I think about things that have happened recently. | True
<input type="checkbox"/> | Uncertain
<input type="checkbox"/> | False
<input type="checkbox"/> |

The Beck Depression Inventory

Instructions to the Beck Depression Inventory

On this questionnaire are groups of statements. Please read each group of statements carefully. Then pick out the one statement in each group which best describes the way you have been feeling for the PAST WEEK, INCLUDING TODAY! Circle the number beside the statement you picked. If several statements in the group seem to apply equally well, circle each one. Be sure to read all the statements in each group before making your choice.

- 41 () 0 I do not feel sad.
1 I feel sad.
2 I am sad all the time and I can't snap out of it.
3 I am so sad or unhappy that I can't stand it.
- 42 () 0 I am not particularly discouraged about the future.
1 I feel discouraged about the future.
2 I feel I have nothing to look forward to.
3 I feel that the future is hopeless and that things cannot improve.
- 43 () 0 I do not feel like a failure.
1 I feel I have failed more than the average person.
2 As I look back on my life, all I can see is a lot of failures.
3 I feel I am a complete failure as a person.
- 44 () 0 I get as much satisfaction out of things as I used to.
1 I don't enjoy things the way I used to.
2 I don't get real satisfaction out of things anymore.
3 I am dissatisfied or bored with everything.
- 45 () 0 I don't feel particularly guilty.
1 I feel guilty a good part of the time
2 I feel guilty most of the time
3 I feel guilty all of the time
- 46 () 0 I don't feel I am being punished.
1 I feel I may be punished.
2 I expect to be punished.
3 I feel I am being punished.
- 47 () 0 I don't feel disappointed in myself.
1 I am disappointed in myself.
2 I am disgusted with myself.
3 I hate myself.
- 48 () 0 I don't feel I am any worse than anybody else.
1 I am critical of myself for my weaknesses or mistakes.
2 I blame myself all the time for my faults.
3 I blame myself for everything bad that happens.
- 49 () 0 I don't have any thoughts of killing myself.
1 I have thoughts of killing myself, but I would not carry them out.
2 I would like to kill myself.
3 I would kill myself if I had the chance.
- 50 () 0 I don't cry anymore than usual.
1 I cry more now than I used to.
2 I cry all the time now.
3 I used to be able to cry, but now I can't cry even though I want to.

- 51 () 0 I am no more irritated now than I ever am.
 1 I get annoyed or irritated more easily than I used to.
 2 I feel irritated all the time now.
 3 I don't get irritated at all by the things that used to irritate me.
- 52 () 0 I have not lost interest in other people.
 1 I am less interested in other people than I used to be.
 2 I have lost most of my interest in other people.
 3 I have lost all of my interest in other people.
- 53 () 0 I make decisions about as well as I ever could.
 1 I put off making decisions more than I used to.
 2 I have greater difficulty in making decisions than before.
 3 I can't make decisions at all anymore.
- 54 () 0 I don't feel I look any worse than I used to.
 1 I am worried that I am looking old or unattractive.
 2 I feel that there are permanent changes in my appearance that make me look unattractive.
 3 I believe that I look ugly.
- 55 () 0 I can work about as well as before.
 1 It takes an extra effort to get started at doing something.
 2 I have to push myself very hard to do anything.
 3 I can't do any work at all.
- 56 () 0 I can sleep as well as usual.
 1 I don't sleep as well as I used to.
 2 I wake 1-2 hours earlier than usual and find it hard to get back to sleep.
 3 I wake up several hours earlier than I used to and cannot get back to sleep.
- 57 () 0 I don't get more tired than usual.
 1 I get tired more easily than I used to.
 2 I get tired from doing almost anything.
 3 I am too tired to do anything.
- 58 () 0 My appetite is no worse than usual.
 1 My appetite is not as good as it used to be.
 2 My appetite is much worse now.
 3 I have no appetite at all anymore.
- 59 () 0 I haven't lost much weight, if any, lately.
 1 I have lost more than 5 pounds.
 2 I have lost more than 10 pounds.
 3 I have lost more than 15 pounds.
- I am purposefully trying to lose weight by eating less. Yes _____ No _____
- 60 () 0 I am no more worried about my health than usual.
 1 I am worried about physical problems such as aches and pains; or upset stomach; or constipation.
 2 I am very worried about physical problems and it's hard to think of much else.
 3 I am so worried about my physical problems, that I cannot think about anything else.
- 61 () 0 I have not noticed any recent change in my interest in sex.
 1 I am less interested in sex than I used to be.
 2 I am much less interested in sex now.
 3 I have lost interest in sex completely.

The Jenkins Activity Survey (Form N)

For each question choose the answer that is true for you, and circle the number to the right of the answer.

- 62 When you are under pressure or stress, do you usually
- | | |
|--|---|
| Do something about it immediately? | 1 |
| Plan carefully before taking any action? | 2 |
- 63 When you listen to someone talking, and this person takes too long to come to the point, how often do you feel like hurrying the person along?
- | | |
|--------------|---|
| Frequently | 1 |
| Occasionally | 2 |
| Almost never | 3 |
- 64 How often do you actually "put words in the person's mouth" in order to speed things up?
- | | |
|--------------|---|
| Frequently | 1 |
| Occasionally | 2 |
| Almost never | 3 |
- 65 If you tell your spouse or a friend that you will meet somewhere at a definite time, how often do you arrive late?
- | | |
|------------------|---|
| Once in a while | 1 |
| Rarely | 2 |
| I am never late. | 3 |
- 66 When you were younger, did most people consider you to be
- | | |
|--|---|
| Definitely hard-driving and competitive? | 1 |
| Probably hard-driving and competitive? | 2 |
| Probably more relaxed and easygoing? | 3 |
| Definitely relaxed and easygoing? | 4 |
- 67 How would your spouse (or closest friend) rate you currently?
- | | |
|--|---|
| Definitely hard-driving and competitive? | 1 |
| Probably hard-driving and competitive? | 2 |
| Probably more relaxed and easygoing? | 3 |
| Definitely relaxed and easygoing? | 4 |
- 68 How would your spouse (or closest friend) rate your general rate of activity?
- | | |
|--|---|
| Too slow. Should be more active. | 1 |
| About average. Is busy much of the time. | 2 |
| Too active. Needs to slow down. | 3 |
- 69 How was your temper when you were younger?
- | | |
|----------------------------|---|
| Fiery and hard to control. | 1 |
| Strong, but controllable. | 2 |
| No problem. | 3 |
| I almost never got angry. | 4 |

70 Would people you know well agree that you have less energy than most people?

- | | |
|-----------------|---|
| Definitely yes. | 1 |
| Probably yes. | 2 |
| Probably no. | 3 |
| Definitely no. | 4 |

71 When you are in a group, how often do people look to you for leadership?

- | | |
|--|---|
| Rarely. | 1 |
| About as often as they look to others. | 2 |
| More often than they look to others. | 3 |

For each of the next questions compare yourself with the average person having the same kind of daily activities and life situation as you (e.g. similar occupation, family responsibilities, outside interests) and mark the most accurate description.

72 In being precise (careful about detail), I am

- | | |
|------------------------|---|
| Much more precise. | 1 |
| A little more precise. | 2 |
| A little less precise. | 3 |
| Much less precise. | 4 |

73 I approach life in general

- | | |
|--------------------------|---|
| Much more seriously. | 1 |
| A little more seriously. | 2 |
| A little less seriously. | 3 |
| Much less seriously. | 4 |

74 When you were at high school, college, or university, did you play in any sports teams?

- | | |
|-------------------------|---|
| No. | 1 |
| Yes, one team. | 2 |
| Yes, two or more teams. | 3 |

Anger Expression (AX) Scale

Directions: a number of statements which people have used to describe themselves when they feel angry or furious are given below. Read each statement and then darken the appropriate circle on the answer sheet to indicate how often you feel or act in the manner described. There are no right or wrong answers. Do not spend too much time on any one statement. For each item darken the circle which seems to best describe how you generally act or feel when you are angry or furious.

<u>WHEN ANGRY OR FURIOUS.....</u>	<u>Almost Never</u>	<u>Some- times</u>	<u>Often</u>	<u>Almost Always</u>
88 I express my anger	1	2	3	4
89 I keep things in	1	2	3	4
90 I pout or snik	1	2	3	4
91 I withdraw from people	1	2	3	4
92 I make sarcastic remarks to others	1	2	3	4
93 I do things like slam doors	1	2	3	4
94 I boil inside, but I don't show it.	1	2	3	4
95 I argue with others	1	2	3	4
96 I tend to harbor grudges that I don't tell anyone about.	1	2	3	4
97 I strike out at whatever infuriates me	1	2	3	4
98 I am secretly quite critical of others	1	2	3	4
99 I am angrier than I am willing to admit	1	2	3	4
100 I say nasty things.	1	2	3	4
101 I am irritated a great deal more than people are aware of.	1	2	3	4
102 I lose my temper.	1	2	3	4
103 If someone annoys me, I am apt to tell him or her how I feel.	1	2	3	4

Acting-out hostility subscale of the Hostility and Direction of Hostility
Questionnaire

Please fill in this form by putting a circle round the "true" or the "false" after each of the following statements. If you find it difficult to decide, ask yourself whether you think the statement is on the whole true or false and put a circle round the appropriate word.

- | | | |
|---|---|---|
| 75 I don't blame anyone for trying to grab everything he can get in this world. | T | F |
| 76 I can easily make other people afraid of me, and sometimes do for the fun of it. | T | F |
| 77 In school I was sometimes sent to the principle for misbehaving. | T | F |
| 78 Sometimes I enjoy hurting persons I love. | T | F |
| 79 Sometimes I feel as if I must injure either myself or someone else. | T | F |
| 80 I sometimes tease animals. | T | F |
| 81 I get angry sometimes. | T | F |
| 82 At times I have a strong urge to do something harmful or shocking. | T | F |
| 83 I am easily downed in an argument. | T | F |
| 84 I easily become impatient with people. | T | F |
| 85 I get angry easily and then get over it soon. | T | F |
| 86 At times I feel like smashing things. | T | F |
| 87 At times I feel like picking a fist fight with someone. | T | F |

APPENDIX B

The Covering Letter, Letter of Informed Consent, Demographic Checklist and Open Ended Question Used in the First Study



City of Johannesburg

CITY HEALTH DEPARTMENT

Medical Officer of Health : Prof Hillard S. Marwitz

Your ref.

Our ref.

Refer to

Civic Centre
Braamfontein
Johannesburg

1477
Johannesburg
2100

☎ 407-6111

Fax 403-1069

Dear Sir/Madam

The Johannesburg City Health Department Cardiac Rehabilitation Centre is currently conducting a research project in conjunction with members of the University of the Witwatersrand's School of Psychology. The aim of the project is to examine aspects of the rehabilitation process in cardiac patients. To achieve this objective we are examining the psychological and physical well being of cardiac patients. The role of the School of Psychology is to analyse and collate the data pertaining to the project. The collated data will be used by a postgraduate student for the purpose of a Masters degree.

Please would you assist us by completing the following questionnaire and returning it in the enclosed envelope as soon as possible. No member of the School of Psychology will have access to patient's records or names. I would further like to emphasise that your participation is voluntary and that your responses will be used for research purposes only.

In accordance with the stipulations set out by the ethics committees of the Johannesburg City Health Department and the University of the Witwatersrand, we are unable to release your anonymous data to the School of Psychology without your written consent. Should you decide to participate in the project, please complete the attached form of informed consent and return it with the questionnaire. Please note that you are free to withhold consent and such action will not effect your status at the unit.

Thank you for your time and cooperation.

Yours sincerely


A.G. Digenio



Your ref.

Civic Centre
Braamfontein
Johannesburg

1477
Johannesburg
2000

Our ref.

☎ 407-8111
Fax 403-1069

Refer to

INFORMED CONSENT

PHYSICAL EXERCISE AS A MODERATOR OF THE TYPE A - STRAIN RELATIONSHIP IN CARDIAC REHABILITATION PATIENTS

I, _____ aged _____ years,
consent to participate in the research project, concerning aspects of the cardiac
rehabilitation process, conducted by the Johannesburg City Health Department
Cardiac Rehabilitation Centre and members of the University of the
Witwatersrand's School of Psychology.

I understand and appreciate that:

a. - by agreeing to participate, I consent to the release of information pertaining
to my physiological and psychological status to members of the University of the
Witwatersrand's School of Psychology.

b. - my name will not be released to the members of the University of the
Witwatersrand's School of Psychology.

c. - the released information will remain confidential.

d. - participation is voluntary.

e. - the information will be used for research purposes only.

(Signature of participant)

(Date)

Witness: _____
(Name)

(Signature of witness)

(Date)

The Cardiac Rehabilitation Questionnaire

The aim of this questionnaire is to assess cardiac rehabilitation patient's physiological and psychological condition. In the following pages, you will find a number of statements with corresponding responses. Please mark the response which is closest to the way you feel. There are no right or wrong responses, so please answer the questions as honestly and accurately as possible. Confidentiality is guaranteed. You are not required to write your name anywhere on this questionnaire, so your responses will remain anonymous.

PLEASE PROVIDE THE FOLLOWING BIOGRAPHICAL INFORMATION:

Present Occupation: _____
Age : _____
Sex : _____
Race : _____
Present weight : _____
Education : (please circle the appropriate response)
1. Std. VIII
2. Matric
3. Technical
4. Professional Qualification
5. University Degree
6. Post Graduate Degree

PLEASE TICK THE APPROPRIATE RESPONSES:

1. Have you experienced any of the following since your last medical check up at the Cardiac rehabilitation Centre:

- a. Myocardial Infarction
- b. Coronary Artery bypass
- c. Angioplasty
- d. Coronary Artery Angiography
- e. Other Cardiac Surgery
- f. At present, do you have any physical complaints (e.g. chest pain, palpitations, shortness of breath, fatigue).
- g. if yes, please specify.

Yes	No
Yes	No
Yes	No
Yes	No
Yes	No
Yes	No

2. Since leaving the Cardiac rehabilitation Centre, have you smoked:

a. Cigarettes

Yes	No	Stopped
-----	----	---------

Date started (year) _____

Date stopped (year, month, day) _____

No. of cigarettes smoked per day (average) _____

b. A Pipe

Yes	No	Stopped
-----	----	---------

Date started (year) _____

Date stopped (year, month, day) _____

c. Cigars

Yes	No	Stopped
-----	----	---------

Date started (year) _____

Date stopped (year, month, day) _____

No. of cigars smoked per day (on average) _____

3a. Are you currently on treatment for Hypertension?

Yes	No
-----	----

b. How long have you been receiving treatment? _____

4a. Have you had your cholesterol level checked within the past six months?

Yes	No
-----	----

b. Were the results positive?

Yes	No
-----	----

5. Have you been diagnosed as having Diabetes Mellitus?

Yes	No
-----	----

6a. Have your parents or siblings ever experienced a coronary event?

Yes	No
-----	----

b. If yes, what was their relationship to you (e.g. Mother, father) _____

c. How old were they when the first event occurred? _____

7a. Are you currently participating in any supervised exercise?

Yes	No
-----	----

b. Where? _____

c. Type of exercise? _____

d. How often do you attend? _____

In order to assess the effectiveness of the Cardiac Rehabilitation Centre's program, we would like you to indicate your reasons for leaving the Centre and any suggestions that you may have for how the program can be improved. The Cardiac Rehabilitation Centre will not be notified of your particular suggestions or reasons for leaving, but will be provided with a summary of all responses.

Reasons for leaving: _____

Suggestions: _____

APPENDIX C

The Type A Component Questionnaire

The Type A Component Questionnaire

For each question choose the answer that is true for you, and circle the number to the right of the answer.

- 1 When you listen to someone talking, and this person takes too long to come to the point, how often do you feel like hurrying the person along?
- | | |
|--------------|---|
| Frequently | 1 |
| Occasionally | 2 |
| Almost never | 3 |
- 2 How often do you actually "put words in the person's mouth" in order to speed things up?
- | | |
|--------------|---|
| Frequently | 1 |
| Occasionally | 2 |
| Almost never | 3 |
- 3 I easily become impatient with people. T F
- 4 When you were younger, did most people consider you to be
- | | |
|--|---|
| Definitely hard-driving and competitive? | 1 |
| Probably hard-driving and competitive? | 2 |
| Probably more relaxed and easygoing? | 3 |
| Definitely relaxed and easygoing? | 4 |
- 5 How would your spouse (or closest friend) rate you currently?
- | | |
|--|---|
| Definitely hard-driving and competitive? | 1 |
| Probably hard-driving and competitive? | 2 |
| Probably more relaxed and easygoing? | 3 |
| Definitely relaxed and easygoing? | 4 |
- 6 I approach life in general
- | | |
|--------------------------|---|
| Much more seriously. | 1 |
| A little more seriously. | 2 |
| A little less seriously. | 3 |
| Much less seriously. | 4 |
- 7 How was your temper when you were younger?
- | | |
|-------------------------------|---|
| Fiercely and hard to control. | 1 |
| Strong, but controllable. | 2 |
| No problem. | 3 |
| I almost never got angry. | 4 |
- 8 I can easily make other people afraid of me, and sometimes do for the fun of it. T F
- 9 Sometimes I enjoy hurting persons I love. T F
- 10 At times I have a strong urge to do something harmful or shocking. T F
- 11 I get angry easily and then get over it soon. T F
- 12 At times I feel like smashing things. T F
- 13 At times I feel like picking a fist fight with someone. T F


WHEN ANGRY OR FURIOUS.....

	<u>Almost Never</u>	<u>Some- times</u>	<u>Often</u>	<u>Almost Always</u>
14 I pout or sulk		2	3	4
15 I withdraw from people	1	2	3	4
16 I boil inside, but I don't show it.	1	2	3	4
17 I tend to harbor grudges that I don't tell anyone about	1	2	3	4
18 I am angrier than I am willing to admit	1	2	3	4
19 I am irritated a great deal more than people are aware of.	1	2	3	4
20 I keep things in.	1	2	3	4
21 I express my anger	1	2	3	4
22 I argue with others	1	2	3	4
23 I strike out at whatever infuriates me	1	2	3	4
24 I say nasty things.	1	2	3	4
25 I lose my temper.	1	2	3	4
26 If someone annoys me, I am apt to tell him or her how I feel.	1	2	3	4


APPENDIX D

Residual Plots used for the Assumption Test of Linearity


Age




Achievement Striving




Impatience Irritability




Anger-in




Anger - out




Hostility




Frequency of Exercise




Duration of Exercise




Age




Achievement Striving




Impatience Irritability




Anger - In




Anger - out




Hostility




Frequency of Exercise



Duration of Exercise




Age




Predicted values of change in VO_{2max}


Achievement Striving




Impatience Irritability




Anger-in




Anger-out




Hostility



Frequency of Exercise




Duration of Exercise




APPENDIX E

**Normal Probability Plots used to Test
the Assumption of Normality**

Normal Probability Plot Achievement Striving




Normal Probability Plot Impatience Irritability




Normal Probability Plot


Anger - In




Normal Probability Plot Anger - Out




Normal Probability Plot Hostility




Normal Probability Plot Exercise Frequency




Normal Probability Plot Exercise Duration




Normal Probability Plot Change in Anxiety




Normal Probability Plot Change in Depression



Normal Probability Plot Change in Maximal Oxygen Uptake



Normal Probability Plot Age





Author: Feldner-Busztin Adrienne.

Name of thesis: Predicting the psychological and physiological prognosis of cardiac rehabilitation patients.

PUBLISHER:

University of the Witwatersrand, Johannesburg

©2015

LEGALNOTICES:

Copyright Notice: All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed or otherwise published in any format, without the prior written permission of the copyright owner.

Disclaimer and Terms of Use: Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.