

https://dspace.org/

Immunoglobulin subtying and quantification in direct antiglobun test positive heamolysis in an HIVprevalent setting

Thirosha Chetty; Nikki Bouwer; Yuen On Wan; Johnny Mahlangu

https://hdl.handle.net/10539/36669
WOS

Downloaded from a DSpace repository. https://dspace.org/

Immunoglobulin subtyping and quantification in direct antiglobulin test: positive haemolysis in an HIV-prevalent setting

Thirosha Chetty ,[1,2] Nikki Bouwer,[1,2] Yuen On Wan,[3] Johnny Mahlangu[1,2]

[1]Department of Molecular Medicine and Haematology, University of the Witwatersrand Faculty of Health Sciences, Johannesburg, South Africa
[2]Haematology, National Health Laboratory Service, Johannesburg, South Africa
[3]Southern Community Laboratories, and Department of Pathology, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand

Correspondence to
Dr Thirosha Chetty, Department of Molecular Medicine and Haematology, University of the Witwatersrand Faculty of Health Sciences, Johannesburg 2000, South Africa; thirosha.chetty@nhls.ac.za

Received 12 September 2020
Revised 29 November 2020
Accepted 30 November 2020
Published Online First 16 December 2020

Check for updates

© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chetty T, Bouwer N, Wan YO, *et al*. *J Clin Pathol* 2022;**75**:117–120.

ABSTRACT

Aims Positive direct antiglobulin tests (DATs) are valuable in identifying the aetiology of autoimmune haemolysis and in guiding therapeutic intervention. However, in HIV-positive individuals with background polyclonal gammopathy, a positive DAT in the absence of haemolysis is common. In this setting, IgG quantification and subtyping may be of value, as this is possible with the recently introduced gel cards. There is paucity of literature evaluating the diagnostic usefulness of IgG subtyping and quantification in HIV-positive individuals who are investigated for autoimmune haemolytic anaemia (AIHA). This study evaluated the usefulness of IgG quantification and subtyping in the diagnostic work-up of AIHA in patients with a positive DAT, with and without HIV infection.

Methods This retrospective, cross-sectional study included patients investigated for AIHA in a quaternary care hospital. Those with a positive DAT had their IgG subtyped and quantified using the ID-Card DAT IgG_1/IgG_3 and IgG-dilution cards (Bio-Rad, Cressier, Switzerland).

Results Ninety patients admitted from December 2019 to March 2020 were investigated for AIHA. Forty-four (49%) patients had a positive DAT of whom 26 (59%) had evidence of haemolysis, and 16 (36%) were HIV positive. Concurrent HIV and haemolysis were present in eight patients, two of whom had IgG_1 although none had an IgG antibody titre >1:30. None of the HIV-positive patients without features of haemolysis had IgG_1/IgG_3 or IgG antibody titres >1:30.

Conclusion In our clinical setting, IgG quantification and subtyping were found to be of limited value in the diagnostic characterisation of AIHA in HIV-positive patients with false-positive DAT.

INTRODUCTION

Haemolysis is premature or accelerated destruction of red blood cells, and anaemia occurs when the rate of red cell breakdown is greater than the red cell production rate by the bone marrow.[1 2] The prevalence of autoimmune haemolytic anaemia (AIHA) in many developing countries is poorly documented. In AIHA, secondary causes can be identified in 50%–70% of all cases, these include drugs, infections, malignancies and autoimmune diseases such as systemic lupus erythematosus (SLE).[1 2]

The direct antiglobulin test and AIHA

In the correct clinical setting, the direct antiglobulin test (DAT) is routinely used to confirm the diagnosis of AIHA. This test detects both red cells coated with immunoglobulin and complement. The clinical utility of this test is dependent on its interpretation in a relevant clinical setting, together with laboratory evidence of red cell breakdown. The 2020 recommendations for the diagnosis and treatment of AIHA in adults define the diagnostic criteria required for AIHA to be the presence of haemolysis, a positive DAT in the absence of an alternative cause.[2] Conditions characterised by raised IgG level are associated with non-specific attachment of IgG to the red blood cells in the absence of haemolysis. These conditions include pregnancy, malignancies, drugs and HIV infection.[3]

HIV infection and AIHA

HIV infection is a common disease in sub-Saharan Africa with an incidence of 2.6/100 person-years reported in the 2011–2015 period.[4] Anaemia is present in 95% of HIV-positive patients and is often multifactorial in origin. Haemolysis is considered to be an important cause due to HIV infection or secondary to treatment.[5]

HIV positivity is an independent risk factor for warm AIHA, and individuals on combination antiretrovirals (cARTs) were noted to have a higher risk compared with HIV-positive individuals who were not on treatment.[6] Yen *et al*[6] reported that HIV-positive patients have a 28-fold greater risk for developing AIHA compared with the control group. Proposed causes for this include molecular mimicry, loss of immune tolerance, dysfunctional B lymphocytes and the production of autoreactive antibodies.[6–8]

DAT positivity in HIV-infected individuals in the absence of haemolysis is a common occurrence that is hypothesised to be secondary to the HIV infection and/or the treatment.[9–11] Lai *et al*[11] showed that DAT positivity, in the absence of overt haemolysis, was associated with lower haemoglobins (Hb), treatment with trimethoprim–sulfamethoxazole and coinfection with hepatitis C.[11] In sub-Saharan Africa, with a high prevalence of HIV infection, there is lack of data describing the pattern of DAT results in relation to other clinical features of haemolysis. Better understanding of the pattern of IgG positivity among the HIV-positive population may translate into a better clinical interpretation of this laboratory finding.

Subtyping IgG and quantifying IgG

The IgG subtypes that are most commonly associated with haemolytic anaemia are IgG_1 and IgG_3,

both of which have increased ability to bind to the Fc receptors on phagocytic cells.[12] Ukita *et al*[12] and Dubarry *et al*[13] showed positive correlation between the presence of IgG_1 and IgG_3 in patients with positive DAT results and clinical/biochemical features of haemolysis. The presence of these IgG subtypes is also proposed to be a predictor of severity of haemolysis.[14–16]

The quantification of IgG in patients with suspected AIHA might help us understand if the amount of IgG attached to the red cell surface is sufficient to cause clinically significant haemolysis.[12 17] In several studies, IgG levels greater than 1:300 were found to correlate with severity of haemolysis.[15 18 19] The gel test for IgG subtyping and quantification is much easier to use and is more cost-effective when compared with flow cytometry, and in fact yields concordant results.[19 20] There is a paucity of literature describing the potential benefit of IgG quantification and subtyping in patients with HIV infection and a positive DAT.

MATERIALS AND METHODS

The study period was from December 2019 to March 2020. The study population comprised all patients who were investigated for haemolysis and included all those where IgG was detected on the DAT (DC screening II, card-monospecific antihuman globulin reagent for IgG and C3d). The DATs were requested as a part of routine work-up in patients with suspected autoimmune haemolysis. Residual blood samples were centrifuged and red cells washed according to the package insert. The washed red cells were then further tested for the presence of IgG_1 or IgG_3 and quantification of the IgG was performed at 1:10, 1:30, 1:100, 1:300 and 1:1000 dilutions by using ID-Card DAT IgG_1/IgG_3 and IgG-dilution cards (Bio-Rad, Cressier, Switzerland). The highest concentration of IgG measured for a single sample was recorded.

For the purpose of this study, haemolysis was defined to be the presence of one or more of the following: reduced haptoglobin, raised reticulocyte count, raised unconjugated bilirubin, raised lactose dehydrogenase, and/or peripheral smear findings that included spherocytes, polychromasia and nucleated red blood cells. In this study, raised biochemical markers of inflammation, namely C reactive protein and/or procalcitonin levels were used as surrogate biochemical markers for sepsis. In this study, anaemia was defined as haemoglobin concentrations lower than the lower limit of laboratory-specific reference ranges for the age and sex of participants. Participants' demographic information was retrieved from the laboratory information system.

Statistical analysis

Qualitative data were described, and quantitative data were tabulated and summarised using standard statistics. Microsoft Excel (Microsoft Corporation, Washington, USA) and Statistica (TIBCO Software Corporation, California, USA) programs were used for data collection and analysis of the DAT IgG-dilution and DAT IgG_1/IgG_3 results.

RESULTS

The demographic data are summarised in table 1. A total of 90 samples were included in this study, 44 samples with positive IgG on DAT and anaemia, and 46 negative control samples which were confirmed to be IgG negative.

Secondary causes of haemolysis were sought in IgG-positive samples with haemolysis (n=26). These causes included sepsis (13/26, 50%), autoimmune conditions (9/26, 35%) and malignancy (3/26, 12%).

Table 1 Demographic baseline characteristics of the study population

	DAT positive for IgG (monospecific)			
	With haemolysis	Without haemolysis	DAT negative*	Study population
Number (n)	26	18	46	90
Median age (years)	33	40	54	42
DAT results				
IgG	26	18		
C3d	3	0		
Gender				
Female (n)	15	15	30	60
Male (n)	11	3	16	30
HIV status				
HIV positive (n)	8	8	9	25
HIV negative (n)	12	6	16	34
Not tested (n)	6	4	21	31
Secondary causes				
Autoimmune (n)	9	4		
Malignancy (n)	3	1		
Sepsis (n)	13	9		
Other (n)	4	5		
Diagnostic parameters†				
Raised LDH (n)	26	–		
Low hapto (n)	17	–		
Raised UBR (n)	21	–		
Reticulocytosis (n)	11	–		
Spherocytosis (n)	13	–		
NRBC (n)	14	–		
Polychromasia (n)	19	–		

*Confirmed to be IgG negative.
†The diagnostic parameters of the patients determined to have haemolysis.
DAT, direct antiglobulin test; hapto, haptoglobin; LDH, lactate dehydrogenase; NRBC, nucleated red blood cells; UBR, unconjugated bilirubin.

The HIV status was recorded in 34 of the 44 cases with IgG-positive anaemia. Of these 34 patients, 16 (47%) were HIV positive and had an IgG-positive anaemia (figure 1). HIV positivity had no statistically significant relationship with warm AIHA (p=0.32, 95% CI). In the 16 HIV-positive patients with IgG-positive DAT, the viral loads were available in 15 patients, and CD4 counts in 11. Information regarding cART therapy use was available for four (25%) of these patients. For the HIV-positive patients without features of haemolysis, there was no information regarding their use of cART. Sepsis was noted in the majority (81%, 13/16) of HIV-positive patients with IgG-positive anaemia, with and without haemolysis.

Six of the cases with haemolysis demonstrated concentrations of IgG_1 (6/26, 23%) (table 2). In 50% (3/6) of these patients the presence of C3d on the DAT was demonstrated. IgG_3 was not detected in any of the 44 patients with and without haemolysis.

Twenty-eight patients in our study (n=90) had concentrations of IgG at ≥1:10 (table 3). Two HIV-negative patients with features of haemolysis demonstrated concentrations of IgG at 1:100 and 1:300. The first was an adult female patient with SLE with the highest concentration measured of IgG (1:1000). The second was a paediatric male patient with a malignancy. Both displayed all features of haemolysis and had severe anaemia (Hb <60g/L).




Figure 1 Summary of HIV-positive population. DAT, direct antiglobulin test.

DISCUSSION

This study highlights the population of patients in a South African setting with DAT-positive anaemia. This includes both HIV-positive and negative patients, with and without features of haemolysis. The median age of our group of patients (34 years) was lower that of Kruhonja Galic *et al*[21] and in the GIMEMA Study.[22] This finding may indicate that haemolysis occurs at a younger median age in a South African setting. The inclusion of paediatric patients (6/44, 14%) in our study is contributory to the younger median age.

In our study there were more women than men in the groups with and without haemolysis. This finding concurs with those reported in the other studies.[14 21–23] Alonso *et al*[23] ascribed the increased female:male ratio to the increased susceptibility of women to autoimmune conditions.

Eighty-five per cent of the patients with features of haemolysis had secondary warm AIHA. This finding concurs with the studies of Alonso *et al*[23] and Kruhonja Galic *et al*.[21] Two of the patients with warm AIHA had SLE. The top two autoimmune conditions associated with secondary AIHA are SLE and antiphospholipid syndrome.[14 23 24] The majority of patients in our study had features of underlying sepsis regardless of HIV status. This serves as a reminder to clinicians to exclude the presence of associated conditions, such as an undiagnosed malignancy or infections, when there is a positive DAT with features of haemolysis.[14]

Table 2 Subtyping of IgG

	IgG positive DAT		
	Patients with haemolysis (n=26)	Patients without haemolysis	Controls (n=46)
IgG_1	6	0	0
IgG_3	0	0	0
IgG_1 and IgG_3	0	0	0
IgG_1 and IgG_3 negative	20	18	46

Breakdown of IgG_1 positive cases (6): HIV-positive with sepsis (2); HIV negative with SLE (2); HIV negative with malignancy (1); HIV negative with unknown (1).
DAT, direct antiglobulin test; SLE, systemic lupus erythematosus.

Table 3 Quantification of IgG

Dilutions of IgG	With haemolysis (n=26)	Without haemolysis (n=18)	Controls (n=46)
1:10	11	6	0
1:30	6	2	0
1:100/1:300	2	0	0
1:1000	1	0	0

Our study population included 16 HIV-positive patients (16/34), 47% of those with known HIV status (figure 1). The presence of non-specific IgG, hyperglobulinaemia and circulating immune complexes attached to red blood cells is often described in the setting of HIV. Half of the HIV-positive patients with IgG-positive anaemia did not demonstrate features of haemolysis and would be classified as false-positive DATs. Our study did not reveal a statistically significant association of HIV infection with haemolysis. Despite this, the appropriate clinical indication to perform specialised laboratory tests, as well as the importance of clinical correlation when interpreting laboratory results, must be emphasised.[11 25]

All of the HIV-positive patients, where the data were available regarding both CD4 count and viral load, had failed to achieve both immunological and virological remission. This is in keeping with the theory of De Angelis *et al*[9] and Lepennec *et al*[26] who suggested that poorly controlled HIV-positive patients were more likely to demonstrate the presence of IgG on the DAT in the absence of haemolysis. The presence of sepsis and failure to achieve remission in the HIV-positive patients (12/16, 75%), with and without haemolysis, could indicate poor control of the underlying immune deficiency syndrome.

Yen *et al*[6] reported that patients on cART, specifically those on efavirenz, had an increased risk of AIHA. Unfortunately, our study population did not have adequate information regarding cART for the majority of the HIV-positive patients included.

In the group of patients with warm AIHA, a small proportion of patients (6/26, 23%) had the subtype IgG_1. Kruhonja Galic *et al*[21] described similar association of IgG_1 positivity in patients with warm AIHA, compared with patients without haemolysis and a positive DAT.

IgG_3 and/or IgG_1 were not detected in any of the patients without haemolysis regardless of HIV status. These findings do not agree with those reported by both Lai *et al*[18] and Kruhonja Galic *et al*[21] of IgG_3 detection in cases without haemolysis.

Although IgG_1 with or without IgG_3 has been reported as commonly associated with AIHA by numerous authors,[15 21 27] correlation of the presence of these IgG subtypes with severity of haemolysis was only described by Das *et al*.[14] The detection of either or both of these IgG subtypes in patients with suspected AIHA has been proposed to be helpful in stratifying the risk of developing severe haemolysis.[20 24] In our study, IgG_1 was only detected in patients with haemolysis. Patients with IgG_1 demonstrated variable severity of haemolytic anaemia. The two HIV-positive patients with haemolysis and IgG_1 were noted to have severe anaemia (Hb <6 g/dL). In HIV-positive patients, the presence of IgG_1 may potentially identify patients with severe AIHA. In addition, the absence of IgG_1 in HIV-positive patients could suggest there is a false-positive DAT.

In our study, three of the patients with IgG_1 and haemolysis were positive for C3d on the DAT (n=26). Two of the three patients were HIV positive, cART naïve and had viral loads >100 000 copies/mL with concurrent features of sepsis. The remaining sample was from an HIV-negative patient with SLE.

Our findings suggest a possible relationship of IgG_1 with poorly controlled HIV and autoimmune haemolysis.[9]

A similar proportion of patients with and without AIHA had IgG concentration of 1:10. These low concentrations do not appear to be clinically significant. Three patients with warm AIHA had concentrations greater than 1:30 (3/26, 12%). Two of the three patients had an underlying diagnosis of SLE and severe anaemia. In HIV-positive patients with or without haemolysis, none of the concentrations of IgG were greater than 1:30. In our study of HIV-positive patients with false-positive DAT, quantification of IgG did not yield more useful information.

In conclusion, our study highlights that the demographics and profile of comorbid diseases of South African patients with IgG-positive DAT anaemia differ from other studies. The presence of DAT positivity in a patient (especially those with HIV) still requires the correlation of both laboratory and clinical findings. Despite the apparent association between absence of IgG_1 and/or IgG_3 and the absence of AIHA in HIV-positive patients, the overall benefit in both subtyping and quantification in HIV-positive patients remains limited.

STUDY LIMITATIONS

This study has the limitations of a retrospective cross-sectional study. Our study had a small sample size. Either due to lack of clinical information or haphazard ordering of biochemical tests by treating clinicians, some data points were not available. This further reduced the amount of usable data.

FURTHER RESEARCH

A prospective study in patients with HIV infection, with and without AIHA, to study the presence of IgG_1 or IgG_3 is required to ascertain if subtyping would be of more benefit in this subpopulation of patients. Additionally, larger, prospective studies in South African hospitals are encouraged to broaden the information on patients with HIV infection, with and without cART and positive DAT results.

Take home messages

- The absence of IgG_1/IgG_3 may be helpful in excluding the presence of severe haemolysis in patients with HIV and a positive DAT.
- The subtyping and quantification of IgG in the patients with a positive DAT and HIV have limited value.

Handling editor Mary Frances McMullin.

Contributors TC—study design, data collection, data analysis and paper writing. NB, YOW and JM—study design, data analysis, critical review and paper writing. All authors approved the submitted and final version of the manuscript.

Funding This work was funded by the National Health Laboratory Service (NHLS) K-project (KNC 180).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This cross-sectional, retrospective study conducted in a quaternary care hospital was reviewed and approved by the Human Research Ethics Committee of the University of the Witwatersrand.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

ORCID iD

Thirosha Chetty http://orcid.org/0000-0002-7749-6169

REFERENCES

1 Hoffbrand VA, Higgs DR, Keeling DM, *et al*. *Postgraduate haematology*. Seventh ed. United Kingdom: John Wiley & Sons Ltd, 2016.
2 Jäger U, Barcellini W, Broome CM, *et al*. Diagnosis and treatment of autoimmune hemolytic anemia in adults: recommendations from the first international consensus meeting. *Blood Rev* 2020;41:100648.
3 Salama A. Clinically and/or serologically misleading findings surrounding immune haemolytic anaemias. *Transfus Med Hemother* 2015;42:311–5.
4 Baisley K, Chimbindi N, Mthiyane N, *et al*. High HIV incidence and low uptake of HIV prevention services: the context of risk for young male adults prior to dreams in rural KwaZulu-Natal, South Africa. *PLoS One* 2018;13:e0208689.
5 Opie J. Haematological complications of HIV infection. *S Afr Med J* 2012;102:465–8.
6 Yen Y-F, Lan Y-C, Huang C-T, *et al*. Human immunodeficiency virus infection increases the risk of incident autoimmune hemolytic anemia: a population-based cohort study in Taiwan. *J Infect Dis* 2017;216:1000–7.
7 Russo S, Lopalco L. Is autoimmunity a component of natural immunity to HIV? *Curr HIV Res* 2006;4:177–90.
8 Tsiakalos A, Routsias JG, Kordossis T, *et al*. Fine epitope specificity of anti-erythropoietin antibodies reveals molecular mimicry with HIV-1 p17 protein: a pathogenetic mechanism for HIV-1-related anemia. *J Infect Dis* 2011;204:902–11.
9 De Angelis V, Biasinutto C, Pradella P, *et al*. Clinical significance of positive direct antiglobulin test in patients with HIV infection. *Infection* 1994;22:92–5.
10 Toy PT, Reid ME, Burns M. Positive direct antiglobulin test associated with hyperglobulinemia in acquired immunodeficiency syndrome (AIDS). *Am J Hematol* 1985;19:145–50.
11 Lai M, d'Onofrio G, Visconti E, *et al*. Aetiological factors related to a positive direct antiglobulin test result in human immunodeficiency virus-infected patients. *Vox Sang* 2006;90:325–30.
12 Ukita M, Takahashi A, Nunotani T, *et al*. IgG subclasses of anti-A and anti-B antibodies bound to the cord red cells in ABO incompatible pregnancies. *Vox Sang* 1989;56:181–6.
13 Dubarry M, Charron C, Habibi B, *et al*. Quantitation of immunoglobulin classes and subclasses of autoantibodies bound to red cells in patients with and without hemolysis. *Transfusion* 1993;33:466–71.
14 Das SS, Nityanand S, Chaudhary R. Clinical and serological characterization of autoimmune hemolytic anemia in a tertiary care hospital in North India. *Ann Hematol* 2009;88:727–32.
15 Singh A, Solanki A, Chaudhary R. Demonstration of IgG subclass (IgG1 and IgG3) in patients with positive direct antiglobulin tests. *Immunohematology* 2014;30:24–7.
16 Garratty G. Effect of cell-bound proteins on the in vivo survival of circulating blood cells. *Gerontology* 1991;37:68–94.
17 Achargui S, Benchemsi N. [A quantitative determination of IgG anti-D subclasses by Elisa in hemolytic disease of the newborn]. *Transfus Clin Biol* 2003;10:284–91.
18 Lai M, Leone G, Landolfi R. Autoimmune hemolytic anemia with gel-based immunohematology tests. *Am J Clin Pathol* 2013;139:457–63.
19 Lynen R, Krone O, Legler TJ, *et al*. A newly developed gel centrifugation test for quantification of RBC-bound IgG antibodies and their subclasses IgG1 and IgG3: comparison with flow cytometry. *Transfusion* 2002;42:612–8.
20 Fabijańska-Mitek J, Lopieńska H, Zupańska B. Gel test application for IgG subclass detection in auto-immune haemolytic anaemia. *Vox Sang* 1997;72:233–7.
21 Kruhonja Galic Z, Jagnjic S, Bingulac-Popovic J, *et al*. Warm red blood cell autoantibodies and clinical diagnoses in patients with or without autoimmune hemolysis. *Transfus Clin Biol* 2020;27:25–9.
22 Barcellini W, Fattizzo B, Zaninoni A, *et al*. Clinical heterogeneity and predictors of outcome in primary autoimmune hemolytic anemia: a GIMEMA study of 308 patients. *Blood* 2014;124:2930–6.
23 Alonso H-C, Manuel A-AV, Amir CGC, *et al*. Warm autoimmune hemolytic anemia: experience from a single referral center in Mexico City. *Blood Res* 2017;52:44–9.
24 Gehrs BC, Friedberg RC. Autoimmune hemolytic anemia. *Am J Hematol* 2002;69:258–71.
25 Zantek ND, Koepsell SA, Tharp DR, *et al*. The direct antiglobulin test: a critical step in the evaluation of hemolysis. *Am J Hematol* 2012;87:707–9.
26 Lepennec PY, Lefrere JJ, Rouzaud AM, *et al*. Red cell autoantibodies in asymptomatic HIV-infected subjects. *Transfusion* 1989;29:465–6.
27 Garratty G, Nance SJ. Correlation between in vivo hemolysis and the amount of red cell-bound IgG measured by flow cytometry. *Transfusion* 1990;30:617–21.