

**DETERMINANTS OF DAY-NIGHT DIFFERENCE IN
BLOOD PRESSURE IN SUBJECTS OF AFRICAN
ANCESTRY**

Joseph Muzi Maseko

**A dissertation submitted to the Faculty of Health Sciences, University of the
Witwatersrand, in fulfilment of the requirements for the degree of
Master of Science**

Johannesburg, 2008

DECLARATION

I, Joseph Muzi Maseko declare that this dissertation is my own work except to the extent indicated in the acknowledgements and references. It is being submitted for the degree of Master of Science in Medicine in the field of Physiology at the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at this or any other university.

Joseph Muzi Maseko.....

.....day of, 2008.

I certify that the studies contained in this dissertation have the approval of the Committee for Research in Human Subjects of the University of the Witwatersrand, Johannesburg. The ethics approval number is M02-04-72

.....

Joseph Muzi Maseko

.....

Gavin R. Norton (supervisor)

.....

Date

.....

Angela J. Woodiwiss (supervisor)

.....

Date

This dissertation is dedicated to my wife Mokgadi and my three children

Thembi, Zinhle and Sibusiso

PUBLICATIONS AND PRESENTATIONS

The following is a list of publications and presentations offered in support of this dissertation.

Muzi J Maseko, Harold Majane, John Milne, Gavin R Norton, Angela Woodiwiss. Salt intake in an urban, developing South African community. *Cardiovascular Journal of South Africa* 2006; 17;186-191.

Muzi Maseko, Angela J Woodiwiss, Lutgarde Thijs, Jan Staessen, Gavin R Norton. Day-night differences in ambulatory blood pressure in subjects of African as compared to European descent. Oral presentation at the annual congress of the Physiology Society of Southern Africa 2005, Cape Town, South Africa 7-9 September.

Muzi Maseko, Angela J Woodiwiss, Lutgarde Thijs, Jan Staessen, Gavin R Norton. Determinants of day-night difference in ambulatory blood pressure in subjects of African ancestry. Poster presentation at the annual congress of the Physiology Society of Southern Africa 2005, Cape Town, South Africa 7-9 September.

ABSTRACT

Hypertension is a major risk factor for cardiovascular disease in both developed and developing countries. Blood pressure normally decreases at night and a number of studies have indicated that a reduced nocturnal decline in blood pressure (BP) increases the risk for cardiovascular disease. Nocturnal decreases in BP are attenuated in subjects of African as compared to European descent, but the mechanisms of this effect require clarity. In the present study I attempted to identify potentially modifiable factors that contribute toward nocturnal decreases in BP in a random sample of 171 nuclear families comprising 438 black South Africans living in Soweto.

Prior studies have suggested that adiposity and salt intake may determine nocturnal decreases in BP. Adiposity and salt intake were considered to be potentially important factors to consider in the present study as 67% of the group studied were either overweight or obese and in 291 subjects that had complete 24-hour urine collections (used to assess salt intake) and BP measurements, Na^+ and K^+ intake was noted to be considerably higher and lower respectively than the recommended daily allowance in the majority of people. Moreover, a lack of relationship between either hypertension awareness and treatment and Na^+ and K^+ intake suggested that current recommendations for a reduced Na^+ intake and increased K^+ intake in hypertensives do not translate into clinical practice in this community.

In order to assess whether adiposity or salt intake are associated with nocturnal decreases in BP in this community, ambulatory BP monitoring was performed using Spacelabs model 90207 oscillometric monitors. Of the 438 subjects recruited, 314 had

ambulatory BP measurements that met pre-specified quality criteria (more than 20 hours of recordings and more than 10 and 5 readings for the computation of daytime and nighttime means respectively). To identify whether adiposity or salt intake are associated with a reduced nocturnal decline in BP, non-linear regression analysis was employed with indices of adiposity and urinary Na^+ and K^+ excretion rates and urine $\text{Na}^+ : \text{K}^+$ ratios included in the regression model with adjustments for potential confounders. Neither body mass index, skin-fold thickness, waist circumference, waist-to hip ratio, urinary Na^+ and K^+ excretion rates, nor urine $\text{Na}^+ : \text{K}^+$ ratios were associated with nocturnal decreases in systolic and diastolic BP. Indices of adiposity were however associated with 24 hour ambulatory systolic and diastolic BP. Unexpectedly, female gender was associated with an attenuated nocturnal decrease in BP.

In conclusion, in the first random, community-based sample with large sample sizes conducted with ambulatory BP monitoring in Africa, I found that neither adiposity nor salt intake are associated with a reduced nocturnal decline in BP. The lack of association between either salt intake or adiposity and nocturnal decreases in BP was despite a high prevalence of excessive adiposity in the community, as well as clear evidence that current recommendations for a reduced Na^+ intake and increased K^+ intake do not translate into clinical practice in this community. Thus, based on this study, the question arises as to whether primordial prevention programs targeting excess adiposity or inappropriate salt intake are likely to modify nocturnal decreases in BP, in urban, developing communities of African ancestry in South Africa. However, unexpectedly I noted that females were more likely to have an attenuated nocturnal decrease in BP. Thus further work is required to explain this finding.

ACKNOWLEDGEMENTS

A special word of thanks goes to my supervisor Professor Gavin Norton whose profound knowledge of pathophysiology led to my completing this dissertation and to my co-supervisor Professor Angela Woodiwiss whose in-depth knowledge of statistics and insight guided me through all of the stages of data analysis. This project would not have been possible without the voluntary participation of nuclear families who were recruited in the framework of the African Project on Genes in Hypertension, which was supported by the International Scientific and Technological Cooperation between South Africa and Flanders (contract number BIL 01/43), the Medical Research Council of South Africa, the National Research Foundation (Women in Research) and the Circulatory Disorders Research Trust. I am also very grateful for the excellent technical assistance of Mthuthuzeli Kiviet, Nkele Maseko and Nomonde Molebatsi.

TABLE OF CONTENTS

	Page
DECLARATION	ii
DEDICATION	iii
PUBLICATIONS AND PRESENTATIONS	iv
ABSTRACT	v
ACKNOWLEDGEMENTS	vii
TABLE OF CONTENTS	viii
LIST OF FIGURES	xi
LIST OF TABLES	xii
LIST OF ABBREVIATIONS	xiv
1. INTRODUCTION	1
1.1 Cardiovascular disease risk factors	1
1.1.1 Ethnic differences in cardiovascular diseases	4
1.1.2 Hypertension as an important cause of cardiovascular disease	6
1.2 How well is hypertension being treated	6
1.2.1 Ethnic differences in the prevalence of hypertension	7
1.2.2 Potential factors accounting for ethnic differences in the prevalence of hypertension	8
1.3 Are blood pressures measured during epidemiological studies appropriate	10
1.3.1 Ambulatory versus conventional blood pressure measurements	11
1.3.2 Nocturnal decreases in BP as a determinant of cardiovascular disease	12

1.3.3	Ethnic differences in ambulatory BP	15
1.3.4	Determinants of nocturnal decreases in BP	18
1.3.5	Potential determinants of ethnic differences in nocturnal decreases in BP	20
1.3.6	Optimal thresholds for salt intake	23
1.3.7	Objectives of the dissertation	25
2	METHODS	27
2.1	Subjects	28
2.2	Questionnaire	30
2.3	Conventional blood pressure measurement	31
2.4	Anthropometric measurements	33
2.5	Blood measurements	35
2.6	Urinary sodium and potassium excretion	35
2.7	Ambulatory blood pressure measurements	36
2.8	Definitions and analysis	39
3	RESULTS	
3.1	Characteristics of the study group	42
3.2	Salt intake	45
3.2.1	General demographic and clinical characteristics of the participants with appropriate 24-hour urine samples grouped according to hypertension status	45
3.2.2	Antihypertensive medication	46
3.2.3	Blood pressures	46

3.2.4	Electrolyte excretion rates	49
3.3	Determinants of 24-hour ambulatory BP and nocturnal decreases in BP	51
3.3.1	Ambulatory BP values in subjects with appropriate urine samples	51
3.3.2	Urinary salt excretion in subjects with appropriate 24-hour BP data	51
3.3.3	Determinants of ambulatory BP	55
3.3.4	Determinants of nocturnal decreases in ambulatory blood pressure	55
4	DISCUSSION AND CONCLUSION	60
4.1	Nocturnal decreases in BP in groups of African descent	61
4.2	Environmental and phenotypic factors other than adiposity or salt intake that may influence nocturnal decreases in BP	63
4.3	Salt intake in the study group	64
4.4	Relationship between salt intake or excess adiposity and ambulatory BP and nocturnal decreases in BP	68
4.5	Relationship between adiposity and ambulatory BP and nocturnal decreases in BP	70
4.6	Relationship between female gender and attenuated nocturnal decreases in BP	72
4.7	Potential limitations and future studies and analysis	73
4.8	Conclusion	75

LIST OF FIGURES	Page
1. Example of BP values obtained using 24 hour ABPM	13
2. Summary of proposed mechanisms responsible for attenuated decrease in nocturnal BP in people of African descent	21
3. Examples of formal and informal dwellings in the suburban regions where people were recruited for this study	29
4. Hypertension clinic in the School of Physiology	34
5. Spacelabs model 90207 ambulatory BP monitors and representative data obtained for analysis purposes	37
6. The relationship between urine sodium excretion and night-day difference	57
7. The relationship between urine potassium excretion and night-day difference	58
8. The relationship between urine sodium to potassium ratio and night-day difference	59
9. Relationship between BMI and night-day difference	60

LIST OF TABLES

Table	Page
1. Summary of studies assessing the impact of ethnicity on nocturnal decrease in BP.	16
2. Demographic and anthropometric characteristics of all study subjects and study subjects with appropriate ambulatory BP monitoring, urinary salt excretion and both ambulatory BP and urinary salt excretion data	43
3. Clinical characteristics of all study subjects and study subjects with appropriate ambulatory BP monitoring, urinary salt excretion and both ambulatory BP and urinary salt excretion data	44
4. Demographic, anthropometric and clinical characteristics of study subjects with appropriate 24-hour urine samples	47
5. Blood pressures and antihypertensive therapy of study subjects with appropriate 24-hour urine samples.	48
6. Urinary values of study subjects	50
7. Ambulatory blood pressures in all subjects with 24-hour ambulatory BP data and subjects with 24-hour urine excretion data	52
8. Indexes of nocturnal decreases in blood pressure in all subjects with 24-hour ambulatory BP data and subjects with 24-hour ambulatory BP and 24-hour urine data	53
9. Urinary values of all study subjects and study subjects with appropriate 24-hour ambulatory BP values.	54

10. Determinants of 24-hour blood pressure in study subjects	56
11. Determinants of day-night differences in blood pressure in study subjects	57
12. Determinants of the ratio of night to day blood pressure in study subjects	58

LIST OF ABBREVIATIONS

24DBP	24-hour Diastolic Ambulatory Blood Pressure
24SBP	24-hour Systolic Ambulatory Blood Pressure
ABP	Ambulatory Blood Pressure
ANOVA	Analysis of variance
APOGH	African Project On Genes in Hypertension
BMI	Body Mass Index
BP	Blood Pressure
DBP	Diastolic Blood Pressure
GLP	Good Laboratory Practice
HbA _{1C}	Glycated Haemoglobin
HDL	High Density Lipoprotein
IHD	Ischemic Heart Disease
LDL	Low Density Lipoprotein
LVM	Left Ventricular Mass
MI	Myocardial Infarction
NHANES	National Health And Nutrition Examination Surveys
NHSL	National Health System Laboratories
PVD	Peripheral Vascular Disease
RDA	Recommended Daily Allowance
SAS	Statistical Analysis Software
SE	Socio-economic

TG	Triacylglycerides
TOD	Target Organ Damage
U _{Na+}	Urine Sodium Concentration
U _{K+}	Urine Potassium Concentration
UK	United Kingdom
USA	United States of America