

https://dspace.org/

Semiproductsproductsand modal predicate logicsSome examples

V.B Shehtman; Dmitry Shkatov

MAIK NAUKA/INTERPERIODICA/SPRINGER
https://hdl.handle.net/10539/41240
WOS

Downloaded from a DSpace repository. https://dspace.org/

ResearchGate

See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/377864772

Semiproducts, Products, and Modal Predicate Logics: Some Examples

Article *in* Doklady Mathematics · January 2024

DOI: 10.1134/S1064562423701296

CITATION
1

READS
10

2 authors, including:



Dmitry Shkatov
University of the Witwatersrand
66 PUBLICATIONS **478** CITATIONS

SEE PROFILE

All content following this page was uploaded by Dmitry Shkatov on 04 February 2024.

The user has requested enhancement of the downloaded file.

Semiproducts, products, and modal predicate logics: some examples*

Valentin Shehtman[a] and Dmitry Shkatov[b]

[a]Steklov Mathematical Institute, Russian Academy of Sciences,
vshehtman@gmail.com
[b]University of the Witwatersrand, Johannesburg, South Africa,
shkatov@gmail.com

Abstract

We study two kinds of combined modal logics, semiproducts and products with **S5**, and their correlation with modal predicate logics. We present examples of propositional modal logics for which semiproducts or products with **S5** are axiomatized in the minimal way (they are called semiproduct- or product-matching with **S5**) and also present counterexamples for these properties. The finite model property for (semi)products, together with (semi)product-matching, allow us to show decidability of corresponding 1-variable modal predicate logics.

Keywords: semiproducts of modal logics, products of modal logics, predicate modal logic.

1 Introduction

We study two kinds of combined modal logics, products and semiproducts (also known as expanding products). Products were introduced in the 1970s to formalise reasoning about multiple independent modalities [1, 2]; a comprehensive investigation of the field was undertaken in [3]; some later developments were presented in [4].

So far the study of semiproducts has been lagging behind the study of products. For example, while there is a general theorem on product-matching [3,

*A prefinal draft of an article to appear in *Doklady. Mathematics*. The published version is available at https://doi.org/10.1134/S1064562423701296

Theorem 5.9], an analogous theorem for semiproducts [3, Theorem 9.10] is much weaker. In many cases, properties of semiproducts are unknown.

In this paper, we are interested in semiproducts and products with **S5**. Our interest is primarily motivated by their connection with modal predicate logics, noted by Fischer-Servi [5]. This connection is, in fact, a bimodal version of the well-known M. Wajsberg's interpretation of **S5** as a single-variable classical predicate logic (see, e.g., [3, Subsection 1.3]).

To study axiomatization of these semiproducts and products and the relationship of these logics with 1-variable modal predicate logics, we introduce a classification of propositional modal logics and give examples for some categories of the proposed classification.

In particular, we consider logics of finite depth, in the sense of [6], with the axiom of thickness corresponding to the Horn condition

$$\forall x, y, z, t\,(xRy \wedge xRz \wedge yRt \to zRt).$$

We show that semiproducts and products of such logics with **S5** are axiomatized in the minimal way and are decidable. Moreover, they have the product and semiproduct finite model property. This implies decidability and the finite model property for corresponding 1-variable predicate logics.

2 Preliminaries

2.1 Propositional modal logics

We consider N-modal propositional formulas constructed from a countable set $PL = \{p_1, p_2, p_3, \dots\}$ of proposition letters, the constant $\bot$, the connective $\to$, and unary modalities $\Box_1, \dots, \Box_N$. In this paper, $N \in \{1, 2\}$.

We use lowercase letters $p, q, r, \dots$ for proposition letters and uppercase letters $A, B, C, \dots$ for formulas. We use the standard abbreviations $\top$, $\neg A$, $A \wedge B$, $A \vee B$, $A \leftrightarrow B$, $\Diamond_i A$ and the iterated modalities $\Box_i^n$ and $\Diamond_i^n$. The modality of the 1-modal language is usually denoted by $\Box$.

A *k-formula* is a formula containing only proposition letters from the set $\{p_1, p_2, \dots, p_k\}$. A 0-formula (i.e., a formula without propositional letters) is called *closed*.

An *N-modal propositional logic* is a set of N-modal formulas containing the Boolean tautologies and formulas of the form $\Box_i(p \to q) \to (\Box_i p \to \Box_i q)$ and closed under Substitution, Modus Ponens, and Necessitation. The smallest such logic is called $\mathbf{K}_N$; also, $\mathbf{K} := \mathbf{K}_1$.

If $\mathbf{\Lambda}$ is an N-modal logic and A an N-modal formula, then $\mathbf{\Lambda} \vdash A$ means the same as $A \in \mathbf{\Lambda}$. The smallest logic including a logic $\mathbf{\Lambda}$ and a set Γ of formulas is denoted by $\mathbf{\Lambda} + \Gamma$; we write $\mathbf{\Lambda} + A$ instead of $\mathbf{\Lambda} + \{A\}$.

The *fusion* $\mathbf{\Lambda}_1 * \mathbf{\Lambda}_2$ of 1-modal logics $\mathbf{\Lambda}_1$ and $\mathbf{\Lambda}_2$ is the $\mathbf{K}_2 + \mathbf{\Lambda}_1 \cup \mathbf{\Lambda}_2^{+1}$, where $\mathbf{\Lambda}_2^{+1}$ is obtained from $\mathbf{\Lambda}_2$ by replacing every occurrence of $\Box_1$ with $\Box_2$.

We use standard definitions from Kripke semantics. An *N-frame* is a tuple $F = (W, R_1, \ldots, R_N)$ where $W \neq \varnothing$ and $R_1, \ldots, R_N \subseteq W^2$; elements of W are called *points*. A *Kripke model over* F is a pair $M = (F, \theta)$ where $\theta \colon PL \to 2^W$. The truth relation between points w of a modal M and formulas is defined by recursion; in particular,

- $M, w \models p_i$ if $w \in \theta(p_i)$;
- $M, w \models \Box_i A_1$ if $M, w' \models A_1$ whenever wR_iw'.

A formula A is (globally) *true in a model* M (in symbols, $M \vDash A$) if $M, w \vDash A$, for every $w \in W$. A formula A is *valid on* a frame F (in symbols, $F \vDash A$) if $M \vDash A$, for every model M over F.

If Γ is a set of formulas, $\mathbf{V}(\Gamma)$ denotes the class of frames validating Γ; if A is a formula, we write $\mathbf{V}(A)$ instead of $\mathbf{V}(\{A\})$. If $\mathbf{\Lambda}$ is a logic, then $\mathbf{V}(\mathbf{\Lambda})$ is said to be the class of *$\mathbf{\Lambda}$-frames*.

By soundness theorem, $\mathbf{V}(\Gamma) = \mathbf{V}(\mathbf{K}_N + \Gamma)$. Also, if F is an N-frame and $\mathcal{C}$ a class of N-frames, then $\mathbf{L}(F) := \{A \mid F \vDash A\}$ and $\mathbf{L}(\mathcal{C}) := \bigcap\{\mathbf{L}(F) \mid \boldsymbol{F} \in \mathcal{C}\}$ are N-modal logics. We say that the logic $\mathbf{L}(\mathcal{C})$ is *determined by* $\mathcal{C}$.

A logic is *Kripke complete* if it is determined by some class of frames. A logic has the *finite model property* (fmp), if it is determined by a class of finite frames.

Lemma 2.1. *Let $\mathbf{\Lambda}_1$ and $\mathbf{\Lambda}_2$ be 1-modal logics and let $F = (W, R_1, R_2)$ be a Kripke frame. Then,*

$$F \vDash \mathbf{\Lambda}_1 * \mathbf{\Lambda}_2 \quad \Longleftrightarrow \quad (W, R_1) \vDash \mathbf{\Lambda}_1 \ \ \& \ \ (W, R_2) \vDash \mathbf{\Lambda}_2.$$

A frame $(W, R_1, \ldots, R_N)$ can also be viewed as a classical first-order model in the signature $\{R_1, \ldots, R_N, =\}$.

Definition 2.2. *A modal logic $\mathbf{\Lambda}$ is* elementary *if the class $\mathbf{V}(\mathbf{\Lambda})$ is definable by a classical first-order sentence. An N-modal formula A and a classical first-order sentence Φ in the signature $\{R_1, \ldots, R_N, =\}$ are* correspondents *if the class $\mathbf{V}(A)$ is definable by Φ.*

Definition 2.3. *A* universal Horn sentence *is a classical first-order sentence in the signature* $\{R_1, \ldots, R_N, =\}$ *of the form*

$$\forall x \, \forall y \, \forall \bar{z} \, (\Phi(x, y, \bar{z}) \to R_i(x, y)),$$

where $\Phi(x, y, \bar{z})$ *is a conjunction of atomic formulas.*

An N*-modal formula* A *is* Horn *if it corresponds to a universal Horn sentence.*

Definition 2.4. *A 1-modal logic* $\mathbf{\Lambda}$ *is* Horn axiomatizable *if* $\mathbf{\Lambda} = \mathbf{K} + \Gamma$*, for some set* Γ *of formulas that are either Horn or closed.*

Definition 2.5. *A* cone *of a frame* $F = (W, R_1, \ldots, R_N)$ *at a point* w*, denoted by* $F{\uparrow}w$*, is the restriction of* F *to the set* $(R_1 \cup \ldots \cup R_N)^*(w)$*, where* S^* *denotes the reflexive transitive closure of a binary relation* S*.*

If $F = F{\uparrow}w$*, then* F *is said to be* rooted at w*.*

The following is well known:

Lemma 2.6. *Let* F *be a Kripke frame with a set* W *of points. Then,*

$$\mathbf{L}(F) \quad = \quad \bigcap_{w \in W} \mathbf{L}(F{\uparrow}w).$$

Definition 2.7. *A 1-frame* (W, R) *is* n-transitive *if* $R^{n+1} \subseteq \bigcup\limits_{m \leqslant n} R^m$*. An* N*-frame* $(W, R_1, \ldots, R_N)$ *is* n-transitive *if the 1-frame* $(W, R_1 \cup \ldots \cup R_N)$ *n-transitive.*

Let $F = (W, R_1 \cup \ldots \cup R_N)$ be an N-frame and $R = R_1 \cup \ldots \cup R_N$. Note that the points from $R^*(w)$ are path-accessible from w. If F is n-transitive, then all these points are accessible from w in at most n steps, i.e.,

$$R^*(w) \quad = \quad \bigcup_{m \leqslant n} R^m(w).$$

Definition 2.8. *A p-morphism from an* N*-frame* $(W, R_1, \ldots, R_N)$ *onto an* N*-frame* $(W', R'_1, \ldots, R'_N)$ *is a surjective map* $f : W \longrightarrow W'$ *satisfying the following conditions:*

- $xR_iy \Rightarrow f(x)R'_if(y)$ *(lift property),*
- $f(x)R'_iz \Rightarrow \exists y \, (f(y) = z \,\&\, xR_iy)$ *(monotonicity).*



Figure 1. Frame F_0 (with universal R_2) and model M_0

We consider the following 1-modal formulas and logics (here, $n \geqslant 1$):

$$\begin{array}{rcl@{\qquad}rcl} \mathit{det} & := & \Diamond p \leftrightarrow \Box p; & \mathit{ref} & := & \Box p \to p; \\ \mathit{sym} & := & \Diamond\Box p \to p; & \mathit{4} & := & \Box p \to \Box\Box p; \\ \mathit{5} & := & \Diamond\Box p \to \Box p; & \mathit{alt}_n & = & \neg \bigwedge\limits_{0\leqslant i\leqslant n} \Diamond(p_i \wedge \bigwedge\limits_{j\neq i} \neg p_j); \\ \mathit{Ath} & := & \Diamond\Diamond p \to \Box\Diamond p. & & & \end{array}$$

$$\begin{array}{rcl@{\qquad}rcl} \mathbf{T} & := & \mathbf{K} + \mathit{ref}; & \mathbf{K4} & := & \mathbf{K} + \mathit{4}; \\ \Box\cdot\mathbf{T} & := & \mathbf{K} + \Box\mathit{ref}; & \mathbf{SL4} & := & \mathbf{K4} + \mathit{det}; \\ \mathbf{K5} & := & \mathbf{K} + \mathit{5}; & \mathbf{K45} & := & \mathbf{K4} + \mathit{5}; \\ \mathbf{S5} & := & \mathbf{K4} + \mathit{ref} + \mathit{sym}; & \mathbf{Alt}_n & := & \mathbf{K} + \mathit{alt}_n; \\ \mathbf{K05} & := & \mathbf{K} + \mathit{Ath}. & & & \end{array}$$

We briefly mention the Kripke semantics of the lesser known of these logics. The logic $\Box\cdot\mathbf{T}$ is determined by the class of frames satisfying $\forall x\forall y(xRy \to yRy)$. The logic **SL4** is determined by the class of transitive and functional frames, and hence by a single frame where an irreflexive point sees a reflexive point (see Fig. 1). The logic $\mathbf{Alt}_n$ is determined by the frames (W, R) where $|R(w)| \leqslant n$ whenever $w \in W$.

Definition 2.9. *A 1-frame (W, R) is* thick *if $R^{-1} \circ R^2 \subseteq R$ or, equivalently,*

$$\forall x, y, z, u\,(xRy \,\&\, xRz \,\&\, yRu \Rightarrow zRu).$$

Lemma 2.10. *The class $\mathbf{V}(\mathit{Ath})(= \mathbf{V}(\mathbf{K05}))$ is the class of thick 1-frames. The logic* **K05** *is Kripke complete; hence, it is determined by the class of thick 1-frames.*

2.2 Products and semiproducts

Definition 2.11. *The* product *of 1-frames $F_1 = (W_1, R_1)$ and $F_2 = (W_2, R_2)$ is the 2-frame $F_1 \times F_2 = (W_1 \times W_2, R_h, R_v)$, where*

$$\begin{array}{rcl@{\quad}rcl} (x,y)R_h(x',y') & \iff & xR_1x' & \& & y = y'; \\ (x,y)R_v(x',y') & \iff & x = x' & \& & yR_2y'. \end{array}$$

A semiproduct *of F_1 and F_2 is a restriction of $F_1 \times F_2$ to some $W \subseteq W_1 \times W_2$ such that $R_h(W) \subseteq W$ (i.e., W is* horizontally closed*).*

Lemma 2.12. *If F is a semiproduct of F_1 and F_2, and x_i is a point of F_i (here, $i = 1, 2$), then $F{\uparrow}(x_1, x_2)$ is a semiproduct of $F_1{\uparrow}x_1$ and $F_2{\uparrow}x_2$.*

If $\mathcal{C}_1$ and $\mathcal{C}_2$ are classes of 1-frames, then we define

$$\begin{array}{lcl} \mathcal{C}_1 \times \mathcal{C}_2 & := & \{F_1 \times F_2 \mid F_1 \in \mathcal{C}_1 \text{ and } F_2 \in \mathcal{C}_2\}, \\ \mathcal{C}_1 \mathbin{\diagup} \mathcal{C}_2 & := & \{F \mid F \text{ is a semiproduct of some frames } F_1 \in \mathcal{C}_1 \text{ and } F_2 \in \mathcal{C}_2\}. \end{array}$$

Definition 2.13. *The* product $\boldsymbol{\Lambda}_1 \times \boldsymbol{\Lambda}$ *and the* semiproduct $\boldsymbol{\Lambda}_1 \mathbin{\diagup} \boldsymbol{\Lambda}_2$ *of 1-modal propositional logics $\boldsymbol{\Lambda}_1$ and $\boldsymbol{\Lambda}_2$ are defined as follows:*

$$\begin{array}{lcl} \boldsymbol{\Lambda}_1 \times \boldsymbol{\Lambda}_2 & := & \mathbf{L}(\mathbf{V}(\boldsymbol{\Lambda}_1) \times \mathbf{V}(\boldsymbol{\Lambda}_2)); \\ \boldsymbol{\Lambda}_1 \mathbin{\diagup} \boldsymbol{\Lambda}_2 & := & \mathbf{L}(\mathbf{V}(\boldsymbol{\Lambda}_1) \mathbin{\diagup} \mathbf{V}(\boldsymbol{\Lambda}_2)). \end{array}$$

We will make use of the following 2-modal formulas and their frame correspondents:

$$\begin{array}{lll} (chr) & \Diamond_2 \Box_1 p \to \Box_1 \Diamond_2 p & R_2^{-1} \circ R_1 \subseteq R_1 \circ R_2^{-1}; \\ (lcom) & \Box_1 \Box_2 p \to \Box_2 \Box_1 p & R_2 \circ R_1 \subseteq R_1 \circ R_2; \\ (rcom) & \Box_2 \Box_1 p \to \Box_1 \Box_2 p & R_1 \circ R_2 \subseteq R_2 \circ R_1. \end{array}$$

Definition 2.14. *We define the* semicommutator $\boldsymbol{\Lambda}_1 \sqsupset \boldsymbol{\Lambda}_2$ *and the* commutator $[\boldsymbol{\Lambda}_1, \boldsymbol{\Lambda}_2]$ *of 1-modal logics $\boldsymbol{\Lambda}_1$ and $\boldsymbol{\Lambda}_2$ as follows:*

$$\begin{array}{lcl} \boldsymbol{\Lambda}_1 \sqsupset \boldsymbol{\Lambda}_2 & := & \boldsymbol{\Lambda}_1 * \boldsymbol{\Lambda}_2 + chr + lcom; \\ [\boldsymbol{\Lambda}_1, \boldsymbol{\Lambda}_2] & := & \boldsymbol{\Lambda}_1 \sqsupset \boldsymbol{\Lambda}_2 + rcom. \end{array}$$

Lemma 2.15. *Let $\boldsymbol{\Lambda}$, $\boldsymbol{\Lambda}_1$, and $\boldsymbol{\Lambda}_2$ be 1-modal logics. Then,*

(1) $\boldsymbol{\Lambda}_1 \sqsupset \boldsymbol{\Lambda}_2 \subseteq [\boldsymbol{\Lambda}_1, \boldsymbol{\Lambda}_2] \subseteq \boldsymbol{\Lambda}_1 \times \boldsymbol{\Lambda}_2$.

(2) $\boldsymbol{\Lambda}_1 \mathbin{\diagup} \boldsymbol{\Lambda}_2 \subseteq \boldsymbol{\Lambda}_1 \times \boldsymbol{\Lambda}_2$.

(3) $\boldsymbol{\Lambda} \sqsupset \mathbf{S5} \subseteq \boldsymbol{\Lambda} \mathbin{\diagup} \mathbf{S5}$.

(4) $\boldsymbol{\Lambda} \sqsupset \mathbf{S5} = \boldsymbol{\Lambda} * \mathbf{S5} + lcom = \boldsymbol{\Lambda} * \mathbf{S5} + chr$.

Definition 2.16. *A semiproduct logic $\boldsymbol{\Lambda}_1 \mathbin{\diagup} \boldsymbol{\Lambda}_2$ has the* semiproduct fmp *if it is determined by a class of finite semiproduct frames. The* product fmp *is defined similarly.*

Remark 2.17. Obviously, the (semi)product fmp implies the fmp. The converse is not always true.

Definition 2.18. *1-modal logics $\boldsymbol{\Lambda}_1$ and $\boldsymbol{\Lambda}_2$ are* product-matching *if $\boldsymbol{\Lambda}_1 \times \boldsymbol{\Lambda}_2 = [\boldsymbol{\Lambda}_1, \boldsymbol{\Lambda}_2]$ and* semiproduct-matching *if $\boldsymbol{\Lambda}_1 \mathbin{\diagup} \boldsymbol{\Lambda}_2 = \boldsymbol{\Lambda}_1 \sqsupset \boldsymbol{\Lambda}_2$.*

The following is well known:

Theorem 2.19. *[3, Theorem 5.9]. If a logic* $\mathbf{\Lambda}$ *is Kripke complete and Horn axiomatizable, then* $\mathbf{\Lambda}$ *and* $\mathbf{S5}$ *are product-matching.*

Theorem 2.20. *[3, Theorem 9.10]. If* $\mathbf{\Lambda} \in \{\mathbf{K}, \mathbf{T}, \mathbf{K4}, \mathbf{S4}\}$*, then* $\mathbf{\Lambda}$ *and* $\mathbf{S5}$ *are semiproduct-matching.*

Note that, while Theorem 2.19 gives infinitely many examples of product-matching logics, Theorem 2.20 gives only four examples of semiproduct-matching logics.

2.3 Monadic modal predicate logics

We refer to monadic fragments of 1-modal predicate logics as *monadic modal predicate logics*. These are logics in the language containing a countable set $\{x_1, x_2, x_3, \dots\}$ of individual variables, a countable set $\{P_1^1, P_2^1, P_3^1, \dots\}$ of monadic predicate letters, a countable set $\{P_1^0, P_2^0, P_3^0, \dots\}$ of nullary predicate letters (i.e., proposition letters), and logical symbols $\bot$, $\to$, $\Box$, and $\forall$. Formulas are defined as usual.

A *monadic modal predicate logic* is a set of monadic modal predicate formulas that includes the propositional logic $\mathbf{K}$ and the monadic classical predicate tautologies and is closed under Predicate Substitution, Modus Ponens, Generalisation, and Necessitation. The minimal such logic will be called[1] $\mathbf{QK}$. If $\mathbf{\Lambda}$ is a propositional 1-modal logic, then $\mathbf{Q\Lambda} := \mathbf{QK} + \mathbf{\Lambda}$ and $\mathbf{Q\Lambda C} := \mathbf{Q\Lambda} + Ba$, where $Ba := \forall x \Box P(x) \to \Box \forall x P(x)$ is the Barcan formula.

A *predicate Kripke frame* over a Kripke frame $F = (W, R)$ is a pair $\boldsymbol{F} = (F, D)$, where $D = (D_w)_{w \in W}$, with $D_w \neq \varnothing$ for all w and with $D_w \subseteq D_v$ whenever wRv.

A *valuation* on $\boldsymbol{F}$ is a family $\xi = (\xi_w)_{w \in W}$ of local valuations: $\xi_w(P_k^1) \subseteq D_w$ and $\xi_w(P_k^0) \in \{0, 1\}$. A *predicate Kripke model* over $\boldsymbol{F}$ is a pair $M = (\boldsymbol{F}, \xi)$, where ξ is a valuation on $\boldsymbol{F}$.

The truth relation $\Vdash$ between points w of a predicate Kripke model M and D_w-sentences (i.e., sentences obtained from formulas by replacing parameters with elements of D_w) is defined by recursion:

- $M, w \Vdash P_k^0$ if $\xi_w(P_k^0) = 1$;
- $M, w \Vdash P_k^1(a)$ if $a \in \xi_w(P_k^1)$;

[1] Usually, $\mathbf{QK}$, $\mathbf{Q\Lambda}$, and $\mathbf{Q\Lambda C}$ denote modal logics in languages with predicates of any arity, but in this paper we use the same notation for logics in languages with only monadic and nullary predicate letters.

- $M, w \Vdash \forall x\, A_1(x)$ if $M, w \Vdash A_1(a)$ whenever $a \in D_w$,

and the clauses for $\bot$, $\to$, $\Box$ are as in the propositional case.

A modal predicate formula A is *true in* M (in symbols, $M \models A$) if $M, u \Vdash \overline{\forall} A$ whenever $w \in W$. The formula A is *valid* on a predicate Kripke frame $\boldsymbol{F}$ (in symbols, $\boldsymbol{F} \models A$) if $M \models A$ whenever M is a Kripke model over $\boldsymbol{F}$. If L is a predicate modal logic, an *L-frame* is a predicate frame $\boldsymbol{F}$ validating all formulas from L; in this case, we write $\boldsymbol{F} \vDash L$.

By Soundness theorem [14, Theorem 3.2.29], $\mathbf{ML}(\boldsymbol{F}) := \{A \mid \boldsymbol{F} \models A\}$ is a modal predicate logic (called *the logic of* $\boldsymbol{F}$). The *modal predicate logic of a class $\mathcal{C}$ of predicate frames* (or the logic *determined by* $\mathcal{C}$) is logic defined as follows:

$$\mathbf{ML}(\mathcal{C}) \quad := \quad \bigcap \{\mathbf{ML}(\boldsymbol{F}) \mid \boldsymbol{F} \in \mathcal{C}\};$$

such logics are said to be *Kripke complete*. Every predicate logic L has the least Kripke complete extension, called the *Kripke completion*, which is the logic $\widehat{L}$ of the class of all L-frames.

3 1-variable predicate modal logics, semiproducts, and products

Let us recall definitions of some classes of monadic predicate modal formulas:

- *1-parametric* formulas contain at most one parameter;
- *1-variable* formulas are monadic containing at most one (fixed) variable x;
- *pure 1-variable* formulas are 1-variable without proposition letters;
- in *monodic* formulas [7, 3] every subformula of the form $\Box A$ is 1-parametric.

Monadic monodic fragments (mm-fragments) of logics **QK**, **QT**, **QK4**, and **QS4** are decidable [7, Theorem 5.1].[2] Even though they are syntactically more restrictive, 1-variable fragments are as expressive as mm-fragments:

Lemma 3.1.

(1) *Every mm-formula is* **QK***-equivalent to a Boolean combination of 1-variable formulas.*

[2]These are probably the largest known decidable fragments of modal predicate logics; most of 2-variable fragments, even in signatures with a single monadic predicate letter, are undecidable [8].

(2) *every 1-parametric mm-formula is* **QK**-*equivalent to a 1-variable formula.*

Moreover, every 1-variable formula A in proposition letters $q_1, q_2, \ldots, q_n$ translates into a pure 1-variable formula

$$A_0 \quad := \quad [\forall x\, Q_1(x), \ldots, \forall x\, Q_n(x)/q_1, \ldots, q_n]\, A,$$

where $Q_1, \ldots, Q_n$ are monadic letters not occurring in A. Since, for every modal predicate logic,

$$L \vdash A \quad \Longleftrightarrow \quad L \vdash A_0,$$

we may assume that all 1-variable formulas are pure.

Furthermore, there exists a validity-preserving bijection $A \mapsto A_*$ between pure 1-variable modal predicate formulas and 2-modal propositional formulas:

$$\begin{array}{lcl} P_i(x)_* & := & p_i; \\ \bot_* & := & \bot; \\ (A \to B)_* & := & A_* \to B_*; \\ (\Box A)_* & := & \Box_1 A_*; \\ (\forall x\, A)_* & := & \Box_2 A_*. \end{array}$$

The *1-variable fragment* of a modal predicate logic L is the set

$$(L{-}1)^* \quad := \quad \{A \in L \mid A \text{ is a pure 1-variable formula}\}.$$

The *propositional counterpart* of $(L{-}1)^*$ is the set

$$L{-}1 \quad := \quad \{A_* \mid A \in L,\ A \text{ is a pure 1-variable formula}\}.$$

Loosely, we sometimes refer to the set $L{-}1$ as the 1-variable fragment of L.

Remark 3.2. The notion of Kripke completeness is also applicable to 1-variable fragments of predicate logics: $(L{-}1)^*$ is said to be Kripke complete if there exists a class $\mathcal{C}$ of predicate frames such that $(\mathbf{ML}(\mathcal{C}){-}1)^* = (L{-}1)^*$, or equivalently, $\mathbf{ML}(\mathcal{C}){-}1 = L{-}1$. Obviously, Kripke completeness of L implies Kripke completeness of $(L{-}1)^*$.

Lemma 3.3. *Let L be a modal predicate logic. Then,*

(1) *$L{-}1$ is a 2-modal propositional logic containing $\mathbf{K} \sqsupset \mathbf{S5}$.*

(2) *If $L \vdash Ba$, then $[\mathbf{K}, \mathbf{S5}] \subseteq L{-}1$.*

Propositon 3.4. *Let $\mathbf{\Lambda}$ be a propositional 1-modal logic. Then,*

(1) $\mathbf{\Lambda} \sqsupset \mathbf{S5} \subseteq \mathbf{Q\Lambda}{-}1 \subseteq \widehat{\mathbf{Q\Lambda}}{-}1 = \mathbf{\Lambda} \nearrow \mathbf{S5}$. *Hence, if* $\mathbf{Q\Lambda}$ *is Kripke complete, then*

$$\mathbf{\Lambda} \sqsupset \mathbf{S5} \subseteq \mathbf{Q\Lambda}{-}1 = \mathbf{\Lambda} \nearrow \mathbf{S5}.$$

(2) $[\mathbf{\Lambda}, \mathbf{S5}] \subseteq \mathbf{Q\Lambda C}{-}1 \subseteq \widehat{\mathbf{Q\Lambda C}}{-}1 = \mathbf{\Lambda} \times \mathbf{S5}$. *Hence, if* $\mathbf{Q\Lambda C}$ *is Kripke complete, then*

$$[\mathbf{\Lambda}, \mathbf{S5}] \subseteq \mathbf{Q\Lambda C}{-}1 = \mathbf{\Lambda} \times \mathbf{S5}.$$

Definition 3.5. *A 1-modal propositional logic* $\mathbf{\Lambda}$ *is called* quantifier-friendly *if* $\mathbf{Q\Lambda}{-}1 = \mathbf{\Lambda} \sqsupset \mathbf{S5}$ *and* Barcan-friendly *if* $\mathbf{Q\Lambda C}{-}1 = [\mathbf{\Lambda}, \mathbf{S5}]$.

Proposition 3.4(1) implies that there exist four possibilities for semiproducts:

(1S) $\mathbf{\Lambda} \sqsupset \mathbf{S5} = \mathbf{Q\Lambda}{-}1 = \mathbf{\Lambda} \nearrow \mathbf{S5}$,

(2S) $\mathbf{\Lambda} \sqsupset \mathbf{S5} = \mathbf{Q\Lambda}{-}1 \subset \mathbf{\Lambda} \nearrow \mathbf{S5}$,

(3S) $\mathbf{\Lambda} \sqsupset \mathbf{S5} \subset \mathbf{Q\Lambda}{-}1 = \mathbf{\Lambda} \nearrow \mathbf{S5}$.

(4S) $\mathbf{\Lambda} \sqsupset \mathbf{S5} \subset \mathbf{Q\Lambda}{-}1 \subset \mathbf{\Lambda} \nearrow \mathbf{S5}$.

(1S) means that $\mathbf{\Lambda}$ and $\mathbf{S5}$ are semiproduct-matching. Some logics $\mathbf{\Lambda}$ of this type are described in Theorem 2.19. Another set of examples is presented in Section 6.

(2S) means that $\mathbf{\Lambda}$ and $\mathbf{S5}$ are not semiproduct-matching, but $\mathbf{\Lambda}$ is quantifier-friendly. Examples are given in Section 4.

Examples for (3S) are the logics $\mathbf{Alt}_n$, as shown in Section 4. Examples for (4S) are not known.

Proposition 3.4(2) implies that there exist four possibilities for products:

(1P) $[\mathbf{\Lambda}, \mathbf{S5}] = \mathbf{Q\Lambda C}{-}1 = \mathbf{\Lambda} \times \mathbf{S5}$,

(2P) $[\mathbf{\Lambda}, \mathbf{S5}] = \mathbf{Q\Lambda C}{-}1 \subset \mathbf{\Lambda} \times \mathbf{S5}$,

(3P) $[\mathbf{\Lambda}, \mathbf{S5}] \subset \mathbf{Q\Lambda C}{-}1 = \mathbf{\Lambda} \times \mathbf{S5}$.

(4P) $[\mathbf{\Lambda}, \mathbf{S5}] \subset \mathbf{Q\Lambda C}{-}1 \subset \mathbf{\Lambda} \times \mathbf{S5}$.

(1P) means that $\mathbf{\Lambda}$ and $\mathbf{S5}$ are product-matching. Examples are well known, see Theorem 2.19. Examples for (3P) are logics $\mathbf{Alt}_n$, as shown in Section 4.

(2P) means that $\mathbf{\Lambda}$ and $\mathbf{S5}$ are not product-matching, but $\mathbf{\Lambda}$ is Barcan-friendly; examples are unknown. Examples for (4P) are also unknown.

4 Logics not semiproduct-matching with S5

The following result was first stated, without a proof, in [9]:

Theorem 4.1. *Let* $\mathbf{\Lambda}$ *be a propositional 1-modal logic such that* $\Box \cdot \mathbf{T} \subseteq \mathbf{\Lambda} \subseteq \mathbf{SL4}$. *Then,*

$$\mathbf{\Lambda} \sqsupset \mathbf{S5} \quad \subset \quad \mathbf{\Lambda} \sqsupset \mathbf{S5} + \Box_1\Box_2 ref_1 \quad \subseteq \quad \mathbf{\Lambda} \nearrow \mathbf{S5},$$

where $ref_1 = \Box_1 p \to p$. *Hence,* $\mathbf{\Lambda}$ *and* $\mathbf{S5}$ *are not semiproduct-matching.*

Proof. Let M_0 be the model from Fig. 1; then, $M_0, u \vDash \Diamond_1\Diamond_2\neg ref$ and $F_0 \vDash \mathbf{SL4} \sqsupset \mathbf{S5}$. Hence, $\Box_1\Box_2 ref_1 \notin \mathbf{SL4} \sqsupset \mathbf{S5}$, which proves that the first inclusion is proper.

In view of Lemma 2.15 (3), to prove the second inclusion, it is enough to show that $\Box_1\Box_2 ref_1 \in \Box \cdot \mathbf{T} \nearrow \mathbf{S5}$. This membership follows from the validity of the formula $\Box_1\Box_2 ref_1$ on every semiproduct of a $\Box \cdot \mathbf{T}$-frame with an $\mathbf{S5}$-frame. ■

Theorem 4.1 implies that the analogue of Theorem 2.19 does not hold for semiproducts (cf. Theorem 2.20); in particular, it gives us the following counterexamples:

Corollary 4.2. *Horn axiomatizable logics* $\Box \cdot \mathbf{T}$, $\mathbf{K5}$, *and* $\mathbf{K45}$ *are not semiproduct-matching with* $\mathbf{S5}$.

Moreover, the following nontrivial result is also true (stated in [10]; the proof is in preparation):

Theorem 4.3. *Every Kripke complete Horn axiomatizable logic is quantifier-friendly.*

Corollary 4.4. *The logics* $\Box \cdot \mathbf{T}$, $\mathbf{K5}$, *and* $\mathbf{K45}$ *satisfy (2S).*

Remark 4.5. A standard modal logic argument shows that there is a continuum of logics between $\Box \cdot \mathbf{T}$ and $\mathbf{SL4}$. Due to Theorem 4.1, this gives us a continuum of logics not satisfying (1S).

Remark 4.6. If $\mathbf{\Lambda}$ is a logic from the statement of Theorem 4.1, then $\mathbf{Q\Lambda}$ is Kripke incomplete [11, Theorem 5.11].

We next recall a well-known property of Jankov–Fine formulas X_G (this property is stated in [12] in a slightly different form):

Proposition 4.7. *Let G be a rooted n-transitive N-frame. Then there exists an N-modal formula X_G such that, for every n-transitive frame F, the following holds: $F \not\Vdash X_G$ iff there exists a p-morphism from some cone of F onto G.*

Theorem 4.8. *If $\mathbf{Alt}_n \subseteq \mathbf{\Lambda} \subseteq \mathbf{Alt}_n + \Box^2\bot$, where $n \geqslant 3$, then $\mathbf{\Lambda}$ and $\mathbf{S5}$ are neither semiproduct- nor product-matching.*

Proof. We sketch the proof for $n = 4$; the general case is argued similarly. Let $G = (W, R, S)$ be the frame depicted in Fig. 2, on the right (S-reflexive accessibilities are not drawn).



Figure 2. Frame G is not a p-morphic image of any $H \in \mathbf{V}(\mathbf{Alt}_4) \nearrow \mathcal{U}$

It is not hard to see that $G \models [\mathbf{Alt}_4 + \Box^2\bot, \mathbf{S5}]$. Hence, G is 3-transitive.[3] Let X_G be the Jankov–Fine formula of G, and let $A := \Box_1^2\bot \to X_G$. Surely, $G \models \Box_1^2\bot$, and, by Proposition 4.7, $G \not\models X_G$. Therefore, $G \not\models A$, and hence $A \notin [\mathbf{Alt}_n + \Box^2\bot, \mathbf{S5}]$.

On the other hand, $A \in \mathbf{Alt}_n \nearrow \mathbf{S5}$, since otherwise, by Proposition 4.7, G is a p-morphic image of a cone $F{\uparrow}(x_1, x_2)$, where F is a semiproduct of $F_1 \vDash \mathbf{Alt}_n$ and $F_2 \vDash \mathbf{S5}$. By Lemma 2.12, this cone is a semiproduct of $F{\uparrow}x_1$ and $F{\uparrow}x_2$.

[3] This fact can also be inferred from Fig. 2.

By Lemma 2.6, $F{\uparrow}x_1$ is an $\mathbf{Alt}_n$-frame. Since $F{\uparrow}x_2$ is a cone in an **S5**-frame, it is a *cluster* (a frame with a universal relation).

However, as we next show, G cannot be a p-morphic of a semiproduct of such frames. Indeed, suppose f is a required p-morphism with $f(x_0, y) = (0, 1)$. By the lift property, there exist points (x_i, y), with $1 \leqslant i \leqslant 4$, and (x_0, y') such that $f(x_i, y) = (i, 1)$, for $i \in \{1, \ldots, 4\}$, and $f(x_0, y') = (0, 2)$ (see Fig. 2). Then, by monotonicity, $f(x_0, y')Rf(x_2, y')$ and $f(x_2, y)S(x_2, y')$; hence, $f(x_2, y') = (1, 2)$. Similarly, $f(x_3, y') = (1, 2)$ and $f(x_4, y') = (4, 2)$. Hence, (x_1, y') is mapped to either $(2, 2)$ or $(3, 2)$, which means that one of these points is not in the range of f, in contradiction with f being a p-morphism. ■

Recall that a modal predicate logic L is *strongly Kripke complete* if every L-consistent theory is satisfiable at a point of a model over an L-frame. By using selective submodels of canonical models (the method described in [11]), we can obtain the following result (for details, see [13]):

Theorem 4.9. *Every logic* $\mathbf{QAlt}_n$ *is strongly Kripke complete.*

Since adding closed propositional formulas preserves strong Kripke completeness, the following is also true:

Corollary 4.10. *Every logic* $\mathbf{QAlt}_n + \Box^m\bot$*, with* $m \geqslant 2$*, is strongly Kripke complete.*

The correspondent of the propositional formula alt_n is a classical first-order universal sentence. Hence, by Tanaka–Ono theorem [14, Theorem 7.4.7], we obtain the following:

Theorem 4.11. *If*

$$\mathbf{\Lambda} \in \{\mathbf{Alt}_n \mid n \geqslant 1\} \cup \{\mathbf{Alt}_n + \Box^m\bot \mid n \geqslant 1, m \geqslant 2\},$$

then the logic $\mathbf{Q\Lambda C}$ *is strongly Kripke complete.*

This implies the following:

Theorem 4.12. *If* $\mathbf{\Lambda} = \mathbf{Alt}_n$ *or* $\mathbf{\Lambda} = \mathbf{Alt}_n + \Box^m\bot$*, with* $n \geqslant 3$ *and* $m \geqslant 2$*, then the logic* $\mathbf{\Lambda} \sqsupset \mathbf{S5}$ *satisfies* (*3S*) *and the logic* $\mathbf{\Lambda} \times \mathbf{S5}$ *satisfies* (*3P*)*.*

Proof. By Theorem 4.8, $\mathbf{\Lambda} \diagup \mathbf{S5} \neq \mathbf{\Lambda} \sqsupset \mathbf{S5}$ and $[\mathbf{\Lambda}, \mathbf{S5}] \neq \mathbf{\Lambda} \times \mathbf{S5}$. By Theorem 4.9, Corollary 4.10, and Proposition 3.4(1), $\mathbf{Q\Lambda} - 1 = \mathbf{\Lambda} \diagup \mathbf{S5}$. Hence, $\mathbf{\Lambda}_1 \sqsupset \mathbf{S5} \neq \mathbf{Q\Lambda} - 1$. Similarly, it follows, by Theorem 4.11 and Proposition 3.4(2), that $\mathbf{Q\Lambda C} - 1 = \mathbf{\Lambda} \times \mathbf{S5}$. Hence, $\mathbf{Q\Lambda C} - 1 \neq [\mathbf{\Lambda}, \mathbf{S5}]$. ■

Problem 4.13. *Suppose that* $\mathbf{\Lambda}$ *is a logic from Theorem 4.11. Axiomatize logics* $\mathbf{Q\Lambda} - 1$ $(= \mathbf{\Lambda} \diagup \mathbf{S5})$ *and* $\mathbf{Q\Lambda C} - 1$ $(= \mathbf{\Lambda} \times \mathbf{S5})$*.*

5 Local tabularity and modal depth

We now recall definitions and facts from [15] about N-modal formulas and logics.

Definition 5.1. *The* modal depth $md(A)$ *of an N-modal propositional formula A is the maximal number of nested occurrences of modal operators in A:*

$$\begin{array}{lcl} md(\bot) & := & 0; \\ md(p_j) & := & 0; \\ md(A \to B) & := & \max(md(A), md(B)); \\ md(\Box_i A) & := & md(A) + 1. \end{array}$$

Definition 5.2. *The* modal depth $md_{\mathbf{\Lambda}}(A)$ of a formula A in an N-modal logic $\mathbf{\Lambda}$ *is defined as follows:*

$$md_{\mathbf{\Lambda}}(A) := \min\{md(B) \mid \mathbf{\Lambda} \vdash A \leftrightarrow B\}.$$

The modal depth $md(\mathbf{\Lambda})$ of a logic $\mathbf{\Lambda}$ *is defined as follows:*

$$md(\mathbf{\Lambda}) := \begin{cases} \max\{md_{\mathbf{\Lambda}}(B) \mid B \text{ is an } N\text{-modal propositional formula}\} & \text{if exists;} \\ \infty & \text{otherwise.} \end{cases}$$

Definition 5.3. *An N-modal logic $\mathbf{\Lambda}$ is* locally tabular *if, for any finite k, there exist only finitely many N-modal k-formulas non-equivalent in $\mathbf{\Lambda}$.*

Propositon 5.4.

(1) *Every locally tabular logic has the fmp.*

(2) *Every propositional modal logic of finite modal depth is locally tabular.*

6 Logics semiproduct-matching with S5

In this section, we show that each logic $\mathbf{K05} + \Box^n\bot$ is semiproduct-matching with **S5** and that the corresponding semiproduct has the semiproduct fmp. We use the following nomenclature for logics:

$$\begin{array}{lcl} \mathbf{\Lambda}_{0n} & := & \mathbf{K05} + \Box^n\bot; \\ \mathbf{\Lambda}_n & := & \mathbf{\Lambda}_{0n} \sqsupset \mathbf{S5}; \\ \mathbf{\Lambda}'_n & := & \mathbf{\Lambda}_n + rcom = [\mathbf{\Lambda}_{0n}, \mathbf{S5}]. \end{array}$$

Theorem 6.1. *If $n \geqslant 1$, then $md(\mathbf{\Lambda}_n) \leqslant 2n - 1$.*

The proof uses bisimulation games; for more details, see [15, 16]. Thus, by Proposition 5.4, we obtain the following:

Corollary 6.2. *The logics $\mathbf{\Lambda}_n$ and $\mathbf{\Lambda}'_n$ have the fmp.*

To prove semiproduct-matching and the semiproduct fmp for $\mathbf{\Lambda}_n$, it suffices to construct p-morphisms from semiproducts of finite $\mathbf{\Lambda}_{0n}$-frames with clusters onto finite $\mathbf{\Lambda}_n$-cones. Similarly, to prove product-matching and the product fmp for $\mathbf{\Lambda}'_n$, it suffices to construct p-morphisms from products of finite $\mathbf{\Lambda}_{0n}$-frames with clusters onto finite $\mathbf{\Lambda}'_n$-cones. We construct the sought p-morphisms in a number of steps (Lemmas 6.6 – 6.10).

Definition 6.3. *Let $F = (W, R_1, \ldots, R_N)$ and $F' = (W', R'_1, \ldots, R'_N)$ be frames. A map $g\colon W \longrightarrow W'$ is a* strong homomorphism *from F to F' if, for every $w, v \in W$ and every i,*

$$wR_iv \iff g(w)R'_ig(v).$$

Lemma 6.4. *Every surjective strong homomorphism is a p-morphism and an elementary equivalence for formulas without equality.*

Thus, if $\mathbf{\Lambda}$ is elementary (with respect to a classical signature without equality), then the class $\mathbf{V}(\mathbf{\Lambda})$ is closed under strong homomorphic pre-images.

Definition 6.5. *Let $F = (W, R_1, R_2)$ be a $\mathbf{K} \sqsupset \mathbf{S5}$-frame.*

- *A* row *in F is an equivalence class under the relation $(R_1 \cup R_1^{-1})^*$.*
- *A* column *in F is an equivalence class under R_2.*
- *A* block *in F is a non-empty intersection of a row and a column.*
- *F is* organized *if, for every row U in F, the frame $(W, R_1) \restriction U$ is rooted.*
- *F is* equalized *if every column in F consists of blocks of the same cardinality.*
- *F is* straight *if all its blocks are singletons.*

Lemma 6.6 (on organizing). *Every finite rooted $\mathbf{\Lambda}_n$-frame is a strong homomorphic image of a finite rooted organized $\mathbf{\Lambda}_n$-frame; a similar fact holds for $\mathbf{\Lambda}'_n$-frames.*

Proof. Let $F = (W, R_1, R_2)$ be a finite rooted $\mathbf{\Lambda}_n$-frame. We say that a point $a \in W$ is R_1-*minimal* if $R_1^{-1}(a) = \varnothing$. We put

$$V := \{(a, x) \mid a \text{ is } R_1\text{-minimal and } aR_1^*x\}$$

and define relations S_i on V:

$$\begin{array}{lcl}(a,x)S_1(b,y) & \iff & a=b \ \&\ xR_1y,\\ (a,x)S_2(b,y) & \iff & xR_2y.\end{array}$$

Then (V,S_1,S_2) is rooted and organized, and the map $(a,x)\mapsto x$ is a required strong homomorphism onto F. ■

Lemma 6.7 (on equalizing). *Every finite rooted organized $\mathbf{\Lambda}_n$-frame is a strong homomorphic image of a finite rooted equalized $\mathbf{\Lambda}_n$-frame; a similar fact holds for $\mathbf{\Lambda}'_n$-frames.*

Proof. To equalize a frame $F=(W,R_1,R_2)$, we add extra points to blocks making the blocks of a column the same size. We use the fact that, for every pair of blocks β and γ in F and each $k\in\{1,2\}$,

$$(1)\qquad \exists x\in\beta\,\exists y\in\gamma\, xR_ky \iff \forall x\in\beta\,\forall y\in\gamma\, xR_ky.$$

Hence, we write $\beta R_k\gamma$ whenever there exist $x\in\beta$ and $y\in\gamma$ such that xR_ky. We replace each block β in F with a block β' whose cardinality is the largest for blocks in the column of β. We put $W':=\{\beta' \mid \beta \text{ is a block in } F\}$ and define, for every $x\in\beta'$, $y\in\gamma'$, and $k\in\{1,2\}$,

$$xR'_ky \iff \beta R_k\gamma.$$

Due to (1), each relation R'_k is well defined. Then the frame $F':=(W',R'_1,R'_2)$ is equalized. A surjective map sending each point of block β' in F' to some point of block β in F is a strong homomorphism. ■

Lemma 6.8 (on straightening). *Every finite rooted equalized $\mathbf{\Lambda}_n$-frame is a strong homomorphic image of a finite rooted straight $\mathbf{\Lambda}_n$-frame; a similar fact holds for $\mathbf{\Lambda}'_n$-frames.*

Proof. To straighten a frame $F=(W,R_1,R_2)$, we first construct a frame whose columns all have the size, say n, of the largest column in F. To that end, we put $W'=W\times n$ and define $R'_1\subseteq W'\times W'$ so that

$$(x,i)R'_1(y,j) \iff xR_1y \ \&\ i=j,$$

and $R'_2\subseteq W'\times W'$ so that, if β and γ are blocks from the same column of F, $x\in\beta$, and $y\in\gamma$, then, for fixed enumerations N_β of β and N_γ of γ,

$$(x,i)R'_2(y,j) \iff N_\beta(x)+i\equiv N_\gamma(y)+j \ (mod\ |\beta|).$$

Then the map $(x,i)\mapsto x$ is a strong homomorphism from $F'=(W',R'_1,R'_2)$ onto F. ■

From Lemmas 6.6–6.8 and 6.4 we immediately obtain the following:

Lemma 6.9. *Every finite rooted $\mathbf{\Lambda}_n$-frame is a p-morphic image of a finite rooted straight $\mathbf{\Lambda}_n$-frame; a similar fact holds for $\mathbf{\Lambda}'_n$-frames.*

Lemma 6.10. *Every finite rooted straight $\mathbf{\Lambda}_n$-frame is isomorphic to a semiproduct of an $\mathbf{\Lambda}_{0n}$-frame and a cluster; a similar fact holds for $\mathbf{\Lambda}'_n$-frames and products.*

Proof. If $F = (W, R_1, R_2)$ is a finite straight frame rooted at x_0, then F is isomorphic to a (semi)product of the frame $(R_1^*(x_0), R_1 \restriction R_1^*(x_0))$ and the cluster whose points are the rows of F. ■

Theorem 6.11.

(1) *The logics* $\mathbf{K05}{+}\Box^n\bot$ *and* $\mathbf{S5}$ *are both semiproduct-matching and product-matching.*

(2) *The logics* $(\mathbf{K05} + \Box^n\bot) \mathbin{\diagup} \mathbf{S5}$ *have the semiproduct fmp, and the logics* $(\mathbf{K05} + \Box^n\bot) \times \mathbf{S5}$ *have the product fmp.*

Proof. Let, as before, $\mathbf{\Lambda}_{0n} := \mathbf{K05} + \Box^n\bot$, $\mathbf{\Lambda}_n := \mathbf{\Lambda}_{0n} \sqsupset \mathbf{S5}$, and $\mathbf{\Lambda}_n := [\mathbf{\Lambda}_{0n}, \mathbf{S5}]$.

(1) Suppose $A \notin \mathbf{\Lambda}_n$. By Corollary 6.2, $\mathbf{\Lambda}_n$ has the fmp. Hence, A is refuted on a finite rooted $\mathbf{\Lambda}_n$-frame. By Lemmas 6.9 and 6.10, this frame is a p-morphic image of a semiproduct of a finite $\mathbf{\Lambda}_{0n}$-frame and a finite cluster. Since p-morphisms preserve validity of modal formulas, $A \notin \mathbf{\Lambda}_{0n} \mathbin{\diagup} \mathbf{S5}$. Thus, $\mathbf{\Lambda}_{0n} \mathbin{\diagup} \mathbf{S5} \subseteq \mathbf{\Lambda}_n$. The converse is given by Lemma 2.15(3).

The proof for $\mathbf{\Lambda}'_n$ is similar.

(2) Since the (semi)product frames obtained in the proof of (1) are finite, the claim follows. ■

Acknowledgments

We thank an anonymous reviewer for comments that helped to improve the paper.

Funding

The work of the first author was carried out at Steklov Mathematical Institute and supported by Russian Science Foundation, project 21-11-00318.

References

1. K. Segerberg. "Two-dimensional modal logic," *Journal of Philosophical Logic*, **2**(1), 77–96 (1973).

2. V. Shehtman. "Two-dimensional modal logics," *Mathematical Notices of the USSR Academy of Sciences*, **23**, 417–424 (1978). (Translated from Russian).

3. D. Gabbay, A. Kurucz, F. Wolter, and M. Zakharyaschev. *Many-Dimensional Modal Logics: Theory and Applications*, volume 148 of *Studies in Logic and the Foundations of Mathematics* (Elsevier, 2003).

4. A. Kurucz. "Combining modal logics," P. Blackburn, J. Van Benthem, and F. Wolter, editors, *Handbook of Modal Logic*, volume 3 of *Studies in Logic and Practical Reasoning*, 869–924 (Elsevier, 2008).

5. G. Fischer-Servi. "On modal logic with an intuitionistic base," *Studia Logica*, **36**, 141-149, (1977).

6. D. Gabbay and V. Shehtman. "Products of modal logics, Part 1," *Logic Journal of the IGPL*, **6**(1), 73–146 (1998).

7. F. Wolter and M. Zakharyaschev. "Decidable Fragments of First-Order Modal Logics," *The Journal of Symbolic Logic*, **66**(3), 1415–1438 (1999).

8. M. Rybakov and D. Shkatov. Undecidability of first-order modal and intuitionistic logics with two variables and one monadic predicate letter. *Studia Logica*, **107**(4), 695–717 (2019).

9. V. Shehtman and D. Shkatov. "On one-variable fragments of modal predicate logics," *Proceedings of SYSMICS 2019*, 129–132 (University of Amsterdam, 2019).

10. V. Shehtman. "Simplicial semantics and one-variable fragments of modal predicate logics," *Abstacts of TACL 2019*, 172–173 (Nice, 2019).

11. V. Shehtman. "On Kripke completeness of modal predicate logics around quantified K5," *Annals of Pure and Applied Logic*, **174**(2), 103202 (2023).

12. M. Kracht. *Tools and Techniques in Modal Logic* (North Holland, 1999).

13. V. Shehtman. and D. Shkatov "Kripke (in)completeness of predicate modal logics with axioms of bounded alternativity," *Proceedings of FOMTL 2023*, 26–29 (ESSLLI, 2023).

14. D. Gabbay, V. Shehtman, and D. Skvortsov. *Quantification in Nonclassical Logic, Volume 1*, volume 153 of *Studies in Logic and the Foundations of Mathematics* (Elsevier, 2009).

15. V. Shehtman. "Segerberg squares of modal logics and theories of relation algebras," S. Odintsov, ed., *Larisa Maksimova on Implication, Interpolation, and Definability*, 245–296. (Springer, 2018).

16. V. Shehtman "Bisimulation games and locally tabular logics" *Russian Mathematical Surveys*, **71**, 979–982 (2016). (Translated from Russian).

View publication stats