

https://dspace.org/

Contributions in the field of palaeopalynology at the Bernard Price Institute, past, present and future

MacRae, C. S.; Aitken, G. A.

1997
Bernard Price Institute for Palaeontological Research
http://hdl.handle.net/10539/16397
None

Downloaded from a DSpace repository. https://dspace.org/

Palaeont. afr., **33**, 37-40 (1997)

CONTRIBUTIONS IN THE FIELD OF PALAEOPALYNOLOGY AT THE BERNARD PRICE INSTITUTE, PAST, PRESENT AND FUTURE

by

C.S. MacRae[1] and G.A. Aitken[2]

[1]P.O. Box 905, Cullinan, 1000, South Africa
[2]Bernard Price Institute for Palaeontological Research
University of the Witwatersrand, Private Bag 3, Wits 2050, South Africa

ABSTRACT

A brief chronological summary of the palynological research carried out by students and past members of the staff at the Bernard Price Institute for Palaeontological Research is presented. The contribution that each of these studies has made to the understanding of stratigraphic relationships in the southern African region is highlighted. A correlation chart of palynological biozones documented from South African localities is presented (Table 1).

KEYWORDS: Palaeopalynology, biostratigraphy, spores, pollen, acritarchs, Karoo Basin.

INTRODUCTION

Over the past 40 years palaeopalynological research has provided a powerful means of correlating sediments in southern Africa. Earlier studies employing this methodology in our country focused on the major coal-bearing beds of Permian age. Here the Karoo Basin, rich in its fossil floras, faunas and mineral deposits, provided an ideal setting for palaeopalynological research. More recently attention has shifted to the younger Cretaceous and Tertiary-aged sediments.

Many of the pioneering individuals who dedicated themselves to southern African palynology have, at one time or another, had links with the Bernard Price Institute for Palaeontological Research. It is through their efforts that future students have a solid foundation on which to build.

HISTORICAL REVIEW

Edna Plumstead was the first person associated with the BPI to become involved in palynological research. She documented the fossil floras of the southern African coal measures by illustrating certain spore genera typical of the coal-bearing strata of the Ecca Group (1957, 1966a, 1966b). Palynological endeavour, however, began in earnest at the BPI with the appointment of George Hart in 1965. A graduate of Sheffield University in the United Kingdom, he had already initiated research on the palynofloras of the Ketewaka-Mchuchuma Coalfield in Tanzania and was able to subdivide the coal-bearing strata there into one transitional and two major florizones. (Hart 1960, 1965a). He later used this scheme to correlate strata and the Songwe-Kiwiri Coalfield of that country (Hart 1965b) and clearly demonstrated the value of palynology in defining equivalent rock strata over a wide geographical area. His first publication on South African material (Hart 1963) documented an assemblage from a borehole in the Orange Free State and he concluded that it contained both Carboniferous and Permian species. His first subdivision of the South African Lower Karoo strata (Hart 1967) resulted in the definition of four zones, with subsequent papers (Hart 1969a, 1969b, 1971) enlarging the stratigraphic subdivision of the Lower Karoo and its equivalents in southern and central Africa. He produced various taxonomic papers (Hart 1964a, 1964b, 1966), but the most ambitious was a systematic revision and account of the distribution of Permian miospores (Hart 1965c).

The appointment of Roger Davey in November 1967 led to research on hystrichospheres of the Cretaceous sediments of Zululand. He demonstrated that stages in evolutionary development in three genera could be used to correlate sections of these strata. Two papers followed (Davey, 1969a, 1969b) documenting dinoflagellate cysts and hystrichospheres from the Upper Cretaceous of Zululand, but his stratigraphic results were not published due to the confidential nature of the project. Richard Pienaar, a student of the BPI at that time, documented the calcareous nannoplankton of these rocks (Pienaar 1969).

John Anderson was appointed in 1968 as a palynologist on the BPI staff, with financial support from SOEKOR (Southern Oil Exploration Corporation). His research on the miospores of the northern Karoo Basin in South Africa led him to propose eight zones and 21 subzones, ranging from the Permo-Carboniferous Dwyka to the *Daptocephalus* Zone (now *Dicynodon* Assemblage Zone) of the Beaufort Group in the Late Permian (Anderson 1977). He also provided a detailed correlation of these units.

Registration of Rosemary Falcon for a masters degree in 1970 led to the production of a thesis documenting the palyno-stratigraphy of the Mid-Zambesi Basin in Rhodesia (now Zimbabwe) (Falcon 1972, see also Falcon 1973, 1975). Continued research for a higher degree extended this correlation scheme to

CHRONOSTRATIGRAPHIC UNITS *vide* Harland *et al.* 1982				SOUTH AFRICA				
				N. Karoo Basin	Waterberg/ Pafuri	Main Basin	Main Karoo Basin	North Main
	EPOCH	AGE	CHRON	Aitken 1998	MacRae 1988	Hart 1967	Anderson 1977	Falcon 1978
TRIASSIC	EARLY TRIASSIC	Scythian		X				
				IX				
				VIII			ZONE 7	
				VII			ZONE 6	
		Tatarian		VI	F		ZONE 5	
	LATE PERMIAN					Striatiti Zone		h'
				V			ZONE 4	
		Kazanian						IV h
				IVB	E			
		Ufimian				Zonati Zone		d g
				IVA				c
			Irenian				ZONE 3	b
PERMIAN		Kungurian						
			Filpovian	III	D			a III f
						Cingulati Zone		
		Artinskian	Baigendzinian					
				II				d II e
	EARLY PERMIAN							
			Aktaskian					d
		Sakmarian	Sterlitamakian				ZONE 2	c
			Tastubian	I	B			b
		Asselian	Krumaian			Cavati Zone		
			Uskalikian					I b
			Surenan					a
CARBONIFEROUS	GZELIAN	Noginskian		Dwyka	A		ZONE 1	a
		Klazminskian						
	KASIMOVIAN	Dorogomilovskian						

TABLE 1.

Correlation chart of zones defined with international chronostratigraphic units and assemblages or biozones documented from South African localities (shaded areas = no information).

include sediments on both sides of the Rhodesian watershed (Falcon 1978). Correlation of these assemblages with similar material from many Gondwanan localities confirmed the Early Permian age of the coal-bearing strata equivalent to our Karoo. Falcon, in collaboration with other researchers, provided a palynological framework for the South African Witbank Basin coal measures and associated sediments, and related this to the Zimbabwe sections (Falcon, Pinheiro and Shepherd, 1984).

Also in 1970, Ian McLachlan registered for a higher degree (part-time) at the BPI. During this time he was still an employee of SOEKOR. He studied the problematic microfossils termed *Anellotubulata*, found in Lower Karoo beds. The term *Anellotubulata* was given to these microfossils as similar forms had been described from the Lias of Germany. Tests using X-ray diffraction and electron microprobe work showed that these microfossils were comprised mainly of iron, phosphorus and a small amount of calcium. Unfortunately they were of limited biostratigraphic use due to their long range. McLachlan (1973) reported similar forms from Cretaceous sediments in Eastern Australia. The nature of their depositional environment was questioned and conclusions indicated that Karoo anellotubulates flourished in fresh rather than marine waters. However, the presence of glauconite bands, spinose acritarchs and sponge spicules suggested that conditions were more saline at times, but the lack of recognisable marine faunas indicated that a connection with the sea was unlikely. In 1977 McLachlan published a palynological assessment of petroleum source rocks of the Mesozoic and Cenozoic on the coastal margin of South Africa.

Research for a masters degree by Colin MacRae, on the coal-bearing Karoo strata near Francistown, Botswana, resulted in the definition of three palynological biozones (MacRae 1978). He correlated these biozones with palynological schemes established for other African and Gondwanan localities. This highlighted the presence of coal seams in strata equivalent to the Lower Ecca Group: previously it was thought that coal seams were restricted to the Middle Ecca. MacRae extended his palynological research to the Waterberg, Pafuri and Springbok Flats Basins of South Africa for his doctorate. Six palynostratigraphic zones were defined and their distribution over a considerable distance correlated (MacRae 1988). These biozones were also compared with similarly defined units Gondwana-wide and an attempt was made to date the sections. His results challenged the generally accepted Early Permian age of the middle Ecca (Vryheid Formation equivalent horizons), favouring a slightly younger age. Broad palaeoenvironmental conclusions were drawn for the various units and some economic aspects and implications for future mining highlighted. Carboniferous sediments in South Africa were documented, and the location of the Carboniferous/Permian boundary in the Waterberg Basin was also identified (MacRae 1991).

Grigor Aitken recently completed palynological studies of the No. 4 and 5 seams (Vryheid Formation, Ecca Group) in the Witbank and Highveld Coalfields (Aitken, 1992, 1994). He included this work as part of a doctoral thesis and extended his research to include Beaufort Group sediments in the northern Karoo Basin. Ten biozones were presented and compared with the work of Hart, Anderson, Falcon and MacRae. A composite biostratigraphic section was proposed for the northern Karoo Basin extending from the Dwyka of Late Carboniferous age to the Beaufort of Late Permian/Triassic age.

Ann Cadman, a staff member of the BPI and Sue de Villiers, a PhD student presently registered at the Institute, are presently studying the Late Cretaceous/ Early Tertiary spore and pollen assemblages from the west coast and adjacent Namaqualand interior. These studies will provide valuable information about climatic and plant regimes during this time. So far, assemblages from the Koingnaas site in Namaqualand indicate a Palaeogene age and show that the climate was humid and subtropical. The vegetation was predominantly ancient rain forest which included gymnospermous elements of araucarian and podocarp affinity, with abundant angiosperm trees, shrubs and herbs (De Villiers 1994). Looking at assemblages from various sites on – and offshore, Ann Cadman was able to conclude that the diverse fern representatives and many angiosperm forms indicated conditions vastly different from present on the west coast of southern Africa (Cadman 1994).

CONCLUSIONS

The scope of palynological work still to be done in southern Africa remains unlimited. It is hoped that future research will continue to focus on the creation and refinement of biozonation schemes and that collaborative work will continue between the palaeontological and geological disciplines. A clearer understanding of the past can only be achieved by applying and integrating the research emanating from the various palaeontological subdivisions. Students and past staff members of the BPI have without doubt significantly contributed to the quest of unraveling the historical development of the earth's crust in southern Africa. It is sincerely hoped that the BPI will continue to foster palynological research and that in time many useful answers will be found in studying the fossil spores and pollen that were 'blowing in the wind' many millions of years ago.

ACKNOWLEDGEMENTS

Dr B Millsteed, Dr A Cadman and a second reviewer who wishes to remain anonymous are thanked for their helpful comments on an earlier draft of this paper.

REFERENCES

AITKEN, G.A. 1992. *Palynology of the Number Four Coal Seam.* Uunpublished BSc (Hons) project, University of the Witwatersrand, Johannesburg.

-------- 1994. Permian palynomorphs from the Number 5 Seam, Ecca Group, Witbank/ Highveld Coalfields, South Africa. *Palaeont. afr.*, **31**, 97-109.

-------- 1998. *A Palynological and Palaeoenvironmental Analysis of Permian and Early Triassic sediments of the Ecca and Beaufort Groups, Northern Karoo Basin, South Africa.* Unpublished PhD thesis, University of the Witwatersrand, Johannesburg.

ANDERSON, J.M. 1977. The biostratigraphy of the Permian and Triassic. Part 3. A review of Gondwana Permian palynology with particular reference to the northern Karoo Basin, South Africa. *Mem. Bot. Surv. S. Afr.*, **41**, 1-132.

CADMAN, A. 1994. *Onshore and off - what the Tertiary pollen of the west coast is telling us.* Abstracts of the 8th Biennial Conference of the Palaeontological Society of South Africa, Grahamstown , p.10.

DAVEY, R.J. 1969a. Some dinoflagellate cysts from the upper Cretaceous of northern Natal, South Africa. *Palaeont. afr.*, **12**, 1-24.

-------- 1969b. The evolution of certain upper Cretaceous hystrichosperes from South Africa. *Palaeont. afr.*, **12**, 25-52.

DE VILLIERS, S.E. (1994*). Palynology of Tertiary sediments from a palaeochannel in Namaqualand, South Africa.* Abstracts of the 8th Biennial Conference of the Palaeontological Society of South Africa, Grahamstown, p.40.

FALCON, R.M.S. 1972. *Preliminary palynology of the Mid-Zambesi Basin, Rhodesia.* Unpublished MSc Dissertation, University of the Witwatersrand, Johannesburg.

-------- 1973. Palynology of the Middle Zambesi Basin. **In:** Bond, G. *The Palaeontology of Rhodesia. Bull. Geol. Surv. Rhodesia*, **70**, 43-70.

-------- 1975. Palyno-stratigraphy of the Lower Karroo Sequence in the Central Sebungwe District, Mid-Zambesi Basin. *Palaeont. afr.*, **18**, 1-29.

-------- 1978. *A palynological comparison of Karoo sediments on opposite sides of the Rhodesian watershed with stratigraphic application.* Unpublished PhD Thesis, University of the Witwatersrand, Johannesburg.

--------, PINHEIRO, H.J. & SHEPERD, P. 1984. The palynostratigraphy of the major coal seams in the Witbank Basin with lithostratigraphic, chronostratigraphic and palaeoclimatic implications. *Comm. Serv. Geol. Portugal*, **70**, 2115-2243.

HART, G.F. 1960. Microfloral investigation of the Lower Coal Measures (K2); Ketewaka-Mchuchuma Coalfield, Tanganyika. *Bull. Geol. Surv. Tanganyika*, **30**, 1-18.

-------- 1963. A probable pre-*Glossopteris* microfloral assemblage from Lower Karroo sediments. *S. Afr. J. Sci.*, **59**, 135-146.

-------- 1964a. A review of the classification and distribution of the Permian miospore: *Disaccate striatiti. C.r. Ve, Congr. Internatl. Strat. Geol. Carb., Paris,* 1171-1199.

-------- 1964b. *Chomotriletes* from the lower Permian of South Africa. *Ann. Geol. Surv. S. Afr.*, **3**, 149-156.

-------- 1965a. Microflora from the Ketewaka-Mchuchuma Coalfield, Tanganyika. *Bull. Geol. Surv. Tanganyika,* **36**, 1-27.

-------- 1965b. Miospore Zones in Karroo Sediments of Tanzania. *Palaeont. afr.*, **9**, 139-150.

-------- 1965c. *Systematics and distribution of Permian miospores.* Witwatersrand Univ. Press, Jhb., 253pp.

-------- 1966. *Vittatina africana*, a new miospore from the Lower Permian of South Africa. *Micropalaeont.*, **12**, 37-42.

-------- 1967. Micropalaeontology of the Karroo deposits in South and Central Africa. **In:** I.U.G.S. Secretariat, Ed., *Reviews for the first symposium on Gondwana stratigraphy,* Haarlem, Netherlands 161-172.

-------- 1969a. Palynology of the Permian Period. **In:** Tschudy R.H & Scott, R.A. Eds., *Aspects of Palynology*, Wiley-Interscience, London, 271-289.

-------- 1969b. The stratigraphic sub-division and equivalents of the Karroo Sequence as suggested by palynology. *Earth Science*, **2**, 23-32.

-------- 1971. The Gondwana Permian palynofloras. *Anais. Acad. Bras., Cienc.*, **43** (suppl.), 145-176.

MACRAE, C.S. (1978). *Palaeozoic palynology of Botswana.* Unpublished MSc thesis, University of the Witwatersrand, Johannesburg.

-------- 1988. Palynostratigraphic correlation between the lower Karoo Sequence of the Waterberg and Pafuri coal-bearing Basins and the Hammanskraal plant macrofossil locality, Republic of South Africa, *Mem. Geol. Surv. S. Afr.*, **75**, 217pp.

-------- 1991. Carboniferous sediments in the Waterberg Basin, South Africa, and the identification of the Carboniferous/Permian boundary. *S. Afr. Jour. Sci.*, **87**, 381-387.

McLACHLAN, I.R. (1973). Problematic microfossils from the Lower Karoo Beds in South Africa. *Palaeont. afr.*, **15**, 1-21.

-------- 1977. Palynological assessment of the petroleum source-rock of the Mesozoic and Cenozoic sediments on the coastal margin of South Africa. *Abstr. 17 Congr. geol. Soc. S. Afr.,* Jhb.

PIENAAR, R.N. 1969. Upper Cretaceous calcareous nannoplankton from Zululand, South Africa. *Palaeont. afr.*, **12**, 75-150.

PLUMSTEAD, E. 1957. *Coal in Southern Africa.* Witwatersrand University Press, Johannesburg, 1-24.

-------- (966a. Recent palaeobotanical advances and problems in Africa. *Symposium on Floristics and Stratigraphy of Gondwanaland, Birbal Sahni Institute of Palaeobotany*, Lucknow, 1-12.

-------- 1966b. The story of Africa's coal. *Optima*, **16**, 186-202.