

https://dspace.org/

A comparative study of the proteins in the body fluids of animals

Coetzee, Naudine

2015-02-09
http://hdl.handle.net/10539/16945

Downloaded from a DSpace repository. https://dspace.org/

ASHTON, N. (1951). Anatomical study of Schlemm's canal and aqueous veins by means of neoprene casts; pt 1. Aqueous veins. Brit. J. Ophth., 35, 291-303.

ASHTON, N. (1952). Anatomical study of Schlemm's canal and aqueous veins by means of neoprene casts; pt 2. Aqueous veins (continued). Brit. J. Ophth., 36, 265-267.

ASHTON, N. & SMITH, R. (1953). Anatomical study of Schlemm's canal and aqueous veins by means of neoprene casts; pt 3. Arterial relations to Schlemm's canal. Brit. J. Ophth., 37, 577-583.

ASHTON, N., BRINI, A. & SMITH, R. (1956). Anatomical studies of the trabecular meshwork of the normal human eye. Brit. J. Ophth., 40, 257-282.

ASTRAND, P. & SALTIN, B. (1964). Plasma and red cell volume after prolonged severe exercise. J. Appl. Physicl., 19, 829-832.

AUKLAND, K. & FADNES, H.O. (1973). Protein concentration of interstitial fluid collected from rat skin by a wick method. Acta physiol. scand., 88, 350-358.

AUKLAND, K. & JOHNSEN, H.M. (1974). Protein concentration and colloid osmotic pressure of rat skeletal muscle interstitial fluid. Acta physiol. scand., 91, 354-364.

AXELSEN, N.H. & SVENDSEN, P.J. (1973). Reversed rocket immunoelectrophoresis. Scand. J. Immun., 1, suppl. 1.

BACH, C. & LEWIS, G.P. (1973). Lymph flow and lymph protein concentration in the skin and muscle of the rabbit hind limb. J. Physiol., 235,477-492.

BAKER, K.J. & BRADLEY, S.E. (1966). Binding of sulfobromophtalein (BSP) sodium by plasma albumin. Its role in hepatic BSP extraction. J. Clin. Invest., 45, 281-287.

BARBOUR, H.G., DAWSON, M.H. & NEUWIRTH, I. (1925). Heat regulation and water exchange. Water, salt and lipoid accumulation in the serum as a preliminary to sweating. Am. J. Physiol., 74, 204-223.

BARCROFT, J. & KATO, T. (1915). Effects of functional activity in striated muscle and the submaxillary. Phil. Trans. B., 207, 149-182.

BARTHOLOMEW, R.J. & DELANEY, A.M. (1966). Proc. Australian Assoc. Clin. Biochem., 1, 214. As quoted by Howorth, P.J.N. (1971).

BASS, D.E. & HENSCHEL, A. (1956). Responses of body fluids compartments to heat and cold. Physiol. Rev., 36, 128-144.

BAUER, K.D. & THOMASSON, W.A. (1966). The effect of urea on histones in polyacrylamide gel electrophoresis. J. Chromatog., 22, 496-498.

BAZETT, H.C., SUNDERMAN, F.W., DOUPE, J. & SCOTT, J.C. (1940). Climatic effects on the volume and composition of blood in man. Am. J. Physiol., 129, 69-83.

BEH, K.J., WATSON, D.L. &LASCELLES, A.K. (1974). Concentrations of immunoglobulins and albumin in lymph collected from various regions of the body of the sheep. Aust. J. exp. Biol. med. Sci., 52, 81-86.

BELL, R.D., KEYL, M.J. & SHRADER, F.R. (1969). Effects of renal medullary concentrating ability on canine renal lymph composition. Am. J. Physiol., 216, 704-706.

BENESCH, H. & BENESCH, R.E. (1956). Heterogeneity of serum albumin. Fed. Proc., 15, 218.

BENNETT, N.B. OGSTON, C.M. & OGSTON, D. (1968). The effect of prolonged exercise on the components of the blood fibrinolytic enzyme system. J. Physiol (Lond.), 198, 479-485.

BENSON, J.A., KIM, K.S. & BOLLMAN, J.L. (1955). Extravascular diffusion of protein. Am. J. Physiol. 182, 217-220.

BERG, K. (1964). Precipitation reactions in agar gel between albumin and beta-lipoprotein of human serum. Acta. path. et microbiol. Scand., 62, 287-298.

BERGMANN, L., DENCKER, S.J., JOHANSSON, B.G. & SVENNERHOLM, L. (1968). Cerebrospinal fluid gamma-globulins in subacute sclerosing leucoencephalitis. J. Neurochem., 15, 781.

BERKMAN, P.M., PASTEWKA, J.V. & PEACOCK, A.C. (1969). Influence of urea on the electrophoretic properties of the proteins of microsomal membranes. Biochim. Biophys. Acta., 181, 159-163.

BERNE, B.H., DRAY, S. & KNIGHT, K.L. (1971). Immunological relationships of serum alpha-macroglobulins in the human, rat and rabbit. Proc. Soc. Exp. Biol. Med., 138, 531-535.

BERTHILLIER, G., GOT, R. & BERTAGNOLIO, G. (1968). Biochemistry of rabbit alpha1-macroglobulin. IV Effect on the esterase activity of trypsin and chymotrypsin. Biochim. Biophys. Acta., 170, 140-151.

BERTLER, A., FALCK, B. & ROSENGREN, E. (1964). The direct demonstration of a barrier mechanism in the brain capillaries. Acta pharmac. tox., 20, 317-321.

BEST, C.H. & TAYLOR, N.B. (1955). The Physiological Basis of Medical Practice. Baltimore, Md. : Williams & Wilkins, pp. 14, 25, 26.

BIGGS, R., MaC FARLANE, R.G. & PHILLING, J. (1947). Observations on fibrinolysis-experimental activity produced by exercise or adrenaline. Lancet, 1, 402-405.

BILLIMORIA, J.D., DRYSDALE, J., JAMES, D.C.O. & MACLAGEN, N.F. (1959). Determination of fibrinolytic activity of whole blood with special reference to the effects on exercise and fat feeding. Lancet, 2, 471-475.

BISHOP, D.H.L., CLAYBROOK, J.R. & SPIEGELMAN, S. (1967). Electrophoretic separation of viral nucleic acids on polyacrylamide gels. J. Mol. Biol., 26, 373-387.

BITO, L.Z. (1977). Inflammatory effects of endotoxin-like contaminants in commonly used protein preparations. Science, 196, 83-85.

BITO, L.Z. & MYERS, R.E. (1970). The ontogenesis of haematoencephalic cation transport processes in the rhesus monkey. J. Physiol., 208, 153-170.

BJERRUM, O.J. (1968). Interaction of bromophenol blue and bilirubin with bovine and human serum albumin determined by gel filtration. Scand. J. Clin. Lab. Invest., 22, 41-48.

BJERRUM, O.J. & LUNDAHL, P. (1974). Crossed immunoelectrophoresis of human erythrocyte membrane proteins. Immunoprecipitation patterns for fresh and stored samples of membranes extensively solubilized with non-ionic detergents. Biochim. et Biophys. Acta, 342, 69-80.

BJORKLUND, A., FALCK, B., HROMEK, F. & OWMAN, C. (1969). An enzymatic barrier mechanism for monoamine precursors in the newly forming brain capillaries following electrolytic or mechanical lesions. J. Neurochem., 16, 1605-1608.

BLAND, J.H. (1963). Clinical Metabolism of Body Water and Electrolytes. Philadelphia, London: Saunders. pp. 33, 400.

BLAUER, G. & KING, T.E. (1970). Interactions of bilirubin with bovine serum albumin in aqueous solution. J. Biol. Chem., 245, 372-381.

BLAUER, G., HARMATZ, D. & SNIR, J. (1972). Optical properties of bilirubin-serum albumin complexes in aqueous solution. Biochim. Biophys. Acta, 278, 68-88.

BLOEMENDAL, H., BONT, W.S., JONGKIND, J.F. & WISSE, J.H. (1962). Purity of alpha-crystallin. Nature, 193, 437-439.

BLOTH, B. CHESEBRO, B. & SVEHAG, S. (1968). Ultrastructural studies of human and rabbit alpha M-globulins. J. Exp. Med., 127, 749-756.

BOCK, E. (1973a). Quantitation of plasma proteins in cerebrospinal fluid. A manual of quantitative immuno-electrophoresis. Methods and applications. pp. 111-117.

BOCK, E. (1973b). Non-plasma protiens in cerebrospinal fluid. A manual of quantitative immuno-electrophoresis. Methods and application. pp. 119-124.

BOJANOVSKY, I. & FILIP, J. (1956). The influence of exhausting graded exercise on the level and composition of blood serum proteins in trained athletes. Rev. Czech. Med., 2, 1-7.

BOMAN, H.G. (1955). Chromatography of serum and some other proteins on an anion-exchange resin. Nature, 175, 898-899.

BOMAN, H.G. & WESTLUND, L.E. (1956). Protein chromatography on an anion-exchange resin. Arch. Biochem. Biophys., 64,217-239.

BÖNING, D., SCHWEIGERT, U., TIBES, U. & HEMMER, B. (1972). In vivo and in vitro investigations on the oxygen dissociation curve of blood of trained and untrained subjects during exercise. Pflügers Arch., 332, suppl. R78.

BORUT, A., HOROWITZ, M. & CASTEL, M. (1972). Blood volume regulation in the spiny mouse: capillary permeability changes due to dehydration. In Comparative Physiology of Desert Animals. (Maloiy, G.M.O., Ed.). Symp. Zool. Soc. Lond., 31, 175-189.

BOYLES, S. LEWIS, G.P. & WESTCOTT, B. (1970). Changes in the composition of hind limb lymph after chemical injury. Br. J. Pharmac., 39, 798-808.

BREWER, J.M. (1967). Artifact produced in disc electrophoresis by ammonium persulfate. Science, 156, 256-257.

BREWER, J.M. & De SA, R.J. (1973). Stopped-flow studies of the N-F transition in bovine serum. Fed. Proc., 32, 1323.

BRIGHTMAN, H.W. & REESE, T.S. (1969). Junctions between intimately apposed cell membranes in the vertebrate brain. J. Cell. Biol., 40, 648-677.

BRIMACOMBE, R., MORGAN, J.M. & COX, R.A. (1971). An improved technique for the analysis of ribonucleoprotein fragments from Escherichia coli 30S Ribosomes. Eur. J. Biochem., 23, 52-60.

BRONTE-STEWART, B., ANTONIS, A., ROSE-INNES, C. & MOODIE, A.D. (1961). An interracial study on the serum protein pattern of adult men in Southern Africa. Am. J. Clin. Nut., 9, 596-605.

BROWN, C.S. & HARDENBERGH, E. (1951). A technique for sampling lymph in unanesthetized dogs by means of an exteriorized duct-venous shunt. Surgery, 29, 502-507.

BROWSE, N.L., RUTT, D.R., SLEELAND, D. & TAYLOR, A (1974). The velocity of lymph flow in the canine thoracic duct. J. Physiol. (Lond.), 237, 401-413.

BULLARD, R.W., DILL, D.B. & YOUSEF, M.K. (1970). Responses of the burro to desert heat stress. J. Appl. Physiol., 29, 159-167.

BUNTING, T.L. (1905). Histology of lymphatic glands; general structure, the reticulum and germ centers. Jour. Anat. Physiol., 39, 56.

CASLEY-SMITH, J.R. (1976). The function and interrelationship of blood capillaries and lymphatics. Experimentia, 32, 1-12.

CASLEY-SMITH, J.R. & BOLTON, T. (1973). The presence of large colloidal osmotic pressure across large pores. Microvasc. Res., 5, 213-216.

CASLEY-SMITH, J.R. & FLOREY, H.W. (1961). The structure of normal small lymphatics. Quart. J. Exp. Physiol., 46, 101-105.

CHALMERS, I.M., PUDIFIN, D.J. & SHEPHARD, E.G. (1977). Rheumatoid factor in non-Rheumatoid black patients. S. Afr. Med. J., 51, 617-620.

CHAPMAN, A.L. & BOPP, W.J. (1970). Electron microscopy of vascular barrier in thymus, tonsil, and lymph node of beagle pups. Am. J. Vet. Res., 31, 1255-1268.

CHRAMBACH, A. & RODBARD, D. (1971). Polyacrylamide gel electrophoresis. Science, 172, 440-451.

CHRAMBACH, A., REISFELD, R.A., WYCKOFF, M. & ZACCARI, J. (1967). A procedure for rapid and sensitive staining of protein fractionated by polyacrylamide gel electrophoresis. Anal. Biochem., 20, 150-154.

CLARK, A.H. (1912). On the fate of the jugulor lymph sacs, and the development of the lymph channels in the neck of the pig. Am. Jour. Anat., 14.

CLARK, E.R. & CLARK, E.L. (1933). Further observations on living lymphatic vessels in the transparent chamber in the rabbit's ear - their relation to the tissue spaces. Am. J. Anat., 52, 273.

CLAUSEN, J. (1961). Proteins in normal cerebrospinal fluid not found in serum. Proc. Soc. Exp. Biol. Med., 107, 170.

CLAUSEN, J. (1971). Immunochemical techniques for the identification and estimation of macromolecules. Edit. by Work, T.S. & Work, E. North-Holland Publishing Co., Amsterdam, London.

CLAUSEN, J., GJEDDE, F. & JORGENSEN, K. (1963). Insulin binding proteins in human serum. In vitro experiments. Proc. Soc. Exp. Biol. Med., 112, 778-781.

CLELAND, W.W. (1964). Dithiothreitol, a new pretective reagent for SH-groups. Biochemistry, 3, 480-482.

CLIFF, W.J. & NICOLL, P.A. (1970). Structure and function of lymphatic vessels of the bat's wing. Quart. J. exp. Physiol., 55, 112-121.

COHEN, S. & PORTER, R.R. (1964). Heterogeneity of the peptide chains of gamma-globulin. Biochem. J., 90, 278-284.

COLLIP. P.J., KAPLAN, S.A., BOYLE, D.C. & SHIMIZU, L.S.N. (1964). Protein bound human growth hormone. Metabolism, 13. 532-538.

DAVIS, G.L., ABILDGAARD, C.F., BERNAUER, E.M. & BRITTON, M. (1976). Fibrinolytic and hemostatic changes during and after maximal exercise in males. J. Appl. Physiol., 40, 287-292.

DAVSON, H. (1953). The penetration of large water-soluble molecules into the aqueous humour. J. Physiol, 122, 10P-11P.

DAVSON, H. (1956). Physiology of the Ocular and Cerebrospinal Fluids. Churchill, London.

DAVSON, H. (1960). Intracranial and intraocular fluids. In: Handbook of Physiology. Edit. by Magoum, H.W. The Williams & Wilkins Co., Sec 1, vol 3, p 1761.

DAVSON, H. (1967). Physiology of the Cerebrospinal Fluid. Little, Brown, Boston.

DAVSON, H. (1970a). The normal protein concentration. In: Physiology of the Cerebrospinal Fluid. Little, Brown and Co., London, pp. 272-273.

DAVSON, H. (1970b). Chemical composition and secretory nature of the fluid. In: Physiology of the Cerebrospinal Fluid. Little, Brown and Co., London. pp. 33-54.

DAVSON, H. (1976). Review lecture. The blood-brain barrier. J. Physiol., 255, 1-28.

DAVSON, H & HOLLINGSWORTH, J.R. (1973). Active transport of ^{131}I across the blood-brain barrier. J. Physiol, 233, 327-347.

DAVSON, H., HOLLINGSWORTH, J.R. & SEGAL, M.B. (1970). The mechanism of the cerebrospinal fluid. Brain, 93, 665-678.

DAVSON, H., DOMER, F. & HOLLINGSWORTH, J.R. (1973). The mechanism of drainage of the cerebrospinal fluid. Brain, 96, 329-336.

DE LANNE, R., BARNES, J.R. & BROUHA, L. (1958). Changes in concentration of plasma protein fraction during muscular work and recovery. J. Appl. Physiol., 13, 97-104.

DE LANNE, R., BARNES, J.R. & BROUHA, L. (1959). Changes in osmotic pressure and ionic concentrations of plasma during muscular work and recovery. J. Appl. Physiol., 14, 804-808.

DENCKER, S.J. & SWAHN, B. (1962). Proteins of central nervous origin present in cerebrospinal fluid. Nature (London), 194, 288.

DENIS, W. & AYER, J.B. (1920). A method for the quantitative determination of protein in cerebrospinal fluid. Arch. Int. Med., 26, 436-442.

DERNOUCHAMPS, J.P., VAERMAN, J.P., MICHIELS, J. & HEREMANS, J.F. (1974). Le dosage des proteines de l'humeur aqueuse et du vitre chez le lapin. Ophthalmologia (Basel), 168, 132-139.

EKELUND, L.G. (1967b). Circulatory and respiratory adaptation during prolonged exercise of moderate intensity in the sitting position. Acta physiol. scand., 69, 327-340.

ELEC, S.D. (1948). The recognition of toxicogenic bacterial strains in vitro. Brit. Med. J., 1, 493-496.

ENGESET, A., HAGER, B., NESHEIM, A. & KOLBENSTVEDT, A. (1973). Studies on human peripheral lymph. Lymphology, 6, 1-5.

ENGESET, A., OLSZEWSKI, W., JAEGER, P.M., SOKOLOWSKI, J. & THEODORSEN, L. (1977). Twenty-four hour variation in flow and composition of leg lymph in normal men. Acta physiol. scand., 99, 140-148.

FAWCETT, J.S. & MORRIS, C.J.O.R. (1966). Separation studies. pp 1,9. (As quoted by Gordon, A.H. in 1975).

FEINSTEIN, A. (1966). Use of charged thiol reagents in interpreting the electrophoretic patterns of immune globulin chains and fragments. Nature, 210, 135-137.

FERRIS, T.G., EASTERLING, R.E. & BUDD, R.E. (1964). Electrophoresis of serum protein in acrylamide gel. V Effect of buffers on separation and mobility. Anal. Biochem., 8, 477-486.

FIELD, M.E., LEIGH, O.C., HEIM, J.W. & DRINKER, C.K. (1934). The protein content and osmotic pressure of blood serum and lymph from various sources in the dog. Am. J. Physiol., 110, 174-181.

FLOCKS, M. (1956). The anatomy of trabecular meshwork as seen in tangential section. Arch. Ophthalmol., 56, 708-718.

FLOREY, H. (1927). Reactions of and absorption by lymphatics with special reference to those of the diaphragm. Bri. J. exp. Path., 8, 479.

FLORINI, J.R. & BRIVIO, R.P. (1969). Disc electrophoresis of myosin and myosin derivatives in dilute polyacrylamide gels. Anal. Biochem., 30, 358-367.

FLOYER, M.A. (1969). Discussion. In: Ciba Foundation Symposium on Circulatory and Respiratory Mass Transport. Edit. by Wolstenholm, G.E. & Knight, J., Churchill: London, p 21.

FORSGREN, A. & SJÖQUIST, J. (1966). Protein A from S. aureus. I. Pseudo-immune reaction with human gamma-globulin. J. Immunol., 97, 822-827.

FOSTER, J.F. (1960). In: The Plasma Proteins: vol 1, 1st ed., pp 179-233, Edit. by Putnam, F.W., Academic Press, New York.

FOSTER, J.F., SOGAMI, M., PETERSEN, H.A. & LEONARD, W.J. (1965). The microheterogeneity of plasma albumins. I. Critical evidence for and description of the microheterogeneity model. J. Biol. Chem., 240, 2495-2502.

GANROT, P.O. & SCHERSTEN, B. (1967). Serum alpha$_2$-macroglobulin concentration and its variation with age and sex. Clin. Chim. Acta, 15, 113-120.

GARLICK, D.G. & RENKIN, E.M. (1970). Transport of large molecules from plasma to interstitial fluid and lymph in dogs. Am. J. Physiol, 219, 1595-1605.

GÄRTNER, K., BANNING, E., VOGEL, G. & ULBRICH, M. (1973). The identity of interstitial fluid and hilar lymph in the kidney. Pfluegers Arch., 343, 331-340.

GERSH, I. & CATCHPOLE, H.R. (1960). The nature of ground substance of connective tissue. Perspectives Biol. Med., 3, 282-319.

GIBSON, H. & GAAR, K.A. (1970). Dynamics of the implanted capsule. Fed. Proc., 29, 319.

GIBSON, A.T. & SEGAL, M.B. (1975). Observations on the composition of pericardial fluid. Proc. Physiol. Soc., 32P-33P.

GIEC, L. (1960). Changes of extra cellular space and the circulatory plasma volume during thermal dehydration. Pol. Arch. Med. Wewnetrznej, 30, 39-48.

GILLIGAN, D.A., ALTSCHULE, M.D. & KATERSKY, E.M. (1943). Physiological intravascular hemolysis of exercise. Hemoglobinemia and hemoglobinuria following cross country runs. J. Clin. Invest., 22, 859-871.

GIRARDET, R.E. & BENNINGHOFF, D.L. (1973). Surgical techniques for long-term study of thoracic duct lymph circulation in the dog. J. Surg. Res., 15, 168-175.

GLICKMAN, N. HICK, F.K., KEFTON, R.W. & MONTGOMERY, M.M. (1941). Blood volume changes in men exposed to hot environmental conditions for a few hours. Am. J. Physiol., 134, 165-176.

GODBILLE, P. (1915). Lymphatic glands in meat producing animals. As quoted by Job, T.T. (1922).

GORDON, A.H. (1975). Electrophoresis of proteins in polyacrylamide and starch gels. Edit. by Work, T.S. & Work, E., North-Holland Publishing Co., Amsterdam, Oxford.

GORNALL, A.G., BARDAWILL, C.J. & DAVID, M.M. (1949). Determination of serum proteins by means of the biuret reaction. J. Biol. Chem., 177, 751-766.

GOT, R., MOURAY, H. & MORETTI, J. (1965). Biochemical study of rabbit alpha$_1$-macroglobulin. I. Preparation and physiochemical characteristics. Biochim. Biophys. Acta, 107, 278-285.

GRABAR, P. & WILLIAMS, C.A. (1953). Méthode permettant l'étude conjugée des propriétés électrophorétiques et immunochimiques d'un mélange de proteines; application au sérum sanguin. Biochim. Biophys. Acta, 10, 193.

GRABAR, P., URIEL, J. & COURCON, J. (1960). L'analyse immuno-électrcphorétique du sérum humain normal. Ann. Inst. Pasteur, 99, 13-27.

GREAVES, D.P. & PERKINS, E.S. (1951). Aqueous veins in rabbits. Brit. J. Ophth., 35, 119-123.

GREGERSEN, M.I. (1959). Blood volume. Physiol. Rev., 39,307-342.

GROTTE, G. (1956). Passage of dextran molecules across the blood-lymph barrier. Acta Chir. Scand., suppl. 211.

GRUBB, A. (1973). Preparation of electroendosmosis-free agarose gel and exemplification of its use in crossed immunoelectrophoresis. Anal. Biochem., 55, 582-592.

GULLAND, G.L. (1894). The development of the lymphatic glands. J. Path. Bact., 2, 447.

GUSTAVSON, G.T., SJÖQUIST, J. & STALENHEIM, G. (1967). Protein A from Staphylococcus aureus. II. Arthus-like reaction produced in rabbits by interaction of protein A and human gamma-globulin. J. Immunol., 98, 1178-1181.

GUTTER, F.J., PETERSON, E.A. & SOBER, H.A. (1957). Sedimentation behaviour of bovine plasma albumin as a function of urea concentration and pH. Arch. Biochem. Biophys., 72, 194-204.

GUYTON, A.C. (1963). A concept of negative interstitial pressure based on pressures in implanted perforated capsules. Cir. Res., 12, 399-414.

GUYTON, A.C. (1965). Pressure-volume relationships in the interstitial spaces. Invest. Ophth., 4, 1075.

GUYTON, A.C. (1976). Textbook of Medical Physiology. 5th ed., W.B. Saunders Co., Philadelphia, London, Toronto.

GUYTON, A.C., GRANGER, H.J. & TAYLOR, A.E. (1971). Interstitial fluid pressure. Physiol. Rev., 51, 527.

GUYTON, A.C., TAYLOR, A.E. & GRANGER, H.J. (1975). Circulatory Physiology. II. Dynamics and Control of the Body Fluids. W.B. Saunders Co., Philadelphia, London, Toronto.

GUYTON, A.C., TAYLOR, A.E. & BRACE, R.A. (1976). A synthesis of interstitial fluid regulation and lymph formation. Fed. Proc., 35, 1881-1885.

HADDEN, D.R. & PROUT, T.E. (1964). A growth hormone binding protein in normal human serum. Nature, 202, 1342-1343.

HAGEN, S. (1920). Die regeneration des kammerwassers im menschlichen auge. Klin. Mbl. Augenheilk., 64, 187-194.

HALJAMäE, H. (1970). Sampling of nanoliter volumes of mammalian subcutaneous tissue fluid and ultra-micro flame photometric analyses of the K and Na concentrations. Acta Physiol. Scand., 78, 1-10.

HALJAMÄE, H., LINDE, A. & AMUNDSON, B. (1974). Comparative analyses of capsular fluid and interstitial fluid. Am. J. Physiol., 227, 1199-1205.

HARALAMBIE, G. (1969). Serum glycoproteins and physical exercise. Clin. Chim. Acta, 26, 287-291.

HARALAMBIE, G., KEUL, J. & THEUMERT, F. (1976). Protein-eisen- und kupferveränderungen im serum bei schwimmern vor und nach höhentraining. Europ. J. Appl. Physiol., 35, 21-31.

HARBOE, N. & INGILD, A. (1973). Immunization, isolation of immunoglobulins, estimation of antibody titre. Scand. J. Immun., 1, Suppl. 1.

HARGENS, A.R. & ZWEIFACH, B.W. (1976). Transport between blood and peripheral lymph in intestine. Microvasc. Res., 11, 89-101.

HARGENS, A.R., TUCKER, B.J. & BLANTZ, R.C. (1977). Renal lymph protein in the rat. Am. J. Physiol., 233, F269-F273.

HARRIS, R.I. & KOHN, J. (1974). The pre-albumin fraction: A useful parameter in the interpretation of routine protein electrophoresis. J. Clin. Path., 27, 986-989.

HATTINGH, J., COETZEE, N. & ROSS, F.P. (1977). Erroneous densitometric albumin estimation as a result of prestaining with certain dyes. Anal. Biochem., 80, 635-638.

HATTINGH, J., COETZEE, N. & ROSS, F.P. (1978). Human plasma proteins: Separation on Sephadex G-25 Polyacrylamide Gels. S.A. J. Science, 74, 437-438.

HATTINGH, J., LAURN, H. & MITCHELL, D. (1979). Fever induced in rabbits by intravenous injection of bovine serum albumin. J. Physiol., (Lond) 209, 69-77.

HAVERBACK, B.J., DYCE, B., BUNDY, H.F., WIRTSCHAFTER, S.K. & EDMONDSON, H.A. (1962). Protein binding of pancreatic proteolytic enzymes. J. Clin. Invest., 41, 972-980.

HAWKINS, J.D. (1961). Possible sites of catabolism of rat serum albumin. Biochem. J., 80, 210-214.

HEMMINGSEN, L. & OTHER, A. (1967). Disc electrophoresis of aqueous humour. Acta ophthal., 45, 359-370.

HENDERSON, E.E. & STARLING, E.H. (1904). The influence of changes in the intra-ocular circulation on the intra-ocular pressure. J. Physiol., 31, 305-319.

HENDERSON, E.E. & STARLING, E.H. (1905). The factors which determine the production of the intra-ocular fluid. Proc. Roy. Soc. London, ser. B, 77, 294-310.

BURGESS, A. & SEGAL, M.B. (1970). Morphological changes associated with inhibition of fluid transport in the rabbit choroid plexus. J. Physiol., 208, 88-91P.

BURNET, F.M. & FERNER, F. (1949). The production of antibodies. Melbourne and London: MacMillian and Co. Ltd.

BURNSTOCK, G. & MERRILLEES, N.C.R. (1964). Structural and experimental studies on autonomic nerve endings in smooth muscle. In: Pharmacologigy of Smooth Muscle, Proceedings of the 2nd International Pharmacology Meeting, 1-17.

CAIN, J.C. GINDLAY, J.H., BOLLMAN, J.L., FOOCK, E.V. & MANN, F.C. (1947). Lymph from liver and thoracic duct. Surg. Gynec. Obstet., 85, 559-562.

CALNAN, J.S., PFLUG, J.J., CHISHOLM, G.D. & TAYLOR, L.M. (1972). Pathophysiology of tissue fluid. Proc. Roy. Soc. Med., 65, 715-719.

CAMPBELL, D.H., GARVEY, J.S., CREMER, N.E. & SUSSDORF, D.H. (1963). Isolation and purification of rabbit antibodies. Methods in Immunology, 114-120.

CANNON, W.B. (1927). A Laboratory Course in Physiology. Cambridge, Mass. : Harvard Univ. Press, 125.

CARLEY, W.D. & MAXWELL, D.S. (1970). Development of the blood vessels and extracellular spaces during postnatal maturation of rat cerebral cortex. J. comp. Neurol., 138, 31-48.

CARPENTER, R.L. (1954). Eye. In: Histology, Edit. by Greep, R.O., N.Y.; Blackiston, pp. 844-889.

CARTWRIGHT, G.E., SMITH, E.L., BROWN, D.M. & WINTROBE, M.M.(1948). Electrophoretic analysis of sera of normal and hypoproteinemic swine. J. Biol. Chem., 176, 585-589.

CASH, J.D. & ALLEN, A.G.E. (1967). The fibrinolytic response to moderate exercise and intravenous adrenaline in the same subjects. Brit. J. Haemtol., 13, 376-383.

CASLEY-SMITH, J.R. (1962). The identification of chylomicra and lipoproteins in tissue sections and their passage into jejunal lacteals. J. Cell. Biol., 15, 259.

CASLEY-SMITH, J.R. (1964). An electron microscopic study of injured and abnormally permeable lymphatics. Ann. N.Y. Acad. Sci., 116, 803.

CASLEY-SMITH, J.R. (1967) The function of the lymphatic system under normal and pathological condition: Its dependence on the fine structure and permeability of the vessels. In: Progress in Lymphology. Edit. by Ruttimann. New York: Hafner, pp. 384-359.

CASLEY-SMITH, J.R. (1973). The lymphatic system in inflammation. In: The Inflammatory Progress. Edit. by Zweifach, B.W., Grant, L. & McCluskey, R.T., 2nd ed., N.Y. and London: Academic Press, p. 189.

CONN, H.J. (1977). Biological Stains. Edit. by Lillie, R.D. The Williams and Wilkins Co., Baltimore.

COOPER, D.W. (1963). A serum protein present in pregnant women. Nature, 200, 892.

CORVOL, P.L., CHRAMBACH, A., RODBARD, D. & BARDIN, C.W. (1971). Physical properties and binding capacity of testosterone estradiol-binding globulin in human plasma determined by polyacrylamide gel electrophoresis. J. Biol. Chem., 246, 3435-3443.

COTMAN, C.W., MAHLER, H.R. & HUGLI, T.E. (1968). Isolation and characterization of insoluble proteins of the synaptic plasma membrane. Arch. Biochem. Biophys., 126, 821-837.

COURTICE, F.C. (1972). The Chemistry of lymph. In: Handbuch der allgemeinen Pathologie. vol. 3, Edit. by Meesen, H., Springer-Verlag, New York, pp309-362.

COURTICE, F.C. & MORRIS, B. (1955). The exchange of lipids between plasma and lymph of animals. Quart. J. exp. Physiol., 40, 138-148.

COURTICE, F.C. & SABINE, M.S. (1966). The effect of changes in local temperature on the transfer of proteins and lipoproteins from plasma to lymph in the normal and injured paw of the hypercholesterolemic rabbit. Austr. J. exp. Biol. med. Sei., 44, 23-36.

COURTICE, F.C., SIMMONDS, W.J. & STEINBECK, A.W. (1951). Some investigations on lymph from a thoracic duct fistula in man. Austr. J. exp. Biol. med. Sci., 29, 201-210.

CRONE, C. (1960). The diffusion of some organic non-electrolytes from blood to brain tissue. Acta physiol. scand., Suppl. 50, 17S.

CRONE, C. (1974). Traek af kapillaer fysiologiens udvikling. Copenhagen.

CROWLE, A.J. (1961). Immunodiffusion. Academic Press, New York.

CSERR, H.F., FENSTERMACHER, J.D. & RALL, D.P. (1972). Brain-Barrier systems in sharks. Comp. Biochem. Physiol., 42, 73-78.

DAHLBERG, A.E., DINGMAN, C.W. & PEACOCK, A.C. (1969). Electrophoretic characterization of bacterial polyribosomes in agarose-acrylamide composite gels. J. Mol. Biol., 41, 139-147.

DAMES, W. MAURER, H.R. & NEUHOFF, V. (1972). Hoppe-Seylers Z. Physiol. Chem., 353, 554-558. (As quoted by Lundahl, P. & Liljas, L., 1975).

DAVIS, B.J. (1964). Disc electrophoresis. II. Method and application to human serum proteins. Ann. N.Y. Acad. Sci.. 121, 404-427.

DEWEY, W.C. (1959). Vascular-extravascular exchange of I^{131} plasma proteins in the rat. Am. J. Physiol., 197, 423-431.

DIETER, W. (1925). Über die oberflächenspannung des kammerwassers und ihre beziehungen zum eiweissgehalt desselben. Arch. Augenheilk., 96, 8-33.

DOEMLING, D.B. & STEGGERDA, F.R. (1960). Chronic thoracic duct-venous shunt preparation in dogs. J. Appl. Physiol., 15, 745-746.

DRINKER, C.K. (1946). Extravascular protein and the lymphatic system. Ann. N.Y. Acad. Sci., 46, 807.

DRINKER, C.K. & FIELD, M.E. (1931). The protein content of mammalion lymph and the relation of lymph to tissue fluid. Am. J. Physiol, 97, 32-39.

DRINKER, C.K. & YOFFEY, J.M. (1941). Lymphatics, Lymph and Lymphoid Tissue. Cambridge, Mass: Harvard University Press.

DUKE-ELDER, W.S. (1927). The biochemistry of the aqueous humour. Biochem. J., 21, 66-77.

DUKE-ELDER, S. (1961). System of Ophthalmology. vol 2, Edit. by Krimpton, H., London, p 360.

DUKE-ELDER, S. & DAVSON, H. (1948). The present position of the problem of the intra-ocular fluid and pressure. Brit. J. Ophth., 32, 555-569.

DVORAK-THEOBALD, G. (1934). Schlemm's canal - its anastomoses and anatomic relations. Tr. Am. Ophthalmol. Soc., 32,574-595.

DVORAK-THEOBALD, G.(1955). Further studies on the canal of Schlemm. Its anastomoses and anatomic relations. Am. J. Ophth., 39, 65-69.

EBERT, R.V. & STEAD, E.A. (1941). Demonstration that in normal man no reserves of blood are mobilized by exercise, epinephrine and hemorrhage. Am. J. Med. Sci., 201, 655-664.

EGEBERG, O. (1963). The effect of exercise on the blood clotting system. Scand. J. Clin. Lab. Invest., 15, 8-13

EIDELBERG, E. FISHMAN, I. & HAMS, M.L. (1967). Penetration of sugars agross the blood-brain barrier. J. Physiol. (Lond.), 191, 47-57.

EISLER, P. (1930). Die Anatomie des menschlichen Auges. In: Handbuch der Ophthalmologie. Edit. by Schieck, F. & Bruckner, A., Berlin, p 157.

EKELUND, L.G. (1967a). Circulatory and respiratory adaptation during prolonged exercise. Acta physiol. scand., 70, suppl. 167, 38.

HENDRICK, J.L. & SMITH, A.J. (1968). Size and charge isomer separation and estimation of molecular weights of proteins by disc gel electrophoresis. Arch. Biochem. Biophys., 126, 155-164.

HERMAN, P.G., YAMAMOTO, I. & MELLINS, H.Z. (1972). Blood microcirculation in the lymph node during the primary immune response. J. exp. Med., 136, 697-714.

HERMANS, P.E., McGUCKIN, W.F., McKENZIE, B.F. & BAYRD, E.D. (1960). Electrophoretic studies of serum proteins in cyanogum gel. Proc. Mayo Clin., 35, 792-808.

HIGGINS, P.J. & RAND, C.S. (1974). A comparative immunochemical study of the serum proteins of several galapagos iguanids. Comp. Biochem. Physiol., 49A, 347-355.

HIRSCHFELD, J. (1960). Shape of the precipitate in immunoelectrophoresis. Nature, 185, 164-165.

HIRSCHFELD, J. & SÖDERBERG, U. (1960). Immuno-electrophoretic demonstration of precipitating components in sera from pregnant women. Nature, 187, 332-333.

HITZIG, W.H. (1961). Das bluteiweissbild im säulingsalter. Bull. Schweiz. Akad. Med. Wiss., 17, 228.

HJERTÉN, S. (1959). Calsium phosphate chromatography of normal human serum and of electrophoretically isolated serum proteins. Biochim. Biophys. Acta, 31, 216-235.

HJERTÉN, S., JERSTEDT, S. & TISSELIUS, A. (1965). Electrophoretic particle sieving in polyacrylamide gels as applied to ribosomes. Anal. Biochem., 11, 211-218.

HOBBS, B.B. & MAY, J.B. (1975). The quantitation of blood flow to and lymph output from single lymph nodes. In: Abstracts of 1st World Congress for Microcirculation. Toronto, 1975, abstract no. 301.

HOLMGREN, A. (1956). Circulatory changes during muscular work in man. With special reference to arterial and central venous pressures in the systemic circulation. Scand. J. Clin. Lab. Invest., 8, suppl. 24, 97.

HOPKINSON, B.R., BORDER, J.R., HEYDEN, W.C. & SCHENK, W.G. (1968). Interstitial fluid pressure changes during hemorrhage and blood replacement with and without hypotension. Surgery, 64, 68.

HOROWITZ, M. & BORUT, A. (1973). Blood volume regulation in dehydrated rodents: plasma colloid osmotic pressure, total osmotic pressure and electrolytes. Comp. Biochem. Physiol., 44A, 1261-1265.

HOROWITZ, M. & BORUT, A. (1975). Blood volume regulation in dehydrated rodents: plasma protein turnover and sedimentation coefficients. Comp. Biochem. Physiol., 51A, 827-831.

HORSTMANN, E. (1959). Beobachtungen zur motorik der lymphgefäbe. Pflügers Arch. ges. Physiol., 269, 511-519.

HOWORTH, P.J.N. (1971). Determination of serum albumin in neonatal jaundice. The albumin saturation index. Clin. Chim. Acta, 32, 271-278.

IKKALA, E., MYLLYLä, G. & SARAJAS, H.S. (1963). Hemostatic changes associated with exercise. Nature, 199, 459-461.

JACOBSSON, S. & KJELLMER, I. (1964). Flow and protein content of lymph in resting and exercising skeletal muscle. Acta Physiol. Scand., 60, 278-285.

JANATOVA, J.,FULLER, J.K. & HUNTER, M.J. (1968). The heterogeneity of bovine albumin with respect to sylfhydryl and dimer content. J. Biol. Chem., 243, 3612-3622.

JARABAK, J., SEEDS, E. & TALALAY, P. (1966). Reversible cold inactivation of a 17 beta-hydroxy steroid dehydrogenase of human placenta: protective effect of glycerol. Biochem., 5, 1269-1279.

JARRETT, I.G. (1972). A polymeric form of haemoglobin-binding protein in sheep following metabolic and hormonal disturbance. Austr. J. Biol. Sci., 25, 941-948.

JASANI, M.K. & LEWIS, G.P. (1971). Lymph flow and changes in intracellular enzymes during healing and rejection of rabbit skit grafts. J. Physiol., 219, 525-554.

JELLINCK, P.H. & FLETCHER, R. (1970). Peroxidase-catalyzed conjugation of (4-^{14}C) estradiol with albumin and thiols. Can. J. Biochem., 48, 1192-1198.

JENNINGS, M.A. & FLOREY, F.R.S. (1967). An investigation of some properties of endothelium related to capillary permeability. Proc. Roy. Soc. (Lond.) Ser. B, 167, 39-63.

JENSEN, K. (1959). Undersoqelser over stafqlococcernes antigenstruktur. Edit. by Munksgaard, E., Copenhagen.

JOB, T.T. (1915). The adult anatomy of the lymphatic system in the common rat. (Epimys norvegicus). Anat. Rec., 9, 447-459.

JOB, T.T. (1922). Studies on lymph nodes. Am. J. Anat., 31, 125-137.

JOHANSSON, B.G. (1972). Agarose gel electrophoresis. Scand. J. Clin. Lab. Invest., 29, suppl. 124, 7-19.

JOHANSSON, B.G. & HJERTÉN, S. (1974). Electrophoresis, crossed immunoelectrophoresis, and isoelectric focussing in agarose gels with reduced electroendosmotic flow. Anal. Biochem., 59, 200-213.

JOHANSSON, B.G. & RYMO, L. (1964). Thin-layer filtration and thin-layer electrophoresis on sephadex gels. Biochem. J., 92, 5P.

JOHANSSON, B.G. & STENFLOW, J. (1971). Polyacrylamide slab electrophoresis followed by electrophoresis into antibody-containing agarose gel. Anal. Biochem., 40, 232-236.

JOHNSON, T.F. & WONG, H.Y.C. (1961). Effects of training and competitive swimming on serum proteins. J. Appl. Physiol., 16, 807-809.

JOLLY, J. (1909). Sur une disposition speciale de la structure des ganglions lymphatiques chex les oiseaux. Soc. de Biol., 66, 499.

JOLLY, J. (1910). Recherches sur les ganglions lymphatiques des oiseaux. Arch. d' Anat. Micro., 11.

JONES, R.T. (1975). Normal values for some biochemical constituents in rabbits. Laboratory Animals, 9, 143-147.

JONES, M.A.S. & SHANNON, A.D. (1972). The effect of staphylococcal antigen on the output of cells from the popliteal lymph node of immunized sheep. New Zealand Veterinary Journal, 20, 19-23.

JONES, C.E., CROWELL, J.W. & SMITH, E.E. (1969). Determination of mean tissue oxygen tensions by implanted perforated capsules. J. Appl. Physiol., 26, 630-633.

JUDAH, J.D., GAMBLE, M. & STEADMAN, J.H. (1973). Biosynthesis of serum albumin rat liver. Evidence of the existence of "proalbumin". Biochem. J., 134, 1083-1091.

KABAT, E.A., MOORE, D.H. & LANDOW, H. (1942). An electrophoretic study of the protein components in cerebrospinal fluid and their relationship to the serum proteins. J. Clin. Invest., 21, 571-577.

KALTREIDER, W.I. & MENEELY, C.R. (1940). The effect of exercise on the volume of the blood. J. Clin. Invest., 19, 627-634.

KAPLAN, L.J. & FOSTER, J.F. (1971). Isoelectric focussing behaviour of bovine plasma albumin, mercaptalbumin, and beta-lactoglobulins A and B. Biochem., 10, 630-636.

KARNOVSKY, M. (1962). Metabolic basis of Phagocytic Activity. Physiol. Rev., 42, 143-168.

KATZMAN, R. & PAPPIUS, H.M. (1973). Cerebrospinal fluid ions. In: Brain Electrolytes and Fluid Metabolism. The William and Wilkens Co., Baltimore, pp75-110

KAWAMURA, S., YASUDA, Y., TAYAMA, H. MOKI, H., TAKASE, K. & OGASAWARA, S. (1974). Cellulose acetate membrane electrophoresis of bovine serum proteins using a barbital calcium buffer. Jap. J. Vet. Sci., 36, 285-290.

KELLER, S. & BLOCK, R.J. (1959). Amino acid composition of the serum proteins. III. The chromatographic isolation of human and bovine serum albumins, and the amino composition of the fractions. Arch. Biochem. Biophys., 85,366-373.

KEYL. M.J., SCOTT, J.B., DABNEY, J.M., HADDY, F.J., HARVEY, R.B., BELL, R.D. & GINN, H.E. (1965). Composition of canine renal hilar lymph. Am. J. Physiol, 209, 1031-1033.

KINSEY, V.E. (1950). A unified concept of aqueous humor dynamics and the maintnance of intra-ocular pressure. Arch. Ophthalmol., 44, 215-235.

KINSEY, V.E. (1951). The chemical composition and the osmotic pressure of the aqueous humor and plasma of the rabbit. J. Gen. Physiol, 34, 389-402.

KIRK, G.R., NEATE, S., McCLURE, R.C. & HUTCHESON, D.P. (1974). Electrophoretic pattern of Equine cerebrospinal fluid. Am. J. Vet. Res., 35, 1263-1264.

KIRK, G.R., HUTCHESON, D.P. & NEATE, S. (1975). Electrophoretic pattern of serum protein in clinically normal horses and ponies with laminitis. Vet. Med. Small Anim. Clin., 70, 337-339.

KJELDAHL, J. (1883). Neue methode zur bestimmung des stickstoffs in organischen körpern. Z. Anal. Chem., 22, 366.

KJELLMER, I. (1964). The effect of exercise on the vascular bed of skeletal muscle. As quoted by Jacobsson, S. & Kjellmer, I. (1964).

KJELLMER, I. (1965). Studies on exercise hyperaemia. Acta Physiol. Scand., 64, suppl. 244, 1-27.

KLATZO, I., MIQUEL, J., FERRIS, P.J., PROKOP, J.D. & SMITH, D.E. (1964). Observations on the passage of the fluorescein labelled serum proteins (FLSP) from the cerebrospinal fluid. J. Neuropath. exp. Neurol., 23, 18-35.

KLEIHAUER, E. & BETKE, K. (1957). Zür hämoglobin bestimmung mittels cyanmethämoglobin nach Betke und Savelberg. Das ärtzl. Lab., 3, 202 -205.

KNIGHT, K.L. & DRAY, S. (1968). Identification and genetic control of two rabbit alpha2-macroglobulins allotypes. Biochem., 7, 1165-1171.

KORNER, P.I., MORRIS, B. & COURTICE, F.C. (1954). An analysis of factors affecting lymph flow and protein composition during gastric absorption of food and fluids, and during intravenous infusion. Austr. J. exp. Biol., 32, 301-320.

KORNGOLD, L. & VAN LEEUWEN, G. (1957). The effect of the molecular weight on the curvature of the precipitin line in the Ouchterlony technique. J. Immunol., 78, 172-177.

KORZHUEV, P.A. (1964). Methods of study of blood in fish. Techniques for the investigation of fish physiology., pp. 2-10.

KOZLOWSKI, S. & SALTIN, B. (1964). Effect of sweat loss on body fluids. J. Appl. Physiol., 19, 1119-1124.

KRAUSE, U. & RAUNIO, V. (1969). Proteins of the normal human aqueous humour. Ophthalmologica, 159, 178-185.

KRONFELD, P.C., LIN, C.K. & LUO, T.H. (1941). The protein content of the reformed aqueous humour in man. Am. J. Ophthal., 24, 264-276.

KUHLMAN, R.E. & KAUFMAN, H.E. (1960). A microchemical study of the aqueous humour enzyme-protein interrelations. Arch. Ophthal., 63, 41-46.

LADD, M.P., KOTTKE, F.J. & BLANCHARD, R.S. (1952). Studies on the effect of massage on the flow of lymph from the fore leg of the dog. Arch. Phys. Med., 38, 604-612.

LADEGAARD-PEDERSEN, H.J. (1970). Measurement of the interstitial pressure in subcutaneous tissue in dogs. Circ. Res., 26, 765.

LADELL, W.S.S. (1948). The measurement of chloride losses in the sweat. J. Physiol, 107, 465-471.

LAMOUREUX, G., JOLICOEUR, R., GIARD, N., ST-HILAIRE, M. & DUPLANTIS, F. (1975). Cerebrospinal fluid proteins in multiple sclerosis. Neurology, 25, 537-546.

LANDIS, E.M. (1931). Capillary pressure and hyperemia in muscle and skin of the frog. Am. J. Physiol., 98, 704-716.

LANDIS, E.M. & PAPPENHEIMER, J.R. (1963). Exchange of substances through the capillary walls. In: Handbook of Physiology. Circulation, vol. 2, Edit. by Hamilton, W.F. and Dow, P., pp. 961-1034, Am. Physiol. Soc., sect. 2, Washington.

LANDIS, E.M., JONAS, L., ANGEVINE, M. & ERB, W. (1932). The passage of fluid and protein through the human capillary wall during venous congestion. J. Clin. Invest., 11, 717-734.

LANGHAM, M.E. (1958). Aqueous humour and control of intra-ocular pressure. Physiol. Rev., 38, 215-242.

LATERRE, E.C. & HEREMANS, J.F. (1963). A note on proteins apparently specific for cerebrospinal fluid. Clin. Chim. Acta, 8, 220-226.

LATERRE, E.C., HEREMANS, J.F. & CARBONARA, A. (1964). Immunological comparison of some proteins found in cerebrospinal fluid, urine and extracts from brain and kidney. Clin. Chim. Acta, 10, 197-209.

LAURELL, C.B. (1965). Antigen-antibody crossed electrophoresis. Anal. Biochem., 10, 358-361.

LAURELL, C.B. (1966). Quantitative estimation of proteins by electrophoresis in agarose gel containing antibodies. Anal. Biochem., 15, 45-52.

LAURELL, C.B. (1968). Orosomucoid and alpha-antitrypsin in maternal and fetal sera at parturition. Scand. J. Clin. Lab. Invest., 21, 136.

LAURELL, C.B. (1972). Composition and variation of the gel electrophoretic fractions of plasma, cerebrospinal fluid and urine. Scand. J. Clin. Lab. Invest., 29, suppl. 124, 71-82.

LAURELL, C.B. & GRÖNVALL, C. (1962). Haptoglobins. Advan. Clin. Chem., 5, 135-172.

LAURELL, C.B. & NILÉHN, J.E. (1966). A new type of inherited serum albumin anomaly. J. Clin. Invest., 45, 1935-1945.

LEAK, L.V. (1971). Studies on the permeability of lymphatic capillaries. J. Cell. Biol., 50, 300-323.

LEAK, L.V. & BURKE, J.F. (1966). Fine structure of the lymphatic capillary and the adjoining connective tissue area. Am. J. Anat., 118, 785.

LEAK, L.V. & BURKE, J.F. (1968). Ultrastructural studies on the lymphatic anchoring filaments. J. Cell. Biol., 36, 129-149.

LEBER, T. (1903). Die cirkulations und erährungs verhältnissc des auges. Graefe-Saemisch Handb. d. ges. Ophthal., 2 aufl., BD2, 207-352.

LEE, D.H.K. & MULDER, A.G. (1935). Some immediate effects on reduced cooling powers upon the water balance and related effects in the human subject. J. Physiol., 84, 410-432.

LEVINSON, A. (1950). Cerebrospinal fluid in infants and children. Med. Clin. N. Am., 34, 107-118.

LEWIS, F.T. (1909). The first lymph glands in rabbit and human embryos. Anat. Res., 3.

LEWIS, G.P. (1969). Changes in the composition of rabbit hind limb lymph after thermal injury. J. Physiol., 205, 619-634.

LEWIS, J.H. (1974). Comparative hematology: studies on elephants, Elephas maximus. Comp. Biochem. Physiol, 49A, 175-181.

LEWIS, G.P. & WESTCOTT, B. (1968). Intracellular enzymes in rabbit hind-limb lymph after injury. J. Physiol., 195, 12-13P.

LEWIS, G.P. & YATES, C. (1972). Flow and composition of lymph collected from the skeletal muscle of the rabbit hind limb. J. Physiol., 226, 57-58P.

LIBERMANN, I.M., GONZÁLEZ, F., BRAZZUNA, H., GARCIA, H. & LABOUNORA, D. (1972). Fluid composition from implanted perforated capsules: an approach to interstitial fluid? J. Appl. Physiol., 33, 751-756.

LIND, K.E., KRAGH-HANSEN, V. & MOLLER, J.V. (1974). Protein binding of small molecules. Biochim. Biophys. Acta, 371, 451-461.

LINK, H. (1967). Immunoglobulin G and low molecular weight proteins in human cerebrospinal fluid. Acta neurol. scand., 43, suppl. 28, 1.

LOENING, U.E. (1967). The fractionation of high-molecular-weight ribonucleic acid by polyacrylamide-gel electrophoresis. Biochem. J., 102, 251-257.

LÖWENSTEIN, A. (1920). Untersuchungen über den stoffwechsel des menschlichen auges. Klin. Mbl. Augenheilk., 65, 654-662.

LOWRY, O.H., ROSEBROUGH, N.J., FARR, A.L. & RANDALL, R.J. (1951). Protein measurement with the Folin phenol reagent. J. Biol. Chem., 193, 256-275.

LUCAS, J. & FLOYER, M.A. (1973). Renal control of changes in the compliance of the interstitial space: a factor in the aetiology of renoprival hypertension. Clin. Sci., 44, 397.

LUCIANI, L. (1901). Tratado Didáctico de Fisiología Humana. vol. 1, Virgilli, Barcelona, 495-497.

LUNDAHL, P. & HJERTÉN, S. (1973). Isoelectric focusing in free ampholine solution and attempts at isoelectric focusing in pH gradients created in ordinary buffers. Ann. N.Y. Acad. Sci., 209, 94-111.

LUNDAHL, P. & LILJAS, L. (1975). Crossed immunoelectrophoresis. Polyacrylamide gel electrophoresis followed by electrophoresis into an agarose gel containing antibodies. Anal. Biochem., 65, 50-59.

MacPHERSON, C.F.C. & COSGROVE, J.B.R. (1961). Immunochemical evidence for a gamma globulin peculiar to cerebrospinal fluid. Canad. J. Biochem., 39, 1567.

MARON, M.B., HORVATH, S.M. & WILKERSON, J.E. (1975). Acute blood biochemical alterations in response to marathon running. Europ. J. appl. Physiol., 34, 173-181.

MARKUS, G., McCLINTOCK, D.K. & CASLELLANI, B.A. (1967). Ligand-stabilized conformations in serum albumin. J. Biol. Chem., 242, 4402-4408.

MARRON, T.U. (1941). Cerebrospinal fluid protein values determined by tyrosine equivalent method. Am. J. Med. Sci., 202, 330-333.

MAYERSON, H.S. (1963). The physiological importance of lymph. In: Handbook of Physiology, vol. 2, Edit. by Hamilton, W.F. and Dow, P., pp 1035-1073, Am. Physiol. Soc., sect. 2, Washington.

MAYERSON, H.S., PATTERSON, R.M., McKEE, A., Le BRIE, S.J. & MAYERSON, P. (1962). Permeability of Lymphatic vessels. Am. J. Physiol., 203, 98-106.

MAYNARD, E.A., SCHULTZE, R.L. & PEASE, D.C. (1957). Electron microscopy of the vascular bed of rat cerebral cortex. Am. J. Anat., 100, 409-433.

MEHL, J.W., O'CONNELL, W. & DE GROOT, J. (1964). Macroglobulin from human plasma which forms an enzymatically active compound with trypsin. Science, 145, 821-822.

MENON, I.S., BURKE, F.D., SMITH, P.A., NEWELL, D.J. & DEWAR, H.A. (1969). A study of possible causes of increased fibrinolytic activity during exercise. Thromb. Diath. Haemorrhag., 21, 287-293.

MERLER, E., REMINGTON, J.S., FINLAND, M. & GITLIN, D. (1962). Differences between urinary albumin and serum albumin. Nature, 196, 1207-1208.

MERRITT, H.H. & FREMONT-SMITH, F. (1937). Cerebrospinal Fluid. W.B. Saunders Co., Philadelphia.

MEYER, B.J., MEYER, A. & GUYTON, A.C. (1968). Interstitial fluid pressure. V. Negative pressure in the lungs. Circ. Res., 22, 263-271.

MICHAELIS, L. (1909). Electrische überfuhrung von fermenten. I. Das inverten. Biochem. Z., 16, 83-86.

MIKATA, A. & NIKI, R. (1971). Permeability of postcapillary venules of the lymph node. An electron microscopic study. Expl. molec. Path., 14, 289-305.

MOORE, D.H. (1945). Species differences in the serum protein patterns. J. Biol. Chem., 161, 21.

MOORE, D.H. & RUSKA, H. (1957). The fine structure of capillaries and small arteries. J. Biophysic. Biochem. Cytol., 3, 457-462.

MORIMOTO, T., SLABOCHOVA, Z., NAMAN, R.K. & SARGENT, F. (1967). Sex differences in physiological reactions to thermal stress. J. Appl. Physiol, 22, 526-532.

Author Coetzee N
Name of thesis A Comparative study of the proteins in the body fluids of animals 1981

PUBLISHER:
University of the Witwatersrand, Johannesburg
©2013

LEGAL NOTICES:

Copyright Notice: All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed, or otherwise published in any format, without the prior written permission of the copyright owner.

Disclaimer and Terms of Use: Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.