DECLARATION

I declare that this dissertation is my work. It is being submitted for the degree of Master of Sciences in Medicine in the field of Nuclear Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination in any other university.

Kayode Adedapo

Date

This research was approved by the committee for Research on Human Subjects, University of the Witwatersrand (protocol M080621) and was carried out at the Nuclear Medicine department of the Johannesburg Hospital in 2008

Dedication

To God almighty, the giver of life and opportunities be all the glory, honour and majesty

Abstract

A 20 years retrospective study of all patients treated at the Johannesburg Hospital with radioactive iodine post thyroidectomy for differentiated thyroid carcinoma from 1986 to 2006 was carried out with the aim of determining the adequate duration and modality of follow-up.

A total of 106 patients (91 female and 15 male) out of 287 patients qualified for inclusion into the study. The mean age of the patients was 45 years (range: 16-81 years). There was a ratio of 6 females to 1 male. The mean ages of incidence in papillary, follicular, Hurthle and mixed papillary and follicular cancers were 40, 49, 53 and 49 years respectively. Only the ages of the patients with papillary cancer differed significantly from the ages of patients with other cancers (p=0.011). Of the 4 histologic types of cancer recorded, papillary thyroid cancer was the most common 58 (55%) followed by follicular 30 (28%), Hurthle cell 10(9%) and the mixed papillary-follicular cancers 8 (8%).

More than half of the patients 58 (53.7%) had total thyroidectomy, while 36(34%), and 12(11.3%) patients had near total thyroidectomy and lobectomy respectively. Majority of the patients with papillary carcinoma (35 out of 58), and Hurthle cell carcinoma (6 out of 10) had total thyroidectomy. All patients had complete 24 months follow-up after a negative whole body iodine scan. A proportion of the patients were followed up to 36 and 60 months after a negative whole body iodine scan.

The mean iodine dose administered by the first 6 months of follow-up was 3.3GBq (88mCi). Following the first iodine treatment, 58 of 101 patients were lodine scan negative. The remaining 43 patients progressed to a negative scan by the fourth treatment.

The histology and the type of surgery did not affect the outcome of iodine therapy in the patients studied.

The proportion of patients with negative whole body iodine scan after radioactive iodine ablation increased progressively from 58 out of 101 (57.4%) at the first 6 months to 99 of 101(98%) by 24 months of follow up.

Twenty four months was found to be the minimum adequate follow-up period, while the modality of follow-up was whole body iodine scan and thyroglobulin determination in concordant patients on and off T_4 suppression.

ACKNOWLEDGEMENTS

I am grateful to my supervisor Prof. MDT Vangu for his deep interest in me, this work and the entire training process for the award of an M.Sc in Nuclear Medicine. Your desire and commitment to see Nuclear Medicine in Nigeria grow to an enviable status in the West African sub region is recognized and fully appreciated.

I am extremely indebted to my head of department and mentor Prof. BOA Osifo for her belief in me and all the support she has given to make this dream of a Master's degree in Nuclear Medicine a reality.

I am grateful to the chief medical director of my hospital University College Hospital (UCH) for making the financial support for this study possible, God bless you sir. Immense gratitude goes to all the consultants in the department of nuclear medicine for showing me the path to follow; Carlos, Sindy and Sonto.

I shall always remember the friendship and camaraderie atmosphere under which I gleaned from the registrars; Nico, Shireen, Kaushica, Hema and Lizzete. Thank you all. I am grateful to Dr Elena Libhaber and Prof. Geoffrey Candy for my data analysis. I could not have made much sense of the study without your expertise. Thank you very much.

I thank my good friend Mr. Bode Akindele and his wife Dr Kemi Akindele for having me to stay with them during the duration of this course in Johannesburg. May the Lord bless and reward your labour of love.

I will miss the friendship of Mr. Maleeme the assistant director of radiography at the Johannesburg Hospital as well as all the other radiographers without which my learning will not be complete. Ma Emily, the departmental secretary is appreciated for her assistance, as well as all other paramedical members of staff.

Finally, I appreciate my one and only true love Dr Adura Adedapo and my lovely children Dami, Ife and Ini for their understanding and love during my period of long absence from the closeness we used to share. You all will remain indelible in my heart and the success of this work is because you all loved me.

TABLE OF CONTENTS

DeclarationiDedicationiiSummaryiiiAcknowledgmentsivTable of ContentsvList of Abbreviationsix			
Chapt	er 1: Introduction1		
1.1	Brief anatomy of the thyroid Gland1		
1.1.1	Description of thyroid gland1		
1.1.2	Blood supply and drainage1		
1.1.3	Nerve supply1		
1.1.4	Lymphatic drainage1		
1.2	Brief physiology1		
1.2.1	Synthesis of thyroid hormones2		
1.2.2.	Storage		
1.2.3	Control of Secretion2		
Chapt	er 2: Thyroid cancer4		

Page

2.1	Epidemiology4
2.2	Histological types4
2.2.1	Papillary thyroid cancer4
2.2.2	Follicular thyroid cancer5
2.2.3	Hurthle (oxyphil or oncocytic) Cell Cancer5
2.2.4	Anaplastic Thyroid Cancer 5
2.2.5	Medullary Thyroid cancer5
2.2.6	Metastatic cancer to the thyroid6
2.2.7	Malignant lymphoma6
2.3	Diagnostic Approach6
2.4	Pathological Staging6
2.4.1	Primary tumour (pT)7
2.4.2	Regional lymph nodes (N)7
2.4.3	Distant metastasis (M)7
2.4.4	Staging Protocol7
2.4.5	Staging for Medullary carcinoma7

Chapter: 3 Manag	ement of differentiated thyroid cancer	8
3.1 Surgery		8
3.2 Prognosis		8
3.3 Follow-up of	differentiated thyroid cancer	9
3.3.1 Radioactive lo	odine (RAI) Ablation	9
3.3.2 Side effect of	radioactive iodine	10
3.3.3 New concepts	s in follow –up of Patients	10
3.3.3.1 Serum Thyro	globulin	11
3.3.3.2 Neck Ultraso	nography	11
3.3.3.3 Recombinant	t TSH	11
3.3.4 Thyroglobulin E	Elevated, Negative Iodine Scan (TENIS)	12
3.3.5 External Bear	m Radiotherapy (EBR)	12
Chapter 4: Method	ds	13
4.1 Objective		13
4.2 Study design.		13

4.3	Study population and sampling13	3
4.3.1	Inclusion criteria13	3
4.4	Ethical issues14	1
4.5	Confidentiality14	1
4. 6	Data management and analysis14	1
Chapte	er 5: Results1	5
5.1 De	mographics1	5
5.2 Su	rgery and Post Surgical follow-up15	5
5.3 Th	yroglobulin level and iodine scan1	5
5.4 Do	ese and number of radioiodine treatments16	3
5.5 Me	etastasis10	5
Chapte	er 6: Discussion2	5
Chapter 7: Conclusion28		
References 29		
Apper	ndix30)

LIST OF ABBREVIATIONS

- DNM- Department of Nuclear medicine
- DTC Differentiated Thyroid Cancer
- PTC Papillary Thyroid Cancer
- FTC Follicular Thyroid Cancer
- HTC Hurthle cell Thyroid Cancer
- MTC Medullary Thyroid cancer
- ATC Anaplastic Thyroid Cancer
- Tg -Thyroglobulin
- MEN Multiple Endocrine Neoplasia
- TNM Tumour Nodes, Metastasis
- RRA Radioactive Remnant Ablation
- RAI Radioactive Iodine
- TSH Thyroid Stimulating Hormone
- WBS Whole Body Scan
- I-131 Iodine -131
- RhTSH- Recombinant Human Thyroid Stimulating Hormone
- FT3 Free Triiodothyronine
- FT4 Free Thyroxine
- LT4 Levothyroxine
- FNAC Fine Needle Aspiration Cytology
- CT Computerized Tomography
- US Ultrasound
- TENIS Thyroglobulin Elevated Negative Iodine Scan
- EBR- External Beam Radiotherapy