

https://dspace.org/


Partial purification and characterization of soluble cyclic nucleotide phosphodiesterases in human and murine tissues

Robinson, Marion Frances

2015-08-18
http://hdl.handle.net/10539/18255

Downloaded from a DSpace repository. https://dspace.org/

Fig. 5:3. Ferguson plot of cyclic nucleotide phosphodieterase from crude cytosol of humar leiomyoma of the uterus. Samples were run, as described in the experimental section, at different gel concentrations ranging from 5% to 8.5%. For each gel concentration, the plot of Rf (logarithmic scale) versus gel concentration (%) was constructed from the results (mean s.d. of four separate runs), by using least-squares linear regression. The plots represent band 1 (●), band 2 (△), band 3 (○) and band 6 (▲). The individual data for each band is given in Table 5:1.




Sucrose gradient ultracentrifugation of crude leiomyoma cytosol.

This method revealed three peaks of cPDE activity (only cAMP tested) with sedimentation coefficients, calculated according to the tables of McEwan (1967), of 3.6 ± 0.35 S, 8.1 ± 0.3 S and 11.8 ± 0.3 S at 1 uM cAMP concentration (Fig. 5:4). Only two peaks at 7.9 S and 3.6 S were observed at 100 uM cAMP. The affinity purified fraction showed a single symmetrical peak of enzyme activity with 1 uM cAMP at 7.8 ± 0.3 S (Fig. 5:4). No activity could be detected at 100 uM cAMP. If one expressed the results as dpm versus fraction number, the crude cytosol also showed a peak in the identical position as that of the purified fraction. This is similar to the findings of Strada *et al.* (1981) who presented their results in the latter way although their reported sedimentation coefficient was less for rat uterus as further detailed in the discussion. It was noted that leiomyoma cytosol, which had been frozen and thawed, showed the majority of specific activity to be associated with the 3.5 S peak (result not shown).

Fig.5:4. Sucrose gradient ultracentrifugation of cPDE from human leiomyoma of the uterus.

Crude cytosol (a) (5 mg/ml protein) was loaded in 200 ul aliquots and the centrifugation and the assay of enzyme activity were performed as described in the Methods. Specific activity of cPDE was measured at 1 uM cAMP (●) and 100 uM cAMP (■). Protein (▲) was measured in mg/ml using the Bio-Rad kit. Purified isoenzymes (b), which were the bands 2-6 observed on native polyacrylamide-gel electrophoresis, were loaded in 200 ul aliquots (0.5 mg/ml protein) after affinity chromatography, ammonium sulphate precipitation and equilibration in buffer A. The enzyme activity was measured at 1 uM cAMP (●) and protein (▲) was determined by the micro-method of the Bio-Rad kit. The results shown are representative ones of three separate runs.



Soluble cPDE activities of eluted bands of leiomyoma cytosols run on non-denaturing gels.

The various activity bands of the leiomyoma were eluted and assayed for cPDE activity (only cAMP tested) and summarized below (Table 5:2)

Table 5:2. Elution of the cPDE forms from the unstained, excised cPDE activity bands of leiomyoma cytosol on 7.5% gels. The lower bands 3-6 were combined into two segments of bands 3-4 and 5-6. The effect of added calmodulin (30 units per assay) was also tested. The units are in pmol/mg protein/min.

Band No.	1 uM cAMP	SD	1uM cAMP + calmodulin	SD
1	465	58	532	72
2	782	72	830	84
3-5	1210	107	1290	116
5-6	449	32	403	33

There did not appear to be increased cPDE activity when the eluted forms were incubated in the presence of added calmodulin plus calcium (present in the assay buffer). This suggested that the calmodulin-bound cPDE forms were still bound to calmodulin after the electrophoresis and elution steps.

Kinetic properties of some of the separated forms of cPDE.


The enzyme forms, designated band 1 and 2 in polyacrylamide-gel electrophoresis (Figs. 5:1a and 5:1b) were obtained by preparative native polyacrylamide electrophoresis followed by elution of the enzyme from the gel. The Km and Vmax values for the forms were calculated from Eadie-Hofstee plots with cAMP as substrate. The plot for form 1, which displayed linear kinetics, is shown (Fig. 5:5). The Km was found to be 5 ± 0.4 uM and the Vmax 26 ± 3.5 nmol/mg/min protein. The band 2 form did not display linear kinetics (Fig. 5:5) and the data was indicative of either two enzyme forms or apparent negative co-operativity. The calculated Km values were 5.8 and 37 uM with respective Vmax values of 12 and 64 nmol/min/mg protein. The band 5 form was also studied after elution and showed non-linear kinetics with an apparent Km range of 17-24 uM and Vmax range of 85-125 nmole/min/mg protein. The anomalous kinetics may, however, have resulted from contamination of this form with those of band 3 or 4 since these shared closer Rf values and, as a result, were difficult to separate when cutting strips from preparative gels.

Table 5:3. cPDE kinetic data for bands 1,2 and 5 of human leiomyoma of the uterus.The bands were eluted from native polyacrylamide gels (7.5%) and assayed for enzyme activity between 0.5 uM and 100 uM cAMP. The data are the mean ± s.d. of three experiments except for band 5 which is expressed as a range.

Band No.	Km (uM)	s.d.	Vmax (nmol/mg/min)	s.d.
1	5.0	0.3	26	3.5
2	5.8	0.95	12	1.6
2	37.0	5.0	64	11.5
5	17-24		85-125	

It appeared that the band 1 form exhibited a higher affinity for cAMP although the Vmax for band 5 was the greatest.


Fig.5:5. Kinetics of cPDE activity of two forms of the enzyme in human leiomyoma of the uterus (cAMP tested).
Enzyme activity was derived from extracted bands 1 and 2 from native polyacrylamide-gel electrophoresis and was assayed at cAMP concentrations between 0.5 uM and 100 uM. Results (a) of a single representative experiment, drawn as an Eadie-Hofstee plot, are shown for band 1. The two kinetically distinct forms of band 2 (b) are presented as an Eadie-Hofstee plot after reconstruction of the raw data according to Spears *et al.* (1971).



Partial purification of the major form of cPDE in cytosols of human leiomyoma.

The specific activity of the high affinity component of the crude leiomyoma supernatant was 0.036 nmol/min/mg protein. Fractionation of the cytosol on DE-52 anion exchange resin between 0.1-0.4 M NaCl followed by theophylline-Sepharose and sucrose gradient ultracentrifugation increased the specific activity to 2.68 nmol/min/mg protein that is a 74 fold purification. It is impossible to calculate an accurate recovery for bands 3-6 because of the presence of the other forms in the crude leiomyoma cytosol. Purification through the same steps described above except that the sucrose gradient centrifugation was replaced by preparative electrophoresis (Fig. 5:6(a)) as described in materials and methods produced enzyme which on SDS gel-electrophoresis yielded the pattern of four bands at Mr 65000 ± 3000, 50000 ± 2000, 44000 ± 2000, 41000 ± 2000 (Fig. 5:6(b) and (c)). The protein standards for SDS electrophoresis were phosphorylase b (P) 94000, albumin (A) 67000, ovalbumin (O) 43000 and carbonic anhydrase (CA) 30000. The molecular masses of the subunits were determined from the method of Weber and Osborn (1969).

Fig. 5:6. (a) Sample of affinity purified fraction of leiomyoma (0.1-0.4 M NaCl eluate from DE-52) was run at 7.5 % gel concentration for cPDE activity depicting band 2 and cluster of bands 3-6. The (b) figure shows the subunits (Coomassie 3DS) of the purified cPDE of leiomyoma in lane 1, a blank with buffer in lane 2 and protein standards in lane 3 (phosphorylase b P, albumin A, ovalbumin O carbonic anhydrase CA) and albumin in lane 4. The (c) figure depicts the same as (b) but the more sensitive silver stain was applied.



Discussion

The initial part of this discussion will compare the present results with purified human and mammalian cPDE enzymes described in other tissues and cells. The second part will summarize the findings on either crude or partially pure preparations from human and mammalian uterine tissues.

Human leiomyoma soluble cPDE activity could be resolved into seven bands of enzyme activity by electrophoresis at lower gel concentrations. This was observed for both cAMP and cGMP at a substrate concentration of 100 uM. Indeed, this pattern, except for band 7, was characteristic in over five separate preparations of this tissue. Homogenizing the tumour in buffers A, B, or C also resulted in the observed pattern of activity for either cAMP or cGMP (Fig. 5:1b). The measured relative mobilities (Rf) for cAMP or cGMP were indistinguishable. Invariably, band 5 exhibited the majority of the staining activity for both cAMP and cGMP. Caution in the interpretation of this observation is necessary since it has been reported that lead ions in the incubation medium inhibit some forms of cPDE more than others (Nemoz *et al.* 1983). I did utilize half the concentration of lead ion, 1.5 mM instead of 3 mM, in an attempt to reduce the possibility of this effect. In addition, elution of cPDE from unstained gel slices, showed the highest activity to be associated with band 5 for both low and high Km forms (1 uM and 100 uM cAMP)(for results at 1 uM cAMP see Table 5:3). Multiple forms, about 5, of cPDE have been described in

rat brain using the technique of isoelectricfocusing on granular gel plates (Nemoz *et al.* 1985).

Ferguson plot analysis of the leiomyoma activity bands revealed four different molecular mass species and a cluster of four charge forms in the group of bands 3-6. Evidence that the Band 1 form was a distinct enzyme might be summarized as follows: 1) This form was eluted at 0.1 M ionic strength from DE-52 cellulose 2) The other 0.1-0.4 M ionic fraction, which contained bands 2-7, when further purified on theophylline-Sepharose did not show interconversion to Band 1 when analyzed on native polyacrylamide-gel electrophoresis (Fig. 5:6(a)). 3) Kinetic analysis of the form indicated linear Michaelis-Menten kinetics from Eadie-Hofstee plots and a higher affinity for cAMP compared with band 2. 4) The form was inhibited by preincubation with EGTA (1 mM). 5) Band 1 exhibited the highest molecular mass of apparent Mr of 229000 ± 4000 and Yo = 2.49 but this higher value was insufficient to be explained as simple aggregation of the lower molecular mass isoenzymes 3-6 which were also EGTA inhibited. There have been reports in the literature, based on different tissues and species, of calcium/calmodulin dependent forms, that is type I cPDE enzymes, hydrolyzing both nucleotides with similar native molecular masses of about 240000 (Hidaka and Asano 1976; Tucker *et al.* 1981) to the band 1 form. The reported Mr of around 200000 for cGMP-stimulated phosphodiesterases in calf liver (Yamomoto *et al.* 1984(a)) (type II cPDE) and 240000 for bovine adrenal and heart tissues (Martins *et al.* 1982)(type II cPDE)

are also not dissimilar to that of the band 1 form. I am, however, unable to comment on the effect of cGMP on the hydrolysis of cAMP by the various enzyme forms in leiomyoma since this was not examined.

Band 2 appeared to be a distinct form of cPDE particularly in the characteristic of being totally impervious to preincubation with EGTA (Fig. 5:1(c) and (d)). It could therefore be regarded as calcium independent and hence calcium/calmodulin independent. Furthermore, the extracted enzyme from non-denaturing polyacrylamide electrophoresis gels did not show increased activity in the presence of exogenous calcium and calmodulin. The anomalous kinetics with approximate Km values for band 2 of 5.8 uM and 37 uM are not dissimilar from those reported for the enzyme(s) in crude cytosols of rat uterus of 3 uM and 20 uM (Strada *et al.* 1981) or values of 2 and 36 uM found for soluble uterine phosphodiesterase from diethylstilbestrol-treated immature rats (d'Auriac and Meyer 1973). The apparent molecular mass reported by Strada and associates (1981) was lower, that is 7.0 S (ca. 130000-140000) compared to my calculated Mr of 186000 $\pm$ 4000 for band 2. An earlier report by Gardner and co-workers (1978) on rat uterus yielded a higher value for the Sedimentation Coefficient of the enzyme namely 7.9 S.

Bands 3 to 6 were characterized as charge species of apparent Mr 174000 $\pm$ 4000. These forms were all largely inhibited by preincubation with 2 mM EGTA, although lower concentrations, <1 mM, did not inhibit forms 3 and 4 very strongly (Fig. 5:1(c)).

This would imply these forms are calcium/calmodulin dependent (Ho *et al.* 1977). Anomalous forms of different Rf were observed in the cytosol incubated with EGTA (Fig. 5:1(c) and (d)). Clearly more detailed research on these species is required. Since it was the bands 3-6 which were partially purified and shown to consist of subunits of apparent Mr 65000 ± 2000, 59000 ± 3000, 45000 ± 2000 and 41000 ± 2000 on SDS electrophoresis, certain inferences may be drawn. I was unable to detect the presence of calmodulin in the purified fraction when subjected to SDS electrophoresis under the conditions used. This would suggest that calmodulin was either lost during the earlier stages of the purification procedure or was tightly bound to the cPDE subunits on the SDS gels, which is unlikely.

The overall purification of the leiomyoma enzyme was only 74 fold because heavy losses in enzyme activity were incurred. This was probably due to the relatively low starting protein concentration (35 g/run) and the very labile nature of the isoenzymes. The two lower molecular mass subunits might be the proteolytic fragments described by Tucker and co-workers (Tucker *et al.* 1981) for the type I(a) cPDE enzyme even though PMSF was present in all solutions. The structure of the native soluble human leiomyoma isoenzymes could theoretically be composed of two subunits of type I cPDE and two calmodulin molecules A_2C_2 (ca. a mass of 160000)(La Porte *et al.* 1979; Sharma *et al.* 1984). I do not consider these forms to be similar to the 175000 form (type IV cAMP PDE) isolated in human platelets (Umekawa *et al.* 1984; Grant and Colman, 1984) as

their reported Km was much lower, that is, <0.4 uM compared to my range of 17-24 uM. In addition, the leiomyoma forms appeared to be equally active when gels were incubated in the presence of either cAMP or cGMP and they appeared to be calcium/calmodulin dependent because of their sensitivity to EGTA (see Chapter 1 for description of type IV cAMP PDE). The cAMP PDE (type II cPDE) isolated in calf liver with Mr of 174000 by Yamomoto and co-workers (1984(a)) more closely resembles the leiomyoma enzyme also exhibiting non-linear kinetics with an apparent low Km component of approximately 2 uM. The authors did not report the Km of the low affinity component. Nonetheless, the leiomyoma isoenzymes appear to be calcium/calmodulin dependent which is not the case with type II cPDE enzymes. It is interesting to speculate that three of the forms of the bands 3-6 could represent the two homodimers and one heterodimer described in bovine brain of a calmodulin-dependent cPDE (type I(a) cPDE (Sharma *et al.* 1984; Hansen and Beavo 1986). The activity form, represented by band 7, was not always present, particularly in the leiomyoma, and might represent a proteolytic fragment which nonetheless was inhibited by EGTA, unlike the fragments described by Tucker and associates (1981) which are not.

The latter part of this discussion centres on research related to uterine cPDE forms and the findings of five different groups will be summarized.

The first group, reported once on the levels of cPDE and adenyl cyclase in rat uterus during the oestrus cycle (Sim and

Chantharaksri 1974). These authors reported that in crude homogenates both enzyme activities were highest at met-oestrus but lowest at pro-oestrus suggesting steroid hormone control of the two enzymes.

The second group, based their studies on cPDE in crude supernatants of rat uterus, where there appeared to be at least two different kinetic forms (only cAMP hydrolysis was measured) both of which were inhibited by ATP (d'Auriac and Meyer 1973). It also appeared that, angiotensin II at levels that could cause contractions of rat uteri, had no effect on the soluble cPDE activities.

A third group have extensively studied cPDE, using both cAMP and cGMP, in monkey uterine tissues at different stages of development (Beatty *et al.* 1979). Beatty and co-workers have reported that the majority of the cPDE activity is associated with the supernatant (about 80%) in agreement with the findings by Bar (1974) in the rat myometrium supernatant and Leroy and associates (1985) in supernatants from human pregnant myometrium. They have also shown that cAMP metabolism in myometrium was altered during the menstrual cycle (Beatty et al. 1977) suggesting regulation of cAMP levels to be under steroid hormone control. The Beatty team published results of non-linear kinetics for cAMP hydrolysis in monkey myometrium but did not characterize the cPDE further or report Km or Vmax values (Beatty *et al.* 1976). This latter team did however provide further characterization of cPDE in their later comparative study of spayed macaque myometrium versus pregnant

myometrium (Beatty *et al.* 1979). The enzyme in crude cytosols of pregnant and spayed myometrium gave anomalous double-reciprocal plots of apparent Km values for cAMP of 4.6 to 18.7 uM and 2.9 to 24.9 uM respectively. The Vmax values for both tissues were very similar and were about 6 nmol/min/mg protein. My Vmax values for the separated soluble cPDE forms of human leiomyoma (Table 5:3) were all considerably higher and ranged from 12 to 125 nmol/mg protein/min. Linear kinetics for cGMP hydrolysis was observed for the enzymes in soluble pregnant myometrium (Km=7.6 uM, Vmax=9.3 nmol/min/mg protein) but not always for the enzyme in spayed monkeys, suggesting more than one form of the enzyme, as would be expected in crude cytosols. Furthermore, cGMP at low concentrations, could stimulate cAMP hydrolysis which is a feature of the type II cPDE form. Finally, the group concluded that levels of cPDE were raised in pregnant myometrium compared to spayed myometrium expressed on the basis of DNA content but the situation was reversed when comparisons were based on either wet tissue mass or nitrogen (protein) content (Beatty *et al.* 1979).

The fourth team have based their studies on supernatants from homogenates of rat uterine tissues and first suggested that in this tissue cAMP PDE (cGMP hydrolysis was not measured) was under the control of steroid hormones (Stancel *et al.* 1975). Their subsequent studies on partial purification of rat uterus cPDE using DEAE-cellulose ion-exchange chromatography followed by sucrose gradient ultracentrifugation consistently revealed a

single form of cPDE of about 7.9 S, as evaluated by sedimentation in the ultracentrifuge, which could hydrolyze both cAMP and cGMP (Gardner *et al.* 1978). This form, however, is in contrast to the report previously mentioned on crude supernatants of rat uterus (d'Auriac and Meyer 1973). A further study by the fourth group, based on limited proteolysis studies of the enzyme using trypsin, indicated the catalytic sites for cGMP and cAMP to be separate (Strada *et al.* 1981). Of particular importance was their finding that limited proteolysis of one of the DEAE-cellulose eluted forms resulted in two forms when rerun on DEAE-cellulose. This report by Strada and colleagues (1981) was disputed by Keravis and colleagues (1981) who reported non-linear kinetics for cAMP of Km 3 to 20 uM and linear kinetics for cGMP with a Km of about 3 uM. These Km values are not dissimilar to those reported by d'Auriac and Meyer (1973)(2 and 36 uM for cAMP) or to those reported by Beatty and associates (1979) or indeed to my own findings. The fourth group did not show stimulation of cAMP hydrolysis by low concentrations of cGMP (see Beatty *et al.* 1979).

The final group have studied cPDE in either human myometrium, human placenta or cultured myometrial cells (Ferre *et al.* 1975; Ferre *et al.* 1977; Vallet-Strouve *et al.* 1984; Leroy *et al.* 1985). At least two cPDE activities were reported for homogenates of human placenta and imidazole was demonstrated to stimulate the lower affinity form (Km = 1.36 mM)(Ferre *et al.* 1975). The high affinity cAMP PDE (cGMP hydrolysis was not

measured), exhibited a Km of 10 uM and Vmax of 0.63 nmol/min/mg protein, and was weakly inhibited by prostaglandins but strongly inhibited by progesterone. They repeated these experiments in cytosols of human pregnant myometrium and, from the crude homogenates, two forms of cAMP PDE were isolated (higher affinity form with Km of 12.6 uM, Vmax of 0.5 nmol/min/mg protein; lower affinity form Km of 90 uM, Vmax of 1.0 nmol/min/mg protein). The stimulatory effect of imidazole for the activity of the lower affinity enzyme was apparent and, as for placenta, indomethacin and progesterone (non physiological concentrations *in vitro*) were found to inhibit the higher affinity form. Of interest, oestradiol 17B and several oestradiol derivatives were shown to have negligible inhibition on cPDE *in vitro* (Ferre *et al.* 1977). Apparently, progesterone has been shown both *in vivo* and *in vitro* to inhibit human uterine contraction (Ferre *et al.* 1977). Subsequent work by this group on myometrial cells in culture, have shown that physiological concentrations of either progesterone or oestradiol inhibited cAMP PDE activity (cGMP hydrolysis was not measured) (Vallet-Strouve *et al.* 1984). This effect has also been reported by others (Stancel *et al.* 1975; Beatty *et al.* 1979). Furthermore, *in vivo* administration of oestrogen decreased cPDE activity in rat uterus (Gardner *et al.* 1978) whereas only progesterone decreased PDE activity in macaque myometrium (Beatty 1978). Recently, a report on murine oocyte maturation has demonstrated that either progesterone or Beta-oestradiol or testosterone inhibited cPDE activity in

oocyte extracts (Kaji *et al.* 1987). Clearly there appear to be some conflicting results regarding the inhibitory effects by progesterone and oestradiol on the cPDE activity. It is possible that the discrepant observations might reflect differences in the oestradiol to progesterone ratio which varies considerably either during the menstrual cycle, or stages of pregnancy or as a result of species differences. Also, discrete forms of cPDE exhibiting differential inhibition by the two hormones might be synthesized either during development, the menstrual cycle, pregnancy or as a result of species differences.

The reported kinetics for cAMP hydrolysis in supernatants of human myometrial cells in culture were non-linear in agreement with the other groups (Stancel *et al.* 1975; Beatty *et al.* 1976; Ferre *et al.* 1977; Beatty *et al.*1977). The apparent Km values of the soluble cPDE in myometrial cells for cAMP were 0.8 uM and 40 uM with respective Vmax values of 0.06 and 1.4 nmoles/min/mg protein. The latter values increased by 100% following several successive subcultures. Futhermore, a decrease in cAMP PDE activity was noted with high densities of cells and *vice versa* (Vallet-Strouve *et al.* 1984). The most recent report by this group has been based on longitudinal sections of human myometrium at the end of pregnancy (Leroy *et al.* 1985). It appears that less than 10% of either cAMP or cGMP PDE activity was associated with the particulate fraction. Two soluble cPDE forms were isolated using DEAE-cellulose chromatography and one form (peak I) most closely resembled the

type I cPDE and the second eluting form (peak II) appeared to be similar to the type IV cAMP PDE. I did not detect the type IV cPDE in the cytosols of human nonpregnant myometrium using my gel method coupled to the cPDE activity stain. That is, I observed the same cPDE activity profile for human myometrium cytosol in the presence of either cAMP or cGMP.

Sucrose gradient ultracentrifugation of the soluble fraction of "end of pregnancy" myometrium (Leroy *et al.* 1985) gave rise to three peaks A,B and C of which, C, appeared to be type IV cAMP PDE. The peaks A and B (hydrolyzing both substrates) could also be resolved on sucrose gradients when only peak I was loaded suggesting at least three distinct forms of cPDE in human pregnant myometrium. Both substrates exhibited non-linear kinetics with apparent Km values for cAMP in crude supernatant of 2.8 uM and 57 uM ; for cGMP of 0.83 uM and 12 uM. The peaks I and II obtained on DEAE chromatography revealed different kinetic properties: Peak I (type I cPDE) exhibited Km values for cAMP of 28 uM and 4200 uM with respective Vmax values of 0.4 and 2.0 nmol/min/mg protein; Km values for cGMP of 1.7 uM and 16 uM with respective Vmax values of 0.04 and 0.22 nmol/min/mg protein; Peak II (type IV cAMP PDE) gave Km values for cAMP of 10 uM and 1000 uM with respective Vmax values of 0.6 and 4.1 nmol/min/mg protein. The estimated molecular masses of A, B and C from sucrose gradient studies suggested 110000 daltons, 47000 daltons and 18000 daltons respectively. It appears that, in agreement with Leroy and associates (1985), in human leiomyoma versus "end of pregnancy" myometrium there are

at least three distinct forms of cPDE and that there is more than one form of type I cPDE. My results, however for cAMP hydrolysis by band 1 form (type I cPDE) and bands 3-6 (type I cPDE) yielded much higher affinities for the substrate and also higher Vmax values in comparison to Leroy and worker's (1985) partially purified "peak I". In fact my values do not differ much from their results reported for crude supernatants of human pregnant myometrium. However, the molecular mass values are radically different. Whether these discrepancies reflect true differences between cPDE forms in human leiomyoma cytosols versus human pregnant myometrium cytosols or are the result of differences in methodology remains to be resolved. It is worth noting, though, based on my own experiences with sucrose gradient ultracentrifugation, that if leiomyoma supernatants, either crude or partially pure, were frozen and rethawed or stored for more than three weeks at -70° C, lower molecular mass forms of cPDE were detected and were regarded as proteolytic but active cPDE enzyme fragments.

Further observations, based on the activity pattern in non-denaturing polyacrylamide electrophoresis of human leiomyoma cyclic nucleotide phosphodiesterase revealed that the cPDE enzyme pattern did not differ for either human myometrium or normal uterus (composed of both myometrium and endometrium). In addition, the myometrium tissue displayed denser staining in all activity bands at the same soluble protein concentrations compared to that of leiomyoma from the same patient (Figs. 5:1a, 5:1b, 5:1c, 5:2b and 5:2c) indicating higher cPDE

activity in myometrium tissue. This finding was most interesting in that when I studied the oestrogen and progesterone receptor content of the two tissues both receptor types were significantly higher in leiomyoma ($p = 0.0002$) (Sadan *et al.* 1987). Therefore, there appears to be an inverse correlation between oestrogen/progesterone receptors and the level of cPDE activity which is in agreement with the findings of Etingof *et al.*(1984) who studied oestrogen receptors and cPDE levels in both human and uterine tissues.

Of interest, a recent report on the kinetic characteristics of cPDE in cruda cytosols of human pregnant and nonpregnant myometrium has shown a markedly lower Vmax in the pregnant tissue (Kofinas *et al.* 1987). This suggests steroid hormonal influences on the soluble cPDE activity in human myometrium during pregnancy. Nonetheless, in view of the multiple forms of cPDE that I found in human myometrium cytosols, studies of cPDE kinetics based on crude myometrium cytosols are difficult to interpret.

Characterization of soluble uterine cyclic nucleotide phosphodiesterases in rat uterus has only revealed one form with anomalous kinetics (Gardner *et al.* 1978; Strada *et al.* 1981). Several recent reports in the literature indicate that there are distinct species differences in the four main types of soluble cyclic nucleotide phosphodiesterases, referred to previously, in either one or more of the following: kinetic parameters, physical properties or regulation by pharmacological agents (Geremia *et al.* 1984; Mullaney and

Clegg, 1984; Mutus *et al.* 1985; Pyne *et al.* 1986).

The diverse molecular mass species and charge forms in the human uterine tissues that I studied might also be ascribed to: 1) the release of membrane-bound forms due to the extremely low temperature of liquid nitrogen (-196^{o} C) (not previously used to my knowledge) and pulverizing of the tissue in this medium. 2) The result of endogenous proteolysis during processing. The very consistent activity pattern observed in all the tissue cytosols would tend to refute this but I cannot discount it entirely. It was found that repeated freezing and thawing of the cytosols considerably lowered the intensity of staining and resulted in poorly resolved bands. In addition, more anionic bands were sometimes observed.

In conclusion, I consider that human leiomyoma possess three distinct soluble cPDE forms and that there also exist isoenzymes. The human myometrium and uterus also contain these three forms. The method of slab gel electrophoresis described here may prove useful in the analysis of cPDE enzymes in tissue available in only small quantities, as is the case in man. It should also be of potential value in the assessment of inhibitor sensitivities in different tissues.

CHAPTER SIX

CONCLUSIONS

Since most of the work in this thesis has been discussed in separate sections, it would be pertinent to consolidate the information in order to gain a clearer perspective of the soluble cPDE enzymes in mammalian tissues and the possible clinical significance of my findings. A brief overview of cPDE forms in more primitive organisms, where some gene sequencing has been achieved, will be summarized prior to my findings of the soluble cPDE enzymes in mammalian tissues.

The cPDE enzymes of primitive eukaryotes.

In view of the overwhelming evidence that cPDE enzymes hold key positions in cellular biochemical regulation and the fact that they are the only specific catabolic enzymes for the ubiquitous substrates, cAMP and cGMP, I would tend to expect the cPDE enzymes to be highly conserved between species. In other words, perhaps only very few mutations would be advantageous and, most in fact, deleterious to the vital function of these regulatory enzymes. Thus reported findings of cPDE enzymes in more primitive organisms ought to answer some of the questions that have arisen from *in vitro* studies based on cPDE enzymes in

mammalian tissues. Certainly, *in vitro* studies of cPDE enzymes in eukaryotes have demonstrated multiple forms of cPDE. In my opinion, however, a disconcerting problem has been the fact that purified cPDE enzymes of the same classification can differ considerably : th respect to kinetic parameters and physical characteristics between mammalian species. A case in point would be the purified soluble type II cPDE in bovine liver (Martins *et al.* 1982) compared to the soluble purified type II cPDE in rat liver (Pyne *et al.* 1986) which represents just one of many dicrepancies (see Introduction). In brief, are the multiple cPDE forms observed *in vitro* between species either the result of: different structural genes; the same gene but considerably variant as a result of accummulated mutations; differential processing of the mRNA's in the nucleus or cytoplasm; differential covalent modification of the translation products; or artifacts arising from different *in vitro* separation techniques? Furthermore, the multiple forms observed within different tissues within the same species could be either: different structural genes that may be differentially expressed or suppressed in specific tissue types; one structural gene that is differentially processed in certain tissues; or artifacts that arise from variations of *in vitro* methodologies.

Recently, a partial or complete DNA sequencing has been attained in *Drosophila melanogaster* (Davis and Kauvar; Chen *et al.* 1986), in yeasts (Sass *et al.* 1986; Nikawa *et al.* 1986) and slime mold (Lacombe *et al.* 1986). It appears that there are two

genetically distinct forms of cPDE in *Drosophila* (Walker and Kiger 1984; Chen *et al.* 1986). The form I enzyme resembles the mammalian type I cPDE in mass but not kinetic parameters and is regulated by a calmodulin-like protein. The form II cPDE, which is associated with the dunce gene mutation, most closely resembles the mammalian type IV cPDE. In the yeast, *Saccharomyces cervisiae*, two genes, PDE 1 and PDE 2, have been cloned and encoded respectively by Nikawa *et al.* ( ) and Sass *et al.* (1986). An interesting study, based on comparisons of partial amino acid sequences between type I cPDE from mammalian brain, type II cPDE from bovine heart with those protein sequences (derived from the nucleotide sequences of the genes) of the yeast PDE 2 gene and the *Drosophila* form II gene, has demonstrated a conserved domain in all four of the cPDE enzymes from these widely differing species (Charbonneau *et al.* 1986). The yeast PDE 1 gene, however, showed no apparent homology with the above four cPDE enzymes but did show homology with the *Dictyostelium discoideum* extracellular cPDE enzyme (Nikawa *et al.* 1987). The latter finding suggests that the slime mold cPDE might represent a structural gene for cPDE not found in higher organisms. It should be noted that the above cPDE genes which have been sequenced in the more primitive eukaryotes might not be the only genes present and it is quite possible that new cPDE genes remain to be identified.

In summary, therefore, there appears to be conservation of amino acid sequences in some of the cPDE enzymes classified as type I, type II and type IV (Charbonneau *et al.* 1986). Evidence

presented by the Charbonneau team (1986) has implied that, since the homology between mammalian type I cPDE and and type II cPDE was rather limited, the respective structural genes were probably discrete. The use of polyclonal antibodies by Sarada and associates (1982) has also demonstrated shared homology between mammalian type I cPDE enzymes with those of mammalian type IV cPDE. One exception to the above was a report by a group (Shenolikar *et al.* 1985) in which they were unable to demonstrate any cross reactivity, using Western blot analysis, between soluble bovine brain calmodulin-stimulated cGMP PDE (type I (c) cPDE, see Introduction) and soluble type IV cPDE of dog kidney. The type III (a) and (b) cPDE enzymes are uniquely associated with the lens of the eye and there is strong evidence, derived from two dimensional tryptic peptide mapping (Takemoto *et al.* 1984), for interspecies conservation of the type III (a) cPDE. Furthermore, Barbehenn and associates (1985) have demonstrated antigenic similarity between type III (a) and type III (b) enzymes. Clearly, evidence thus far, seems to favour the theory that the four main cPDE classifications, type I, type II, type III and type IV cPDE represent discrete structural genes, some of which share a degree of sequence homology.

The findings outlined above, however, have still not satisfactorily resolved the reported differences in properties within a cPDE type either between species or within different tissues of the same organism. It is indeed likely that many of these differences can be attributed to artifacts arising from

in vitro methodology rather than to true physical variations in the cPDE forms. Some of the major problems, not the least of which are the facts that the cPDE enzymes are both highly active and present in small amounts, include: the sensitivity of cPDE enzymes to either endogenous or exogenous proteolysis and the ability of fragments to remain catalytically active, such as the irreversible activation of type I cPDE by proteolysis with concomitant loss of responsiveness to calcium/calmodulin (review, Krinks *et al.* 1984); either similar or differential sensitivity of cPDE enzymes to an immense array of agents ranging from endogenous cations, sulphydryl compounds, small peptides, phospholipids and hormones to exogenous effectors such as imidazole and many exogenous inhibitors (see Introduction); the effect of pH which irreversibly inactivates cPDE under acidic conditions (pH < 5) (Strada *et al.* 1978); hydrophobic interactions (Van Inwegen *et al.* 1976); and temperature sensitivity (Strada *et al.* 1978).

The above difficulties explain, in large part, why so few studies of the cPDE enzymes in human and murine tissues have been published. Since the study of these enzymes in human and murine tissues is only possible with small tissue specimens, it was necessary to adapt the method of non-denaturing polyacrylamide gel electrophoresis coupled to the specific cPDE activity stain. This technique can enable as little as 0.5 mg of soluble protein to effect a separation of the soluble cPDE enzymes.

Comparative findings of the cPDE forms in murine and human tissue cytosols.

My studies, which have dealt mainly with the characterization and partial purification of the soluble cPDE enzymes in murine tissue cytosols and human mammary and uterine cytosols, consistently demonstrated a significant portion of the total soluble cPDE activity to be calcium/calmodulin dependent. This was proved by inhibition to the following; EDTA, a calcium chelator; EGTA, also a calcium ion chelator; and the specific inhibitor, trifluoroperazine. Furthermore, when these cPDE activity forms were excised and eluted from gels, the enzymes were not further activated in the presence of exogenous calcium/calmodulin which suggested that nd PAGE did not separate calmodulin from the type I cPDE forms. Of those cPDE activity forms that were shown to be calmodulin/dependent (type I cPDE), there appeared to be activity bands that were common to all the tissue cytosols I studied. In the murine tissue cytosols, the Rf range of a common calcium/calmodulin sensitive cPDE activity band was 0.28-0.35 at 7.5% gel concentration (Table 2:4, repeated below). In human mammary tumour cytosols (Fig. 4:1a (c)) the EGTA-sensitive cPDE activity bands, 2 to 4, exhibited an Rf range of 0.28-0.34 at 7.5% gel concentration which is within the same range as that found for the common calmodulin/dependent cPDE activity band in all the murine tissue cytosols. Similarly, the normal human mammary cytosol bands 2 and 3 had an Rf range of 0.31-0.34 at 7.5% gel

concentration. Lastly, the human uterine cytosol bands 3 to 5, also EGTA-sensitive (Fig. 5:1(a)), exhibited an Rf range of 0.29-0.35 (results summarized below in Tables 2:4, 6:1 and 6:2). The two faster migrating cPDE bands observed only in murine mammary tumour cytosol (5 to 6, Rf range 0.34-0.41 and EDTA-sensitive) were, however, apparent in all the human cytosols, except for the fastest migrating cPDE activity form in normal human mammary cytosol, and are summarized: human mammary tumour cytosol (bands 5 to 4, Rf range 0.35-0.40); normal human mammary cytosol (band 4, Rf range 0.35-0.36); uterus, myometrium and leiomyoma cytosols (bands 6 to 7, Rf range 0.36-0.41) (see Tables 2:4, 6:1 and 6:2.)

Unfortunately, I am unable to comment on the comparative molecular masses of the soluble cPDE forms, calculated by Ferguson plot, which were found to be higher for all the murine tissue cytosols. The reason for this is that I used protein standards that were optimized for running on nd PAGE at the start of the human tissue soluble cPDE studies and the new standards appeared to yield lower molecular masses. Nonetheless, my findings did reveal that the faster migrating cPDE activities appeared to be type I cPDE isoenzymes based on the facts that they were either inhibited by EDTA or EGTA and also the forms exhibited nearly identical molecular masses but possesed discrete charge densities (Yo) within the different tissue cytosols.

Another soluble cPDE activity band that was consistently demonstrated, using non-denaturing polyacrylamide gel

electrophoresis coupled to the cPDE activity stain, with the exception of murine brain cytosol, was the EDTA/EGTA-insensitive cPDE enzyme. When this cPDE activity form was excised and eluted from gels, the cPDE activity was not increased in the presence of exogenous calcium/calmodulin and I considered this cPDE activity to be calcium/calmodulin independent. In murine tissue cytosols, except for murine brain and the EDTA-insensitive band 1 of murine heart, the EDTA-insensitive cPDE activity bands in murine cytosols had an Rf range of 0.24-0.28. This same characteristic cPDE activity band had an Rf range of 0.25-0.27 in human mammary cytosols and an Rf range of 0.26-0.27 in human uterine cytosols (see Tables 2:4, 6:1 and 6:2 below). Kinetic analysis of this EGTA-insensitive form from human mammary tumour cytosols and human leiomyoma cytosols was achieved by eluting the band of interest (note: not exposed to lead ions) from polyacrylamide gels and then performing the cPDE radioassay. The kinetic parameters of the calcium/calmodulin independent cPDE enzyme in human mammary tumour cytosol (for cAMP, Km values of 3.8 uM and 50 uM with respective Vmax values of 4.5 and 25 nmol/mg protein/min) were not dissimilar from those reported for the same enzyme in human leiomyoma cytosol (for cAMP, Km values of 5.3 and 37 uM with respective Vmax values of 12 and 64 nmol/mg protein/min). In both tissues, the cPDE enzyme exhibited complex kinetics with apparent negative cooperativity. This suggests that the EGTA-insensitive cPDE activity band is either indicative of more than one cPDE enzyme, possibly not fully

separated on nd PAGE, or that the enzyme is allosteric with different catalytic binding sites.

The calcium/calmodulin cPDE activity band with the characteristic Rf range of 0.24-0.28 (molecular mass range of 168000 to 186000 daltons in human cytosols) which I observed in all the tissue cytosols, except murine brain, could be regarded as similar to either the soluble type II cPDE described in bovine heart (Martins *et al.* 1982) or calf liver (Yamomoto *et al.* 1983(a)) or a soluble type IV cPDE purified from human platelets (Umekawa *et al.* 1984). The soluble type II cPDE from heart has been reported to have a molecular mass of 240000 daltons and Km for cAMP of 30 uM (Vmax not given) (Martins *et al.* 1982). The similar enzyme in calf was demonstrated to have a mass of 201000 daltons, Km for cAMP of about 30 uM and a Vmax of 170 umol/mg protein/min (Yamomoto *et al.* 1983(a)). The soluble type IV cPDE has been shown to have a mass of 175000 daltons, Km of 0.34 uM (with 0.11 uM for cGMP) and a Vmax of 85 nmol/mg protein/min (Umekawa *et al.* 1984). The very high Vmax (one thousand fold higher, in umol as averse to nmol/mg protein/min) reported for the calf liver cPDE would seem to suggest a closer resemblance of my form to the unusual soluble type IV cPDE. The soluble type IV enzyme in human platelets is unusual in that it has been demonstrated to exhibit an almost equal hydrolysis rate for both cAMP and cGMP (Umekawa *et al.* 1984) whereas the type IV cPDE enzymes tend to be cAMP-specific (see Introduction section on type IV cPDE's). Thus the dichotomy still remains as to why the soluble cPDE types vary

so considerably with respect to both physical and kinetic parameters in different species. Only further biochemical studies, using several different separation techniques, will possibly resolve this question.

Table 2:4. Relative mobility (Rf range) of the cPDE activity bands of murine tissue cytosols observed in 7.5% nd PAGE coupled to the cPDE activity stain as shown in Fig. 2:4a for both cAMP and cGMP. The results are the mean of four experiments.

	Rf range from top of gel	Tissue and band position (Fig. 2:4)
A.	0.095-0.11	kidney (1)
B.	0.12 -0.14	heart (1)
C.	0.24 -0.28	tumour (1), heart (2), liver (1), kidney (2),
D.	0.29 -0.35	bovine heart forms, heart (3), brain (1), liver(2), kidney (3), tumour (2)
E.	0.34 -0.36	tumour (3)
F.	0.37 -0.41	tumour (4)

Table 6:1. Relative mobilities (Rf range) of the cPDE activity bands of human mammary cytosols observed in 7.5 % nd PAGE coupled to the cPDE activity stain as shown in Fig. 4:1a (a). The calculated Rf values were the same for either cAMP or cGMP. The results are the mean of three experiments.

	Rf range from top of gel	Tissue and band position (Fig. 4:1a (a))
A.	0.25-0.27	normal breast(1), breast tumour (1)
B.	0.29-0.30	breast tumour (2)
C.	0.31-0.32	normal breast (2), breast tumour (3)
D.	0.33-0.34	normal breast (3), breast tumour (4)
E.	0.35-0.36	normal breast (4), breast tumour (5)
F.	0.39-0.40	breast tumour (6)

T.ble 6:1. Relative mobilities (Rf range) of the cPDE activity bands of human mammary cytosols observed in 7.5 % nd PAGE coupled to the cPDE activity stain as shown in Fig. 4:1a (a). The calculated Rf values were the same for either cAMP or cGMP. The results are the mean of three experiments.

	Rf range from top of gel	Tissue and band position (Fig. 4:1a (a))
A.	0.25-0.27	normal breast(1), breast tumour (1)
B.	0.29-0.30	breast tumour (2)
C.	0.31-0.32	normal breast (2), breast tumour (3)
D.	0.33-0.34	normal breast (3), breast tumour (4)
E.	0.35-0.36	normal breast (4), breast tumour (5)
F.	0.39-0.40	breast tumour (6)

Table 6:2. Relative mobility (Rf range) of the cPDE activity bands of human uterine cytosols observed in 7.5% nd PAGE coupled to the cPDE activity stain. The calculated Rf values were the same for either cAMP or CGMP. The results represent the mean of four experiments.

	Rf range from top of gel	Tissue and band position (Fig. 5:1(a))
A.	0.18-0.19	uterus (1), myometrium (1), leiomyoma (1)
B.	0.26-0.27	uterus (2), myometrium (2), leiomyoma (2)
C.	0.29-0.30	uterus (3), myometrium (3), leiomyoma (3)
D.	0.31-0.32	uterus (4), myometrium (4), leiomyoma (4)
E.	0.33-0.35	uterus (5), myometrium (5), leiomyoma (5)
F.	0.36-0.38	uterus (6), myometrium (6), leiomyoma (6)
G.	0.40-0.41	uterus (7), myometrium (7), leiomyoma (7)

From the above Tables it can be seen that the murine kidney cytosol band 1 cPDE form, murine heart band 1 and the human uterine band 1 represent unique cPDE forms with their own characteristic relative mobility under controlled conditions. It is possible that the murine kidney band 1 cPDE could represent an aggregate of murine kidney band 3 (both EGTA-sensitive) where the mass of band 1 was 395000 daltons and band 3 was 245000 daltons (Table 2:12). This would not appear

to be the case for either the murine heart soluble EDTA-insensitive cPDE forms (band 1 = 246000 daltons compared to band 2 = 227000 daltons) or for the EGTA-sensitive cPDE enzymes of human uterine cytosols (band 1 = 229000 daltons compared to bands 3-7 = 175000 daltons).

Biochemical considerations.

My experiments based on nd PAGE coupled to the cPDE activity stain showed identical cPDE profiles for either cAMP or cGMP in all the tissue cytosols I studied. The staining intensity was similiar for both substrates when incubated with either 50 to 300 uM cAMP or cGMP. At lower substrate concentrations (10 uM), however, there was evidence for increased cPDE activity in the presence of 10 uM cGMP compared to 10 uM cAMP when I studied this effect in murine heart cytosols (Figs. 2:5e and 2:5f). It is necessary to reiterate, though, that no new cPDE activity bands were observed to appear at different substrate concentrations. This has important implications: 1. It is highly unlikely that there are discrete cPDE enzymes for the hydrolysis of either cAMP or cGMP. In other words, the catalytic binding site(s) probably reside(s) in the same enzyme. 2. It is unlikely that there are discrete high affinity and low affinity cPDE enzymes.

For the type I cPDE, which seemed to be the predominant soluble cPDE activity in the tissue cytosols I studied, there is controversy as to whether cAMP and cGMP are hydrolyzed by

separate catalytic sites as suggested by Geremia *et al.* (1984) or by a common catalytic site (Sullivan *et al.* 1987). I am, at present, unable to offer any definitive proof although my observations are consistent with a common catalytic site for both cAMP and cGMP with possibly a discrete high affinity cGMP site that becomes active at low substrate concentrations. Also, my findings tend to discredit the still popular practice of reporting high affinity cPDE and low affinity cPDE as if they were separate enzymes. This is usually done by studying the cPDE enzyme kinetics in crude extracts and then attributing the non-linear plots to low and high affinity enzymes (see Kofinas *et al.* 1987; Kurtz *et al.* 1987; Moses *et al.* 1987). My findings unequivocally demonstrated at least three or more cPDE forms in all the tissue cytosols I studied (see results of gradient nd PAGE on murine brain, kidney and liver Fig. 2:7) and these cPDE profiles were identical for each tissue cytosol from 50 uM (10 uM in murine heart) to 300 uM substrate concentrations. This multiplicity of both cPDE enzymes and isoenzymes makes it difficult to determine the relative contributions of the specific cPDE forms in crude enzymes or extracts. Therefore some form of separation technique of the cPDE forms, prior to attempting enzyme kinetics analysis, seems to be a necessary prerequisite. As I stated previously (Multiple Forms section in the Introduction), DEAE-cellulose anion-exchange chromatography has almost always been the method of choice for the separation of cPDE forms, on the basis of net protein charge, prior to further purification. My experience with this technique,

indicated that large losses of cPDE activity were incurred during the DEAE c.romatography stage in addition to the overnight dialysis and concentraion steps. Thus, it is not a suitable method for small tissue samples. The adaptation and modification of the method of nd PAGE coupled to the cPDE activity stain to slab gels, has enabled the separation of cPDE forms to be attained with very little soluble protein (0.1 to 0.5 mg soluble protein). Furthermore, several samples from different tissue cytosols can be assessed on a comparative basis for their cPDE enzymes. Also, by means of preparative nd PAGE, it is possible to excise the separated cPDE bands (note, in the absence of lead ions), elute them from the polyacrylamide gel and then calculate the kinetic parameters of these separated cPDE forms. Another important consideration of the nd PAGE method is the fact that there was likely to be less breakdown of the cPDE forms by proteolysis as: 1. The tissues were processed in the presence of the proteolytic inhibitor, PMSF. 2. The tissue cytosols were not exposed to excessive dilution as occurs with DEAE-anion exchange and dialysis. 3. The slab gel electrophoresis runs were performed at a very low current at 4^{o} C in a period of only five hours.

It is interesting to note that, at no time did I detect any low molecular mass cPDE forms using the gel method (molecular masses > 162000 daltons). Thus, I did not detect the soluble cAMP-specific type IV cPDE of native molecular mass 60000 daltons (Thompson *et al.* 1979 (a)). This low molecular mass enzyme has also been described in the soluble fraction of human

lung (Moore and Schroedter 1982) although these latter authors speculated on the possibility of their enzyme being a catalytic subunit of a larger native enzyme. Of course, it cannot be entirely discounted that my gel method was not sensitive enough to detect the presence of this soluble type IV cPDE. When I performed sucrose ultracentrifugation (18 hour runs) on crude leiomyoma cytosols I detected three cPDE forms (Fig 5:4), one of which was a low molecular mass enzyme of 3.6 S (c.f. 46000 daltons). The seven forms that I detected, using the gel method, in leiomyoma cytosols, however, all had molecular masses of greater than 162000 daltons. It is also worth noting, based on my own experiences with sucrose gradient ultracentrifugation, that if leiomyoma supernatants, either crude or partially pure, were frozen and rethawed or stored for more than three weeks at - 70° C, lower molecular mass forms of cPDE were detected and regarded as proteolytic but active cPDE enzyme fragments.

Finally, the nd PAGE method might prove useful in the assessment of inhibitor/activator sensitivities of the cPDE activity forms of different tissues to a variety of compounds.

Clinical considerations.

From a clincal standpoint, I had hoped to be able to detect a soluble cPDE activity band that might be specific for tumour mammary cytosols. Indeed, the finding of two faster migrating cPDE forms in murine mammary tumour cytosols compared to other

normal murine tissue cytosols seemed promising. The work on human mammary cytosols also indicated two possible tumour associated cPDE forms with Rf ranges of 0.28-0.30 and 0.39-0.40. The subsequent observation of these two migrating cPDE forms in both normal myometrium and benign uterine tumours, however, suggested that these lower cPDE activity bands, might in fact, be type I cPDE isoenzymes associated with steroid hormone target tissues. An interesting finding, though, was the relatively reduced cPDE activity staining in leiomyoma cytosols compared to their normal myometrium cytosol counterparts. I have demonstrated the oestrogen and progesterone receptor content to be significantly higher in leiomyoma compared to myometrium ($p < 0.0002$) (Sadan *et al.* 1987). Therefore, there appears to be an inverse correlation between levels of oestrogen/progesterone receptors and the level of cPDE activity which is in agreement with the findings of Etingof *et al.* (1984). Also, this finding concurs well with that of Cho-Chung and associates (1978) who reported that cPDE activity was increased in regressing DBMA-induced rat mammary tumours. DMBA-induced tumours are known to posses high levels of functional oestrogen and progesterone receptors and regression of growth of these tumours is achieved by ovariectomy of the host which leads to withdrawal of steroid hormones and thus reduction of receptors. Of concern, therefore, is the fact that many breast cancer patients who have detectable levels of oestrogen/progesterone receptors in their tumours are often treated with the anti-oestrogen,

tamoxifen. This drug, however, has also been shown to be a type I cPDE inhibitor (Lam 1984) which would, at first, appear to be a disadvantageous effect in view of the fact that cPDE levels are observed to be increased in regressing DBMA-induced rat mammary tumours (Cho-Chung *et al.* 1978). Nonetheless, it is important to note that the levels of adenyl cyclase and cAMP are also increased in these regressing tumours (Cho-Chung *et al.* 1978). It seems, therefore, that despite the increased cPDE activity level, the overall concentration of cAMP is higher in regressing tumours. On the basis of this observation and assuming the increased cAMP levels to be associated with growth inhibition, it is interesting to speculate that tamoxifen in conjunction with either a drug that stimulates adenyl cyclase (a forskolin-like compound) or the use of site-specific cAMP analogues that have a much higher affinity for the cAMP receptor protein (Tagliaferri *et al.* 1988) might further increase the efficacy of tamoxifen in causing tumour regression. Alternatively, it should prove worthwhile to search for an anti-oestrogen that is inhibitory to all the cPDE types and is not solely specific for type I cPDE, as is the case with the calmodulin antagonist tamoxifen.

The assessment of the inhibitory effects of such an anti-oestrogen on soluble cPDE enzymes in mammary tumour cytosols could easily be performed with the nd PAGE technique coupled to the cPDE activity stain. The drug therapies proposed above might also be of value in the treatment of leiomyoma which are rapidly growing, steroid receptor positive benign tumours of the

uterus. This latter condition is usually treated surgically (Sadan *et al.* 1987).

Although I was unable to detect a specific tumour-associated soluble cPDE activity in mammary tissue cytosols, this was not the case for the nuclear-associated particulate cPDE in malignant mammary tissues (Chapter 3). This membrane-bound cPDE (only cAMP tested) was studied by using an ultracytochemical technique coupled to the cPDE activity stain. The nuclear-associated cPDE was strikingly observed in malignant mammary cells, to a lesser extent in fibrocystic mammary cells but was not apparent in normal human breast epithelial cells. This finding has important implications for possible therapeutic therapy against the mammary cancer cells. If the nuclear-associated cPDE could be characterized and a specific inhibitor identified then it might be possible to induce tumour regression by using the inhibitor to cause increased cellular cAMP concentration. This would be of particular value in those patients who have oestrogen/receptor negative tumours that are usually associated with more aggressive disease (Robinson et al. 1985). In short, anti-oestrogen therapy is ineffective against steroid receptor negative tumours and thus an inhibitor directed specifically against the nuclear-associated cPDE that I detected in all the malignant breast tumours, might be able to target those hormone-independent tumours. Of possible relevance to my findings is the recent report by Lichtner and associates (1987), who have demonstrated that the antimetastatic action of the pyrimido-pyrimidine analogue RA

233 (Rapenton) on tumour cell lines, is not merely the result of inhibition of platelet function but also the alteration in cAMP metabolism because of inhibition by RA 233 of type IV cPDE (Lichtner *et al.* 1987). It would be useful to perform further ultracytochemical experiments to establish whether this RA 233 inhibitor inhibits the nuclear-associated cPDE.

At present, I am studying a murine model which exhibits spontaneous hepatomas. There is sufficient normal liver for comparative studies between the normal liver and hepatoma tissues. I am using the methodology described in this thesis to study the soluble liver cPDE forms and I am developing new techniques to study the membrane-bound cPDE forms.

REFERENCES

Ahluwalia GS, Rhoads A Proceedings of the National Academy of Science. USA. Selective inhibition of cyclic AMP cyclic GMP phosphodiesterases of cardiac nuclear fraction. *Biochem Pharmacol* 1982;31:665-669

Ahluwalia GS, Rhoads AR, Lulla M. Particulate cyclic 3',5'-nucleotide phosphodiesterase and calmodulin of cardiac muscle. *Int J Biochem* 1984;16:483-488

Aitchison R, West, DW, Clegg RA. Insulin-stimulated high affinity cyclic AMP phosphodiesterase in rat mammary acini. *FEBS Lett* 1984;167:25-28

Andrenyak DM, Epstein PM. Regulation of a high affinity cyclic nucleotide phosphodiesterase from chick heart by proteases, phospholipids and calmodulin. *Fed Proc* 1982;41:1728-1728

Appleman MM, Thompson WJ, Russel TR. Cyclic nucleotide phosphodiesterases. *Adv Cyclic Nucleotide* 1973;3:65-98

Appleman MM, Terasaki WL. Regulation of cyclic nucleotide phosphodiesterase. *Adv Cyclic Nucleotide Res* 1975;5:153-162

Appleman MM, Allan EH, Ariano MA, Ong KK, Tunang CA, Weber HW, Whitson RH. Insulin control of cyclic AMP phosphodiesterase. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:149-158

Arch JRS, Newsholme EA. Activities and some properties of adenylate cyclase and phosphodiesterase in muscle, liver and nervous tissues from vertebrates and invertebrates in relation to the control of the concentration of adenosine 3':5'-cyclic monophosphate. *Biochem J* 1976;153:603-622

Ashman DF, Lipton R, Melican MM, Price TD. Isolation of adenosine 3',5'-monophosphate and guanosine 3',5'-monophosphate from rat urine. *Biochem Biophys Res Commun* 1963;11:330-334

d'Auriac AG, Meyer P. 3'5'-cyclic phosphodiesterases in cytosol of uterine smooth muscle; effects of angiotensin. *Life* 1973;12:233-239

Ayad SR, Tobia AS. Multiple interconvertible forms of cAMP phosphodiesterases in normal, malignant and hybrid mammalian cells. *Exp Cell Res* 1980;128:257-265

Azhar S, Menon KMJ. Cyclic nucleotide phosphodiesterases from rat anterior pituitary. *Eur J Biochem* 1977;73:73-82

Azhar S, Menon KMJ. Regulatory role of cyclic nucleotide phosphodiesterases in intracellular adenonsine 3':5'-cylic monophosphate turnover and gonadotropin-stimulated progesterone production by rat ovarian cells. *Biochem J* 1979;180:201-211

Baehr W, Devlin MJ, Applebury ML. Isolation and characterization of cGMP phosphodiesterase from bovine rod outer segments. *J Biol Chem* 1979;254:11669-11677

Ball EH, Narindrasorasak S, Sanwal BD. The cyclic adenosine monophosphate phosphodiesterase of myoblasts, fibroblasts and their somatic cell hybrids. *Can J Biochem* 1979;57:1220-1228

Bar HP. Cyclic nucleotides and smooth muscle. *Adv Cyclic Nucleotide Res* 1974;4:195-237

Barber R, Butcher RW. The quantitative relationship between intracellular concentration and egress of cyclic AMP from cultured cells. *Mol Pharmacol* 1981;19:38-43

Barbehenn EK, Wiggert B, Lee L, Kapoor CL, Zonnenberg BA, Redmond TM, Passoneau JV, Chader GJ. Extracellular cGMP phosphodiesterase related to the rod outer segment isolated from bovine and monkey retinas. *Biochemistry* 1985;24:1309-1316

Beatty CH, Bocek RM, Young MK, Novy MJ. Effects of indomethacin on cyclic AMP phosphodiesterase activity of myometrium from pregnant rhesus monkeys. *Prostaglandins* 1976;11:713-725

Beatty CH, Herrington PT, Bocek RM. Regulation of cyclic nucleotide phosphodiesterase activity in rhesus fetal muscle. *Biol Neonate* 1977;31:33-42

Beatty CH, Bocek RM, Herrington PT. Regulation of cyclic nucleotide phosphodiesterase activity in myometrium from pregnant and spayed rhesus monkeys. *J Reprod Fertil* 1979;55:391-400

Beavo JA, Hardman JG, Sutherland EW. Hydrolysis of cyclic guanosine and adenosine 3',5'-monophosphate by rat and bovine tissues. *J Biol Chem* 1970;245:5649-5655

Beavo JA, Hardman JG, Sutherland EW. Stimulation of adenosine 3',5'-monophosphate hydrolysis by guanosine 3',5'-monophosphate. *J Biol Chem* 1971;246:3841-3846

Beavo JA, Hansen RS, Harrison SA, Hurwitz RL, Martins TJ, Mumby MC. Identification and properties of cyclic nucleotide phosphodiesterases. *Mol Cell Endocrinol* 1982;28:387-410

Benedeczky I, Rozsa KS. Cytochemical demonstration of cyclic 3',5' AMP phosphodiesterase in different tissues of migratory locust. *Histochemistry* 1981;70:189-197

Benelli C, Desbuquois B, De Galle B. Acute in vivo stimulation of low-Km cyclic AMP phosphodiesterase activity by insulin in rat-liver Golgi fractions. *Eur J Biochem* 1986;156:211-220

Bergstrand H, Lundquist B. Partial purification and characterization of adenosine- and guanosine-3',5'-monophosphate phosphodiesterase from human lung tissue. *Biochemistry* 1976;15:1727-1735

Bergstrand H, Lundquist B. Partial purification and characterization of cyclic nucleotide phosphodiesterases from human bronchial tissue. *Mol Cell Biochem* 1978;21:9-15

Berridge MJ. The interaction of cyclic nucleotides and calcium in the control of cellular activity. *Adv Cyclic Nucleotide Res* 1975;6:1-98

Berridge MJ, Irvine RF. Inositol triphosphate, a novel second messenger in cellular signal transduction. *Nature(London)* 1984(a);312:315-321

Berridge MJ. Cellular control through interactions between cyclic nucleotides and calcium. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984(b);17:329-335

Best L, Dunlop M, Mlaisse WJ. Phospholipid metabolism in pancreatic islets. *Experentia* 1984;40:1085-1091

Bhatnagar SK, Anand SR. Association of cyclic nucleotide phosphodiesterase of buffalo spermatozoa with sperm chromatin. *Biochim Biophys Acta* 1981;674:212-224

Bhatnagar SK, Anand SR. Cyclic nucleotide phosphodiesterase activity in midpiece and tail of buffalo spermatozoa and its role in sperm motility. *Biochim Biophys Acta* 1982;716:133-139

Billingsley M, Khun D, Velletri PA, Kincaid R, Lovenberg W. Carboxymethylation of phosphodiesterase attenuates its activitation by Ca^{2+}/calmodulin. *J Biol Chem* 1984;259:6630-6635

Birnbaum RJ, Head JF. Studies on the soluble phosphodiesterases of chicken gizzard smooth muscle. *Biochem J* 1983;215:627-636

Boldicke TW, Schubert-Rheberg K, Wombacher H. Purification of an enzyme aggregate containing 3',5'-cyclic nucleotide phosphodiesterase and nucleotidase. *FEBS Lett* 1986;204:125-128

Boudreau RJ, Drummond GI. A modified assay of 3':5'-cyclic-AMP phosphodiesterase. *Anal Biochem* 1975;63:388-399

Bourne HR, Tomkins GM, Dion S. Regulation of phosphodiesterase synthesis: requirement for cyclic adenosine monophosphate-dependent protein kinase. *Science* 1973;181:952-954

Bourne HR, Brothers VM, Kaslow HR, Groppi V, Walker N, Steinberg F. Genetic analysis of cyclic nucleotide phosphodiesterases in S49 mouse lymphoma cells. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:185-194

Boynton AL, Whitfield JF. The role of cyclic AMP in cell proliferation: a critical assessment of the evidence. *Adv Cyclic Nucleotide Res* 1983;15:193-294

Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248-254

Braumann T, Erneux C, Petridis G, Stohrer W-D, Jastorff B. Hydrolysis of cyclic nucleotides by a purified cGMP-stimulated phosphodiesterase: structural requirements for hydrolysis. *Biochim Biophys Acta* 1986;871:199-206

Broadus AE. Clinical cyclic nucleotide research. *Adv Cyclic Nucleotide Res* 1977;8:509-546

Brooker G, Thomas LJ, Appleman MM. The assay of adenosine 3',5'-cyclic monophosphate and guanosine 3',5'-cyclic monophosphate in biological materials by enzymatic radioisotopic displacement. *Biochemistry* 1968;12:4177-4181

Brothers VM, Walker N, Bourne HR. Increased cyclic nucleotide phosphodiesterase activity in mutant S49 lymphoma cell. *J Biol Chem* 1982;257:9349-9355

Bryan JK. Molecular weight of protein multimers from ployacrylamide gel electrophoresis. *Anal Biochem* 1977;78:513-519

Butcher RW, Sutherland EW. Adenosine 3',5'-phosphate in biological materials. I. Purification and properties of cyclic 3',5'-nucleotide phosphodiesterase and use of this enzyme to characterize adenosine 3',5'-phosphate in human urine. *J Biol Chem* 1962;237:1244-1250

Butcher RW. Cyclic 3':5'-nucleotide phosphodiesterase from bovine heart. *Methods Enzymol* 1974;38:218-223

Butcher RW. Phosphodiesterase after twenty years:an introduction. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:1-12

Butcher RW, Barber R, Goka TJ. cAMP metabolism in an S49 mouse lymphoma variant with diminished phosphodiesterase activity. *J Cyclic Nucleotide Protein Phosphor Res* 1986;11:275-290

Campbell MT, Oliver IT. 3':5'-cyclic nucleotide phosphodiesterase in rat tissues. *Eur J Biochem* 1972;28:30-37

Capito K, Hedeskov CJ, Thams P. Cyclic AMP phosphodiesterase activity in mouse pancreatic islets: effects of calmodulin and phospholipids. *Acta Endocrinol* 1986;111:533-538

Cercek B, Houslay MD. Submitochondrial localization and asymmetric disposition of two peripheral cyclic nucleotide phosphodiesterases. *Biochem J* 1982;207:123-132

Charbonneau H, Beier N, Walsh K, Beavo J. Identification of a conserved domain among cyclic nucleotide phosphodiesterases from diverse species. *Proc Natl Acad Sci(USA)* 1986;83:9308-9312

Chasin M, Harris DN. Inhibitors and activators of cyclic nucleotide phosphodiesterase. *Adv Cyclic Nucleotide Res* 1976;7:225-264

Chassy BM. Cyclic nucleotide phosphodiesterase in Dictyostelium discoideum: interconversion of two enzyme forms. *Science* 1972;175:1016-1018

Chatterjee SK, Kim U. Adenosine-3',5'-cyclic monophosphate levels and adenosine-3',5'-cyclic monophosphate phosphodiesterase activity in metastasizing and nonmetastasizing rat mammary carcinomas. *J Natl Cancer Inst* 1975;54:181-186

Chen C-N, Denome S, Davis RL. Molecular analysis of cDNA clones and the corresponding genomic coding sequences of the Drosophila dunce gene, the structural gene for cAMP phosphodiesterase. *Proc Natl Acad Sci(USA)* 1986;83:9313-9317

Cheung WY. Cyclic 3',5'nucleotide phosphodiesterase: demonstration of an activator. *Biochem Biophys Res Commun* 1970;38:533-538

Cheung WY. Cyclic 3',5'nucleotide phosphodiesterase: evidence for and properties of a protein activator. *J Biol Chem* 1971;246:2859-2869

Cheung WY, Lynch TJ, Wallis RW. An endogenous calcium dependent activator of brain adenylate cyclase and cyclic nucleotide phosphodiesterase. *Adv Cyclic Nucleotide Res* 1978;9:233-251

Cho-Chung YS, Newcomer SF. Adenylate cyclase, cyclic adenosine 3':5'-monophosphate phosphodiesterase, and regression of Walker 256 mammary carcinoma. *Cancer Res* 1977;37:4493-4499

Cho-Chung YS, Bodwin JS, Clair T. Cyclic AMP-binding proteins: inverse relationship with estrogen-receptors in hormone-dependent mammary tumor regression. *Eur J Biochem* 1978;86:51-60

Cho-Chung YS. On the mechanism of cyclic AMP-mediated growth arrest of solid tumors. *Adv Cyclic Nucleotide Res* 1980;12:111-121

Cho-Chung YS. Potential application of dibutyryl-cAMP + L-arginine in breast cancer therapy. In : *Endocrine therapy of breast cancer*, edited by U Eppenberger, D Fabbro and P Schafer 1986;23:44-45. Karger, Basel

Chuang NN, Newby AC, Luzio JP. Characterization of different molecular forms of 5'-nucleotidase in normal serum and serum from cholestatic patients and bile-duct-ligated rats. *Biochem J* 1984;224:689-695

Clark JF, Morris HP, Weber G. Cyclic adenosine 3',5'-monophosphate phosphodiesterase activity in normal, differentiating, regenerating, and neoplastic liver. *Cancer Res* 1973;33:356-361

Clark RB. Desensitization of hormonal stimuli coupled to regulation of cyclic AMP levels. *Adv Cyclic Nucleotide Protein Phosphor Res* 1986;20:151-210

Clayberger CA, Goodman DBP, Rasmussen H. Regulation of cyclic AMP metabolism in the rat erythrocyte during chronic B-adrenergic stimulation. Evidence for calmodulin-mediated alteration of membrane- bound phosphodiesterase activity. *J Membr Biol* 1981;58:191-201

Coffey RG, Hadden EM, Lopez C, Hadden JW. cGMP and calcium in the initiation of cellular proliferation. *Adv Cyclic Nucleotide Res* 1978;9:661-676

Cohen LA, Chan P-C. Intracellular cAMP levels in normal rat mammary glands and adenocarcinoma. *Life Sci* 1975;16:107-115

Cohen LA, Straka D, Chan P-C. Cyclic nucleotide phosphodiesterase activity in normal and neoplastic rat mammary cells grown in monolayer culture. *Cancer Res* 1976;36:2007-2012

Collier HOJ, Butt NM, Saeed SA. Endogenous peptides that inhibit brain cyclic AMP phosphodiesterase. *J Neurochem* 1982;38:275-277

Collings JR, Levin J, Savige N. Racial differences in oestrogen receptor and peroxidase status of human breast cancer tissue. *S Afr Med J* 1980;57:444-446

Conti M, Toscano MV, Geremia R, Stefanini M. Follicle-stimulating hormone regulates in vivo testicular phosphodiesterase. *Mol Cell Endocrinol* 1983;29:79-89

Conti M, Kasson BG, Hsueh AJW. Hormonal regulation of 3',5'-adenosine monophosphate phosphodiesterase in cultured rat granulosa cells. *Endocrinol* 1984;114:2361-2368

Coquil J-F, Franks DJ, Wells JN, Dupuis M, Hamet P. Characteristics of a new binding protein distinct from the kinase for guanosine 3',5'-monophosphate in rat platelets. *Biochim Biophys Acta* 1980;631:148-165

Couchie D, Erneux C, Dumont JE. Characterization of a rat liver cyclic GMP-activated phosphodiesterase by chromatography on hexyl-agarose. *Biochem J* 1981;199:441-446

Couchie D, Petridis G, Jastorff B, Erneux C. Characterization of phosphodiesterase catalytic sites by means of cyclic nucleotide derivatives. *Eur J Biochem* 1983;136:571-575

Coulson AS, Kennedy LA. Lymphocyte membrane enzymes. II. Cyclic 3',5'-adenosine monophosphatase located on unstimulated human small lymphocyte nuclear membranes. *Blood* 1971;38:485-490

Cox JA,Malnoe A, Stein EA. Regulation of brain cyclic nucleotide phosphodiesterase by calmodulin. *J Biol Chem* 1981;256:3218-3222

Davis BJ. Disc electrophoresis - II Method and application to human serum proteins. *Ann NY Acad Sci* 1964;121:404-427

Davis CW, Kuo JF. Purification and characterization of guanosine 3',5' monophosphate specific phosphodiesterase from the guinea pig lung. *J Biol Chem* 1977;252:4078-4084

Davis RL, Kauvar LM. Drosophila cyclic nucleotide phosphodiesterases. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:393-402

Dedman JR, Potter JD, Means AR. Biological cross-reactivity of rat testis phosphodiesterase activator protein and rabbit skeletal muscle troponin-C. *J Biol Chem* 1977;252:2437-2440

Dedman JR, Jackson RL, Schreiber WE, Means AR. Sequence homology of the Ca2+-dependent regulator of cyclic nucleotide phosphodiesterase from rat testis with other Ca2+-binding proteins. *J Biol Chem* 1978;253:343-346

De Jong D. Gluteraldehyde activation of nuclear acid phosphatase in cultured plant cells. *Science* 1967;155:1672-1674

DeRubertis FR, Chayoth R, Field JB. The content and metabolism of cyclic adenosine 3',5'-monophosphate and cyclic guanosine 3',5'-monophosphate in adenocarcinoma of the human colon. *J Clin Invest* 1976;57:641-649

DeRubertis FR, Craven PA. Cyclic nucleotides in carcinogenesis: activation guanylate cyclase-cyclic GMP system by chemical carcinogens. *Adv Cyclic Nucleotide Res* 1980;12:99-109

Donnelly TE Jr. Reconsideration of the multiple cyclic nucleotide phosphodiesterases in bovine heart. *Biochem Biophys Res Commun* 1978;82:964-970

Donnelly TE Jr., Barron B. Effects of divalent metals on the specificity of inhibitors of the cyclic nucleotide phosphodiesterases from bovine heart. *Biochim Biophys Acta* 1981;657:168-178

Doskeland SO, Ogreid D. Binding proteins for cyclic AMP in mammalian tissues. *Int J Biochem* 1981;13:1-19

Duma A, Moraczewski J. Ultracytochemistry of cyclic 3',5'-nucleotide phosphodiesterase activity in the planarian Dugesia lugubris (O.Schmidt). *Histochemistry* 1980;66:211-220

Egrie JC, Siegel FL. Adrenal medullary cyclic nucleotide phosphodiesterase: subcellular distribution, partial purification and regulation of enzyme activity. *Biochim Biophys Acta* 1977;483:348-366

Elks ML, Manganiello VC, Vaughan M. Hormone-sensitive particulate cAMP phosphodiesterase activity in 3T3-L1 adipocytes. *J Biol Chem* 1985;258:8582-8587

Eppelen JT, Engel W, Schmidtke J. Patterns of cyclic AMP phosphodiesterases during rat testis development. *Hum Genet* 1980;53:393-399

Epstein PM, Mills JS, Strada SJ, Hersh EM, Thompson WJ. Increased cyclic nucleotide phosphodiesterase activity associated with proliferation and cancer in human and murine lymphoid cells. *Cancer Res* 1977;37:4015-4023

Epstein PM, Pledger WJ, Gardner EA, Stancel GM, Thompson WJ, Strada SJ. Activation of mammalian cyclic AMP phosphodiesterases by trypsin. *Biochim Biophys Acta* 1978;527:442-455

Epstein PM, Fiss K, Hachisu R, Andrenyak DM. Interaction of calcium antagonists with cyclic AMP phosphodiesterases and calmodulin. *Biochem Biophys Res Commun* 1982(a);105:1142-1149

Epstein PM, Strada SJ, Sarada K, Thompson WJ. Catalytic and kinetic properties of purified high-affinity cyclic AMP phosphodiesterase from dog kidney. *Arch Biochem Biophys* 1982(b);218:119-133

Epstein PM, Hachisu R. Cyclic nucleotide phosphodiesterase in normal and leukemic human lymphocytes and lymphoblasts. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:303-324

Epstein PM, Andrenyak DM, Smith CJ, Pappano AJ. Ontogenic changes in adenylate cyclase, cyclic AMP phosphodiesterase and calmodulin in chick ventricular myocardium. *Biochem J* 1987;243:525-531

Erneux C, Couchie D, Dumont JE. Characterization of horse thyroid cyclic nucleotide phosphodiesterases. *Eur J Biochem* 1980;104:297-304

Erneux C, Couchie D, Dumont JE, Baraniak J, Stec WJ, Garcia Abbad E,Petridis G, Jastorff B. Specificity of cyclic GMP activation of a multi-substrate cyclic nucleotide phosphodiesterase from rat liver. *Eur J Biochem* 1981;115:503-510

Erneux C, Miot F, Boeynaems J-M, Dumont JE. Paradoxical stimulation by 1-methyl-3-isobutylxanthine of rat liver cyclic AMP phosphodiesterase activity. *FEBS Lett* 1982;142:251-254

Erneux C, Miot F, Van Haastert PJM, Jastorff B. The binding of cyclic nucleotide analogs to a purified cyclic GMP-stimulated phosphodiesterase from bovine adrenal tissue. *J Cyclic Nuclectide Protein Phosphor Res* 1985;10:463-472

Etingof RN, Dumler IL, Garnovskaja MN, Kalinia SN. Estradiol receptor and cyclic nuclectide phosphodiesterase: funtional relationship, possible role of guanine nucleotide binding proteins. *Biochem Int* 1984;9:229-236

Exton JH. Mechanisms involved in calcium-mobilizing agonist responses. *Adv Cyclic Nucleotide Protein Phosphor Res* 1986;20:211-262

Farnham CJM. Cytochemical localization of adenylate cyclase and 3',5'-nucleotide phosphodiesterase in Dictyostelium. *Exp Cell Res* 1975;91:36-46

Fell DA. Theoretical analyses of the functioning of the high- and low-Km cyclic nucleotide phosphodiesterases in the regulation of the concentrations of adenosine 3',5'-cyclic monophosphate in animal cells. *J Theor Biol* 1980;84:361-385

Ferguson KA. Starch-gel electrophoresis application to the classification of pituitary proteins and polypeptides. *Metabolism* 1964;13:985-1001

Ferre F, Breuiller M, Cedard L. Human placental cAMP phosphodiesterase activity, kinetic properties and sensitivity to some drugs and hormones. *FEBS Lett* 1975;52:295-299

Ferre F, DeParIente D, Breuiller M, Cedard L. Inhibition of human myometrial cyclic AMP phosphodiesterase by uterine relaxant drugs. *Biochem Pharmacol* 1977;27:1294-1296

Florendo NT, Barnett RJ, Greengard P. Cyclic 3',5'-nucleotide phosphodiesterase: cytochemical localization in cerebral cortex. *Science* 1971;173:745-747

Fougier S, Nemoz G, Prigent AF, Marivet MD, Bourguignon JJ, Wermuth C, Pacheco H. Purification cAMP-specific phosphodiesterase from rat heart by affinity chromatography on immobilized rolipram. *Biochem Biophys Res Commun* 1986;138:205-214

Francis SH, Lincoln TM, Corbin JD. Characterization of a novel cGMP binding protein from rat lung. *J Biol Chem* 1980;255:620-626

Francis SH. Effectors of rat lung cGMP binding protein-phosphodiesterase. *Curr Top Cell Regul* 1985;26:247-262

Franke J, Kessin RH. The cyclic nucleotide phosphodiesterase inhibitory protein of Dictyostelium discoideum. *J Biol Chem* 1981;256:7628-7637

Friedman DL. Role of cyclic nucleotides in cell growth and differentiation. *Physiol Rev* 1976;56:652-708

Fukunaga K, Yamamoto H, Tanaka H, Iwasa T, Miyamoto E. Phosphorylation and activation of calmodulin-sensitive cyclic nucleotide phosphodiesterase by a brain Ca2+,calmodulin-dependent protein kinase. *Life Sci* 1984;35:493-499

Galoyan AA, Gurvitz BY. Regulation of multiple forms of cyclic nucleotide phosphodiesterase from bovine hypothalamus: new factors modulating enzyme activity. *Neurochem Res* 1985;10:1467-1481

Gardner EA, Thompson WJ, Strada SJ, Stancel GM. Characterization of soluble uterine cyclic nucleotide phosphodiesterase. *Biochemistry* 1978;17:2995-3000

Geremia R, Rossi P, Pezotti R, Conti M. Cyclic nucleotide phosphodiesterase in developing rat testis: identification of somatic and germ cell forms. *Mol Cell Endocrinol* 1982;28:37-53

Geremia R, Rossi P, Mocini D, Pezzotti R, Conti M. Characterization of a calmodulin-dependent high-affinity cyclic AMP and cyclic GMP phosphodiesterase from male mouse germ cells. *Biochem J* 1984;217:693-700

Gill DL. Receptors coupled to calcium mobilization. *Adv Cyclic Nucleotide Protein Phosphor Res* 1985;19:307-324

Goff AK, Armstrong DT. Stimulatory action of gonadotropins and prostaglandins on adenosine-3',5'-monophosphate production by rat isolated granulosa cells. *Endocrinol* 1977;101:1461-1467

Goldberg ND, Dietz SB, O'Toole AG. Cyclic guanosine 3',5'-monophosphate in mammalian tissues and urine. *J Biol Chem* 1969;16:4458-4466

Goldberg ND, O'Dea RF, Haddox MK. Cyclic GMP. *Adv Cyclic Nucleotide Res* 1973;3:155-223

Goldberg ND, Haddox MK, Nicol SE, Glass DB, Sanford CH, Kuehl FA, Estensen R. Biologic regulation through opposing influences of cyclic GMP and cyclic AMP: the Yin Yang hypothesis. *Adv Cyclic Nucleotide Res* 1975;5:307-330

Goldberg ND, Walseth TF, Eide SJ, Krick TP, Kuehn BL, Gander JE. Cyclic AMP metabolism in intact platelets determined by 18 O incorporation into adenine nucleotide alpha-phosphoryls. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:363-379

Goren EN, Hirsch AH, Rosen OM. Activity stain for the detection of cyclic nucleotide phosphodiesterase separated by polyacrylamide gel electrophoresis and its application to the cyclic nucleotide phosphodiesterase of beef heart. *Anal Biochem* 1971;43:156-161

Goren EN, Rosen OM. Purification and properties of a cyclic nucleotide phosphodiesterase from bovine heart. *Arch Biochem Biophys* 1972;153:384-397

Goridis C, Virmaux N, Cailla HL, Delaage MA. Rapid light-induced changes of retinal cyclic GMP levels. *FEBS Lett* 1974;49:167-169

Grant PG, Colman RW. Purification and characterization of a human platelet cyclic nucleotide phosphodiesterase. *Biochemistry* 1984;23:1801-1807

Guan WR, Cheung WY. Stimulation of rat liver cyclic 3',5'-cyclic nucleotide phosphodiesterase by cyclic GMP is dependent on enzyme concentration. *Arch Biochem Biophys*
Guan WR, Cheung WY. Stimulation of rat liver cyclic 3',5'-cyclic nucleotide phosphodiesterase by cyclic GMP is dependent on enzyme concentration. *Arch Biochem Biophys* 1980;201:191-198

Guerinot F, De Larue JC, Contesso G, Bohuon C. Adenosine 3',5'-cyclic monophosphate and guanosine 3',5'-cyclic monophosphate levels in human breast cancer tissue. *Oncol* 1977;34:261-263

Hagiwara M, Endo T, Hidaka H. Effects of vinpocetine on cyclic nucleotide metabolism in vascular smooth muscle. *Biochem Pharmacol* 1984;33:453-457

Hait WN, Weiss B. Cyclic nucleotide phosphodiesterase of normal and leukemic lymphocytes. *Mol Pharmacol* 1979;16:851-864

Hamet P, Coquil JF. Cyclic GMP binding and cyclic GMP phosphodiesterase in rat platelets. *J Cyclic Nucleotide Res* 1978;4:281-290

Hamet P, Coquil J-F, Bousseau-Lafortune S, Franks DJ, Tremblay J. Cyclic GMP binding and phosphodiesterase: implication for platelet function. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:119-136

Hansen RS, Beavo JA. Purification of two calcium/calmodulin-dependent forms of cyclic nucleotide phosphodiesterase by using conformation-specific monoclonal antibody chromatography. *Proc Natl Acad Sci(USA)* 1982;79:2788-2792

Hansen RS, Beavo JA. Differential recognition of calmodulin-enzyme complexes by a conformation-specific anti-calmodulin antibody. *J Biol Chem* 1986;261:14363-14645

Harrison SA, Reifsnyder DH, Cadd GC, Beavo JA. Isolation and characterization of a bovine cardiac muscle cyclic GMP inhibited phosphodiesterase: a receptor for new cardiotonic drugs. *Mol Pharmacol* 1986;29:506-514

Hedrick JL, Smith AJ. Size and charge isomer separation and estimation of molecular weights of proteins by disc gel electrophoresis. *Arch Biochem Biophys* 1968;126:155-164

Heidrick ML, Ryan WL. Cyclic nucleotides on cell growth in vitro. *Cancer Res* 1970;30:376-378

Heidrick ML, Ryan WL. Adenosine 3',5'-cyclic monophosphate and contact inhibition. *Cancer Res* 1971;31:1313-1315

Helfman DM, Shoji M, Kuo JF. Purification to homogeneity and general properties of a novel phosphodiesterase hydrolyzing cyclic CMP and cyclic AMP. *J Biol Chem* 1981;256:6327-6334

Heyworth CM, Rawal S, Houslay MD. Guanine nucleotides can activate the insulin-stimulated phosphodiesterases in liver plasma membranes. *FEBS Lett* 1983(a);154:87-90

Heyworth CM, Wallace AV, Houslay MD. Insulin and glucagon regulate the activation of two distinct membrane-bound cyclic AMP phosphodiesterases in hepatocytes. *Biochem J* 1983(b);214:99-110

Heyworth CM, Grey AM, Wilson SR, Hanski E, Houslay MD. The action of islet activating protein (pertussis toxin) on insulin's ability to inhibit adenylate cyclase and activate cyclic AMP phosphodiesterases in hepatocytes. *Biochem J* 1986;235:145-149

Hickie RA, Walker CM, Datta A. Increased activity of low-Km cyclic adenosine 3':5'-monophosphate phosphodiesterase in plasma membranes of Morris hepatoma 5123tc (h). *Cancer Res* 1975;35:601-605

Hickie RA, Thompson WJ, Strada SJ, Couture-Murillo J, Morris HP, Robison GA. Comparison of cyclic adenosine 3':5'-monophosphate and cyclic guanosine 3':5'-monophosphate levels, cyclases and phosphodiesterases in Morris hepatomas and liver. *Cancer Res* 1977;37:3599-3606

Hidaka H, Asano T. Human blood platelet 3':5'-cyclic nucleotide phosphodiesterase: isolation of low-Km and high-Km phosphodiesterase. *Biochim Biophys Acta* 1976;429:485-497

Hidaka H, Endo T. Selective inhibitors of three forms of cyclic nucleotide phosphodiesterase: basic and potential clinical applications. *Adv Cyclic Nucleotide Protein Phosphr Res* 1984;16:245-260

Ho HC, Wirch E, Stevens FC, Wang JH. Purification of a Ca2+/activatable cyclic nucleotide phosphodiesterase from bovine heart by specific interaction with its Ca2+/dependent modulator protein. *J Biol Chem* 1977;252:43-50

Holtz A, Brennen RG, Batista D, Terner C. Androgen control of an inhibitor modulator of phosphodiesterase in rat epididymus and prostate. *Endocrinology* 1981;108:1538-1544

Houslay MD, Wallace AV, Marchmont RJ, Martin BR, Heyworth CM. Insulin controls intracellular cyclic AMP concentrations in hepatocytes by activating specific cyclic AMP phosphodiesterases: phosphorylation of the peripheral plasma membrane enzyme. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:159-176

Hovi T, Keski-Oja J, Vahen A. Growth control in chick embryo fibroblasts; no evidence for a specific role for cyclic nucleotides. *Cell* 1974;2:235-240

Huang CY, Chau V, Chock PB, Wang JH, Sharma Rk. Mechanism of activation of cyclic nucleotide phosphodiesterase: requirement of the binding of four Ca2+ to calmodulin for activation. *Proc Natl Acad Sci(USA)* 1981;78:871-874

Huang CY, King MM. Mode of calcium activation of calmodulin-regulated enzymes. *Curr Top Cell Regul* 1985;27:437-446

Hunt NH, Martin TJ. Hormone receptors and cyclic nucleotides: significance for growth and function of tumors. *Mol Aspects Med* 1980;3:59-118

Hurley JB, Stryer L. Purification of the y regulatory subunit of the cyclic GMP phosphodiesterase from retinal rod outer segments. *J Biol Chem* 1982;257:11094-11099

Hurwitz RL, Bunt-Milan AH, Beavo JA. Immunologic characterization of the photoreceptor outer segment cyclic GMP phosphodiesterase. *J Biol Chem* 1984(a);259:8612-8618

Hurwitz RL, Hansen RS, Harrison SA, Martins TJ, Mumby MC, Beavo JA. Immunologic approaches to to the study of cyclic nucleotide phosphodiesterases. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984(b);16:89-106

Hurwitz RL, Bunt-Milan AH, Chann ML, Beavo JA. cGMP phosphodiesterase in rod and core outer segments of the retina. *J Biol Chem* 1985;260:568-573

Ip C, Dao TL. Regulation of cyclic adenosine monophosphate in relation to growth of dimethylbenzanthracene -induced mammary tumours in rats. *Cancer Lett* 1980;8:227-253

Iwasa Y, Yonemitsu K, Miyamoto E. A calcium-dependent cyclic nucleotide phosphodiesterase from Escherichia coli. *FEBS Lett* 1981;124:207-209

Johnson GS, Friedman RM, Pastan I. Restoration of several morphological characteristics of normal fibroblasts in sarcoma cells treated with adenosine 3',5'-cyclic monophosphate and its derivatives. *Proc Natl Acad Sci(USA)* 1971;68:425-429

Kakiuchi S, Yamazaki R. Calcium dependent phosphodiesterase activity and its activating factor (PAF) from brain. *Biochim Biophys Res Commun* 1970;41:1104-1110

Kakuichi S, Yamazaki R, Teshima Y, Yasuda S, Kashiba A, Sobue K, Ohshima M, Nakajima T. Membrane-bound protein modulator and phosphodiesterase. *Adv Cyclic Nucleotide Res* 1978;253-254

Kalderon AE, Ravanshenas SF. Localization of 3',5'-cyclic nucleotide phosphodiesterase activity in isolated thyroid cells and intact thyroid tissue. *Histochemistry* 1974;39:229-242

Kalderon AE. Cyclic 3',5'-nucleotide phosphodiesterase. In: *Electron microscopy of enzymes.* Edited by MA Hayat 1977;4:1-25. Van Nostrand Reinhold Company.

Kaji E, Bornslaeger EA, Schultz RM. Inhibition of mouse oocyte cyclic AMP phosphodiesterase by steroid hormones: a possible mechanism for steroid hormone inhibition of oocyte maturation. *J Exp Zool* 1987;243:489-493

Kelly SJ, Butler LG. Enzymic hydrolysis of phosphate esters. Reaction mechanism of intestinal 5'-nucleotide phosphodiesterase. *Biochemistry* 1977;16:1102-1104

Keravis TM, Wells JN, Hardman JG. Cyclic nucleotide phosphodiesterase activity from pig coronary arteries: lack of interconvertibility of forms. *Biochim Biophys Acta* 1980;613:116-129

Keravis TM, Duemler BH, Wells JN. Calmodulin sensitive phosphodiesterase of porcine cerebral cortex: kinetic behaviour, calmodulin activation and stability. *J Cyclic Nucleotide Protein Phosphor Res* 1987; 11:365-372

Kiger JA, Davis RL, Salz H, Fletcher T, Bowling M. Genetic analysis of cyclic nucleotide phosphodiesterases in Drosophila melanogaster. *Adv Cyclic Nucleotide Res* 1981;14:273-288

Kincaid RL, Vaughan M. Sequential adsorbtion-electrophoresis: combined procedure for purification of calcium-dependent cyclic nucleotide phosphodiesterase. *Proc Natl Acad Sci(USA)* 1979;76:4901-4907

Kincaid RL, Manganiello VC, Vaughan M. Calmodulin-activated cyclic nucleotide phosphodiesterase from brain. *J Biol Chem* 1981;256:11345-11350

Kincaid RL, Vaughan M. Affinity chromatography of brain cyclic nucleotide phosphodiesterase using 3-(2-pyridyldithio)propionyl-substituted calmodulin linked to thio-sepharose. *Biochemistry* 1983;22:826-830

Kincaid RL, Manganiello VC, Odya CE, Osborne Jr JC, Stith-Coleman IE, Danello MA, Vaughan M. Purification and properties of calmodulin-stimulated phosphodiesterase from mammalian brain. *J Biol Chem* 1984;259:5158-5166

King LE, Solomon SS, Hashimoto K. Characterization of a human melanoma cell line (KHm-1) containing phophodiesterase activity only for cyclic adenosine 3':5'-monphosphate. *Cancer Res* 1978;38:3879-3885

Kishimoto A, Takai Y, Mori T, Kikkawa U, Nishizuka Y. Activation of calcium and phospholipid-dependent protein kinase by diacylglycerol, its possible relationship to phosphoinositol turnover. *J Biol Chem* 1980;;255:2273-2276

Klee CB, Krinks MH. Purification of cyclic 3',5'-nucleotide phosphodiesterase inhibitory protein by affinity chromatography on activator protein coupled to sepharose. *Biochemistry* 1978;17:120-126

Klee CB, Crouch TH, Krinks MH. Subunit structure and catalytic properties of bovine brain Ca2+/dependent cyclic nucleotide phosphodiesterase. *Biochemistry* 1979;18:722-729

Klein C, Darmon M. Effects of cyclic AMP pulses on adenylate cyclase and the phosphodiesterase inhibitor of D. discoideum. *Nature (London)* 1977;268:76-78

Klein DM, Loizzi RF. Enhancement of R3230AC rat mammary tumor growth and cellular differentiation by dibutyryl cyclic adenosine monophosphate. *J Natl Cancer Inst* 1977;58:813-818

Knecht M, Ranta T, Catt KJ. Hormonal regulation of a plasma membrane phosphodiesterasein differentiating granulosa cells. *J Biol Chem* 1983;258:12420-12426

Kofinas AD, Rose JC, Meiss PJ. Changes in cyclic adenosine monophosphate- phosphodiesterase activity in nonpregnant and pregnant human myometrium. *Am J Obstet Gynecol* 1987;157:733-737

Kono T, Robinson FW, Sarver JA. Insulin-sensitive phosphodiesterase: its localization, hormonal stimulation and oxidative stabilization. *J Biol Chem* 1975;250:7826-7835

Krinks MH, Haiech J, Rhoads A, Klee CB. Reversible and irreversible activation of cyclic nucleotide phosphodiesterase: separation of the regulatory and catalytic domains by limited proteolysis. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:31-47

Kung W, Bechtel E, Geyer E, Salokangas A, Preisz J, Huber P, Torhorst J, Jungmann RA, Talmadge K, Eppenberger U. Altered levels of cyclic nucleotides, cyclic AMP phosphodiesterase and adenylyl cyclase activities in normal, dysplastic and neoplastic human mammary tissue. *FEBS Lett* 1977;82:102-106

Kuo JF, Greengard P. Cyclic-nucleotide dependent protein kinases. *Proc Natl Acad Sci(USA)* 1969;64:1349-1355

Kuo JF, Shoji M, Brackett NL, Helfman DM. Effects of phosphodiesterase inhibitors, imidazole and phosphate on cyclic GMP phosphodiesterase are different from those on cyclic AMP and cyclic GMP phosphodiesterases. *J Cyclic Nucleotide Res* 1978;4:463-474

Kuribayashi N, Makino H, Fujimoto S, Kanatsuka A, Yoshida S. Insulin-sensitive phosphodiesterase in fat cells from human subcutaneous adipose tissue. *Horm Metabol Res* 1987;19:341-342

Kurtz MJ, Edwards RB, Schmidt SY. Cyclic nucleotide phosphodiesterases in cultured normal and RCS rat pigment epithelium: kinetics of cyclic AMP and cyclic GMP hydrolysis. *Exp Eye Res* 1987;45:67-75

Lacombe M-L, Podgorski GJ, Franke J, Kessin RH. Molecular cloning and developmental expression of the cyclic nucleotide phosphodiesterase gene of Dictyostelium discoideum. *J Biol Chem* 1986;261:16811-16817

Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature(Lond)* 1970;227:680-685

Lam H-YP. Tamoxifen is a calmodulin antagonist in the activation of cAMP phosphodiesterase. *Biochem Biophys Res Comm* 1984;118:27-32

LaPorte DC, Toscana WA, Storm DR. Cross-linking of iodine-125-labelled, calcium-dependent regulatory protein to the Ca2+-sensitive phosphodiesterase purified from bovine heart. *Biochemistry* 1979;18:2820-2825

Larner EH, Rutherford CL. Implementation of micromethods to resolve problems of human breast tumor heterogeneity in analysis of cyclic 3':5'-nuclectide phosphodiesterase. *Cancer Res* 1982;42:1661-1665

Leroy M-J, Pichard A-L, Cabrol D, Ferre F. Cyclic 3':5'-nucleotide phosphodiesterase in human myometrium at the end of pregnancy:partial purification and characterization of the different soluble isoenzymes. *Gynecol Obstet Invest* 1985;20:27-36

Lichtner RB, Goka TJ, Butcher RW, Nicolson GL. Direct effects of the pyrimido-pyrimidine derivative RA 233 (Rapenton) on rat 13762NF mammary tumor cell clones in vitro. *Cancer Res* 1987;47:1870-1877

Lincoln TM, Corbin JD. Characterization and biological role of the cGMP-dependent protein kinase. *Adv Cyclic Nucleotide Res* 1983;15:139-192

Lineweaver H, Burk D. The determination of enzyme dissociation constants. *J Am Chem Soc* 1934;56:658-666

Lohmann SM, Walter U. Regulation of the cellular and subcellular concentrations and distribution of cyclic nucleotide-dependent protein kinases. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;18:63-118

Londesborough J, Jonkakari L. Low Km cyclic AMP Phosphodiesterase of yeast may be bound to ribosomes associated with the nucleus. *Mol Cell Biochem* 1982;46:65-71

Londesborough J. Evidence that the peripheral cyclic AMP phosphodiesterase is a metalloenzyme. *Biochem J* 1985;225:143-147

Loten EG, Sneyd JGT. An effect of insulin on adipose-tissue adenosine 3':5'-cyclic monophosphate phosphodiesterase. *Biochem J* 1970;120:187-193

Loten EG. Stimulation of a low Km phosphodiesterase from liver by insulin and glucagon. *J Biol Chem* 1978;253:746-757

Loten EG. Proteolytic solubilization and modification hormone-sensitive cyclic nucleotide phosphodiesterase. *J Biol Chem* 1980;255:7838-7844

Loten EG. Detergent solubilisation of rat liver particulate cyclic AMP phosphodiesterase. *Int J Biochem* 1983;15:923-928

Lynch TJ, Cheung WY. Underestimation of cyclic 3',5'-nucleotide phosphodiesterase activity by a radioisotopic assay using an anionic-exchange resin. *Anal Biochem* 1975;67:130-138

Macauley SL, Kiechle FL, Jarett L. Phospholipid modulation of low-km cyclic AMP phosphodiesterase activity on rat adipocyte microsomes. *Arch Biophys* 1983;225:130-136

Macphee CH, Harrison SA, Beavo JA. Immunological identification of the major platelet low-Km cAMP phosphodiesterase: probable target for anti-thrombitic agents. *Proc Natl Acad(USA)* 1986;83:6660-6663

Major GN, Loten EG, Sneyd JGT. Partial purification of protease-solubilized low Km cyclic nucleotide phosphodiesterase from liver. *Int J Biochem* 1983;15:217-223

Makino H, Buschiazzo PM, Pointer RH, Jordan JE, Kono T. Characterization of insulin-sensitive phosphodiesterase in fat cells. *J Biol Chem* 1980;255:7845-7849

Makino H, Osegawa M, Kanatsuka A. Hypotonic activation of insulin-sensitive phosphodiesterase in fat cells. *Mol Cell Endocrinol* 1982(a);28:313-320

Makino H, Kanatsuka A, Osegawa M, Kumagi A. Effects of dithiothreitol on insulin-sensitive phosphodiesterase in rat fat cells. *Biochim Biophys Acta* 1982(b);704:31-36

Manalan AS, Klee CB. Calmodulin. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;18:227-278

Manganiello V, Vaughan M. Prostaglandin E1 effects on adenosine 3':5'-cyclic monophosphate concentration and phosphodiesterase activity in fibroblasts. *Proc Natl Acad Sci(USA)* 1972;69:269-273

Manganiello V, Vaughan M. An effect of insulin on cyclic adenosine 3':5'-monophosphate phosphodiesterase activity in fat cells. *J Biol Chem* 1973;248:7164-7170

Manganiello VC, Yamomoto T, Lin MC, Vaughan M. Regulation of specific forms of cyclic nucleotide phosphodiesterases in cultured cells. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:291-301

Marchmont RJ, Houslay MD. Insulin triggers cyclic AMP-dependent activation and phosphorylation of a plasma membrane cyclic AMP phosphodiesterase. *Nature(London)* 1980(a);286:904-906

Marchmont RJ, Houslay MD. A peripheral and an intrinsic enzyme constitute the cyclic AMP phosphodiesterase activity of rat liver plasma membranes. *Biochem J* 1980(b);187:381-392

Marchmont RJ, Ayad SR, Houslay MD. Purification and properties of the insulin-stimulated cyclic AMP phosphodiesterase from rat liver plasma membranes. *Biochem J* 1981(a);195:645-652

Marchmont RJ, Houslay MD. Characterization of the phosphorylated form of the insulin-stimulated cyclic AMP phosphodiesterase from rat liver plasma membranes. *Biochem J* 1981(b);195:653-660

Marcus R, Grant BF. Parathyroid hormone induces a calmodulin-dependent alteration in phosphodiesterase activity of rat kidney. *Endocrinol* 1983;112:1065-1069

Martin RG, Ames BN. A method for determining the sedimentation behaviour of enzymes:application to protein mixtures. *J Biol Chem* 1961;236:1372-1379

Martins TJ, Mumby MC, Beavo JA. Purification and characterization of a cyclic GMP-stimulated cyclic nucleotide phosphodiesterase from bovine tissues. *J Biol Chem* 1982;257:1974-1979

Mason M, Narindrasorasak S, Sanwal BD. Regulation of cyclic 3',5'-adenosine monophosphate phosphodiesterase in Friend erythroleukemia cells. *J Cyclic Nucleotide Res* 1985;10:129-142

Matusik RJ, Hilf R. Relationship of adenosine 3',5'-cyclic monophosphate and guanosine 3',5'-cyclic monophosphate to growth of dimethylbenzanthracene -induced mammary tumors in rats. *J Natl Cancer Inst* 1976;56:659-661

Mayer RM, Diamond EJ, Hollander VP. Hormone, serum and methylxanthine interactions in the regulation of DNA synthesis organ-cultured rat mammary tumors. *J Natl Cancer Inst* 1980;65:1033-1038

McEwen CR. Tables for estimating sedimentation through linear concentration gradients of sucrose solution. *Anal Biochem* 1967;20:114-149

Merril CR, Harrington MG. Ultrasensitive silver stains: their use exemplified in the study of normal human cerebrospinal fluid proteins separated by two-dimensional electrophoresis. *Clin Chem* 1984;30:1938-1942

Miki N, Barban JM, Kierns JJ, Boyce JJ, Bitensky MW. Purification and properties of the light-activated cyclic nucleotide phosphodiesterase of rod outer segments. *J Biol Chem* 1975;250:6320-6327

Minton JP, Wisenbaugh T, Matthews RH. Elevated cyclic AMP levels in human breast-cancer tissue. *J Natl Cancer Inst* 1974;53:283-284

Minton JP, Mathews RH, Wisenbaugh TW. Elevated adenosine 3',5'-cyclic monophosphate levels in human and animal tumors in vivo. *J Natl Cancer Inst* 1976;57:39-41

Miot F, Dumont JE, Erneux C. The involvement of a calmodulin-dependent phosphodiesterase on the negative control of carbamycholine on cyclic AMP levels in dog thyroid slices. *FEBS Lett* 1983;151:310-319

Miot F, Van Haastert PJM, Erneux C. Specificity of cGMP binding to a purified cGMP-stimulated phosphodiesterase from bovine adrenal tissue. *Eur J Biochem* 1985;149:59-65

Miyamoto M, Takayanagi I, Ohkubo H, Takagi K. Actions of papaverine on intestinal smooth muscle and its inhibition of cyclic AMP and cyclic GMP phosphodiesterases. *Japan J Pharmacol* 1976;26;114-120

Monahan TM, Marchand NW, Fritz RR, Abell CW. Cyclic adenosine 3':5'-monophosphate levels and activities of related enzymes in normal and leukemic lymphocytes. *Cancer Res* 1975;35:2540-2547

Monn E, Christiansen RO. Adenosine 3',5'-monophosphate phosphodiesterase: multiple molecular forms. *Science* 1971;173:540-542

Moore Jr JB, Schroedter DE. Purification and characterization of human lung calmodulin-independent cyclic AMP phosphodiesterase. *Arch Biochem Biophys* 1982;213:267-287

Morrill ME, Thompson ST, Stellwagen E. Purification of a cyclic nucleotide phosphodiesterase from bovine brain using blue dextran-sepharose chromatography. *J Biol Chem* 1979;254:4371-4374

Moses HJ, Rosenthal AS. Pitfalls in the use of lead ion for histochemical localization of nucleotide phosphateses. *Histochem Cytochem* 1968;16:530-539

Moses MA, Szabo EI, Terner C. Control of low Km cyclic adenosine 3',5'-monophosphate phosphodiesterase in prostate and heart of adult and aged rats. *Mec Ageing Dev* 1987;39:103-112

Moss J, Manganiello VC, Vaughan M. Substrate and effector specificity of a guanosine 3':5'-monophosphate phosphodiesterase from rat liver. *J Biol Chem* 1977;252:5211-5215

Moss J, Manganiello VC, Vaughan M. Effects of chymotrypsin and a calcium-dependent protein activator on guanosine 3',5'-monophosphate phosphodiesterase from rat liver. *Biochim Biophys Acta* 1978;541:279-287

Mullaney I, Clegg RA. Cyclic AMP phosphodiesterase and cyclic GMP phosphodiesterase activities of rat mammary tissue. *Biochem J* 1984;219:801-809

Mumby MC, Martins TJ, Chang ML, Beavo JA. Identification of cGMP-stimulated cyclic nucleotide phosphodiesterase in lung tissue with monoclonal antibodies. *J Biol Chem* 1982;257:13283-13293

Munjaal RP, Dedman JR, Means AR. Isolation of the stuctural gene for calmodulin. *Ann NY Acad Sci* 1980;356:110-117

Murad F, Arnold WP, Mittal CK, Braughter JM. Properties and regulation of guanylate cyclase and some proposed functions for cyclic GMP. *Adv Cyclic Nucleotide Res* 1979;11:175-204

Mutus B, Wang K, Sharma RK, Lam H-Y, Wang JH. A fluorescent probe and monoclonal antibody for studies of the interaction between calmodulin and cyclic nucleotide phosphodiesterase. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;16:65-76

Mutus B, Karppiah N, Sharma RK, MacManus JP. The differential activation of brain and heart cyclic-AMP phosphodiesterase by oncomodulin. *Biochem Biophys Res Commun* 1985;131:500-506

Nagasaka A, Ohkubo S, Hidaka H. 3':5'-cyclic-nucleotide phosphodiesterase in the bovine pituitary gland. *Biochim Biophys Acta* 1983;755:481-487

Narindrasorasak S, Tan LU, Seth PK, Sanwal BD. Regulation of cyclic adenosine 3':5'-monophosphate phosphodiesterase. *J Biol Chem* 1982;257:4618-4626

Nasu S, Shioji K, Ueda I, Tanigawa Y, Shimogama M. Ca2+/protein modulator-dependent and -independent cyclic GMP phosphodiesterase from hog heart. *J Biochem* 1978;83:1449-1458

Nemoz G, Prigent A-F, Pageaux JF, Pacheco H. Isoelectric-focusing patterns of cyclic nucleotide phosphodiesterase from rat heart. *Biochem J* 1981;199:113-119

Nemoz G, Prigent A-F, Pacheco H. Analysis of cyclic nucleotide phosphodiesterase by isoelectric focusing coupled to an activity stain. *Anal Biochem* 1983;133:296-301

Nemoz G, Prigent A-F, Moueqqit M, Fougier S, Macovischi O. Pacheco H. Selective inhibition of one of the cyclic AMP phosphodiesterase from rat brain by the neurotropic compoumd rolipram. *Biochem Pharmacol* 1985;34:2997-3000

Nikawa J-I, Sass P, Wigler M. Cloning and characterization of the low affinity cyclic AMP phosphodiesterase gene of Saccharomyces cerevisiae. *Mol Cell Biol* 1987;7:3629-3636

Niles RM, Loewy B. Endogenous inhibitor of cyclic nucleotide phosphodiesterase in cultured human epithe ial cells. *Biochem Biophys Res Commun* 1981;99:236-242

Nishizuka Y, Takai Y, Kishimoto A, Hashimoto E, Inove M, Yamomoto M, Criss WE, Kuroday Y. A role of calcium in the activation of a new protein kinase system. *Adv Cyclic Nucleotide Res* 1978;9:209-220

Onali P, Strada SJ, Chang L, Epstein PM, Hersh EM, Thompson WJ. Purification and characterization of high-affinity cyclic adenosine 5'-monophosphate phosphodiesterase from human acute myelogenous leukemic cells. *Cancer Res* 1985;45:1384-1391

Orellana O, Jedlicki E, Allende CC, Allende JE. Properties of a cyclic nucleotide phosphodiesterase of amphibian oocytes that is activated by calmodulin and calcium by tryptic proteolysis, and by phospholipids. *Arch Biochem Biophys* 1984;231:345-354

Ovchinnikov YA, Lipkin VM, Kumarev VP, Gubanov VV, Khramstov NV, Akhmedov NB, Zagranichny VE, Muradov KG. Cyclic GMP phosphodiesterase from cattle retina. *FEBS Lett* 1986;204:288-292

Palmer WK, Kane TA, Bach F, Doukas S. Effect of cold exposure on liver and muscle cAMP content and cAMP phosphodiesterase activities. *J Appl Physiol* 1985;58:211-216

Patwardhan VV, Lanthnier A. Cyclic GMP phosphodiesterase and guanylate cyclase activities in rabbit ovaries and the effect of in-vivo stimulation with LH. *J Endocrinol* 1984;101:305-310

Pledger WJ, Stancel GM, Thompson WJ, Strada SJ. Separation of multiple forms of cyclic nucleotide phosphodiesterases from rat brain by isoelectricfocusing. *Biochim Biophys Acta* 1974;370-242-248

Pichard A-L, Cheung WY. Cyclic 3',5'-nucleotide phosphodiesterase: interconvertible multiple forms and their effects on enzyme activity and kinetics. *J Biol Chem* 1976;254:5726-5736

Podgorski GJ, Franke J, Kessin RH. Isolation of a cDNA encoding a portion of the cyclic nucleotide phosphodiesterase of Dictyostelium discoideum. *J Gen Microbiol* 1986;132:1043-1050

Pober JS, Bitensky MW. Light-regulated enzymes of vertebrate retinal rods. *Adv Cyclic Nucleotide Res* 1979;11:265-290

Prasad KN. Role of cyclic nucleotides in the expression of malignancy and differentiated functions in nerve cells. *Adv Med Oncol Res Educ* 1979;4:57:73

Prasad KN. Involvement of cyclic nucleotides in transformation. In: *The transformed cell*, edited by IL Cameron and TB Pool 1981;235-236. Academic Press, New York

Purvis K, Olsen A, Hansson V. Calmodulin-dependent cyclic nucleotide phosphodiesterases in the immature rat testis. *J Biol Chem* 1981;256:11434-11441

Pyne NJ, Cooper ME, Houslay MD. Identification and characterization of both the cytosolic and particulate forms of cyclic GMP-stimulated cyclic AMP phosphodiesterase from rat liver. *Biochem J* 1986;234:325-334

Pyne NJ, Cooper ME, Houslay MD. The insulin- and glucagon-stimulated "dense vesicle" high-affinity cyclic AMP phosphodiesterase from rat liver. *Biochem J* 1987;242:33-42

Rall TW, Sutherland EW. Adenyl cyclase :II. The enzymatically catalyzed formation of adenosine 3',5'-phosphate and inorganic pyrophosphate from adenosine triphosphate. *J Biol Chem* 1962;237:1228-1231

Ralph RK. Cyclic AMP, calcium and control of cell growth. *FEBS Lett* 1983;161:1-7

Rasmussen H, Jensen P, Lake W, Friedman N, Goodman DBP. Cyclic nucleotides and cellular calcium metabolism. *Adv Cyclic Nucleotide Res* 1975;5:375-393

Rasmussen H, Goodman DPB. Relationships between calcium and cyclic nucleotides in cell activation. *Physiol Rev* 1977;57:421-50.

Rasmussen H, Kojima I, Kojima K, Zawalich W, Apfeldorf W. Calcium as intracellular messenger: sensitivity modulation, C-kinase pathway, and sustained cellular response. *Adv Cyclic Nucleotide Protein Phosphor Res* 1984;18:159-193

Rasmussen H. The calcium messenger system: part I. *New Engl J Med* 1986(a);314:1094-1101

Rasmussen H. The calcium messenger system: part II. *New Engl J Med* 1986(b);314:1164-1170

Reeves ML, Leigh BK, England PJ. The identification of a new cyclic nucleotide phosphodiesterase in human and guinea-pig cardiac ventricle. *Biochem J* 1987;241:535-541

Rhoads AR, Morris HP, West WL. Cyclic 3',5'-nucleotide monophosphate phosphodiesterase activity in hepatomas of different growth rates. *Cancer* Res 1972;32:2651-2655

Ridgway AAG, De Vouge MW, Mukherjee BB. Dibutyryl cAMP inhibits expression of transformation-related properties in Kirsten murine sarcoma virus transformed Balb/c-3T3 cells despite continued presence of p21(v-Ki-ras). *Biochem Cell Biol* 1988;66:54-65

Rillema JA. Cyclic nucleotides, adenylate cyclase and cyclic AMP phosphodiesterase in mammary glands from pregnant and lactating mice (39299). *Proc Soc Exp Biol Med* 1976;151:748-751

Rillema JA, Mulder JA, Anderson LD. Cyclic nucleotides and their associated enzymes in 9,10-dimethyl-1, 2-benzanthracene-induced mammary tumors of rats. *Cancer Res* 1978;38:741-744

Rillema JA. Calcium requirement for prolactin actions on ribonucleic acid and casein synthesis in mouse mammary gland explants. *Endocrinol* 1980;106:1360-1364

Robinson MF, Fok G, Kundig H, Savage N. Cytosolic cyclic nucleotide phosphodiesterases from a spontaneous mouse mammary tumour. *S Afr Cancer Bull* 1983;27:171-172

Robinson MF, Savage N, Esser JD, Levin J. Retrospective survey of the clinical significance of cytosolic oestrogen receptor status in 287 patients with primary breast cancer. *S Afr J Surg* 1985;23:107-109

Robinson MF, Levin J, Savage N. Characterization of soluble cyclic AMP phosphodiesterases and partial purification of a major form in human leiomyoma of the uterus. *Clin Physiol Biochem* 1987;5:249-260

Robison GA, Butcher RW, Sutnerland EW. *Cyclic AMP.* Academic Press, New York 1971:17-28

Rodbard D, Chrambach A. Estimation of molecular radius, free mobility and valence using polyacrylamide gel electrophoresis. *Anal Biochem* 1971;40:95-134

Rosenfeld MG, O'Malley BW. Steroid hormones: effects on adenyl cyclase-activity and adenosine 3',5'-monophosphate in target tissues. *Science* 1970;168:253-255

Rosenthal A, Moses H, Beaver D, Seschewan S. Lead ion and phosphodiesterase histochemistry: nonenzymatic hydrolysis of nucleotide phosphates by lead ion. *J Histochem Cytochem* 1966;14:698-707

Ross EM, Gilman AG. Biochemical properties of hormone-sensitive adenylate cyclase. *Ann Rev Biochem* 1980;49:533-564

Rossi P, Pezzotti R, Conti M, Geremia R. Cyclic nucleotide phosphodiesterase in somatic and germ cells of mouse semniferous tubules. *J Reprod Fertil* 1985;74:317-327

Russell TR, Terasaki WL, Appleman MM. Separated phosphodiesterases for the hydrolysis of cyclic adenosine 3',5'-monophosphate and cyclic guanosine 3',5'-monophosphate in rat liver. *J Biol Chem* 1973;243:1334-1340

Rutten WJ, Schoot BM, De Pont JHM. Adenosine 3',5'-monophosphate phosphodiesterase assay in tissue homogenates. *Biochim Biophys Acta* 1973;315:378-383

Ryan WL, Heidrick ML. Inhibition of cell growth in vitro by adenosine 3',5'-monophosphate. *Science* 1968;162:1484-1485

Ryan WL, Heidrick ML. Role of cyclic nucleotides in cancer. *Adv Cyclic Nucleotide Res* 1974;4:81-116

Sabatine JM, Coffee CJ. The identification and characterization of two cyclic nucleotide phosphodiesterases from bovine adrenal medulla. *Arch Biochem Biophys* 1986;249:95-105

Sadan O, Van Iddekinge B, Van Gelderen CJ, Savage N, Becker PJ, Van Der Walt LA, Robinson MF. Oestrogen and progesterone receptor concentrations in leiomyoma and normal myometrium. *Ann Clin Biochem* 1987;24:263-267

Saitoh Y, Hardman JG, Wells JN. Differences in the association of calmodulin with cyclic nucleotide phosphodiesterase in relaxed and contracted arterial strips. *Biochemistry* 1985;24:1613-1618

Sakai T, Makino H, Tanaka R. Increased activity of cyclic AMP phosphodiesterase from frozen-thawed rat liver: a role of lysosomal protease in enzyme activation. *Biochim Biophys Acta* 1978;522:477-490

Saltiel AR, Steigerwalt RW. Purification of putative insulin-sensitive cAMP phosphodiesterase or its catalytic domain from rat adipocytes. *Diabetes* 1986;35:698-704

Saltiel AR. Insulin generates an enzyme modulator from hepatic plasma membranes: regulation of adenosine 3',5'-monophosphate phosphodiesterase, pyruvate dehydrogenase and adenylate cyclase. *J Cyclic Nucleotide Protein Phosphor Res* 1987;11:345-354

Sapag-Hagar M, Greenbaum AL. The role of cyclic nucleotides in the development and function of rat mammary tissue. *FEBS Lett* 1974;46:180-183

Sapag-Hagar M, Greenbaum AL. Adenosine 3',5'-monophosphate and hormone interrelationships in the mammary gland of the rat during pregnancy and lactation. *Eur J Biochem* 1974;47:303-312

Sarada K, Epstein PM, Strada SJ, Thompson WJ. Analysis of cyclic nucleotide phosphodiesterase(s) by radioimmunoassay. *Arch Biochem Biophys* 1982;215:183-198

Sass P, Field J, Nikawa J, Toda T, Wigler M. Cloning and characterization of the high-affinity cAMP phosphodiesterases of Saccharomyces cervisiae. *Proc Natl Acad Sci(USA)* 1986;83:9303-9307

Savage N, Robinson MF, Silberman K, Kundig H, Levin J. Cyclic 3'-5'-AMP phosphodiesterase in a spontaneous mammary tumour in the mouse. *J Steroid Biochem* 1983;19:125

Scatchard G. The attractions of proteins for small molecules and ions. *Ann NY Acad Sci* 1949;51:660-672

Schmidtke J, Meyer H, Epplen JT. Cyclic AMP phosphodiesterases of the rat ovary. oestrous cycle dependent activity change of high affinity form. *Acta Endocrinol* 1980;404-411

Seth PK, Rogers J, Narindrasorasak S, Sanwal BD. Regulation of cyclic adenosine 3':5'-monophosphate phosphodiesterases: altered pattern in transformed myoblasts. *J Cell Physiol* 1983;116:336-344

Author Robinson M F
Name of thesis Partial Purification and Characterization of Soluble Cyclic Nucleotide Phosphodiesterases in Human and Murine tissues 1988

PUBLISHER:
University of the Witwatersrand, Johannesburg
©2013

LEGAL NOTICES:

Copyright Notice: All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed, or otherwise published in any format, without the prior written permission of the copyright owner.

Disclaimer and Terms of Use: Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.