

A NEW SPECIES OF *PSAMMOBATES* (REPTILIA: TESTUDINIDAE) FROM THE EARLY PLEISTOCENE OF SOUTH AFRICA.

by

Donald G. Broadley

Research Associate, Natural History Museum of Zimbabwe

Present address: Biodiversity Foundation for Africa, P.O. Box FM 730, Famona, Bulawayo, Zimbabwe

ABSTRACT

A new species of *Psammobates* Fitzinger is described from early Pleistocene cave deposits at Sterkfontein in the Gauteng Province of South Africa. It seems to be most closely related to *P. oculiferus* (Kuhl), which today occurs to the west and north of the type locality, but the proportions of its depressed carapace come closest to *Homopus femoralis* Boulenger.

KEYWORDS: *Psammobates*, Pleistocene, South Africa, Drimolen.

INTRODUCTION

Until recently almost all the fossil tortoises recorded from Africa were large forms belonging to the genus *Geochelone* (Meylan & Auffenberg 1986), apart from some *Chersina* fossils reported from the Pliocene of Langebaanweg (Hendey (1973, 1981) and from upper Pleistocene and Holocene deposits throughout the southern Cape Province (Klein & Cruz-Urbe 1983). Meylan & Auffenberg (1986) recorded *Kinixys erosa* Schweigger and a new genus and species *Impregnochelys pachytectis* from the Miocene of Kenya and *Chersina* sp. from the Miocene of South Africa (Arrisdrift on the Orange River), also noting the presence of *Chersina* among Pleistocene material from Hopefield. Subsequently *Homopus fenestratus* was described from Carlisle Bridge in the Eastern Cape Province (geological horizon unknown) (Cooper & Broadley 1990).

Braun (1981: p.184) reported fragments of unidentified tortoise carapace and plastron from Sterkfontein Member 5, Swartkrans Member 2 and Kromdraai A and B, but most of these would probably be *Geochelone*, which is the common tortoise in cave deposits at Makapansgat (Broadley 1962) and in Zimbabwe (Broadley in prep.).

In the circumstances, the recent discovery of a partial shell of one of the smaller South African genera is of considerable interest.

MATERIAL & METHODS

Comparative osteological material of all African tortoise genera was available in the Herpetology Department of the Natural History Museum of Zimbabwe in Bulawayo (NMZB). The osteology of the shells of the Malagasy genus *Pyxis* and its subgenus *Acinixys* is well illustrated by Bour (1981).

Nomenclature for bones follows Zangerl (1969) and the convention for neural formulae is that of Auffenberg (1974).

SYSTEMATIC PALAEOONTOLOGY

Class Reptilia

Order Testudines

Family Testudinidae

Psammobates antiquorum sp. nov

Diagnosis: A species of *Psammobates* apparently closest to *P. oculiferus*, but different in the more depressed carapace, with a step slop anteriorly, the wider anterior peripherals and the very broad posterior lobe of the plastron. The anterior neural formula is

? ? <6 <6 4.

Etymology: The specific name is derived from *antiquorum* (Latin = of old times), as this is the first extinct species of *Psammobates* to be described.

Holotype: DN803, an adult, collected by Dr. A. Keyser of the Palaeo-anthropological Research Group, University of the Witwatersrand, housed in the collections of the Bernard Price Institute for Palaeontological Research, University of the Witwatersrand, Johannesburg. The specimen consists of a partial shell, missing the portion of the carapace posterior to the seventh peripherals, the fifth (left) or sixth (right) costals and the sixth neural. The carapace has gaped open between the second and third neurals and costals post mortem; the left half of the nuchal, most of first and second neurals and the first and second left costals are missing. There is minor damage to the margins of the carapace and the anterior lobe of the plastron, while much of the left side of the plastron is missing

Type locality and horizon: Drimolen, a Plio-Pleistocene dolomitic cave site located on the farm Sterkfontein 519JQ, Krugersdorp District, Gauteng Province, South Africa. From Plio-Pleistocene sediments estimated at 1.6 to 2 M.a.

Comparisons and generic assignment: As the specimen is incomplete, its generic assignment was determined by a process of elimination. Despite its small size, the specimen has a fully ossified shell with complete sutures between all bones, thus ruling out *Malacochersus*. The lack of any indication of a carapacial hinge and the alternate widening and narrowing of costals 3 to 5 eliminates *Kinixys* from consideration. The depressed carapace and $<6 <6 <4$ formula for neurals 3 to 5 indicates that the specimen cannot be assigned to *Geochelone*, *Testudo*, *Chersina* or *Pyxis*, which all have at least the fourth neural octagonal (Meylan & Auffenberg 1986; Bour 1981), while *Acinixys* has a $6 > 6 > 6 > 6 >$ neural formula (Bour 1981). The sulcus on the left epiplastron indicates that the gulars were at least as long as broad – a diagnostic character separating *Psammobates*

from *Homopus* (Loveridge & Williams 1957). In their cladogram for the Testudinidae, Gaffney & Meylan (1988) treat *Psammobates* and *Homopus* as sister genera.

DESCRIPTION

Measurements

Plastron length 98 mm; maximum shell width (at sutures of peripherals 4 and 5) 80 mm; maximum shell height (at anterior edge of neural 3) 48 mm.

Carapace

Probably oval in outline, somewhat depressed (Table 1). From the front of the first neural the nuchal slopes at an angle of 45° to the horizontal. The first neural is missing (apart from an anterior fragment with an interdigitating suture) and the second is represented only by the right posterior portion, which has parallel outer borders. The third neural is almost square, but the anterior corners are cut off, making it hexagonal like the fourth neural, which is wider anteriorly, while the fifth is tetragonal and narrower caudad, the shape of the damaged sixth

TABLE 1.
Shell proportions for African Species of Testudinidae

		Plastron Length	Shell Width	Shell Height	H/W	Ratios L/W	H/L
<i>Geochelone pardalis</i>	NMZB 6751	103.0	77.5	61.5	0.79	1.33	0.60
<i>Testudo graeca</i>	NMZB-UM 32982	152.0	126.6	87.0	0.69	1.20	0.57
<i>Kinixys natalensis</i>	NMZB 1107	99.0	83.0	50.5	0.61	1.19	0.51
<i>Psammobates antiquorum</i>	DLN - LI-P2-1	98.0	80.0	49.0	0.61	1.22	0.50
<i>Psammobates aocculiferus</i>							
NMZB 6104	No locality	94.0	80.0	58.0	0.72	1.17	0.62
NMZB - UM 9823	Botswana	93.0	78.5	57.0	0.73	1.18	0.61
NMZB-UM 12756	Botswana	81.0	71.0	51.0	0.72	1.14	0.63
NMZB-UM12757	Botswana	91.0	80.0	59.0	0.74	1.14	0.65
NMZB-UM12758	Botswana	84.0	76.0	49.0	0.64	1.11	0.58
NMZB-UM16109	Botswana	89.0	75.0	52.0	0.69	1.19	0.58
NMSB-UM 23226	Botswana	75.0	66.0	44.5	0.67	1.14	0.59
<i>Psammobates tentorius</i>							
NMZB 5531	Beaufort West	108.0	92.5	66.0	0.71	1.17	0.61
NMZB 5534	Hanover	93.0	87.0	55.0	0.63	1.07	0.59
NMZB 6650	Beaufort West	103.0	88.0	68.5	0.78	1.17	0.67
NMZB 7173	Springbok-Augrabies	75.0	68.0	45.5	0.67	1.10	0.61
NMZB-UM 33636	Victoria West	76.0	69.0	42.0	0.61	1.10	0.55
<i>Psammobates geometricus</i>							
NMZB 6563	Gordons Bay-Strand	94.0	79.0	60.0	0.76	1.19	0.64
<i>Homopus femoralis</i>							
NMZB 840	Middelburg	70.5	59.0	35.0	0.59	1.19	0.50
NMZB 1260	Cradock District	119.0	104.0	55.0	0.53	1.14	0.53
NMZB 7069	Neuweldberg	70.0	56.0	32.5	0.58	1.25	0.46
<i>Homopus areolatus</i>							
NMZB 1261	Port Elizabeth	76.5	64.5	35.5	0.55	1.19	0.46
NMZB 6460	Port Elizabeth	89.0	75.5	50.0	0.66	1.18	0.56
<i>Homopus boulengeri</i>							
NMZB 7179	Sutherland	81.0	68.5	37.5	0.55	1.18	0.46
<i>Homopus signatus</i>							
NMZB 6468 10Km	S of Steinkopf	83.5	71.0	41.0	0.58	1.18	0.49
<i>Chersina angulata</i>							
NMZB-UM 32979	Port Elizabeth	136.5	97.0	67.5	0.70	1.41	0.49

neural cannot be determined. The sutures between the first, second and third costals are almost straight and parallel, the third and fifth costals are wider proximally and narrower distally, while this condition is reversed in the fourth costal. The first four peripherals continue the slope of the first two costals, with no tendency to recurve; the fifth and sixth become more steeply sloping and the seventh (left) flares outward. It is unfortunate that the

posterior portion of the carapace is missing, because the three Recent species of *Psammobates* have diagnostic suprapygal patterns (Broadley 1997).

Plastron

The left epiplastron shows traces of a gular sulcus, indicating that the gulars were at least as long as broad, while the endoplastron is similar in size and shape to that of *P. oculiferus*. The right hyoplastron




Figure 1. Left - *Psammobates antiquorum*, holotype: (A) dorsal, (B) lateral and (C) ventral views. KEY: C = costal; Ento = entoplastron; Epi = epiplastron; Hypo = hyoplastron; Hypo = hypoplastron; L = left; N = neural; Nu = nuchal; P = peripheral; R = right; Xiphi = xiplastron. The heavy stipple indicates the sandy matrix, the light stipple indicates plaster filler. Right - *Psammobates oculiferus* (NMZB-UM 9823 10km west of Nata, Botswana): (D) dorsal, (E) lateral and (F) ventral views. The line represents 1 cm to scale.




Figure 2. Anterior views of the shells of (A) *Psammobates oculiferus* (NMZB-UM 9823); (B) *Psammobates antiquorum*, holotype; (C) *Homopus femoralis* (NMZB 1260 - Cradock District, Eastern Cape Province, South Africa). The line represents 1 cm to scale.

and the damaged hypoplastra closely match those of *P. oculiferus*. The xiphiplastra are much broader than those of Recent *Psammobates* or *Homopus* and look more like those of *Testudo*, although the margins are too damaged to show the development of notches at the femoro-anal sulci and the anal notch.

Anterior aspect

The depressed shell is most noticeable from the front, because the nuchal and peripherals 1 and 2 are not recurved, the opening for head and limbs is very low, comparable with that of *Homopus femoralis* rather than any species of *Psammobates* (Figure 2).

ZOOGEOGRAPHY

No Recent species of *Psammobates* occurs in Gauteng Province today, but *P. oculiferus* occurs in the North-West Province and western Free State, reaching its western limit in the Northern Province. This species inhabits arid grassland and open *Acacia* savanna on deep Kalahari sands (Kalahari Thornveld), a region which has cold dry winters (Boycott & Branch 1989). Krugersdorp District lies in the Bankveld, a sour grassveld, but the area was probably covered with bushveld vegetation up to 500 years ago (Acocks 1988) and this habitat is occupied by *P. oculiferus* in Pietersburg District.

ACKNOWLEDGEMENTS

I thank Dr A. Keyser for making the fossil tortoise available for study and Wulf Haacke (Transvaal Museum) for arranging to send it to me on loan.

REFERENCES

- ACOCKS, J.P.H. 1988. *Veld types of South Africa. Mem. Bot. Survey S. Afr.*, No. 57.
- AUFFENBERG, W. 1974. Checklist of fossil land tortoises (Testudinidae). *Bull. Florida State Mus., Biol. Sci.* **18**, 121-251.
- BOYCOTT, R.C. & BRANCH, W. 1989. *Psammobates oculifer*. In: Swingland, I.R. & Klemens, M.W. Eds. *The conservation biology of tortoises*, pp -- 88- 90. (Occ. Pap. IUCN Species Survival Commission No. 5) Gland: IUCN
- BOUR, R. 1981. Etude systematique du genre endemique Malagache *Pyxis* Bell, 1827 (Reptilia, Chelonii). *Bull. Soc. Linn. Lyon*, **50**(4), 132-176.
- BRAIN, C.K. 1981. *The hunters or the hunted. An introduction to African cave taphonomy*. Chicago, University of Chicago Press.
- BROADLEY, D.G. 1962. Some fossil Chelonian fragments from Makapansgat. *Nature*, **194** (4830), 791-792.
- 1997. Diagnostic suprapygal patterns in the genus *Psammobates* Fitzinger (Chelonii: Testudinidae). *Arnoldia Zimbabwe*, **10** (12), 121-126.
- COOPER, M.R. & BROADLEY, D.G. 1990. A new species of fossil *Homopus* (Cryptodira: Testudinidae) from South Africa. *Studia Palaeocheloniologica* **3** (4), 41-55.
- GAFFNEY, E.S. & MEYLAN, P.A. 1988. A phylogeny of turtles. In: Benton, M.J., Ed. *The phylogeny and classification of the Tetrapods, Vol. I: Amphibians, reptiles, birds*, 157-219. Oxford, Clarendon Press.
- GREIG, J.C. & BURDETT, P.D. 1976. Patterns in the distribution of southern African terrestrial tortoises (Cryptodira: Testudinidae). *Zoologica Africana*, **11** (2), 249-273.
- HENDEY, Q. B. 1973. Fossil occurrences at Langebaanweg, Cape Province. *Nature* **244**, 13-14.
- 1981. Palaeoecology of the Late Tertiary fossil occurrences in "E" Quarry, Langebaanweg, South Africa, and a reinterpretation of their geological context. *Ann. S. Afr. Mus.*, **84**, 1-104.
- KLEIN, R.G. & CRUZ-URIBE, K. 1983. Stone age population numbers and average tortoise size at Byneskranskop Cave 1 and Die Kelders Cave 1, Southern Cape Province, South Africa. *S. Afr. archaeol. Bull.*, **38**, 26-30.
- LOVERIDGE, A. & WILLIAMS, E.E. 1957. Revision of the African tortoises and turtles of the suborder Cryptodira. *Bull. Mus. comp. Zool. Harv.*, **115**, 163-557.
- MEYLAN, P. & AUFFENBERG, W. 1986. New land tortoises (Testudines: Testudinidae) from Miocene of Africa. *Zool. J. Linn. Soc.*, **86**, 279- 307.
- ZANGERL, R. 1969. The Turtle Shell. In: Gans, C., Bellairs, A. d'A. & Parsons, T.S. *Biology of the Reptilia. I. Morphology A.*, pp 311- 339. London/ New York, Academic Press