

https://dspace.org/

Nature based solutions for coastoal resilience in South Africa

Jasper Knight

https://hdl.handle.net/10539/41551
WOS

Downloaded from a DSpace repository. https://dspace.org/



South African Geographical Journal

ISSN: (Print) (Online) Journal homepage: www.tandfonline.com/journals/rsag20

Nature-based solutions for coastal resilience in South Africa

Jasper Knight

To cite this article: Jasper Knight (2024) Nature-based solutions for coastal resilience in South Africa, South African Geographical Journal, 106:1, 21-50, DOI: 10.1080/03736245.2023.2193565

To link to this article: https://doi.org/10.1080/03736245.2023.2193565

Published online: 23 Mar 2023.

Submit your article to this journal

Article views: 416

View related articles

View Crossmark data

Citing articles: 1 View citing articles

Full Terms & Conditions of access and use can be found at
https://www.tandfonline.com/action/journalInformation?journalCode=rsag20

SOUTH AFRICAN GEOGRAPHICAL JOURNAL
2024, VOL. 106, NO. 1, 21–50
https://doi.org/10.1080/03736245.2023.2193565

Routledge
Taylor & Francis Group

Check for updates

Nature-based solutions for coastal resilience in South Africa

Jasper Knight

School of Geography, Archaeology & Environmental Studies, University of the Witwatersrand, Johannesburg, South Africa

ABSTRACT

Coastlines globally are sensitive to the effects of sea-level rise, increased coastal storminess and changes in coastal sediment supply and sediment dynamics in the Anthropocene. Coastlines are also influenced by land use change, urbanization and development of built infrastructure. These changes can affect the dynamics of coastal landforms, weaken coastal resilience and make coasts more sensitive to climate hazards. This study critically examines the properties of coastlines that contribute to coastal biophysical resilience in South Africa, highlighting their relativerates of change and dynamic behaviour in response to physical and human forcing factors. Coastal landforms can be considered as 'green infrastructure' that can buffer the effects of climate change as well as providing ecosystem and environmental services in their own right. Viewing coastal landforms as green infrastructure provides a 'nature-based solution' to mitigate against climate change impacts that can work with – not against – the natural geomorphic, sedimentary and ecological processes of coastlines. Coastal landforms can also contribute to socioecological resilience, where they provide environmental and ecosystem services. The green infrastructure approach to coastal resilience has not been well developed in South Africa but is more effective in supporting coastal sustainable development.

ARTICLE HISTORY
Received 11 February 2022
Accepted 9 March 2023

KEYWORDS
Anthropogenic climate change; coastal processes; green infrastructure; nature-based solutions; sandy beaches

1. Introduction

Coastlines lie at the interface of terrestrial and marine environments and, as such, are sensitive to hazards such as sea-level rise and extreme storms (Kesavan & Swaminathan, 2006; Kron, 2013; Luijendijk et al., 2018; Mather & Stretch, 2012; Ramesh et al., 2015). Understanding the impacts that these and other hazards can have on coastlines is important for hazard impact mitigation, management, and sustainable development. Hazard impacts on coasts have been extensively studied worldwide at different spatial and temporal scales, and using many different dataset types, from numerical models of future climate or wave regimes, to geological evidence for the imprints of past hazards, and to behavioural analysis of affected coastal populations (e.g. Gaillard et al., 2009; Kazemi et al., 2017; Pile et al., 2018). The resilience of coastlines to withstand hazards is related to the nature of the climate events themselves (their magnitude, frequency, duration); the properties of the affected coastline (sandy vs rocky coasts, nearshore

CONTACT Jasper Knight jasper.knight@wits.ac.za
© 2023 The Society of South African Geographers

bathymetry and coastal hinterland topography, vegetation); the nature of human activity and infrastructure in the affected region (population exposed, socioeconomic factors, urban development, port/harbour facilities, geoengineering such as sea walls); and any mitigating actions that may reduce hazard impacts to the natural or to the built or human environments (Glavovic et al., 2015; Ruckelshaus et al., 2020). In South Africa, several studies on coastal hazards and their impacts have been undertaken, but these have tended to focus on urban areas (e.g. Colenbrander & Bavinck, 2017; Colenbrander et al., 2015; Mather & Stretch, 2012). There has been less attention paid to open rocky and sandy coastlines, and in understanding their morphodynamics and responses to climate forcing (Botha et al., 2018; La Cock et al., 1992; Mitchell et al., 2005; Musekiwa et al., 2015). This is important as these rocky, sandy and urbanized coastlines, with different landforms and dynamics, respond differently to climate forcing and may have different abilities to withstand climate forcing (i.e. their resilience). In a recent analysis, Williams et al. (2022) evaluated the 'intactness' of global coastlines based on the anthropogenic footprint present within the coastal hinterland, using data points spaced 1 km apart. For South Africa (number of data points = 2606), the authors calculated that 97.5% of the coastline was exposed to high human pressure arising mainly from demographic factors (based on GPD, GINI coefficient, regulatory quality), compared to 47.9% globally. This may highlight the potential sensitivity – and thus low resilience – of the South African coastline to climate change impacts and coastal hazards.

The concept of coastline resilience provides a useful and integrated framework to consider the effects of climate forcing as it describes the net *outcomes* of forcing on a coastline rather than the detailed factors or properties that give rise to those outcomes (Nichols et al., 2019; Flood & Schechtman, 2014; Masselink & Lazarus, 2019). *Resilience* is useful because it considers the nature of the feedbacks that exist between the human environment (geoengineering, socioeconomic and cultural factors) and the physical environment, through the ways in which coastal landforms are used, valued and managed (Knight, 2022a). Measuring coastal resilience, however, is not easy because it requires (1) establishing the state of the system that exists before a forcing event(s) takes place; (2) understanding the nature of the forcing event(s) itself, including its magnitude, characteristics and spatial and temporal extent; and (3) evaluating its impacts on the physical and human environments, and any changes in the workings of physical and human systems that have taken place afterwards, bearing in mind that the spatial and temporal extent of any impacts may not be the same as the original forcing (Brown et al., 2016; Schultz & Smith, 2016). These sequential steps and this kind of *post hoc* analysis is in reality difficult to achieve. This is because it requires monitoring of coastal and hinterland sediment systems and geomorphic changes over long (10^2–10^4 year) timescales, which is often unachievable, and it assumes a simple forcing – response relationship along a coastline that does not change. An alternative approach to evaluating coastal resilience is by monitoring and measuring coastal morphodynamic responses to forcing by wind, waves and/or sea-level changes that take place over decadal or shorter timescales, where field and remote sensing datasets are more likely to be available. This has been the focus of most coastal research undertaken in South Africa, based on field or remote sensing methods from individual coastal stretches (e.g. Corbella & Stretch, 2012a; Harris et al., 2011a; Henrico et al., 2020; Knight & Burningham, 2019, 2021; Lubke & Webb, 2016; Mitchell et al., 2005). The advantage of this approach is that it is not

dependent on *a priori* understanding of the state of the system and fits with the spatial and temporal scales of coastal morphodynamic behaviour and coastal management (Cowell & Thom, 1994). However, there has been less work on considering how data from such studies can be used to evaluate the vulnerability of different coastal stretches in South Africa, and how vulnerability and risk arise as a result of climate forcing and human activity in combination (Colenbrander et al., 2015; Musekiwa et al., 2015).

Sandy coastlines are particularly vulnerable to the effects of climate change because of the relative ease with which sand grains can be moved by wind and water. The vulnerability of sandy coasts to climate change has been noted worldwide (Knight & Harrison, 2009; Luijendijk et al., 2018; Ranasinghe & Stive, 2009). Sandy coasts make up 38% of South Africa's coast, with 29% of the remainder being rocky and 32% being mixed rocky-sandy (Wepener & Degger, 2019), and this likely means that these coasts will be differentially affected by climate forcing. The aim of this study is to critically examine the resilience of different landforms present along the South African coast. This is achieved through consideration of their dynamic behaviour, based on examples of specific landform types. The study highlights that different geomorphic features along coastlines can be considered as 'green infrastructure' that provides natural resilience to the coastline. Further, certain management strategies can increase landform resilience as well as the provision of environmental and ecosystem services along the coast, thereby also increasing societal resilience (Passeri et al., 2021). Focusing on the natural resilience provided by coastal landforms is a nature-based solution to address climate change impacts along coastlines globally, and treats coastlines as integrated biophysical and socioecological systems. This is an integrated and progressive approach to climate change risk management (e.g. Knight, 2021a, 2022b).

2. The definition and scope of resilience

Pachauri & Meyer (2014, p. 127) define resilience as the capacity of social, economic and environmental systems to cope with a hazardous event or trend or disturbance, responding or reorganizing in ways that maintain their essential function, identity and structure, while also maintaining the capacity for adaptation, learning and transformation. In a coastal context, Masselink & Lazarus (2019, p. 1) define resilience as 'the capacity of the socioeconomic and natural systems in the coastal environment to cope with disturbances, induced by factors such as sea level rise, extreme events and human impacts, by adapting whilst maintaining their essential functions'. This therefore considers resilience as having both physical and human elements, termed biophysical and socioecological resilience, respectively (Poe et al., 2014; Silva et al., 2019), but that there may be complex relationships between them.

Resilience can arise as a result of the properties of coastal landforms themselves (their size, shape, location, geology, ecosystems) and their dynamical behaviour. Here, rapid temporal and spatial changes in landform processes and properties (i.e. highly dynamic behaviour) suggest that the landform has high sensitivity (low resilience) to forcing; landforms that do not change significantly over time or space exhibit more subdued dynamic behaviour and are therefore considered to have low sensitivity (high resilience) to forcing. Resilience can be broadly considered as the opposite of vulnerability (Angus & Hansom, 2021; Masselink & Lazarus, 2019), so when resilience is high vulnerability is

low, and *vice versa*. Both terms have been used in the context of coastal physical systems and human activity along coasts (Alexandrakis & Poulos, 2014; Angus & Hansom, 2021; Ballesteros & Esteves, 2021) although societal resilience to coastal events and hazards (which is particularly important in the developing world) has not been well explored in southern Africa (Cabral et al., 2017; DST, 2017; Palmer et al., 2011).

2.1. Resilience along the South African coast

Different landforms present along the South African coast should in theory exhibit different degrees of resilience (and its converse, vulnerability) to forcing. The key landform types under consideration in this study are indicated in Table 1. These landforms are identified based on their commonness along the South African coast (Dardis & Grindley, 1988). These are discussed in detail below. Overall, it can be seen that coastal landforms of different types can provide physical and topographic buffers against sea-level rise and wave erosion and flooding (Hanley et al., 2014; Kim et al., 2021), as these are the major mechanisms of geomorphic change and coastal impacts that are issues along the South African coast (Holmes et al., 2016). Threats and vulnerabilities arise from climate change through sea level and wave forcing, and from different types of human activity along the coast. Both of these sources of vulnerability are amplified along sandy coasts where unconsolidated sand is easily eroded, transported and deposited, resulting in rapid geomorphic change and thus higher sensitivity to forcing (Hanley et al., 2014; Knight, 2022a; Luijendijk et al., 2018). Rocky shorelines exhibit completely different styles of vulnerability, influenced by geologic structure and elevation of the shore platform (Payo et al., 2015).

3. Examples of the dynamics of coastal landforms in South Africa

Every coastal landscape comprises different landforms that may exhibit different controls, dynamics and thus varying levels of resilience to forcing (Hapke et al., 2013; Schultz & Smith, 2016). Examples of the dynamics of the different major landforms found along the South African coast are now discussed (Table 1), with the aim of relating their dynamic behaviour to their potential for coastal resilience.

3.1. Coastal sand dunes

Coastal sand dune locations in South Africa have been mapped by Tinley (1985, p. 19) who recognized a number of dune types: bare; vegetated; related to topographic barriers; associated with temporary wetlands; and relict/gully eroded dunes. These are located in different positions around the coast, in particular in the southern Cape where headland – embayment systems dominate. However, in detail, these mapped locations are often incorrect and dune types are incorrectly assigned (Knight, 2021b). A morphological classification of coastal dunes in South Africa, however, identifies seven types: estuary barrier, headland bypass, mainland beach, prograding beach ridge, transverse, parabolic/migrating and climbing (Jackson et al., 2014). Rates of coastal dune migration have been examined in only a few studies, mainly from unvegetated transverse dunes located in the supratidal zone of sandy beaches, where these dunes respond dynamically to seasonal

Table 1. Key landform types and their contributions to resilience along the South African coast.

Landforms	Contribution of these landforms to resilience	Threats and vulnerabilities
Coastal sand dunes	• Buffer of coastal erosion as a topographic barrier • Sediment store for potential reworking between the dunes and beach • Substrate for plant and animal ecosystems • Store of organic carbon in vegetation and soil • Groundwater store and buffer for landward diffusion of salt water • Alteration of microclimate, as windbreak or firebreak for areas inland • Provision of ecosystem and environmental services for local communities and tourists	• Sea-level rise and wave erosion of the foredune foot • Infrastructure development, increased foot/vehicle traffic • Decreased indigenous species biodiversity • Agriculture/overgrazing • Deforestation, introduction of invasive species (e.g. marram grass *Ammophila arenaria*, sea buckthorn *Hippophae rhamnoides*, rhododendron *Rhododendron* spp.) • Fire (natural or caused by people) • Wind erosion and development of blowouts • Sand mining • Development of tourist facilities (car parking, caravan parks, campsites, toilets, cafes) within the dunes • Changes in groundwater position/recharge of dune slacks and ponds
Sandy beaches	• Dissipate wave energy • Buffer against sea-level rise and coastal flooding • Role as sediment store for longshore drift • As sites of recreation/tourism, contributing to societal socioeconomic and cultural resilience	• Sea-level rise and transgressive wave erosion of the shoreface • Storms and the effects of storm waves as extreme events • Sand mining on the beach or adjacent sand dunes, estuaries or nearshore • Changes in river mouth location • Geoengineering of sea walls, revetments, groynes, and development of the backbeach environment (tourist facilities, walkways, apartments) • Updrift developments that may impact on longshore sediment supply • Human overuse, pollution/trash • Human health risks – drowning, rip currents, sharks/jellyfish attacks in the nearshore • Sand dune development in the backbeach/supratidal zone, contributing to beach squeeze
River mouths and estuaries	• Sediment supply to sandy beaches • Barrier protects the river mouth/lagoon from wave erosion • River mouth/lagoon as a store of blue and green carbon • Sites of high biodiversity for plants (see Mangroves, below), animals and birds (including protection as Ramsar sites and Biosphere Reserves)	• Changes in river water and river sediment yield from the catchment inland • Decreased water quality as a result of industrial/agricultural pollution/runoff from within the catchment • Infrastructure development around the river mouth, leading to increased sedimentation/lagoon infilling, or increased erosion • Barrier breaching and increased exposure of the river mouth to wave erosion/increased salt water overwash into the lagoon • Changes in longshore sediment transport; can result in the barrier becoming thinner and thus more vulnerable to wave overwashing • Sea-level rise and barrier overtopping • Changes in barrier opening/closing behaviour, with impacts on lagoon processes, water quality and ecosystems

(*Continued*)

Table 1. (Continued).

Landforms	Contribution of these landforms to resilience	Threats and vulnerabilities
Mangroves	• Areas of higher sedimentation; buffer of wave erosion • Blue and green carbon store • Substrate for associated ecosystems, especially fish and invertebrates	• Water pollution or decreased water quality from incoming rivers or within lagoons, related to industry and agriculture • Sea-level rise and salt water transgression inland • Wave/storm erosion of the mangrove front • Invasive species • Climate change (water temperature, dissolved oxygen content) • Infrastructure development or encroachment into the mangrove area • Agriculture, livestock grazing, deforestation, wood harvesting on the mangrove fringe
Rocky shorelines	• Buffer of sea-level rise • Dissipation of wave energy across the rock platform	• Possibility of sudden cliff collapse • Vulnerable to sea-level rise, increased wave height and energy and enhanced shoreface erosion • Sea-level rise and migration of the intertidal zone, driving ecosystem responses • Effects of wave bores • Invasive rocky shoreline species • Pollution runoff from the adjacent land

changes in wind strength and direction. Tinley (1985, p. 207) gave backbeach dune migration rates in the Algoa Bay dune field of 4 m yr^{-1}, with migration rates in the inland dune field varying between undisturbed (0.1 m yr^{-1}) and disturbed areas (1.3 m yr^{-1}). Examples of calculated rates from different locations and different dune types along the South Africa coasts are 2.9 m yr^{-1} (La Cock et al., 1992), 3.2–12.3 m yr^{-1} (Knight & Burningham, 2019), 3.5 m yr^{-1} (Lubke & Sugden, 1990), 4.44–1.50 m yr^{-1} (Knight & Burningham, 2021), and 7.23–9.00 m yr^{-1} (Henrico et al., 2020). However, these values are calculated over different timescales and employ different methodologies (field, remote sensing), meaning that results from these studies cannot be easily compared. There are also statistically significant differences ($p < 0.01$) between summer (3.99–11.51 m yr^{-1}) and winter (3.55–9.94 m yr^{-1}) transverse dune migration rates (Knight & Burningham, 2021), which indicate the role of seasonal climate forcing in dune dynamics (Burkinshaw & Rust, 1993). A key element influencing dune migration rates is the relationship that the monitored dunes have to surrounding landforms in different coastal contexts. Sand dunes found adjacent to river mouths respond to variations in fluvial sediment supply (Knight & Burningham, 2022; Olivier & Garland, 2003), and this shows that dune dynamics may reflect other controls apart from wind climate. Lucrezi et al. (2014) showed that infrastructure development in South Africa decreases dune width and biodiversity, and thus the vulnerability of the dune system to climate or wave erosion. Sand dune rehabilitation strategies in South Africa have focused on encouraging vegetation growth, monitored through changes in biodiversity and soil properties (van Aarde et al., 1996, 1998; Ott & van Aarde, 2014), and mainly in response to disturbance by mining rather than for mitigation of climate impacts (Mentis & Ellery, 1994; Lubke et al., 1996). Previously, dune stabilization had been achieved through planting of the European marram grass *Ammophila arenaria* (Hellström, 1996), although now indigenous species are preferred as they can result in higher biodiversity and carbon sequestration potential (de Villiers et al., 1999; van Rooyen et al., 2013). A further

outcome of vegetation growth is stabilization of the land surface, reduced dune movement and sediment yield from erosion (Hellström, 1996), and thus increased resilience to erosion (Hanley et al., 2014; Jackson et al., 2019).

3.2. Sandy beaches

There are few studies on the morphodynamics of sandy beaches in South Africa, but dissipative, reflective and intermediate beach states have all been identified (Harris et al., 2011a). Several studies highlight the role of storm waves as the driver of coastal erosion, in particular by steepening the shoreface and decreasing beach width (Smith et al., 2010; Corbella & Stretch, 2012a, b). Shoreline change is most significantly related to wave peak period and wave direction, and these attributes are influenced by episodic storms (Smith et al., 2013; Corbella & Stretch, 2012a). There are very little field-based data on the dynamics of sandy beaches, including short-term changes in beach profile and volume, although low-resolution but long term (1973–2010) profile data suggest a beach recovery response time of two years after storm events with a volume flux of 90 m^3 m^{-1} yr^{-1} (Corbella & Stretch, 2012b), reflecting the timescale of system resetting and thus the degree of resilience of such coastlines. It is not clear to what extent the dynamic behaviour of different beaches may vary spatially or temporally, or between different beach types (Harris et al., 2011a). Some previous work highlights the close coupling within integrated beach – dune systems in South Africa, where sediment moves from beach to dune storage and back again (Hellström, 1996), and where storm-forced reductions in beach width lead to reduced accommodation space for supratidal dune development (Knight & Burningham, 2021). However, volume fluxes and the rates of morphological change of the beach systems have not been widely presented. Systematic mapping of sandy beach width along the South African coast does not show consistent variations in gross morphological properties between different coastal stretches. However, it highlights how bedrock outcrops can act as pinning points, stabilizing the beach front and thus beach width variability, and reducing beach dynamics (Knight & Burningham, 2022). These bedrock-pinned beaches are therefore sites of higher resilience along sandy coasts.

Sandy beach dynamics may refer to variations in their volume, surface slope, intertidal width, supratidal width, surficial landforms (embryo dunes, beach cusps) or longshore volume fluxes. Different studies may measure one or several of these elements using different methods and this means that their results may not be comparable. At a site scale, monitoring of changes in beach properties over time has been undertaken in several studies in South Africa (e.g. Guastella & Smith, 2014; Mitchell et al., 2005). The most detailed beach morphological study is along different sectors of the KwaZulu-Natal coast, which describes the results of near-monthly beach surveys between 2011 and 2020 (Leuci et al., 2022). Seasonal variations in swell wave frequency and longshore processes give rise to variations in beach aggradation/erosion behaviour. Reduced fluvial sediment input and downstream sediment starvation from the development of the Port of Durban results in net beach erosion, with increased sediment yield and net aggradation in a down-drift direction. Such long-term variations in beach state have implications for coastline resilience, with beaches being more vulnerable where they are narrower and sediment-starved. Aeolianite and beachrock are also found extensively along sandy stretches of the eastern

South African coast (Cawthra & Uken, 2012; Wiles et al., 2018). Progressive development of these lithified coastal sediment through the late Quaternary has 'hardened' the coastline, increasing coastal resilience and providing a fixed substrate that can reduce net longshore sediment transport and promote differential erosion on 'harder' and 'softer' sandy beaches. There has also been consideration of how beach properties and dynamics can influence population viability and species richness of marine/intertidal macrofauna (Dye et al., 1981; Harris et al., 2013,). This highlights that beach dynamics can also influence ecological resilience (Kombiadou et al., 2019) and that this can arise as a result of different beach and environmental properties (beach geometry, sediment grain size, tidal/wave regime, seasonal/interannual volume flux) (Harris et al., 2013,).

3.3. River mouths and estuaries

The dynamics of river mouths in South Africa have been mainly investigated in terms of changes in the size, shape and open/closed status of the lagoons or small estuaries that are commonly found there. These water areas are impounded behind a sandy beach bar or barrier located at the end of the river mouth that separates the river from the sea (e.g. Cooper, 2001; Garden & Garland, 2005; Green et al., 2013; Magoro et al., 2020). Variations in river discharge cause changes in the position of the river channel, changes in river sediment supply and river/coastal geomorphic activity (Cooper et al., 1990; Cooper, 1989; Fernandes & Adams, 2016; Magoro et al., 2020; Marker, 2004; Pillay et al., 2003). A beach bar or barrier that extends across the river mouth serves to hold back freshwater in the lagoon at the river mouth, and separates the functioning of the river system from the coastal system that is influenced by waves and tides (Bate et al., 2017; Cooper, 2001). The development or erosion of this barrier by coastal longshore processes means that river mouths may be variously open, closed or partly open at different times, according to phases of wave overwashing and barrier breaching (Garden & Garland, 2005; Green et al., 2013; Magoro et al., 2020). Although measurements of rates of change of different elements around river mouths and tidal inlets can be made, these do not necessarily inform on system resilience where the major morphological control is the fronting barrier system (Green et al., 2013). Here, phases of barrier overwashing (wave forcing) and lagoon overspilling (river discharge forcing) reflect the exceedance of a critical threshold in system behaviour. It is likely that this behaviour is cyclic (Marker, 2000), by patterns of sediment redistribution by river channel migration, or by seasonal wave climate (Lubke & Webb, 2016; Pillay et al., 2003). This means that barrier resilience is also cyclic and in response to both fluvial and marine forcing.

3.4. Mangroves

Mangroves can offer significant coastal resilience as they can buffer coastal erosion and sea-level rise (Barbier, 2016; Blankespoor et al., 2017; Gedan et al., 2011). They can also be considered as a coastal 'landform' where their roots stabilize the substrate and encourage sediment aggradation, especially around the margins of river mouths (Adams, 2020; Charrua et al., 2020). Bedin (2001) showed a rapid rate of mangrove colonization across a developing delta as part of the Mhlathuze River estuary, KwaZulu-Natal, at initial rates of 20–55 ha yr^{-1} (1976–1982) and then at a steady 5.4 ha yr^{-1} (1986–1999). Mangrove

ecosystems are found in river mouths and estuaries at 41 locations on the east coast of South Africa with around half of these sites being less than 1 ha in area (Steinke, 1999). Several studies have examined recent changes in mangrove area along the South African coast. For example, in the period 1982–1999 mangrove area in the Eastern Cape Province decreased by 6.5% with local extinction at three estuary sites (Adams et al., 2004). Analysis of changes in mangrove forest area at 17 sites in the Eastern Cape Province (1982–2012) shows a decrease at 9 sites and increase at 7 sites (one site shows no change) (Hoppe-Speer et al., 2015). Overall, however, there is a decrease in total mangrove area in the region by 12% over this time period. Rajkaran et al. (2009) showed that mangroves along the KwaZulu-Natal coast were only present in river estuaries that remained open to the sea for more than 56% of the time, in the period 1982–1999. When closed from the sea by a barrier, river mouths experience environmental stresses (higher water salinity and pore-water content) that result in a decrease in mangrove survival (Charrua et al., 2020; Hoppe-Speer et al., 2013). This is particularly the case for micro-estuaries formed at the mouths of small rivers (Bate et al., 2017) which, by virtue of their size, are sensitive to changes in rainfall inland (driving river discharge), lagoon water properties, and wave climate (driving longshore drift and lagoon barrier dynamics) (Charrua et al., 2020). Vertical and horizontal erosion and/or accretion rates in South African mangrove systems have not been evaluated, but rates elsewhere are highly variable, with averaged lateral erosion rates of 0.27–0.96 m yr^{-1} along mangrove coasts in Bangladesh, but with short-term storm-forced rates of 5 m yr^{-1} (Mullick et al., 2020). In India, decadal-scale shoreline erosion rates are 0.91–3.12 m yr^{-1} (Jayanthi et al., 2018). Vertical sediment accretion in Senegal is 0.2–0.3 cm yr^{-1} (Sakho et al., 2011). Thus, mangrove 'landforms' can change in both area and height. The impacts of climate change on the distributions and productivity of mangrove ecosystems in South Africa have been examined based on trajectories of historical data (Hoppe-Speer et al., 2015; Peer et al., 2018) and on future climate and ecological modelling (Yang et al., 2014). Both these approaches suggest a poleward spread of mangrove distributions but a net decrease in area, mainly as a result of human activity (agriculture, pollution, urbanization) rather than sea-level rise and coastal squeeze. However, likely sites at which these changes may take place have not yet been identified (Peer et al., 2018).

3.5. Rocky shorelines

Rocky shorelines include the morphological elements of rock cliffs and rock shore platforms. Despite being relatively common in South Africa (Emery & Kuhn, 1982; Illgner, 2008), rocky shoreline morphology and dynamics have not been examined in detail, either along extensive shore platforms or as part of headland – embayment systems of sandy shorelines (Knight & Grab, 2015). Different rock landforms along the Eastern Cape coast include prominent sea cliffs whose heights are geologically controlled, in addition to caves, arches and stacks (Illgner, 2008; Knight & Grab, 2015). Rock headlands can provide accommodation space within their intervening embayments for sediment accumulation and can therefore protect these deposits from erosion by the process of wave refraction (Shaw et al., 2001). Coastal caves in South Africa are key sites for the preservation of geological records of Quaternary coastal environmental, archaeological and sea-level changes (Hendey & Volman, 1986; Jacobs et al., 2003), and based on different radiogenic dating methods are over 1 million years old (Pickering et al.,

2013). This may in turn indicate the resilience of rock coast features in the landscape, and their survival over multiple Quaternary climate cycles.

Most work along South African rock coasts has been undertaken on shore platforms where microscale variations in rock surface hardness and weathering features reflect differential exposure to subaerial weathering (which softens the bedrock surface) and wave erosion (which removes weathered rock revealing a harder bedrock surface) (Knight & Burningham, 2020a, b). Wave detachment of bedrock blocks takes place in the lower part of the platform below mean high water spring tide positions; wetting and drying associated with wave-splash around highest astronomical tide positions; and land surface runoff in a landward position on rock platforms above highest astronomical tide (Knight & Burningham, 2020a). Bioerosion takes place mainly in seaward locations. Rock platform erosion rates have not been calculated locally, but long term (1973–2003) microerosion monitoring in New Zealand yields a rate of 1.09 mm yr^{-1} (Stephenson et al., 2010). However, such an averaged rate should be considered as highly spatially and temporally variable (Naylor & Stephenson, 2010). Rock cliff collapse or rock platform erosion can potentially protect the rocks behind from wave attack, reducing future erosion.

Different molluscan and macrophyte ecosystems are also found along rocky shorelines in South Africa, and these have been examined in terms of their relationships to bedrock properties and wave regime (Bustamante & Branch, 1996; Dye, 1998; Tucker et al., 2017). Sink et al. (2005) also identified different rocky shoreline intertidal ecosystems in KwaZulu-Natal, thus different biogeographical regions, according to water temperatures and therefore upwelling conditions. The growth of organisms on rock surfaces can also decrease wave velocity and protect against rock erosion (Baxter et al., 2022; Gowell et al., 2015). Ecological resilience along rocky shorelines therefore arises where the shoreline itself provides suitable habitats for certain species, enhancing coastal biodiversity (Malherbe & Samways, 2014), and where ecosystem development contributes to coastal protection (Mead et al., 2013).

4. Coastal landforms as 'green infrastructure'

This survey of different coastal landform types in South Africa highlights their potential to exhibit resilience to coastal forcing (Table 1). As such, these landforms can be considered as 'green infrastructure' that acts as a buffer against coastal processes such as wave erosion and sea-level rise (Ruckelshaus et al., 2016; Sohn et al., 2021; Sutton-Grier et al., 2018), and can thus reduce hazard risk and impacts, and promote resilience (Kombiadou et al., 2019; Monteiro & Ferreira, 2020; Saleh & Weinstein, 2016). The concept of green infrastructure emphasizes how the three-dimensional nature of coastal landforms can be used in the same way as hard engineering structures to reduce coastal hazard impacts (Cohn et al., 2022; Pontee et al., 2016; Schoonees et al., 2019). The role of coastal landforms in buffering coastal change and providing the context for sustainable coastal management has been examined in several studies globally (Chang & Mori, 2021; Silva et al., 2019; Sohn et al., 2021). Most commonly, such green infrastructure has been framed in the context of providing a physical barrier that helps protect against negative coastal change – for example, a sand dune ridge that has a greater height than the height of any incoming waves (Kombiadou et al., 2019; Ranasinghe & Stive, 2009). This is,

however, a very narrow definition of the meaning and application of green infrastructure. This is because coastal landforms do not just provide resilience against *physical* change (e.g. beach width, volume), they can also promote the resilience of the *properties* of the coastline, such as its environmental and ecosystem services (Spalding et al., 2014). The ability of coastal properties to maintain or enhance these environmental and ecosystem services (Knight, 2021a) is important in the context of climate change. For example, clearing invasive plants, maintaining boardwalks, planting unstable sand surfaces and fencing off areas such as dune slacks are management strategies that can enhance coastal dune ecosystems, protect endangered species, encourage carbon storage, and reduce erosion (Itzkin et al., 2020; Keesstra et al., 2018; Morris et al., 2019). These are all strategies that can promote dune system resilience to climate change (e.g. Farrell & Connolly, 2019; Itzkin et al., 2020), independent of a dune system existing as a geomorphic barrier. This highlights that considering the natural resilience of coastlines, and then identifying ways in which management strategies can strengthen this resilience, can help reduce coastal hazard risk and impacts (Hanley et al., 2014; Knight & Goff, 2016; Schultz & Smith, 2016). Thus, coastal landforms can provide 'nature-based solutions' to environmental and climate changes, both locally and globally. The potential of coastal landforms in South Africa to be used as 'green infrastructure' as part of coastal management strategies is now examined (Table 2). However, most work on coastal management policies and practices has focused on the legislative context of coastal management, the nature of engagement with stakeholder communities and developing frameworks for management decision-making (e.g. Colenbrander, 2019; Goble et al., 2017, 2019, 2020; Taljaard et al., 2019). There have been few studies that have considered the management of coastal landscapes, their properties and dynamics, or that have employed ongoing monitoring or modelling of their behaviour to evaluate the success of management strategies. In South Africa, these activities have been undertaken in three major contexts, as described in the literature.

(1) *Estuary and river mouth* studies show how anthropogenic changes in river dynamics can impact on estuary sedimentation. For example, damming along the Umgeni River (KwaZulu-Natal) from the 1980s onwards resulted in a decrease in river discharge, decreased downstream sediment yield, and fining of sediment reaching the river mouth (Garland & Moleko, 2000). Land use change in the Salt River catchment (Eastern Cape) has influenced sedimentation in the lower part of the estuary, and channel migration patterns (Marker, 2004). The sensitivity of estuary ecosystems to pollution and other anthropogenic changes has motivated different management strategies, especially in estuaries whose mouths may be periodically closed to the sea, which can negatively impact on water quality, benthic and planktic organisms, and sediment dynamics (Claassens et al., 2020; Whitfield et al., 2012). As a result, controlled artificial breaching, as was done in the Umdloti River estuary (KwaZulu-Natal) in February 2004, can be used to drain anoxic lagoonal water. Artificial breaching also worked successfully on the Nkongweni River mouth (Margate, KwaZulu-Natal) where lagoon drainage relieved pressure on coastal infrastructure (Guastella et al., 2014). However, a similar breaching strategy used in the St Lucia estuary system from the mid-1950s to 2001 resulted in seawater incursion and hypersaline conditions, causing

Table 2. Management strategies that can increase the resilience of coastal landforms, drawn from the literature worldwide but with specific reference to South Africa.

Landforms	Actions and their likely outcomes
Coastal sand dunes	• Banning new sand mining and requiring full ecological and geomorphological rehabilitation of existing/closing sand mines • Restrict groundwater extraction • Revegetation/planting for increased dune surface stability • Sand or brushwood fences, to reduce wind speed and encourage sand deposition • Planting and wind fencing in the supratidal zone, to encourage embryo dune formation • Moving existing beach sand to engineer an artificial dune, which is then planted and stabilised • Clearance of scrub and invasive species, to promote indigenous plant species growth • Maintain dune slacks and ponds by shallow excavation, to encourage habitat diversity especially for birds and amphibians • Construct and maintain boardwalks/fences to keep pedestrians from wandering • Fencing off areas seasonally for ground-nesting birds and flowering of rare plants • Provide litter bins and information signs to guide good visitor behaviour
Sandy beaches	• Beach nourishment by dredging/pumping to increase beach width • Building groynes to obstruct or slow longshore drift • Planting of the supratidal zone to encourage embryo dune growth and to increase the surface elevation of the supratidal zone • Build or maintain offshore shoals/surf breaks to decrease nearshore wave height • Where the beach is narrow or with high wave energy: stabilize the landward position of the beach with a sloping seawall, gabion, tetrapods, riprap • Build and maintain a landward barrier to the beach to prevent storm wave overwashing or sand blow inland, especially where urban development gets very close to the beach. This barrier may include (hard engineering): a seawall, revetment, gabion, riprap (soft engineering): engineered beach ridge, planted low dune or sand fences • Support beach users by providing car parking, toilets and other facilities, places to dispose of trash, accessible entry to the beach environment, information and safety/signage boards
River mouths and estuaries	• Manage any river mouth barrier, by artificial breaching/dredging (to drain lagoon water or to encourage seawater ingress) or building up the barrier (to keep water in the lagoon in dry periods) • River mouth stabilization by jetties/training walls • Ensure a low-elevation riparian fringe around river/lagoon margins, to encourage plant succession, reduce river flow velocity, promote carbon storage and support other plants/animals • Manage and increase water quality within the river system, in particular from agriculture, industry and wastewater treatment plants • Monitor water quality within the lagoon to reduce the likelihood of anoxic conditions and water stratification in hot/dry periods, removal of invasive species (e.g. water hyacinth *Pontederia crassipes*) • Sea defences around tidal flats/saltmarsh around estuaries can be breached to allow for tidewater ingress, which increases tidal flat elevation through enhanced deposition, and increases biodiversity and carbon storage • Banning sand mining within estuaries and river mouths • Ensuring that roads/bridges/railways spanning the river mouth or estuary have pilings that do not interfere with water or sediment flow
Mangroves	• Replant areas of lost vegetation • Decrease pollution from upstream industrial and agricultural sources • Maintain tidal creeks to prevent them from closing up due to sediment aggradation • Manage the activities of other mangrove users (e.g. farmers, wood harvesters) • Remove any invasive plant species • Clear overgrown mangrove patches to encourage fresh regrowth and provide softground habitats for fish/invertebrates • Restrict infrastructure development that would either reduce the mangrove area or cause enhanced erosion of the mangrove fringe

(*Continued*)

Table 2. (Continued).

Landforms	Actions and their likely outcomes
Rocky shorelines	• Rock armouring on the landward edge of rock platforms, to reduce wave impacts on rock cliffs or infrastructure • Understand the landward limit of extreme tide/wave events and do not allow infrastructure development seaward of this limit • Monitor the stability of rock cliffs or caves, warn users to keep away from these areas • Support unstable rock slopes and reduce mass movements by rock bolts, rock nets/mesh, geotextiles, planting of soft-rock cliffs, grading slopes. This can also support plant biodiversity and provide bird nest sites • Reduce risk of waves to users by providing safety equipment and safety warning signage • Support intertidal ecosystems by ensuring that rock pools are maintained, and removing invasive macrophyte and mollusc species • Reduce land-derived water runoff onto the rock platform

die-back of shore vegetation (Taylor et al., 2006). Dredging has also been done to maintain the river mouth channel during low rainfall periods: 600,000 m^3 of sediment accumulated at the St Lucia estuary mouth between February 1988 and November 1989, of which 466,000 m^3 was subsequently dredged, although this has not resolved ongoing problems of sediment aggradation (Wright & Mason, 1993). Floods and droughts also cause problems of estuary mouth stability and infrastructure erosion in other estuary systems (Schumann, 2021). Catchment land use changes, in particular agricultural practices and invasive species, can lead to macrophyte and invertebrate habitat loss as well as changes in estuary water quality and the potential for coastal erosion (Adams, 2020; Fernandes & Adams, 2016; de Villiers et al., 2021). For example, pollution from sugarcane plantations has reduced natural habitat area and species richness in the uMkhomazi and Mvoti estuaries (KwaZulu-Natal) (Fernandes & Adams, 2016), and this can be considered to have resulted in estuary system resilience (e.g. Claassens et al., 2020; Rajkaran et al., 2009).

(2) *Sand dune revegetation and afforestation* has taken place along some South African coasts over the last ~100 years, undertaken as a management strategy to stabilize dune surfaces and reduce dune and blown sand migration (Hertling & Lubke, 1999; Lubke, 2022). The causes of sand dune destabilization in South Africa include overgrazing, deforestation/vegetation clearance, sand mining, fire, population pressure through tourism and development, coastal erosion and the development of new dunes by river mouth migration (Hellström, 1996; Lubke & Sugden, 1990; Mentis & Ellery, 1998). Sand dune or blown sand movement is seen as a problem where it impacts on people or infrastructure, by blocking roads or silting up drainage channels (Nichols, 1996; La Cock & Burkinshaw, 1996; Lucrezi et al., 2014). Dune revegetation and planting is a recognized standard coastal management strategy and has been undertaken along many sandy coasts worldwide (e.g. Hayes & Kirkpatrick, 2012; Lithgow et al., 2013). Although its impacts are most clearly evaluated in terms of reduced dune migration and sediment transport (Nichols, 1996), planting of non-native and often invasive species such as European marram grass (*Ammophila arenaria*) and *Acacia* spp. can have significant impacts on plant and animal biodiversity (Ott & van Aarde, 2014; van Aarde et al., 1996). Lubke (2022) reviewed the history of marram grass

revegetation strategies and successes on sand dunes along the temperate Eastern Cape coast, highlighting that it helps in dune stabilization and does not tend to outcompete indigenous species. This should also be considered in combination with sand fences, which reduce windspeed and encourage sand deposition to form embryo dunes (Itzkin et al., 2020). Along the KwaZulu-Natal coast under subtropical conditions, dune revegetation appears to be a less common strategy, but here the major driver of sand dune degradation is sand mining, followed by piecemeal environmental rehabilitation (Lubke et al., 1996; Mentis & Ellery, 1998; van Aarde et al., 1998). Stands of high-density secondary succession and climax dune forests in KwaZulu-Natal stabilize much older backdune systems (Goble et al., 2014) and thus can be considered to increase the resilience of the dune cordon as a whole.

(3) *Beach and shoreline manipulation* includes a number of management strategies. Beach nourishment focuses on maintaining sediment supply to sandy beaches, thus maintaining beach width and promoting resilience, and often as a direct response to the effects of inappropriate coastal management updrift (Castelle et al., 2009). Infrastructure development and building of groynes/piers around the Port of Durban in the period 1982–1988 led to downdrift sediment starvation to KwaZulu-Natal beaches north of Durban. After 1990, around 265,000 m^3 yr^{-1} of marine sand was pumped onto the beaches in order to broadly maintain shoreline position following sediment redistribution by longshore processes (Rautenbach & Theron, 2018). The continual need for artificial beach maintenance along this coastal stretch, caused by Port of Durban development, has led to this coastline being deemed to be on 'life support', where it cannot survive without regular human intervention (Cooper & Jackson, 2019). Coastal reclamation, by extending the land surface seaward by port, harbour or urban development, can also be considered as a strategy to develop coastal cities and to increase the resilience of the coastline by stabilizing it with built structures. Coastal reclamation has been done extensively in Cape Town and also around the ports of Durban and Richards Bay (Nichols, 1996), and there is a government White Paper on this topic (2017). However, such development can create further problems with changing coastal erosion/deposition patterns, and may increase flood risk and the vulnerability of infrastructure to extreme events (e.g. Griggs & Patsch, 2019). Coastal setback, which is based on a coastal buffer zone of varying width (or height above mean sea level) within which development is restricted, is also promoted in South Africa as a means to manage coastal risk (Colenbrander et al., 2015; Desportes & Colenbrander, 2016). However, this is based on risk aversion rather than working with coastal properties to increase resilience and reduce climate change impacts.

As outlined above, these three major approaches to coastal management in South Africa only apply to certain coastal types like sand dunes or estuaries. This is not to say, however, that coastal management has not been undertaken in other ways or in different coastal contexts. However, it is notable that many other coastal types (like rock coasts) and many low-population rural coasts have not been systematically evaluated for their biophysical dynamics, the human activities that take place there, or their management needs. There is simply a lack of data on many coasts. As a result, Table 2 lists a range of potential management interventions that can help increase coastal resilience. These are drawn from practices globally but focusing in particular on the physical landforms and

properties the different coastline types found in South Africa that are discussed herein, and that are presented in Table 1. Many of the suggested interventions (Table 2), however, have not been done extensively or even at all in South Africa, but these can form part of a menu of options for increasing coastal resilience, and working with – not against – the natural biophysical processes that take place along coastlines on different scales (Cohn et al., 2022; Hamin et al., 2018; Morris et al., 2019; Spalding et al., 2014).

5. Discussion

5.1. Resilience and coastal systems

Resilience is a useful concept when applied to coastal landforms and systems, because it describes the nature of changes to landform properties in response to forcing by climate and/or human activity. Such forcing – response relationships are the basis of understanding coastline responses to climate change in the Anthropocene (Kittinger & Ayers, 2010; Luijendijk et al., 2018; Martínez et al., 2017). Despite these advantages, the focus of coastline analysis in South Africa has been on mapping spatial vulnerability, not resilience, and consideration of the role of human activity along different coastal stretches (Musekiwa et al., 2015; Palmer et al., 2011). These few studies are also very generalized and do not explore in detail either the physical or human environments along these coasts. However, socioecological vulnerability has been well described in the literature (e.g. Bevacqua et al., 2018; Lloyd et al., 2013) and is especially important in the developing world where coastal populations may be poorer as well have closer relationships to the environmental and ecosystem services provided by the coast (Ballesteros & Esteves, 2021). It is very likely that this is also significant in South Africa although this has not been well explored, especially in rural areas (Palmer et al., 2010). Reconciling and considering the interplay of biophysical and socioecological resilience (vulnerability) along coasts, as has been done in other environments such as mountains (Knight, 2022b), should be considered as a key future research aim (Nichols et al., 2019; Glaser & Glaeser, 2014; Martínez et al., 2017; Sutton-Grier et al., 2018).

A key factor in resilience is the role of coastal geoengineering (sea walls, revetments, groynes, river mouth jetties, port/harbour construction) and changes in land use in the coastal hinterland (affecting runoff and sediment yield). These can affect the properties, dynamics and therefore the resilience of coastal landforms (e.g. Brown et al., 2016; Floerl et al., 2021; Griggs & Patsch, 2019; Hamin et al., 2018; Kittinger & Ayers, 2010). Overwhelmingly, geoengineering has resulted in negative impacts on coastal landforms and ecosystems, in which system integrity, connectedness and service provision have been degraded, resulting in decreased resilience (Berry et al., 2014; Schoonees et al., 2019; Tejada et al., 2007). However, more recent approaches to coastal (hard and soft) geoengineering such as managed realignment, dune afforestation, sandscaping and beach nourishment can increase coastline resilience when used appropriately (Nichols et al., 2019; Hamin et al., 2018; Masselink & Lazarus, 2019). These can also contribute to the morphological development of coastal landforms, by encouraging dune and beach growth and ecosystem change, thus making these landforms more 'green infrastructure' that can better withstand the impacts of climate forcing.

5.2. Nature-based solutions and green infrastructure to increase coastal resilience

There has not been a detailed examination of how green coastal infrastructure can be engineered or managed to provide nature-based solutions to coastal change impacts (Cohn et al., 2022; Floerl et al., 2021; King et al., 2021; Saleh & Weinstein, 2016). Studies have mainly focused on managing the properties of specific coastal landforms such as sand dunes (De Jong et al., 2014; Eichmanns et al., 2021) rather than coastal landscapes as a whole. However, understanding the sediment system dynamics of sandy coastlines (Knight & Goff, 2016) can better inform on not only their resilience but also how changes in the properties of dynamics of one landform can impact on other landforms or coastal properties (Floerl et al., 2021) (Figure 1). The concept of green (or ecological) infrastructure has also not been widely discussed in South Africa (e.g. Malan & Swart, 1997; Mbopha et al., 2021) although the ideas of riparian vegetation or footslope wetlands as a flood buffer, or bioremediation of water pollution in riparian wetlands, are well established in the literature in South Africa (e.g. Abiye et al., 2018; Belle et al., 2018), albeit without using the term 'green infrastructure'. Table 2 describes the management activities that can be employed to increase the resilience of specific coastal landforms as green coastal infrastructure. This highlights the potential of these activities to positively impact on both the maintenance of the coastal landforms themselves (in providing morphological resilience), and landform properties through their environmental and ecosystem services. National- and regional-scale coastal management strategies employing nature-based solutions and green infrastructure have been developed in the USA, the Netherlands, the UK and Australia, and these have focused on both specific landforms and coastal systems as a whole (e.g. Cohn et al., 2022; Davis et al., 2022; Morris et al., 2019). As such, there is an appreciation of coastal landforms as green infrastructure, and this is embedded as part of (some) coastal management strategies. However, these concepts have not really been considered in South African coastal management strategies at all, although the importance of landforms and ecosystems generally is implicit rather than explicit throughout.

5.3. Promoting green infrastructure and nature-based solutions for coastal management in South Africa

In South Africa, coastal management is undertaken at different levels by national, provincial and municipal governments under the Integrated Coastal Management Act (Act No. 24 of 2008), which is also informed by the Natural Environment Management Act (Act No. 107 of 1998) and other relevant instruments including the Coastal Management Policy Programme (1992–97), the National Water Act (Act No. 36 of 1998), and the Biodiversity Act (Act No. 10 of 2004), amongst others. Goble et al. (2014, pp. 168–169) present a table outlining the roles and responsibilities of different spheres of government in terms of the Integrated Coastal Management Act, and this highlights the multiple contexts in which coastal management takes place in South Africa (Celliers et al., 2015; Colenbrander et al., 2015; Desportes & Colenbrander, 2016; Guastella et al., 2014). Although this provides the legislative and policy framework for coastal management in the country, the strategies enacted in any one place may be highly variable, related to local expertise, finances and management priorities (Goble et al.,



Figure 1. Schematic model of how coastal resilience can be enhanced through working with the 'green infrastructure' of existing coastal landforms and their properties.

2020). The outcome is that this can lead to inconsistencies and contradictions in coastal management practices.

A management approach that works with – not against – the natural sedimentary and ecological processes of coasts (Hanley et al., 2014; Knight & Goff, 2016; Sutton-Grier et al., 2018) can more effectively increase biophysical resilience and reduce hazard impacts whilst also promoting sustainable socioeconomic development of local communities and thus socioecological resilience (Floerl et al., 2021). Coastal resilience in its totality in South Africa and elsewhere can be enhanced following two main approaches (see examples in Table 2):

- Supporting the continuation or the enhancement of the natural physical and ecological processes that take place in specific coastal environments (biophysical resilience). In allowing sediments to move from one location to another within the coastal zone (by wind, waves, tides, river flow), sedimentary landforms can naturally readjust to changing environmental conditions (De Jong et al., 2014; Hanley et al., 2014). In this way, they can be incorporated as soft (rather than hard) engineering structures (Gijsman et al., 2021; Pontee et al., 2016; Schoonees et al., 2019) that can contribute to coastline protection. This has been done specifically, for example, on mangroves (Gijsman et al., 2021), wetlands (Liu et al., 2021; Van Coppenolle & Temmerman, 2019), sand dunes (De Jong et al., 2014) and beaches/barriers (Hanley et al., 2014; Orchard & Schiel, 2021). Coastal ecosystems can be maintained and enhanced where indigenous species exist in areas that are protected from development or human overuse, and where invasive species are removed (De Battisti, 2021; Lithgow et al., 2013). This includes working with existing ecosystem succession and ecological conditions in order to achieve habitat rehabilitation (Ott & van Aarde, 2014; van Aarde et al., 1996). These ecosystems can contribute to coastal risk reduction by their stabilizing effects on the land surface (Spalding et al., 2014) as well as their contribution to carbon storage, biodiversity and ecosystem services (Lithgow et al., 2013; van Rooyen et al., 2013).

- Building the adaptive capacity of coastal communities (socioecological resilience). This focuses on understanding the relationship between coastal users and the coastline ecosystem and environmental services that contribute to socioeconomic and cultural activities and human wellbeing. In the developing world, coastal communities may be impoverished and have greater vulnerability, lower adaptive capacity and lower socioecological resilience (Almutairi et al., 2020; Ferro-Azcona et al., 2019). They may also rely on coastal properties and resources for livelihoods (through agriculture, fishing, tourism). Enhancing these coastal resources while also diversifying socioeconomic activity can better safeguard coastal communities against coastal change.

Biophysical and socioecological resilience are connected to each other, and studies highlight how one can enhance the other (e.g. Arkema et al., 2017; McFadden, 2010; Sutton-Grier et al., 2018; Trégarot et al., 2021). The nature of these interconnections is summarized in Figure 1, which describes the interplay between coastal biophysical and socioecological properties and dynamics, and how these can contribute to resilience (Silva et al., 2019). This means that effective and sustainable coastal management focusing on the integrity and dynamics of coastal landforms as an integrated system can help enhance the physical landscape and provide environmental and ecosystem services, and so in turn lead to wider societal and socioeconomic benefits.

6. Conclusions and future research directions

Resilience is a contested and multifaceted topic, but in different physical environments, biophysical and socioecological resilience (and its flip side, vulnerability) can be identified. In coastal environments, biophysical and socioecological properties and processes are strongly interlinked through the provision of ecosystem and environmental services and resources. However, changes in coastal environments globally – caused by ongoing climate change as well as increasing human development, land use change and pollution – are decreasing the resilience of natural coastal processes and landforms (Figure 1). This is a key issue when making use of coastal landforms as 'green infrastructure' (Hanley et al., 2014). These landforms also support the ecosystem and environmental services that contribute to socioecological resilience; thus, the effective management of coastal landforms can lead to positive impacts for coastal communities. This is particularly the case in the developing world such as South Africa.

Relationships between coastal resilience and risk management under climate change is important for the sustainable development of coastlines and coastal communities (Saja et al., 2021; Wood et al., 2021). To adequately address coastal resilience in countries such as South Africa, three systemic requirements are needed:

(1) An integrated Earth Systems approaches to coastal risk, vulnerability and morphodynamics, that is evidence-based and founded on an understanding of the dynamics of coastal biophysical systems (Figure 1). This is currently lacking in many areas of the world, including many South African coasts, and this means that (when done) managers are having to make decisions without the best information being available;

(2) Routine and ongoing monitoring of coastal properties and dynamics across different coastal sectors and different coastline types. This can include both biophysical and socioecological elements of coastal landscapes and include a combination of field and remote data types;
(3) Engagement with stakeholder communities, not just at the outset but as co-partners in coastal management at the local level. This is currently missing in many locations and contexts but is also important in terms of education, community empowerment and governance.

Employing these approaches as well as specific management strategies appropriate to particular coastal types (Table 2) can result in positive outcomes for both coastal landforms and communities.

Acknowledgments

I thank two anonymous reviewers for their comments on a previous version of this paper.

Disclosure statement

No potential conflict of interest was reported by the author.

References

Abiye, T., Mkansi, S., Masindi, K., & Leshomo, J. (2018). Effectiveness of wetlands in retaining metals from mine water, South Africa. *Water and Environment Journal*, *32*(2), 259–266. https://doi.org/10.1111/wej.12323

Adams, J. B. (2020). Salt marsh at the tip of Africa: Patterns, processes and changes in response to climate change. *Estuarine, Coastal and Shelf Science*, *237*, 106650. https://doi.org/10.1016/j.ecss.2020.106650

Adams, J. B., Colloty, B. M., & Bate, G. C. (2004). The distribution and state of mangroves along the coast of Transkei, Eastern Cape Province, South Africa. *Wetlands Ecology and Management*, *12*(5), 531–541. https://doi.org/10.1007/s11273-005-5165-0

Alexandrakis, G., & Poulos, S. E. (2014). An holistic approach to beach erosion vulnerability assessment. *Scientific Reports*, *4*(1), 6078. https://doi.org/10.1038/srep06078

Almutairi, A., Mourshed, M., & Ameen, R. F. M. (2020). Coastal community resilience frameworks for disaster risk management. *Natural Hazards*, *101*(2), 595–630. https://doi.org/10.1007/s11069-020-03875-3

Angus, S., & Hansom, J. D. (2021). Enhancing the resilience of high-vulnerability, low-elevation coastal zones. *Ocean & Coastal Management*, *200*, 105414. https://doi.org/10.1016/j.ocecoaman.2020.105414

Arkema, K. K., Griffin, R., Maldonado, S., Silver, J., Suckale, J., & Guerry, A. D. (2017). Linking social, ecological, and physical science to advance natural and nature-based protection for coastal communities. *Annals of the New York Academy of Sciences*, *1399*(1), 5–26. https://doi.org/10.1111/nyas.13322

Ballesteros, C., & Esteves, L. S. (2021). Integrated assessment of coastal exposure and social vulnerability to coastal hazards in East Africa. *Estuaries & Coasts*, *44*(8), 2056–2072. https://doi.org/10.1007/s12237-021-00930-5

Barbier, E. B. (2016). The protective service of mangrove ecosystems: A review of valuation methods. *Marine Pollution Bulletin*, *109*(2), 676–681. https://doi.org/10.1016/j.marpolbul.2016.01.033

Bate, G. C., Nunes, M., Escott, B., Mnikathi, A., & Craigie, J. (2017). Micro-estuary – a new estuary type recognised for South African conditions. *Transactions of the Royal Society of South Africa*, *72*(1), 85–92. https://doi.org/10.1080/0035919X.2016.1259689

Baxter, T. I., Coombes, M. A., & Viles, H. A. (2022). The bioprotective properties of the blue mussel (*Mytilus edulis*) on intertidal rocky shore platforms. *Marine Geology*, *445*, 106734. https://doi.org/10.1016/j.margeo.2022.106734

Bedin, T. (2001). The progression of a mangrove forest over a newly formed delta in the Umhlatuze Estuary, South Africa. *South African Journal of Botany*, *67*(3), 433–436. https://doi.org/10.1016/S0254-6299(15)31160-1

Belle, J. A., Collins, N., & Jordaan, A. (2018). Managing wetlands for disaster risk reduction: A case study of the Eastern Free State, South Africa. *Jàmbá: Journal of Disaster Risk Studies*, *10*(1). https://doi.org/10.4102/jamba.v10i1.400

Berry, A. J., Fahey, S., & Meyers, N. (2014). Boulderdash and beachwalls – the erosion of sandy beach ecosystem resilience. *Ocean & Coastal Management*, *96*, 104–111. https://doi.org/10.1016/j.ocecoaman.2014.05.006

Bevacqua, A., Yu, D., & Zhang, Y. (2018). Coastal vulnerability: Evolving concepts in understanding vulnerable people and places. *Environmental Science & Policy*, *82*, 19–29. https://doi.org/10.1016/j.envsci.2018.01.006

Blankespoor, B., Dasgupta, S., & Lange, G. -M. (2017). Mangroves as a protection from storm surges in a changing climate. *Ambio*, *46*(4), 478–491. https://doi.org/10.1007/s13280-016-0838-x

Botha, G. A., Porat, N., Haldorsen, S., Duller, G. A. T., Taylor, R., & Roberts, H. M. (2018). Beach ridge sets reflect the late Holocene evolution of the St Lucia estuarine lake system, South Africa. *Geomorphology*, *318*, 112–127. https://doi.org/10.1016/j.geomorph.2018.06.001

Brown, J. M., Prime, T., Phelps, J. J. C., Barkwith, A., Hurst, M. D., Ellis, M. A., Masselink, G., & Plater, A. J. (2016). Spatio-temporal Variability in the Tipping Points of Coastal Defense. *Journal of Coastal Research*, *75*(sp1), 1042–1046. https://doi.org/10.2112/SI75-209.1

Burkinshaw, J. R., & Rust, I. C. (1993). Aeolian dynamics on the windward slope of a reversing transverse dune, Alexandria coastal dunefield, South Africa. In K. Pye & N. Lancaster (Eds.), *Aeolian sediments, ancient and modern* (pp. 13–21). Blackwell, Oxford.

Bustamante, R. H., & Branch, G. M. (1996). Large scale patterns and trophic structure of southern African rocky shores: The roles of geographic variation and wave exposure. *Journal of Biogeography*, *23*(3), 339–351. https://doi.org/10.1046/j.1365-2699.1996.00026.x

Cabral, P., Augusto, G., Akande, A., Costa, A., Amade, N., Niquisse, S., Atumane, A., Cuna, A., Kazemi, K., Mlucasse, R., & Santha, R. (2017). Assessing Mozambique's exposure to coastal climate hazards and erosion. *International Journal of Disaster Risk Reduction*, *23*, 45–52. https://doi.org/10.1016/j.ijdrr.2017.04.002

Castelle, B., Turner, I. L., Bertin, X., & Tomlinson, R. (2009). Beach nourishments at Coolangatta Bay over the period 1987–2005: Impacts and lessons. *Coastal Engineering*, *56*(9), 940–950. https://doi.org/10.1016/j.coastaleng.2009.05.005

Cawthra, H., & Uken, R. (2012). Modern beachrock formation in Durban, KwaZulu-Natal. *South African Journal of Science*, *108*(7/8), 27–32. https://doi.org/10.4102/sajs.v108i7/8.935

Celliers, L., Colenbrander, D. R., Breetzke, T., & Oelofse, G. (2015). Towards increased degrees of integrated coastal management in the City of Cape Town, South Africa. *Ocean & Coastal Management*, *105*, 138–153. https://doi.org/10.1016/j.ocecoaman.2014.11.005

Chang, C. -W., & Mori, N. (2021). Green infrastructure for the reduction of coastal disasters: A review of the protective role of coastal forests against tsunami, storm surge, and wind waves. *Coastal Engineering Journal*, *63*(3), 370–385. https://doi.org/10.1080/21664250.2021.1929742

Charrua, A. B., Bandeira, S. O., Catarino, S., Cabral, P., & Romeiras, M. M. (2020). Assessment of the vulnerability of coastal mangrove ecosystems in Mozambique. *Ocean & Coastal Management*, *189*, 105145. https://doi.org/10.1016/j.ocecoaman.2020.105145

Claassens, L., Barnes, R. S. K., Wasserman, J., Lamberth, S. J., Miranda, N. A. F., van Niekerk, L., & Adams, J. B. (2020). Knysna Estuary health: Ecological status, threats and options for the future. *African Journal of Aquatic Science*, *45*(1–2), 65–82. https://doi.org/10.2989/16085914.2019.1672518

Cohn, J. L., Copp Franz, S., Mandel, R. H., Nack, C. C., Brainard, A. S., Eallonardo, A., & Magar, V. (2022). Strategies to work towards long-term sustainability and resiliency of nature-based solutions in coastal environments: A review and case studies. *Integrated Environmental Assessment and Management*, *18*(1), 123–134. https://doi.org/10.1002/ieam.4484

Colenbrander, D. (2019). Dissonant discourses: Revealing South Africa's policy-to-praxis challenges in the governance of coastal risk and vulnerability. *Journal of Environmental Planning and Management*, *62*(10), 1782–1801. https://doi.org/10.1080/09640568.2018.1515067

Colenbrander, D., & Bavinck, M. (2017). Exploring the role of bureaucracy in the production of coastal risks, City of Cape Town, South Africa. *Ocean & Coastal Management*, *150*, 35–50. https://doi.org/10.1016/j.ocecoaman.2016.11.012

Colenbrander, D., Cartwright, A., & Taylor, A. (2015). Drawing a line in the sand: Managing coastal risks in the City of Cape Town. *South African Geographical Journal*, *97*(1), 1–17. https://doi.org/10.1080/03736245.2014.924865

Cooper, J. A. G. (1989). Fairweather versus flood sedimentation in Mhlanga Lagoon, Natal: Implications for environmental management. *South African Journal of Geology*, *92*, 279–294.

Cooper, J. A. G. (2001). Geomorphological variability among microtidal estuaries from the wave-dominated South African coast. *Geomorphology*, *40*(1–2), 99–122. https://doi.org/10.1016/S0169-555X(01)00039-3

Cooper, J. A. G., & Jackson, D. W. T. (2019). Coasts in Peril? A shoreline health perspective. *Frontiers in Earth Science*, *7*, 260. https://doi.org/10.3389/feart.2019.00260

Cooper, J. A. G., Mason, T. R., Reddering, J. S. V., & Illenberger, W. K. (1990). Geomorphological effects of catastrophic flooding on a small subtropical estuary. *Earth Surface Processes and Landforms*, *15*(1), 25–41. https://doi.org/10.1002/esp.3290150104

Corbella, S., & Stretch, D. D. (2012a). Decadal trends in beach morphology on the east coast of South Africa and likely causative factors. *Natural Hazards and Earth System Sciences*, *12*(8), 2515–2527. https://doi.org/10.5194/nhess-12-2515-2012

Corbella, S., & Stretch, D. D. (2012b). Shoreline recovery from storms on the east coast of Southern Africa. *Natural Hazards and Earth System Sciences*, *12*(1), 11–22. https://doi.org/10.5194/nhess-12-11-2012

Cowell, P. J., & Thom, B. G. (1994). Morphodynamics of coastal evolution. In R. W. G. Carter & C. D. Woodroffe (Eds.), *Coastal evolution* (pp. 33–86). Cambridge University Press.

Dardis, G. F., & Grindley, J. R. (1988). Coastal geomorphology. In B. P. Moon & G. F. Dardis (Eds.), *The geomorphology of Southern Africa* (pp. 141–174). Southern Book Publishers.

Davis, J., Whitfield, P., Szimanski, D., Golden, B. R., Whitbeck, M., Gailani, J., Herman, B., Tritinger, A., Dillon, S. C., & King, J. (2022). A framework for evaluating island restoration performance: A case study from the Chesapeake Bay. *Integrated Environmental Assessment and Management*, *18*(1), 42–48. https://doi.org/10.1002/ieam.4437

De Battisti, D. (2021). The resilience of coastal ecosystems: A functional trait-based perspective. *The Journal of Ecology*, *109*(9), 3133–3146. https://doi.org/10.1111/1365-2745.13641

De Jong, B., Keijsers, J. G. S., Riksen, M. J. P. M., Krol, J., & Slim, P. A. (2014). Soft engineering vs. a dynamic approach in coastal dune management: A case study on the North Sea barrier island of Ameland, the Netherlands. *Journal of Coastal Research*, *30*, 670–684.

Desportes, I., & Colenbrander, D. R. (2016). Navigating interests, navigating knowledge: Towards an inclusive setback delineation along Cape Town's coastline. *Habitat International*, *54*, 124–135. https://doi.org/10.1016/j.habitatint.2015.12.016

de Villiers, N. M., Harasti, D., Hodgson, A. N., & Claassens, L. (2021). A comparison of the fauna in eelgrass and erosion control structures in a warm temperate Southern African estuary. *Regional Studies in Marine Science*, *44*, 101757. https://doi.org/10.1016/j.rsma.2021.101757

de Villiers, A. J., van Rooyen, M. W., Theron, G. K., & van Rooyen, N. (1999). Vegetation diversity of the Brand-se-Baai coastal dune area, west coast, South Africa: A pre-mining benchmark survey for rehabilitation. *Land Degradation & Development*, *10*(3), 207–224. https://doi.org/10.1002/(SICI)1099-145X(199905/06)10:3<207:AID-LDR333>3.0.CO;2-0

DST. (2017). *South African risk and vulnerability Atlas, 2nd Ed.* Department of Science and Technology.

Dye, A. H. (1998). Dynamics of rocky intertidal communities: Analyses of long time series from South African Shores. *Estuarine and Coastal Shelf Science*, *46*(2), 287–305. https://doi.org/10.1006/ecss.1997.0251

Dye, A. H., McLachlan, A., & Wooldridge, T. (1981). The ecology of sandy beaches in Natal. *South African Journal of Zoology*, *18*(4), 200–209. https://doi.org/10.1080/02541858.1981.11447758

Eichmanns, C., Lechthaler, S., Zander, W., Pérez, M. V., Blum, H., Thorenz, F., & Schüttrumpf, H. (2021). Sand trapping fences as a nature-based solution for coastal protection: An international review with a focus on installations in Germany. *Environments*, *8*(12), 135. https://doi.org/10.3390/environments8120135

Emery, K. O., & Kuhn, G. G. (1982). Sea cliffs: Their processes, profiles, and classification. *Geological Society of America Bulletin*, *93*(7), 644–654. https://doi.org/10.1130/0016-7606(1982)93<644:SCTPPA>2.0.CO;2

Farrell, E., & Connolly, N. (2019). Historic and contemporary dune inventories to assess dune vulnerability to climate change impacts. *Irish Geography*, *52*, 1–21.

Fernandes, M., & Adams, J. (2016). Quantifying the loss of and changes in estuary habitats in the uMkhomazi and Mvoti estuaries, South Africa. *South African Journal of Botany*, *107*, 179–187. https://doi.org/10.1016/j.sajb.2016.04.009

Ferro-Azcona, H., Espinoza Tenorio, A., Calderón-Contreras, R., Ramenzoni, V. C., Gómez País, M. D., & Mesa-Jurado, M. A. (2019). Adaptive capacity and social-ecological resilience of coastal areas: A systematic review. *Ocean & Coastal Management*, *173*, 36–51. https://doi.org/10.1016/j.ocecoaman.2019.01.005

Floerl, O., Atalah, J., Bugnot, A. B., Chandler, M., Dafforn, K. A., Floerl, L., Zaiko, A., & Major, R. (2021). A global model to forecast coastal hardening and mitigate associated socioecological risks. *Nature Sustainability*, *4*(12), 1060–1067. https://doi.org/10.1038/s41893-021-00780-w

Flood, S., & Schechtman, J. (2014). The rise of resilience: Evolution of a new concept in coastal planning in Ireland and the US. *Ocean & Coastal Management*, *102*, 19–31. https://doi.org/10.1016/j.ocecoaman.2014.08.015

Gaillard, J. C., Maceda, E. A., Stasiak, E., Le Berre, I., & Espaldon, M. V. O. (2009). Sustainable livelihoods and people's vulnerability in the face of coastal hazards. *Journal of Coastal Conservation*, *13*(2–3), 119–129. https://doi.org/10.1007/s11852-009-0054-y

Garden, S. E., & Garland, G. G. (2005). Spit development in the Mdloti River estuary, KwaZulu-Natal. *South African Journal of Geology*, *108*(2), 257–270. https://doi.org/10.2113/108.2.257

Garland, G., & Moleko, L. (2000). Geomorphological impacts of Inanda Dam on the Mgeni estuary, north of Durban, South Africa). *Bulletin of Engineering Geology and the Environment*, *59*(2), 119–126. https://doi.org/10.1007/s100640000057

Gedan, K. B., Kirwan, M. L., Wolanski, E., Barbier, E. B., & Silliman, B. R. (2011). The present and future role of coastal wetland vegetation in protecting shorelines: Answering recent challenges to the paradigm. *Climatic Change*, *106*(1), 7–29. https://doi.org/10.1007/s10584-010-0003-7

Gijsman, R., Horstman, E. M., van der Wal, D., Friess, D. A., Swales, A., & Wijnberg, K. M. (2021). Nature-based engineering: A review on reducing coastal flood risk with Mangroves. *Frontiers in Marine Science*, *8*, 702412. https://doi.org/10.3389/fmars.2021.702412

Glaser, M., & Glaeser, B. (2014). Towards a framework for cross-scale and multi-level analysis of coastal and marine social-ecological systems dynamics. *Regional Environmental Change*, *14*(6), 2039–2052. https://doi.org/10.1007/s10113-014-0637-5

Glavovic, B. C., Limburg, K., Liu, K. -K., Emeis, K. -C., Thomas, H., Kremer, H., Avril, B., Zhang, J., Mulholland, M. R., Glaser, M., & Swaney, D. P. (2015). Living on the margin in the Anthropocene: Engagement arenas for sustainability research and action at the ocean–land interface. *Current Opinions in Environmental Sustainability*, *14*, 232–238. https://doi.org/10.1016/j.cosust.2015.06.003

Goble, B. J., Hill, T. R., & Phillips, M. R. (2017). An assessment of integrated coastal management governance and implementation using the DPSIR Framework: KwaZulu-Natal, South Africa. *Coastal Management*, *45*(2), 107–124. https://doi.org/10.1080/08920753.2017.1278144

Goble, B. J., Hill, T. R., & Phillips, M. R. (2020). Building institutional knowledge for effective integrated coastal management decision making: KwaZulu-Natal, South Africa. *Journal of Coastal Research*, *95*(sp1), 808–813. https://doi.org/10.2112/SI95-157.1

Goble, B. J., MacKay, C. F., & Hill, T. R. (2019). Design, development and implementation of a decision support info-portal for integrated coastal management, KwaZulu-Natal, South Africa. *Environmental Management*, *64*(1), 27–39. https://doi.org/10.1007/s00267-019-01173-8

Goble, B. J., van der Elst, R. P., & Oellermann, L. K. (Eds.). (2014). *Ugu Lwethu – Our Coast. A profile of coastal KwaZulu-Natal*. KwaZulu-Natal Department of Agriculture and Environmental Affairs and the Oceanographic Research Institute.

Gowell, M. R., Coombes, M. A., & Viles, H. A. (2015). Rock-protecting seaweed? Experimental evidence of bioprotection in the intertidal zone. *Earth Surface Processes and Landforms*, *40*(10), 1364–1370. https://doi.org/10.1002/esp.3736

Green, A. N., Cooper, J. A. G., & LeVieux, A. (2013). Unusual barrier/inlet behaviour associated with active coastal progradation and river-dominated estuaries. *Journal of Coastal Research*, *SI69*, 35–45. https://doi.org/10.2112/SI_69_4

Griggs, G., & Patsch, K. (2019). The protection/hardening of California's coast: Times are changing. *Journal of Coastal Research*, *35*(5), 1051–1061. https://doi.org/10.2112/JCOASTRES-D-19A-00007.1

Guastella, L. A., & Smith, A. M. (2014). Coastal dynamics on a soft coastline from serendipitous webcams: KwaZulu-Natal, South Africa. *Estuarine, Coastal and Shelf Science*, *150*, 76–85. https://doi.org/10.1016/j.ecss.2013.12.009

Guastella, L. A., Smith, A. M., & Breetzke, T. (2014). Coastal management and mis-management: Comparing successes and failures at two lagoon outlets in KwaZulu-Natal, South Africa. *Journal of Coastal Research*, *SI70*, 513–520. https://doi.org/10.2112/SI70-087.1

Hamin, E. M., Abunnasr, Y., Dilthey, M. R., Judge, P. K., Kenney, M. A., Kirshen, P., Sheahan, T. C., DeGroot, D. J., Ryan, R. L., McAdoo, B. G., Nurse, L., Buxton, J. A., Sutton-Grier, A. E., Albright, E. A., Marin, M. A., & Fricke, R. (2018). Pathways to coastal resiliency: The adaptive gradients framework. *Sustainability*, *10*(8), 2629. https://doi.org/10.3390/su10082629

Hanley, M. E., Hoggart, S. P. G., Simmonds, D. J., Bichot, A., Colangelo, M. A., Bozzeda, F., Heurtefeux, H., Ondiviela, B., Ostrowski, R., Recio, M., Trude, R., Zawadzka-Kahlau, E., & Thompson, R. C. (2014). Shifting sands? Coastal protection by sand banks, beaches and dunes. *Coastal Engineering*, *87*, 136–146. https://doi.org/10.1016/j.coastaleng.2013.10.020

Hapke, C. J., Kratzmann, M. G., & Himmelstoss, E. A. (2013). Geomorphic and human influence on large-scale coastal change. *Geomorphology*, *199*, 160–170. https://doi.org/10.1016/j.geomorph.2012.11.025

Harris, L., Holness, S., Nel, R., Lombard, A. T., & Schoeman, D. (2013). Intertidal habitat composition and regional-scale shoreline morphology along the Benguela coast. *Journal of Coastal Conservation*, *17*(1), 143–154. https://doi.org/10.1007/s11852-012-0226-z

Harris, L., Nel, R., & Schoeman, D. (2011a). Mapping beach morphodynamics remotely: A novel application tested on South African sandy shores. *Estuarine and Coastal Shelf Science*, *92*(1), 78–89. https://doi.org/10.1016/j.ecss.2010.12.013

Harris, L., Nel, R., Smale, M., & Schoeman, D. (2011b). Swashed away? Storm impacts on sandy beach macrofaunal communities. *Estuarine, Coastal and Shelf Science*, *94*(3), 210–221. https://doi.org/10.1016/j.ecss.2011.06.013

Hayes, M., & Kirkpatrick, J. B. (2012). Influence of *Ammophila arenaria* on half a century of vegetation change in eastern Tasmanian sand dune systems. *Australian Journal of Botany*, *60*(5), 450–460. https://doi.org/10.1071/BT12071

Hellström, G. B. (1996). Preliminary investigations into recent changes of the goukamma nature reserve frontal dune system, South Africa — with management implications. *Landscape and Urban Planning*, *34*(3–4), 225–235. https://doi.org/10.1016/0169-2046(95)00223-5

Hendey, Q. B., & Volman, T. P. (1986). Last interglacial sea levels and coastal caves in the Cape Province, South Africa. *Quaternary Research*, *25*(2), 189–198. https://doi.org/10.1016/0033-5894(86)90056-6

Henrico, I., Ledwaba, T., & van Zyl, G. (2020). Measuring the effect of wind-driven processes on coastal dunes: A study of the Atlantis and Geelbek dune fields along the West Coast of South Africa. *Spatial Information Research*, *28*(5), 569–577. https://doi.org/10.1007/s41324-020-00317-x

Hertling, U. M., & Lubke, R. A. (1999). Use of *Ammophila arenaria* for dune stabilization in South Africa and its current distribution?Perceptions and problems. *Environmental Management*, *24* (4), 467–482. https://doi.org/10.1007/s002679900247

Holmes, P. J., Grab, S. W., & Knight, J. (2016). South African geomorphology: Current status and new challenges. *South African Geographical Journal*, *98*(3), 405–416. https://doi.org/10.1080/03736245.2016.1208581

Hoppe-Speer, S. C. L., Adams, J. B., & Bailey, D. (2015). Present state of mangrove forests along the Eastern Cape coast, South Africa. *Wetlands Ecology and Management*, *23*(3), 371–383. https://doi.org/10.1007/s11273-014-9387-x

Hoppe-Speer, S. C. L., Adams, J. B., & Rajkaran, A. (2013). Response of mangroves to drought and non-tidal conditions in St Lucia Estuary, South Africa. *African Journal of Aquatic Science*, *38*(2), 153–162. https://doi.org/10.2989/16085914.2012.759095

Illgner, P. M. (2008). Coastal processes and associated landforms. In C. A. Lewis (Ed.), *Geomorphology of the Eastern Cape* (2nd ed., pp. 63–84). NISC.

IPCC. (2014). Annex II: Glossary. In Pachauri, R. K., & Meyer, L. A. (Eds.), *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 117–130).

Itzkin, M., Moore, L. J., Ruggiero, P., & Hacker, S. D. (2020). The effect of sand fencing on the morphology of natural dune systems. *Geomorphology*, *352*, 106995. https://doi.org/10.1016/j.geomorph.2019.106995

Jackson, D. W. T., Cooper, J. A. G., & Green, A. N. (2014). A preliminary classification of coastal sand dunes of KwaZulu-Natal. *Journal of Coastal Research*, *SI70*, 718–722. https://doi.org/10.2112/SI70-121.1

Jackson, D. W. T., Costas, S., González-Villanueva, R., & Cooper, A. (2019). A global 'greening' of coastal dunes: An integrated consequence of climate change? *Global and Planetary Change*, *182*, 103026. https://doi.org/10.1016/j.gloplacha.2019.103026

Jacobs, Z., Wintle, A. G., & Duller, G. A. T. (2003). Optical dating of dune sand from Blombos Cave, South Africa: I—multiple grain data. *Journal of Human Evolution*, *44*(5), 599–612. https://doi.org/10.1016/S0047-2484(03)00048-4

Jayanthi, M., Thirumurthy, S., Samynathan, M., Duraisamy, M., Muralidhar, M., Ashokkumar, J., & Vijayan, K. K. (2018). Shoreline change and potential sea level rise impacts in a climate hazardous location in southeast coast of India. *Environmental Monitoring and Assessment*, *190* (1), 51. https://doi.org/10.1007/s10661-017-6426-0

Kazemi, K., Mlucasse, R., & Santha, R. (2017). Assessing Mozambique's exposure to coastal climate hazards and erosion. *International Journal of Disaster Risk Reduction*, *23*, 45–52.

Keesstra, S., Nunes, J., Novara, A., Finger, D., Avelar, D., Kalantari, Z., & Cerdà, A. (2018). The superior effect of nature based solutions in land management for enhancing ecosystem services. *The Science of the Total Environment*, *610–611*, 997–1009. https://doi.org/10.1016/j.scitotenv.2017.08.077

Kesavan, P. C., & Swaminathan, M. S. (2006). Managing extreme natural disasters in coastal areas. *Philosophical Transactions of the Royal Society A – Mathematical Physical and Engineering Sciences*, *364*(1845), 2191–2216. https://doi.org/10.1098/rsta.2006.1822

Kim, M., Song, K., & Chon, J. (2021). Key coastal landscape patterns for reducing flood vulnerability. *The Science of the Total Environment*, *759*, 143454. https://doi.org/10.1016/j.scitotenv.2020.143454

King, J., Holmes, R., Burkholder, S., Holzman, J., & Suedel, B. (2021). Advancing nature-based solutions by leveraging engineering with Nature® strategies and landscape architectural practices in highly collaborative settings. *Integrated Environmental Assessment and Management*, *18* (1), 108–114. https://doi.org/10.1002/ieam.4473

Kittinger, J. N., & Ayers, A. L. (2010). Shoreline armoring, risk management, and coastal resilience under rising seas. *Coastal Management*, *38*(6), 634–653. https://doi.org/10.1080/08920753.2010.529038

Knight, J. (2021a). Environmental services: A new approach towards addressing sustainable development goals in sub-Saharan Africa. *Frontiers in Sustainable Food Systems*, *5*, 687863. https://doi.org/10.3389/fsufs.2021.687863

Knight, J. (2021b). The late Quaternary stratigraphy of coastal dunes and associated deposits in South Africa. *South African Journal of Geology*, *124*(4), 995–1006. https://doi.org/10.25131/sajg.124.0032

Knight, J. (2022a). Presence, absence, transience: The spatiotemporalities of sand. *Geographies*, *2*(4), 657–668. https://doi.org/10.3390/geographies2040040

Knight, J. (2022b). Scientists' warning of the impacts of climate change on mountains. *PeerJ*, *10*, e14253. https://doi.org/10.7717/peerj.14253

Knight, J., & Burningham, H. (2019). Sand dunes and ventifacts on the coast of South Africa. *Aeolian Research*, *37*, 44–58. https://doi.org/10.1016/j.aeolia.2019.02.003

Knight, J., & Burningham, H. (2020a). Bedrock hardness values and morphological zonation of a shore platform in South Africa. *Transactions of the Royal Society of South Africa*, *75*(1), 40–53. https://doi.org/10.1080/0035919X.2019.1670751

Knight, J., & Burningham, H. (2020b). What controls bedrock shore platform hardness? A field study from South Africa. *Journal of Coastal Research*, *95*(sp1), 537–541. https://doi.org/10.2112/SI95-104.1

Knight, J., & Burningham, H. (2021). The morphodynamics of transverse dunes on the coast of South Africa. *Geo-Marine Letters*, *41*(4), 47. https://doi.org/10.1007/s00367-021-00717-4

Knight, J., & Burningham, H. (2022). A morphological classification of coastal forelands, with examples from South Africa. *Geomorphology*, *415*, 108410. https://doi.org/10.1016/j.geomorph.2022.108410

Knight, J., & Goff, J. R. (2016). Coastal science for post-tsunami reconstruction. *The Holocene*, *26*(8), 1334–1340. https://doi.org/10.1177/0959683616638438

Knight, J., & Grab, S. W. (2015). Rocky Shorelines of the Wild Coast. In S. W. Grab & J. Knight (Eds.), *Landforms and Landscapes of South Africa* (pp. 55–61). Springer.

Knight, J., & Harrison, S. (2009). Sediments and future climate. *Nature Geoscience*, *2*(4), 230. https://doi.org/10.1038/ngeo491

Kombiadou, K., Costas, S., Carrasco, A. R., Plomaritis, T. A., Ferreira, Ó., & Matias, A. (2019). Bridging the gap between resilience and geomorphology of complex coastal systems. *Earth-Science Reviews*, *198*, 102934. https://doi.org/10.1016/j.earscirev.2019.102934

Kron, W. (2013). Coasts: The high-risk areas of the world. *Natural Hazards*, *66*(3), 1363–1382. https://doi.org/10.1007/s11069-012-0215-4

La Cock, G. D., & Burkinshaw, J. R. (1996). Management implications of development resulting in disruption of a headland bypass dunefield and its associated river, Cape St Francis, South Africa. *Landscape and Urban Planning*, *34*(3–4), 373–381. https://doi.org/10.1016/0169-2046(95)00226-X

La Cock, G. D., Lubke, R. A., & Wilken, M. (1992). Dune movement in the Kwaaihoek region of the Eastern Cape, South Africa, and its bearing on future developments of the region. *Journal of Coastal Research*, *8*, 210–217.

Leuci, R., Wiles, E., Thackeray, Z., & Vella, G. (2022). Trends in sandy beach variability EThekwini Municipality, South Africa. *Journal of Sea Research*, *179*, 102149. https://doi.org/10.1016/j.seares.2021.102149

Lithgow, D., Martínez, M. L., Gallego-Fernández, J. B., Hesp, P. A., Flores, P., Gachuz, S., Rodríguez-Revelo, N., Jiménez-Orocio, O., Mendoza-González, G., & Álvarez-Molina, L. L. (2013). Linking restoration ecology with coastal dune restoration. *Geomorphology*, *199*, 214–224. https://doi.org/10.1016/j.geomorph.2013.05.007

Liu, Z., Fagherazzi, S., & Cui, B. (2021). Success of coastal wetlands restoration is driven by sediment availability. *Communications Earth & Environment*, *2*(1), 44. https://doi.org/10.1038/s43247-021-00117-7

Lloyd, M. G., Peel, D., & Duck, R. W. (2013). Towards a social–ecological resilience framework for coastal planning. *Land Use Policy*, *30*(1), 925–933. https://doi.org/10.1016/j.landusepol.2012.06.012

Lubke, R. A. (2022). Current state of *Ammophila arenaria* (Marram Grass) distribution in the Eastern Cape, South Africa, and the possible effect of the grass on the dune system dynamics. *Plants*, *11*(17), 2260. https://doi.org/10.3390/plants11172260

Lubke, R. A., Avis, A. M., & Moll, J. B. (1996). Post-mining rehabilitation of coastal sand dunes in Zululand, South Africa. *Landscape and Urban Planning*, *34*(3–4), 335–345. https://doi.org/10.1016/0169-2046(95)00227-8

Lubke, R. A., & Sugden, J. (1990). Short-term changes in mobile dunes at Port Alfred, South Africa. *Environmental Management*, *14*(2), 209–220. https://doi.org/10.1007/BF02394038

Lubke, R. A., & Webb, C. (2016). The interaction between the dunes systems and the lower estuary at the Bushmans River Mouth, Eastern Cape over the past 60 years. *South African Journal of Botany*, *107*, 148–159. https://doi.org/10.1016/j.sajb.2016.05.020

Lucrezi, S., Saayman, M., & van der Merwe, P. (2014). Influence of infrastructure development on the vegetation community structure of coastal dunes: Jeffreys Bay, South Africa. *Journal of Coastal Conservation*, *18*, 193–211. https://doi.org/10.1007/s11852-014-0307-2

Luijendijk, A., Hagenaars, G., Ranasinghe, R., Baart, F., Donchyts, G., & Aarninkhof, S. (2018). The State of the World's Beaches. *Scientific Reports*, *8*(1), 6641. https://doi.org/10.1038/s41598-018-24630-6

Magoro, M. L., Perissinotto, R., Wooldridge, T. H., & Whitfield, A. K. (2020). Micro-estuaries and micro-outlets as incipient estuarine systems – Does size and coastal connectivity count? *The Science of the Total Environment*, *703*, 134707. https://doi.org/10.1016/j.scitotenv.2019.134707

Malan, D. E., & Swart, D. H. (1997). South African integrated coastal management and engineering: Converging towards collision or collaboration? *Transactions of the Royal Society of South Africa*, *52*(1), 227–252. https://doi.org/10.1080/00359199709520622

Malherbe, H., & Samways, M. (2014). Rocky shores of a major southern African Marine Protected Area are almost free from intertidal invertebrate alien species. *Koedoe*, *56*(1), 1206. https://doi.org/10.4102/koedoe.v56i1.1206

Marker, M. E. (2000). A descriptive account of sand movement in the Knysna Estuary. *Transactions of the Royal Society of South Africa*, *55*(2), 129–139. https://doi.org/10.1080/00359190009520439

Marker, M. E. (2004). Changes in the Salt River, tributary to the Knysna Estuary, South Africa: 1936–1998. *South African Geographical Journal*, *86*(2), 131–140. https://doi.org/10.1080/03736245.2004.9713817

Martínez, M. L., Taramelli, A., & Silva, R. (2017). Resistance and resilience: Facing the multi-dimensional challenges in coastal areas. *Journal of Coastal Research*, *SI77*, 1–6. https://doi.org/10.2112/SI77-001.1

Masselink, G., & Lazarus, E. D. (2019). Defining Coastal Resilience. *Water*, *11*(12), 2587. https://doi.org/10.3390/w11122587

Mather, A. A., & Stretch, D. D. (2012). A perspective on sea level rise and coastal storm surge from Southern and Eastern Africa: A Case Study Near Durban, South Africa. *Water*, *4*(1), 237–259. https://doi.org/10.3390/w4010237

Mbopha, M. S., Marais, C., Kleynhans, T. E., & Esler, K. J. (2021). Unlocking and securing ecological infrastructure investments: The needs and willingness to invest and institutional support mechanisms used. *South African Journal of Science*, *117*(9/10), 8666. https://doi.org/10.17159/sajs.2021/8666

McFadden, L. (2010). Exploring system interactions for building resilience within coastal environments and communities. *Environmental Hazards*, *9*(3), 266–283. https://doi.org/10.3763/ehaz.2010.0056

Mead, A., Griffiths, C. L., Branch, G. M., McQuaid, C. D., Blamey, L. K., Bolton, J. J., Anderson, R. J., Dufois, F., Rouault, M., Froneman, P. W., Whitfield, A. K., Harris, L. R., Nel, R., Pillay, D., & Adams, J. B. (2013). Human-mediated drivers of change — impacts on

coastal ecosystems and marine biota of South Africa. *African Journal of Marine Science*, *35*(3), 403–425. https://doi.org/10.2989/1814232X.2013.830147

Mentis, M. T., & Ellery, W. N. (1994). Post-mining rehabilitation of dunes on the north-east coast of South Africa. *South African Journal of Science*, *90*, 69–74.

Mentis, M. T., & Ellery, W. N. (1998). Environmental effects of mining coastal dunes: Conjectures and refutations. *South African Journal of Science*, *94*, 215–222.

Mitchell, J., Jury, M. R., & Mulder, G. J. (2005). A study of Maputaland beach dynamics. *South African Geographical Journal*, *87*(1), 43–51. https://doi.org/10.1080/03736245.2005.9713825

Monteiro, R., & Ferreira, J. C. (2020). Green infrastructure planning as a climate change and risk adaptation tool in Coastal Urban Areas. *Journal of Coastal Research*, *95*(sp1), 889–893. https://doi.org/10.2112/SI95-173.1

Morris, R. L., Strain, E. M. A., Konlechner, T. M., Fest, B. J., Kennedy, D. M., Arndt, S. K., & Swearer, S. E. (2019). Developing a nature-based coastal defence strategy for Australia. *Australian Journal of Civil Engineering*, *17*(2), 167–176. https://doi.org/10.1080/14488353.2019.1661062

Mullick, M. R. A., Islam, K. M. A., & Tanim, A. H. (2020). Shoreline change assessment using geospatial tools: A study on the Ganges deltaic coast of Bangladesh. *Earth Science Informatics*, *13*(2), 299–316. https://doi.org/10.1007/s12145-019-00423-x

Musekiwa, C., Cawthra, H. C., Unterner, M., & van Zyl, F. W. (2015). An assessment of coastal vulnerability for the South African coast. *South African Journal of Geomatics*, *4*(2), 123–137. https://doi.org/10.4314/sajg.v4i2.5

Naylor, L. A., & Stephenson, W. J. (2010). On the role of discontinuities in mediating shore platform erosion. *Geomorphology*, *114*(1–2), 89–100. https://doi.org/10.1016/j.geomorph.2008.12.024

Nichols, G. R. (1996). Preliminary observations on managing and reclaiming frontal dunes with the Durban municipal area. *Landscape and Urban Planning*, *34*(3–4), 383–388. https://doi.org/10.1016/0169-2046(95)00230-8

Nichols, C. R., Wright, L. D., Bainbridge, S. J., Cosby, A., Hénaff, A., Loftis, J. D., Cocquempot, L., Katragadda, S., Mendez, G. R., Letortu, P., Le Dantec, N., Resio, D., & Zarillo, G. (2019). Collaborative science to enhance coastal resilience and adaptation. *Frontiers in Marine Science*, *6*, 404. https://doi.org/10.3389/fmars.2019.00404

Olivier, M. J., & Garland, G. G. (2003). Short-term monitoring of foredune formation on the east coast of South Africa. *Earth Surface Processes and Landforms*, *28*(10), 1143–1155. https://doi.org/10.1002/esp.549

Orchard, S., & Schiel, D. R. (2021). Enabling nature-based solutions for climate change on a peri-urban sandspit in Christchurch, New Zealand. *Regional Environmental Change*, *21*(3), 66. https://doi.org/10.1007/s10113-021-01791-1

Ott, T., & van Aarde, R. J. (2014). Coastal dune topography as a determinant of abiotic conditions and biological community restoration in northern KwaZulu-Natal, South Africa. *Landscape and Ecological Engineering*, *10*(1), 17–28. https://doi.org/10.1007/s11355-013-0211-1

Palmer, B. J., McGregor, G. K., Hill, T. R., & Paterson, A. W. (2010). A spatial assessment of coastal development and land use change in the Eastern Cape, South Africa. *South African Geographical Journal*, *92*, 117–128.

Palmer, B. J., Van der Elst, R., Mackay, F., Mather, A. A., Smith, A. M., Bundy, S. C., Thackeray, Z., Leuci, R., & Parak, O. (2011). Preliminary coastal vulnerability assessment for KwaZulu-Natal, South Africa. *Journal of Coastal Research*, *SI64*, 1390–1395.

Passeri, D. L., Bilskie, M. V., Hagen, S. C., Mickey, R. C., Dalyander, P. S., & Gonzalez, V. M. (2021). Assessing the effectiveness of nourishment in Decadal Barrier Island Morphological Resilience. *Water*, *13*(7), 944. https://doi.org/10.3390/w13070944

Payo, A., Hall, J. W., Dickson, M. E., & Walkden, M. J. A. (2015). Feedback structure of cliff and shore platform morphodynamics. *Journal of Coast Conservation*, *19*(6), 847–859. https://doi.org/10.1007/s11852-014-0342-z

Peer, N., Rajkaran, A., Miranda, N. A. F., Taylor, R. H., Newman, B., Porri, F., Raw, J. L., Mbense, S. P., Adams, J. B., & Perissinotto, R. (2018). Latitudinal gradients and poleward

expansion of mangrove ecosystems in South Africa: 50 years after Macnae's first assessment. *African Journal of Marine Science*, *40*(2), 101–120. https://doi.org/10.2989/1814232X.2018.1466728

Pickering, R., Jacobs, Z., Herries, A. I. R., Karkanas, P., Bar-Matthews, M., Woodhead, J. D., Kappen, P., Fisher, E., & Marean, C. W. (2013). Paleoanthropologically significant South African sea caves dated to 1.1–1.0 million years using a combination of U–Pb, TT-OSL and palaeomagnetism. *Quaternary Science Reviews*, *65*, 39–52. https://doi.org/10.1016/j.quascirev.2012.12.016

Pile, J., Gouramanis, C., Switzer, A. D., Rush, B., Reynolds, I., & Soria, J. L. A. (2018). Can the risk of coastal hazards be better communicated? *International Journal of Disaster Risk Reduction*, *27*, 439–450. https://doi.org/10.1016/j.ijdrr.2017.11.008

Pillay, S., Lutchmiah, J., Gengan, R. M., & Lindsay, P. (2003). Morphology and sediment dynamics of the ephemeral Mfolozi estuary, KwaZulu-Natal, South Africa. *South African Geographical Journal*, *85*(2), 158–163. https://doi.org/10.1080/03736245.2003.9713796

Poe, M. R., Norman, K. C., & Levin, P. S. (2014). Cultural dimensions of socioecological systems: key connections and guiding principles for conservation in coastal environments. *Conservation Letters*, *7*(3), 166–175. https://doi.org/10.1111/conl.12068

Pontee, N., Narayan, S., Beck, M. W., & Hosking, A. H. (2016). Nature-based solutions: Lessons from around the world. *Maritime Engineering*, *169*(1), 29–36. https://doi.org/10.1680/jmaen.15.00027

Rajkaran, A., Adams, J., & Taylor, R. (2009). Historic and recent (2006) state of mangroves in small estuaries from Mlalazi to Mtamvuna in KwaZulu-Natal, South Africa. *Southern Forests: A Journal of Forest Science*, *71*(4), 287–296. https://doi.org/10.2989/SF.2009.71.4.6.1033

Ramesh, R., Chen, Z., Cummins, V., Day, J., D'elia, C., Dennison, B., Forbes, D. L., Glaeser, B., Glaser, M., Glavovic, B., Kremer, H., Lange, M., Larsen, J. N., Le Tissier, M., Newton, A., Pelling, M., Purvaja, R., & Wolanski, E. (2015). Land–ocean interactions in the coastal zone: Past, present & future. *Anthropocene*, *12*, 85–98. https://doi.org/10.1016/j.ancene.2016.01.005

Ranasinghe, R., & Stive, M. J. F. (2009). Rising seas and retreating coastlines. *Climatic Change*, *97*(3–4), 465–468. https://doi.org/10.1007/s10584-009-9593-3

Rautenbach, C., & Theron, A. K. (2018). Study of the Durban Bight shoreline evolution under schematised climate change and sand-bypassing scenarios. *Journal of the South African Institution of Civil Engineering*, *60*(4), 2–15. https://doi.org/10.17159/2309-8775/2018/v60n4a1

Ruckelshaus, M. H., Guannel, G., Arkema, K., Verutes, G., Griffin, R., Guerry, A., Silver, J., Faries, J., Brenner, J., & Rosenthal, A. (2016). Evaluating the benefits of green infrastructure for Coastal Areas: Location, Location, Location. *Coastal Management*, *44*(5), 504–516. https://doi.org/10.1080/08920753.2016.1208882

Ruckelshaus, M., Reguero, B. G., Arkema, K., Guerrero Compéan, R., Weekes, K., Bailey, A., & Silver, J. (2020). Harnessing new data technologies for nature-based solutions in assessing and managing risk in coastal zones. *International Journal of Disaster Risk Reduction*, *51*, 101795. https://doi.org/10.1016/j.ijdrr.2020.101795

Saja, A. M. A., Teo, M., Goonetilleke, A., & Ziyath, A. M. (2021). A critical review of social resilience properties and pathways in disaster management. *International Journal of Disaster Risk Science*, *12*(6), 790–804. https://doi.org/10.1007/s13753-021-00378-y

Sakho, I., Mesnage, V., Deloffre, J., Lafite, R., Niang, I., & Faye, G. (2011). The influence of natural and anthropogenic factors on mangrove dynamics over 60 years: The Somone Estuary, Senegal. *Estuarine and Coastal Shelf Science*, *94*(1), 93–101. https://doi.org/10.1016/j.ecss.2011.05.032

Saleh, F., & Weinstein, M. P. (2016). The role of nature-based infrastructure (NBI) in coastal resiliency planning: A literature review. *Journal of Environmental Management*, *183*, 1088–1098. https://doi.org/10.1016/j.jenvman.2016.09.077

Schoonees, T., Gijón Mancheño, A., Scheres, B., Bouma, T. J., Silva, R., Schlurmann, T., & Schüttrumpf, H. (2019). Hard structures for coastal protection, towards greener designs. *Estuaries & Coasts*, *42*(7), 1709–1729. https://doi.org/10.1007/s12237-019-00551-z

Schultz, M. T., & Smith, E. R. (2016). Assessing the resilience of coastal systems: A probabilistic approach. *Journal of Coastal Research*, *32*, 1032–1050. https://doi.org/10.2112/JCOASTRES-D-15-00170.1

Schumann, E. H. (2021). Floods, sedimentation and tidal exchanges in the Keurbooms Estuary, South Africa. *Geo-Marine Letters*, *41*(3), 34. https://doi.org/10.1007/s00367-021-00709-4

Shaw, A., Holmes, P., & Rogers, J. (2001). Depositional landforms and environmental change in the headward vicinity of Dias Beach, Cape Point. *South African Journal of Geology*, *104*(2), 101–114. https://doi.org/10.2113/1040101

Silva, R., Chávez, V., Bouma, T. J., van Tussenbroek, B. I., Arkema, K. K., Martínez, M. L., Oumeraci, H., Heymans, J. J., Osorio, A. F., Mendoza, E., Mancuso, M., Asmus, M., & Pereira, P. (2019). The incorporation of biophysical and social components in Coastal Management. *Estuaries & Coasts*, *42*(7), 1695–1708. https://doi.org/10.1007/s12237-019-00559-5

Sink, K. J., Branch, G. M., & Harris, J. M. (2005). Biogeographic patterns in rocky intertidal communities in KwaZulu-Natal, South Africa. *African Journal of Marine Science*, *27*(1), 81–96. https://doi.org/10.2989/18142320509504070

Smith, A., Guastella, L. A., Mather, A. A., Bundy, S. C., & Haigh, I. D. (2013). KwaZulu-Natal coastal erosion events of 2006/2007 and 2011: A predictive tool? *South African Journal of Science*, *109*(3/4), 0025. https://doi.org/10.1590/sajs.2013/20120025

Smith, A. M., Mather, A. A., Bundy, S. C., Cooper, J. A. G., Guastella, L. A., Ramsay, P. J., & Theron, A. (2010). Contrasting styles of swell-driven coastal erosion: Examples from KwaZulu-Natal, South Africa. *Geological Magazine*, *147*(6), 940–953. https://doi.org/10.1017/S0016756810000361

Sohn, W., Bae, J., & Newman, G. (2021). Green infrastructure for coastal flood protection: The longitudinal impacts of green infrastructure patterns on flood damage. *Applied Geography*, *135*, 102565. https://doi.org/10.1016/j.apgeog.2021.102565

Spalding, M. D., Ruffo, S., Lacambra, C., Meliane, I., Hale, L. Z., Shepard, C. C., & Beck, M. W. (2014). The role of ecosystems in coastal protection: Adapting to climate change and coastal hazards. *Ocean & Coastal Management*, *90*, 50–57. https://doi.org/10.1016/j.ocecoaman.2013.09.007

Steinke, T. (1999). Mangroves in South African estuaries. In B. Allanson & D. Baird (Eds.), *Estuaries of South Africa* (pp. 119–140). Cambridge University Press.

Stephenson, W. J., Kirk, R. M., Hemmingsen, S. A., & Hemmingsen, M. A. (2010). Decadal scale micro erosion rates on shore platforms. *Geomorphology*, *114*(1–2), 22–29. https://doi.org/10.1016/j.geomorph.2008.10.013

Sutton-Grier, A. E., Gittman, R. K., Arkema, K. K., Bennett, R. O., Benoit, J., Blitch, S., Burks-Copes, K. A., Colden, A., Dausman, A., DeAngelis, B. M., Hughes, A. R., Scyphers, S. B., & Grabowski, J. H. (2018). Investing in Natural and Nature-Based Infrastructure: Building Better Along Our Coasts. *Sustainability*, *10*, 523. https://doi.org/10.3390/su10020523

Sutton-Grier, A. E., Wowk, K., & Bamford, H. (2018). Future of our coasts: The potential for natural and hybrid infrastructure to enhance the resilience of our coastal communities, economies and ecosystems. *Environmental Science & Policy*, *51*, 137–148. https://doi.org/10.1016/j.envsci.2015.04.006

Taljaard, S., van Niekerk, L., & Weerts, S. P. (2019). The legal landscape governing South Africa's coastal marine environment – Helping with the 'horrendogram'. *Ocean & Coastal Management*, *178*, 104801. https://doi.org/10.1016/j.ocecoaman.2019.05.003

Taylor, R., Adams, J. B., & Haldorsen, S. (2006). Primary habitats of the St Lucia Estuarine System, South Africa, and their responses to mouth management. *African Journal of Aquatic Science*, *31*(1), 31–41. https://doi.org/10.2989/16085910609503869

Tejada, M., Malvarez, G. C., & Navas, F. (2007). A New Environmental Indicator for Coastal Artificialisation and Resilience Mapping. *Journal of Coastal Research*, *SI50*, 67–71.

Tinley, K. L. (1985). *Coastal Dunes of South Africa*. CSIR, Pretoria.

Trégarot, E., Catry, T., Pottier, A., El-Hacen, E. -H.M., Cheikh, M. A. S., Cornet, C. C., Maréchal, J. P., & Failler, P. (2021). Coastal protection assessment: A tradeoff between ecological, social, and economic issues. *Ecosphere*, *12*(2), e03364. https://doi.org/10.1002/ecs2.33641

Tucker, L., Griffiths, C. L., Schroeter, F., & Vetter, H. D. (2017). Boulder shores in South Africa – a distinct but poorly documented coastal habitat type. *African Journal of Marine Science*, *39*(2), 193–202. https://doi.org/10.2989/1814232X.2017.1329167

Van Aarde, R. J., Ferreira, S. M., & Kritzinger, J. J. (1996). Successional changes in rehabilitating coastal dune communities in northern KwaZulu/Natal, South Africa. *Landscape and Urban Planning*, *34*(3–4), 277–286. https://doi.org/10.1016/0169-2046(95)00237-5

Van Aarde, R. J., Smit, A. -M., & Claassens, A. S. (1998). Soil Characteristics of Rehabilitating and Unmined Coastal Dunes at Richards Bay, KwaZulu-Natal, South Africa. *Restoration Ecology*, *6*(1), 102–110. https://doi.org/10.1046/j.1526-100x.1998.06113.x

Van Coppenolle, R., & Temmerman, S. (2019). A global exploration of tidal wetland creation for nature-based flood risk mitigation in coastal cities. *Estuarine, Coastal and Shelf Science*, *226*, 106262. https://doi.org/10.1016/j.ecss.2019.106262

Van Rooyen, M. W., van Rooyen, N., & Stoffberg, G. H. (2013). Carbon sequestration potential of post-mining reforestation activities on the KwaZulu-Natal coast, South Africa. *Forestry*, *86*(2), 211–223. https://doi.org/10.1093/forestry/cps070

Wepener, V., & Degger, N. (2019). South Africa. In C. Sheppard (Ed.), *World Seas: An Environmental Evaluation, the Indian Ocean to the Pacific* (Vol. *vol* II, 2nd ed., pp. 101–119). Elsevier.

Whitfield, A. K., Bate, G. C., Adams, J. B., Cowley, P. D., Froneman, P. W., Gama, P. T., Strydom, N. A., Taljaard, S., Theron, A. K., Turpie, J. K., van Niekerk, L., & Wooldridge, T. H. (2012). A review of the ecology and management of temporarily open/closed estuaries in South Africa, with particular emphasis on river flow and mouth state as primary drivers of these systems. *African Journal of Marine Science*, *34*(2), 163–180. https://doi.org/10.2989/1814232X.2012.675041

Wiles, E., Green, A. N., & Cooper, J. A. G. (2018). Rapid beachrock cementation on a South African beach: Linking morphodynamics and cement style. *Sedimentary Geology*, *378*, 13–18. https://doi.org/10.1016/j.sedgeo.2018.10.004

Williams, B. A., Watson, J. E. M., Beyer, H. L., Klein, C. J., Montgomery, J., Runting, R. K., Roberson, L. A., Halpern, B. S., Grantham, H. S., Kuempel, C. D., Frazier, M., Venter, O., & Wenger, A. (2022). Global rarity of intact coastal regions. *Conservation Biology*, *2022*, e13874. https://doi.org/10.1111/cobi.13874

Wood, E., Sanders, M., & Frazier, T. (2021). The practical use of social vulnerability indicators in disaster management. *International Journal of Disaster Risk Reduction*, *63*, 102464. https://doi.org/10.1016/j.ijdrr.2021.102464

Wright, C. I., & Mason, T. R. (1993). Management and sediment dynamics of the St. Lucia Estuary mouth, Zululand, South Africa. *Environmental Geology*, *22*(3), 227–241. https://doi.org/10.1007/BF00767408

Yang, S. -C., Riddin, T., Adams, J. B., & Shih, S. -S. (2014). Predicting the spatial distribution of mangroves in a South African estuary in response to sea level rise, substrate elevation change and a sea storm event. *Journal of Coastal Conservation*, *18*(4), 459–469. https://doi.org/10.1007/s11852-014-0331-2