

https://dspace.org/

Linear combinations of factorial and Sunit in a ternary recurrence sequence with a double root

Florian Luca; Armand Noubissie

AKADEMIAI KIADO RT
https://hdl.handle.net/10539/39212
WOS

Downloaded from a DSpace repository. https://dspace.org/

DOI 10.1007/s10986-023-09604-0
Lithuanian Mathematical Journal, Vol. 63, No. 3, July, 2023, pp. 249–253

On the arguments of the roots of the generalized Fibonacci polynomial*

Adel Alahmadi [a], **Oleksiy Klurman** [b], **Florian Luca** [c,d,1], **and Hatoon Shoaib** [a]

[a] Research Group in Algebraic Structures and its Applications, King Abdulaziz University, P.O. Box 80200, Postal Code 21589, Jeddah, Saudi Arabia

[b] School of Mathematics, University of Bristol, Fry Building, Woodland Road, Bristol, BS8 1UG, United Kingdom

[c,] School of Maths, Wits University, 1 Jan Smuts Ave, Braamfontein, Johannesburg, 2000, South Africa

[d] Centro de Ciencias Matemáticas, UNAM, Morelia, Mexico

(e-mail: analahmadi@kau.edu.sa; lklurman@gmail.com; florian.luca@wits.ac.za; hashoaib@kau.edu.sa)

Received January 29, 2021; revised June 6, 2022

Abstract. We revisit the classical subject of equidistribution of the roots of Littlewood-type polynomials. More precisely, we show that the roots of the family of polynomials $\Psi_k(z) = z^k - z^{k-1} - \cdots - 1$, $k \geqslant 1$, are uniformly distributed around the unit circle in the strong quantitative form, confirming a conjecture from [C.-A. Gómez and F. Luca, *Commentat. Math. Univ. Carol.*, On the distribution of roots of $z^k - z^{k-1} - \cdots - z - 1$, 62(3):291–296, 2021].

MSC: primary 11C08; secondary 11B39 12D10 30C15

Keywords: zeros of generalized Fibonacci polynomials

1 Introduction

Location of zeros of polynomials with restricted coefficients is a central subject in analysis with vast literature; see Ambrose [1], Bloch and Polya [2], Bombieri and Vaaler [3], Erdős and Turan [5], Borwein and Erdeley [4], Littlewood [7], Schur [9], Odlyzko and Poonen [8], and many others. In particular, if we restrict attention to the polynomials with coefficients ± 1, the so-called Littlewood polynomials, we expect that for a "typical polynomial" in this family, the zeros become "evenly distributed around the unit circle." Erdős and Turan [5] made this statement precise by proving their beautiful theorem. See also a recent paper of Soundararajan [10] with an new Fourier-analytic proof of their result. It is however a rather nontrivial task to show the equidistribution of zeros *for any particular* family of polynomials. One example when this task is essentially obvious is the family of polynomials

$$\psi_n(z) = z^n + z^{n-1} + \cdots + 1 = \frac{z^{n+1} - 1}{z - 1}.$$

* This project was funded by the Deanship of Scientific Research, King Abdulaziz University, under grant No. DF-595-130-1441.

[1] Corresponding author.

0363-1672/23/6303-0249 © 2023 The Author(s)

In this case the corresponding roots are the roots of unity $\alpha_n^k = \mathrm{e}^{2\pi \mathrm{i} k/(n+1))}$ with $k = \overline{1, n+1}$, which are evenly distributed around the circle.

Gómez and Luca [6] looked at the roots of the Littlewood polynomials $\Psi_k(z) = z^k - z^{k-1} - \cdots - 1$ for $k \geqslant 1$. Sometimes these polynomials are called the k-generalized Fibonacci polynomials since they are the characteristic polynomials of any k-length Fibonacci recurrence $\{F_n^{(k)}\}_{n \geqslant 0}$ with $F_{n+k}^{(k)} = F_{n+k-1}^{(k)} + \cdots + F_n^{(k)}$ for $n \geqslant 0$. Since $F_n^{(k)}$ is generally given by a Binet-type formula, which is a linear combination of the nth powers of the roots of $\Psi_k(z)$, it is important to understand the behavior of these roots if we want to answer Diophantine questions with k-generalized Fibonacci numbers. Let these roots be labeled $\alpha_1, \ldots, \alpha_k$. It is known that such a polynomial has exactly one real root larger than 1, which belongs to the interval $(2-1/2^k, 2)$. The other roots are inside the disc $|z| < 1$. In fact, a bit more is is known about their modulus, namely that they satisfy the inequality $1 - (\log 3)/k < |z| < 1 - 1/(2^8 k^3)$. This inequality was proved in [6]. As for the arguments of these roots, based on numerical evidence, Alexis Gómez, one of the authors of [6] conjectured the following:

Conjecture 1. For each $h \in \{0, 1, \ldots, k-1\}$, there is a root $\alpha_j = \rho_j \mathrm{e}^{\mathrm{i}\theta_j}$ of $\Psi_k(z)$ with $\theta_j \in [0, 2\pi)$ and

$$\left| \theta_j - \frac{2h\pi}{k} \right| < \frac{1}{k}.$$

The conjecture has been verified numerically by Gómez for all $k \in [2, 1000]$. Here we prove Conjecture 1 up to a small constant factor.

Theorem 1. *For $k \geqslant 2$ and every $h \in \{0, 1, \ldots, k-1\}$, there is θ_j such that*

$$\left| \theta_j - \frac{2h\pi}{k} \right| < \frac{\pi}{k}.$$

Furthermore, it appears that in the above inequality the right-hand side $O(1/k)$ improves to $O(1/k^2)$ as roots get closer to the imaginary axis.

Our proof avoids complex analysis. It only uses real analysis, more precisely, the fact that a continuous function of a real variable changing sign in an interval must have a zero in that interval.

2 Proof

Consider the related polynomial

$$g_k(z) = (z-1)\Psi_k(z) = z^{k+1} - 2z^k + 1.$$

Let $z = \rho \mathrm{e}^{\mathrm{i}\theta}$, $0 < \rho < 1$, and let $\theta \in [0, \pi)$ be a root of $g_k(z)$ in the upper half-plane. Rewrite the equation $g_k(z) = 0$ as

$$z + z^{-k} = 2$$

and consider the real and imaginary parts of both sides:

$$\rho \cos\theta + \rho^{-k} \cos(k\theta) = 2, \tag{2.1}$$

$$\rho \sin\theta - \rho^{-k} \sin(k\theta) = 0. \tag{2.2}$$

From (2.2) we obtain

$$\rho^{k+1} = \frac{\sin(k\theta)}{\sin\theta}. \tag{2.3}$$

Multiplying Eqs. (2.1) and (2.2) by $\sin(k\theta)$ and $\cos(k\theta)$, respectively, and then adding them, we obtain

$$\rho(\cos\theta\sin(k\theta)+\sin\theta\cos(k\theta))=2\sin(k\theta)$$

or

$$\rho\sin\big((k+1)\theta\big)=2\sin(k\theta), \tag{2.4}$$

which gives

$$\rho=\frac{2\sin(k\theta)}{\sin((k+1)\theta)}. \tag{2.5}$$

Elimination of ρ using (2.3) and (2.5) yields

$$\frac{2^{k+1}\sin^{k+1}(k\theta)}{\sin^{k+1}((k+1)\theta)}=\frac{\sin(k\theta)}{\sin\theta}. \tag{2.6}$$

By (2.2) and (2.4), $\sin\theta$, $\sin(k\theta)$, and $\sin((k+1)\theta)$ are all nonzero for $\theta\notin\{0,\pi\}$. Then by (2.6) θ must be one of the zeros of the function

$$H(x)=2^{k+1}\sin x\cdot\sin^k(kx)-\sin^{k+1}\big((k+1)x\big).$$

In the interval $x\in(0,\pi)$, $\sin(kx)$ vanishes at the points $\beta_\ell=\pi\ell/k$, $1\leqslant\ell\leqslant k-1$, whereas $\sin((k+1)x)$ vanishes at the points $\alpha_m=\pi m/(k+1)$, $1\leqslant m\leqslant k$. In view of the inequalities

$$\frac{j}{k+1}<\frac{j}{k}<\frac{j+1}{k+1}<\frac{j+1}{k},$$

these vanishing points are interlaced:

$$0<\alpha_1<\beta_1<\cdots<\alpha_j<\beta_j<\alpha_{j+1}<\beta_{j+1}\cdots<\beta_{k-1}<\alpha_k<\pi.$$

Since $\sin((k+1)x)>0$ for $x\in(\alpha_{2j},\alpha_{2j+1})$, by interlacing we obtain

$$H(\beta_{2j})=-\sin^{k+1}\big((k+1)\beta_{2j}\big)<0.$$

Likewise, $\sin(kx)>0$ for $x\in(\beta_{2j},\beta_{2j+1})$, whereas $\sin x>0$ in $(0,\pi)$. By interlacing

$$H(\alpha_{2j+1})=2^{k+1}\sin(\alpha_{2j+1})\sin^k(k\alpha_{2j+1})>0.$$

This means that $H(x)$ changes sign in each interval $(\beta_{2j},\alpha_{2j+1})$, and therefore it has at least one zero there, say, $\theta_j\in(\beta_{2j},\alpha_{2j+1})$. By interlacing the functions $\sin(kx)$ and $\sin((k+1)x)$ are both positive precisely in the intervals $(\beta_{2j},\alpha_{2j+1})$. If $H(\theta_j)=0$, then Eq. (2.6) holds for $\theta=\theta_j$. Consequently, the positive real number $\rho_j=\rho(\theta_j)$ defined by Eq. (2.4) satisfies Eqs. (2.3) and (2.4). This means that the pair (θ_j,ρ_j) is a solution to the system of equations (2.1)–(2.2). Therefore the complex number $z_j=\rho_j\mathrm{e}^{\mathrm{i}\theta_j}$ is the root of the polynomial $g_k(z)$. Conversely, by Eqs. (2.3) and (2.5) each nonreal root z_j with positive imaginary part must have an argument $\theta_j\in(\beta_{2j},\alpha_{2j+1})$ for some j in the range $1\leqslant j\leqslant\lfloor(k-1)/2\rfloor$. Since the number of such intervals $\lfloor(k-1)/2\rfloor$ matches the number of complex roots of $g_k(z)$, namely $(k-2)/2$ when k is even and $(k-1)/2$ when k is odd, there is a one-to-one correspondence between the set of roots and the set of intervals. Now let z be an arbitrary nonreal root of $g_k(z)$. By replacing z with its complex conjugate if necessary, we can

find an integer j in $[1, \lfloor (k-1)/2 \rfloor]$ such that $z = \rho_j e^{i\theta_j}$ with $\rho_j < 1$ and $\theta_j \in (\beta_{2j}, \alpha_{2j+1})$, as it was shown above. So the distance between β_{2j} to θ_j is at most

$$0 \leqslant \theta_j - \beta_{2j} \leqslant \alpha_{2j+1} - \beta_{2j} = \frac{\pi(2j+1)}{k+1} - \frac{\pi(2j)}{k} = \frac{\pi(k-2j)}{k(k+1)} \leqslant \frac{\pi(k-2)}{k(k+1)} \leqslant \frac{\pi}{k+1}. \tag{2.7}$$

This concludes the proof.

Remark. Note that as j approaches $\lfloor (k-1)/2 \rfloor$, the approximations get even better until the error decreases to about $2\pi/k^2$ near the imaginary axis. Namely, substituting $j = \lfloor (k-1)/2 \rfloor$ in (2.7) yields

$$0 \leqslant \theta_j - \beta_{2j} \leqslant \frac{\pi(k - 2\lfloor (k-1)/2 \rfloor)}{k(k+1)} \leqslant \frac{2\pi}{k(k+1)} < \frac{2\pi}{k^2}.$$

Acknowledgments. We thank the referee for supplying the present proof. Our original proof was longer, more complicated, and only yielded a slightly weaker result with a right-hand of $18/k$ in Theorem 1 instead of π/k. The authors gratefully acknowledge the Deanship of Scientific Research, King Abdulaziz University, Jeddah, for technical and financial support.

Funding. Open access funding provided by University of the Witwatersrand.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/

References

1. F. Amoroso, Sur le diamètre transfini entier d'un intervalle réel, *Ann. Inst. Fourier*, **40**(4):885–911, 1990, https://doi.org/10.5802/aif.1240.
2. A. Bloch and G. Pólya, On the Roots of Certain Algebraic Equations, *Proc. Lond. Math. Soc. (2)*, **33**(2):102–114, 1931, https://doi.org/10.1112/plms/s2-33.1.102.
3. E. Bombieri and J.D. Vaaler, Polynomials with low height and prescribed vanishing, in *Analytic Number Theory and Diophantine Problems. Proceedings of a Conference at Oklahoma State University, 1984*, Prog. Math., Vol. 70, Birkhäuser, Boston, MA, 1987, pp. 53–73, https://doi.org/10.1007/978-1-4612-4816-3_3.
4. P. Borwein and T. Erdélyi, On the zeros of polynomials with restricted coefficients, *Ill. J. Math.*, **41**(4):667–675, 1997, http://projecteuclid.org/euclid.ijm/1256068987.
5. P. Erdős and P. Turán, On the distribution of roots of polynomials, *Ann. Math. (2)*, **51**:105–119, 1950, https://doi.org/10.2307/1969500.
6. C.-A. Gómez and F. Luca, On the distribution of roots of $z^k - z^{k-1} - \cdots - z - 1$, *Commentat. Math. Univ. Carol.*, **62**(3):291–296, 2021, https://doi.org/10.14712/1213-7243.2021.025.
7. J.E. Littlewood, *Some Problems in Real and Complex Analysis*, D.C. Heath & Co, Lexington, MA, 1968.
8. A.M. Odlyzko and B. Poonen, Zeros of polynomials with $0, 1$ coefficients, *Enseign. Math. (2)*, **39**(3–4):317–348, 1993.

9. J. Schur, Affektlose Gleichungen in der Theorie der Laguerreschen und Hermiteschen Polynome, *J. Reine Angew. Math.*, **165**:52–58, 1931, https://doi.org/10.1515/crll.1931.165.52.

10. K. Soundararajan, Equidistribution of zeros of polynomials, *Am. Math. Mon.*, **126**(3):226–236, 2019, https://doi.org/10.1080/00029890.2019.1546078.