



DSPACE

<https://dspace.org/>

A new GIS-based biozone map of the Beaufort Group (Karoo Supergroup), South Africa

van der Walt, Merrill; Day, Michael; Rubidge, Bruce; Cooper, Antony K.; Netterberg, Inge

2010-12

Bernard Price Institute for Palaeontological Research, University of the Witwatersrand
<http://hdl.handle.net/10539/13819>

A new GIS-based biozone map of the Beaufort Group (Karoo Supergroup), South Africa

Merrill van der Walt^{1*}, Michael Day², Bruce Rubidge², Antony K. Cooper³ & Inge Netterberg⁴

¹Origins Centre, University of the Witwatersrand, Private Bag 3, WITS, 2050 South Africa

²Bernard Price Institute for Palaeontological Research, School of Geosciences, University of the Witwatersrand, Private Bag 3, WITS, 2050 South Africa

³Council for Scientific and Industrial Research (CSIR) Built Environment, P.O. Box 395, Pretoria, 0001 South Africa

⁴SAHA International, 39 Lytton Road, Observatory, Cape Town, 7925 South Africa

Received 16 November 2010. Accepted 15 December 2010

The stratigraphic subdivision of the Karoo Supergroup of South Africa has been addressed for more than a century by various geoscience studies. A lack of good lithostratigraphic markers in the Beaufort Group and the fact that the succession preserves a rich diversity of therapsid fossils for which there is a robust taxonomic scheme, make biostratigraphy the best option for correlation. This is applicable both within the basin and when comparing the Karoo succession to other coeval depositional basins. Published biostratigraphic maps of the Beaufort Group were compiled based on rough estimates of the distribution of zone-defining tetrapod fossil genera throughout the basin. None of these simultaneously utilized all the databases of South African museum collections. The recent application of GIS technology to integrate the vertebrate fossil databases of all South African museums has resulted in a far more precise biozone map which can be continuously updated as new information is received. This digital map introduces an entirely new way of representing the geographical distribution of fossil data, and thus can potentially enhance basin development and tetrapod biogeographic studies.

Keywords: Karoo Supergroup, fossil tetrapods, biostratigraphy.

INTRODUCTION

Biostratigraphy

The rocks of the Beaufort Group (Adelaide and Tarkastad subgroups) of the Karoo Supergroup cover approximately 60% of the surface of South Africa and comprise an approximately 3000 m thick succession of predominantly sedimentary rocks that are richly fossiliferous. Absolute ages for the Beaufort Group are not well constrained, with current dates based mainly on faunal correlations. The oldest stratigraphic units are considered Middle Permian (Kazanian) (Rubidge 1995a) and the uppermost strata as Middle Triassic (Anisian) (Ochev & Shishkin 1989; Hancox *et al.* 1995; Hancox & Rubidge 1996; Hancox 1998).

The global importance of the Beaufort Group is largely due to its remarkable assemblage of tetrapod fossils representing one of the best preserved ecological assemblages of pre-mammalian (therapsid) terrestrial vertebrates in the world (Keyser & Smith 1979), and the succession is held by many to be the global biostratigraphic standard for the non-marine Permo-Triassic (e.g. Shishkin *et al.* 1995; Lucas 1998). Furthermore, the rocks of the Beaufort Group preserve one of the most complete and best-studied non-marine Permo-Triassic (PT) boundary sequences globally (Smith 1995; Ward *et al.* 2000; Smith & Botha 2005; Botha & Smith 2006). These boundary sections preserve the terrestrial record of the Permo-Triassic extinction event – the greatest mass extinction event experienced on Earth (Erwin 1993, 1994, 2006).

Tetrapod fossils (particularly therapsids) have long been used for biostratigraphic subdivision and correlation of the Group which is especially useful because of the scarcity of basin-wide lithostratigraphic marker beds (Broom

1907a, 1909; Haughton 1924, 1963, 1969; Kitching 1970, 1972, 1977, 1984; Keyser & Smith 1979; Keyser 1979; Rubidge 1995b; SACS 1980). The few major lithological boundaries that have been traced throughout the basin have been shown to be diachronous (Keyser 1979), whereas the biozones are considered to closely approximate time lines (Rubidge 2005). Dicynodonts, the dominant herbivores of the Permian and early Triassic, are the most abundant fossil tetrapod taxon in the Beaufort Group and for this reason they are used as index fossils for the majority (five of the eight) of the currently recognized vertebrate biozones (Hancox & Rubidge 1997). Moreover, due to their abundance, dicynodonts have great potential for global correlation and have therefore been used in a number of previous biogeographic studies and bio-chronological schemes (Cooper 1982; Cruickshank 1985, 1986; Lucas 1990, 1993, 1995, 1998; Shishkin & Ochev 1992; De Fauw 1993).

Following is a summary of the history of work on the biostratigraphy of the Beaufort Group (Table 1) since Seeley (1892) proposed the first biostratigraphic subdivision of the Karoo which included three zones. Broom (1907b, 1909) refined the biozonation of Seeley by recognizing six subdivisions. Watson (1914a) agreed with Broom's subdivision and was the first to indicate the distribution of these units (which he termed 'zones') on a map. Later he replaced the *Pareiasaurus* Zone with the *Tapinocephalus* Zone (Watson 1914b), and an updated biozone distribution map was published in 1940 by von Huene. Hotton & Kitching (1963) pointed out that *Procolophon* occurs throughout the *Lystrosaurus* Zone and suggested that a separate overlying *Procolophon* Zone was not valid. Kitching (1970, 1977) introduced the *Daptocephalus* Zone to encompass the strata between the level where *Cistecephalus* became

*Author for correspondence. E-mail: merrill@origins.org.za

Table 1. Table showing the historical progression of Beaufort Group biozonation.

Beds	Zones		Assemblage Zones		
	Watson (1914 a, modified 1914b)	Kitching (1970, 1977)	Keyser & Smith (1979)	Keyser (1979) SACS (1980)	Current Biozonation Rubidge (1995b)
Broom (1906)					
<i>Cynognathus</i>	<i>Cynognathus</i>	<i>Cynognathus</i>	<i>Kannemeyeria</i>	<i>Kannemeyeria</i> <i>Diademodon</i>	<i>Cynognathus</i>
<i>Procolophon</i>	<i>Procolophon</i>	<i>Lystrosaurus</i>	<i>Lystrosaurus</i>	<i>Lystrosaurus</i> <i>Thrinaxodon</i>	<i>Lystrosaurus</i>
<i>Lystrosaurus</i>	<i>Lystrosaurus</i>				
<i>Kistecephalus</i>	<i>Cistecephalus</i>	<i>Daptocephalus</i>	<i>Dicynodon lacerticeps</i>	<i>Dicynodon lacerticeps</i> <i>Whaitsia</i>	<i>Dicynodon</i>
		<i>Cistecephalus</i>	<i>Aulacephalodon bairi</i>	<i>Aulacephalodon</i> <i>Cistecephalus</i>	<i>Cistecephalus</i>
<i>Endothiodon</i>	<i>Endothiodon</i>		<i>Tropidostoma microtrema</i>	<i>Tropidostoma</i> <i>Endothiodon</i>	<i>Tropidostoma</i>
<i>Pareiasaurus</i>	<i>Tapinocephalus</i>	<i>Tapinocephalus</i>	<i>Pristerognathus/Diictodon</i>	<i>Pristerognathus/Diictodon</i>	<i>Pristerognathus</i>
			<i>Dinocephalian</i>	<i>Dinocephalian</i>	<i>Tapinocephalus</i>
					<i>Eodicynodon</i>

extinct and the first appearance of *Lystrosaurus*, and discarded Broom's *Endothiodon* and *Procolophon* Zones. Kitching (1977) also produced a map showing zone-defining fossil localities. Keyser & Smith (1979) proposed a more refined vertebrate biozonation (and accompanying biozone map) for the Beaufort Group in the western part of the basin and linked their assemblage zones to the litho-stratigraphic scheme proposed by SACS (1980). The discovery of a tetrapod fauna from below the *Tapinocephalus* Assemblage Zone (Rubidge 1984; Rubidge *et al.* 1983; Rubidge *et al.* 1994) led to the erection of the *Eodicynodon* Assemblage Zone, which is currently the oldest vertebrate biozone of the Beaufort Group (Rubidge 1990, 1995a).

Following the recommendations of the International Stratigraphic Guide (ISSC 1976), the names of the assemblage zones that were accepted by SACS (1980) incorporated two genera. The most recent biostratigraphic scheme of the Beaufort Group which recognized eight biozones (Rubidge 1995b) follows the most recent nomenclatorial practice of the ISSC (1994) and uses only one taxon name in defining a biozone. This biozonation was represented in map form (Rubidge 1995b) and has since been the standard reference for the biozones of the Beaufort Group. The method used to create this map was manual drawing by enhancing the biozone maps of Kitching (1977) and Keyser & Smith (1979). This was essentially an exercise in outlining diagnostic fossil localities on an A3-sized map with a consequent low-level resolution. Over the past 15 years many more fossils from the Beaufort Group with quantifiable provenance data have been accessioned into museum collections and provide the impetus for a reassessment of biozone boundaries. In addition, since 2007 a GIS database has been built up incorporating data relating to all fossil specimens from the Beaufort Group which are curated in South African

palaeontological collections (Nicolas 2007; Nicolas & Rubidge 2009) and is an excellent tool to produce an updated biozone map.

MATERIALS AND METHODS

To produce the refined biozonation map from the GIS locality data, each biozone had to be defined in terms of key or 'marker' genera. Filtering for key genera allowed for biozone boundaries from the database to be defined. Ideally the lowermost boundary of a biozone is marked by the first appearance datum (FAD) of a key taxon, but in practise this is not always practical in the Beaufort biozonation scheme, which is based on assemblage zones, as the stratigraphic range of some biozone defining fossils are not necessarily restricted to the Assemblage Zone named after them. The diagnostic taxa utilized to define the individual biozones were taken from data presented in Rubidge (1995b), but in order to circumscribe the lower boundary of each biozone on the map the criteria set out in Table 2 were used.

Although ArcGIS allows for complex spatial analyses, numerous obstacles relating to data quality were encountered that presented barriers to the generation of biozone boundaries. These were:

- 1) *Quality of the locality data.* For older collections this often constitutes a farm name only. These are represented on the GIS map as farm centroids (Nicolas 2007). A farm covering two or more biozones may thus group genera from different biozones at a single point. Problematic localities were checked and topography examined to deduce the cause and nature of such apparent overlaps.
- 2) *The relationship between geology and topography.* At localities of high relief, the surface area of each biozone represented on the map is much smaller than when




Figure 1. Geographic distribution of the vertebrate biozones of the Beaufort Group based on a GIS database.

Table 2. Criteria used to circumscribe lower boundaries of biozones shown on the map.

Assemblage Zone	Criteria used to circumscribe lower boundary
<i>Eodicynodon</i>	Ecce-Beaufort boundary
<i>Tapinocephalus</i>	Last appearance of <i>Eodicynodon</i> , <i>Australosyodon</i> and <i>Tapinocaninus</i>
<i>Pristerognathus</i>	Last appearance of dinocephalians, presence of <i>Pristerognathus</i>
<i>Tropidostoma</i>	First appearance of <i>Tropidostoma</i> with presence of <i>Endothiodon</i>
<i>Cistecephalus</i>	First appearance of <i>Aulacephalodon</i> with presence of <i>Cistecephalus</i>
<i>Dicynodon</i>	Presence of <i>Dicynodon</i> , <i>Daptocephalus</i> absence of <i>Cistecephalus</i>
<i>Lystrosaurus</i>	Last appearance of <i>Dicynodon</i> , <i>Daptocephalus</i>
<i>Cynognathus</i>	First appearance of <i>Trirachodon</i> and <i>Cynognathus</i>

exposed on flat ground. The orientation of topographic features, such as escarpments, relative to dip is also important as this defines whether the zone remains at a constant level or alters in height laterally. This meant that without sophisticated three-dimensional modelling the accurate mapping of biozone boundaries was not possible, even if all locality input data was exact. As a result, biozone boundaries were drawn digitally utilising overlays of fossil genus localities, as well topographic and geological maps. In areas of great fossil abundance accuracy is highest, but for many areas, particularly in the Free State and eastern portion of the Eastern Cape provinces, this could not be achieved as there is a dearth of collecting localities. In these places, where fortuitously the strata are relatively horizontal, biozone boundaries were established through extrapolation by following mapped contour lines from one locality to the next. In a few instances where fossil localities are separated by tens of kilometres and the topography is complex as a result of folding of the strata or dolerite intrusions, biozone boundaries were traced by assessing lithostratigraphic boundaries. It should be noted that neither topography nor geology are always reliable features to follow because of the dip of beds and the diachroneity of lithostratigraphic boundaries. However, in the absence of fossil data, this method provides the best approximation.

There are some spurious specimen localities in the database that are aberrant compared to surrounding data points and cannot be explained by the farm centroid effect. Possible causes include the presence of an outlier of another biozone, specimen misidentification, the specimen not being *in situ* or incorrect cataloguing. In areas without physical deformation and flat topography, the former could be ruled out with confidence. Specimens with apparently anomalous localities that could not be explained by relief or physical deformation had their provenance and description data checked. If they were incorrectly situated then they were either moved to the correct locality or, if their true provenance could not be determined, ignored. Any specimens that were correctly situated but the authors remained unconvinced of their identity, were ignored and listed to be reidentified at a later date.

DISCUSSION

The new, GIS-based biozone map (Fig. 1) shows relatively few large-scale changes compared to its published precursors (Rubidge 1995b). The termination of the *Tapinocephalus* AZ close to the town of De Aar is an edu-

cated estimation because of the absence of dinocephalian fossils further north. The exact position of the boundary here is unknown due to a lack of collecting and the Ecce-Beaufort boundary at the nearest site to the north at Somersfontein near Philippolis is known to be *Pristerognathus* AZ (Welman *et al.* 2001). Also, small biozone outliers which usually coincide with high hills, have been acknowledged where fossil evidence illuminates their position. Previously these occurrences were too small-scale to be mapped, but the far higher resolution of the new digital map now makes this possible.

Despite the caveats discussed above, it is clear that the use of GIS greatly increases the achievable accuracy and ultimately the utility of the resulting digital map. It can be viewed at both small-scale, for the observation of broad overall patterns, and at large-scale for the assessment of smaller areas. It will be useful to identify areas where collecting has been sparse and the need for further research is greatest, as well as those locations where the boundary is best visible. One of the foremost applications, at least in the early stages, will be to draw attention to misidentified specimens, or specimens identified long ago using outdated characters.

This is an ongoing project. The GIS database, which is housed and curated at the BPI Palaeontology (University of the Witwatersrand, Johannesburg), will be updated continuously to include nomenclatorial modifications of existing specimens as well as the inclusion of new acquisitions to collections. The map is available at <http://web.wits.ac.za/Academic/Science/GeoSciences/BPI/>. Updating metadata will increase the accuracy of the map and provide an accessible record in 2D and ultimately in 3D. Use of this kind of spatial modelling will revolutionize the way biozones are mapped and will ensure that all new data are recorded in a systematic and centralized manner. The final product will serve as an up-to-date representation of the surface expression of vertebrate fossil assemblage zones of the main Karoo Basin in South Africa.

Setting up the Karoo fossil GIS database has been a long-term project which would not have been possible without the cooperation and enthusiasm of the curators of all the Karoo fossil collections in South Africa. We record our gratitude to Sheena Kaal and Roger Smith (Iziko South African Museum), Billy de Klerk (Albany Museum), Richard, Robert and Marion Rubidge (Rubidge Collection), Elize Butler and Jennifer Botha-Brink (National Museum), Bernhard Zipfel (BPI Palaeontology), Heidi Fourie (Transvaal Museum) and Johann Neveling and Ellen de Kock (Council for Geoscience). We are greatly indebted to Cynthia Kemp for the many hours she spent painstakingly editing and updating records. We also record our gratitude to the Palaeontological Scientific Trust (PAST), the Department of Science and Technology (DST), and the (National Research Foundation) NRF for providing funding. We are grateful to Jennifer Botha-Brink and an anonymous reviewer for improving the manuscript.

REFERENCES

- BOTHA, J. & SMITH, R.M.H. 2006. Rapid recuperation in the Karoo Basin of South Africa following the End-Permian extinction. *Journal of African Earth Sciences* **45**, 502–514.
- BROOM, R. 1906. On the Permian and Triassic faunas of South Africa. *Transactions of the Geological Society of London* **7**, 175–192.
- BROOM, R. 1907a. On the geological horizons of the fossil vertebrates of the Karoo Formation. *Records of the Albany Museum* **2**, 156–163.
- BROOM, R. 1907b. On some new fossil reptiles from the Karoo beds of Victoria West, South Africa. *Transactions of the South African Philosophical Society* **18**, 31–42.
- BROOM, R. 1909. An attempt to determine the horizons of the fossil vertebrates of the Karoo. *Annals of the South African Museum* **7**, 285–289.
- COOPER, M.R. 1982. A Mid-Permian to Earliest Jurassic tetrapod biostratigraphy and its significance. *Arnoldia Zimbabwe* **7**, 77–103.
- CRUICKSHANK, A.R.I. 1985. Biostratigraphic significance of kane-meyeriid Evolution. *Terra Cognita* **5**, 136.
- CRUICKSHANK, A.R.I. 1986. Biostratigraphic and classification of a new Triassic Dicynodont from East Africa. *Modern Geology* **10**, 121–131.
- DE FAUW, S.L. 1993. The Pangean dicynodont *Rechnisaurus* from the Triassic of Argentina. In: Lucas, S.G. & Morales, M. (eds), The non-marine Triassic. *New Mexico Museum of Natural History and Science Bulletin* **3**, 101–105.
- ERWIN, D.H. 1993. *The Great Paleozoic Crises: Life and Death in the Permian*. New York, Columbia University Press.
- ERWIN, D.H. 1994. The Permo-Triassic extinction. *Nature* **367**, 231–236.
- ERWIN, D.H. 2006. *Extinction: How Life on Earth Nearly Ended 250 Million Years Ago*. Princeton, New Jersey, Princeton University Press.
- HANCOX, P.J. 1998. A stratigraphic, sedimentological and palaeo-environmental synthesis of the Beaufort-Molteno contact in the Karoo Basin. Unpublished Ph.D. thesis, University of the Witwatersrand, Johannesburg.
- HANCOX, P.J., SHISHKIN, M.A., RUBIDGE, B.S. & KITCHING, J.W. 1995. A threefold subdivision of the *Cynognathus* Assemblage Zone (Beaufort Group, South Africa) and its palaeogeographical implications. *South African Journal of Science* **91**, 143–144.
- HANCOX, P.J. & RUBIDGE, B.S. 1996. The first specimen of the Mid-Triassic Dicynodont *Angonisaurus* from the Karoo of South Africa: implications for the dating and biostratigraphy of the *Cynognathus* Assemblage Zone, Upper Beaufort Group. *South African Journal of Science* **92**, 391–392.
- HANCOX, P.J. & RUBIDGE, B.S. 1997. The role of fossils in interpreting the development of the Karoo Basin. *Palaeontologia africana* **33**, 41–54.
- HAUGHTON, S.H. 1924. A bibliographic list of Pre-Stormberg Karoo Reptilia, with a table of horizons. *Transactions of the Royal Society of South Africa* **12**(2), 51–104.
- HAUGHTON, S.H. 1963. Note on the distribution of fossil reptiles of Karoo age. *Palaeontologia africana* **7**, 1–11.
- HAUGHTON, S.H. 1969. *Geological History of Southern Africa*, 348–526. Johannesburg, Geological Society of South Africa.
- HOTTON, N. & KITCHING, J.W. 1963. Speculations on upper Beaufort deposition. *South African Journal of Science* **59**, 254–258.
- ISSC: International Subcommittee on Stratigraphic Classification, 1976. Hedberg, H.D. (ed.), *International Stratigraphic Guide*, 200. New York, Wiley-Interscience.
- ISSC: International Subcommittee on Stratigraphic Classification, 1994. *International Stratigraphic Guide* (2nd edn). Trondheim, International Union of Geological Sciences, and Boulder, Geological Society of America.
- KEYSER, A.W. 1979. A review of the biostratigraphy of the Beaufort Group in the Karoo basin of South Africa. Abstracts from Geocongress, Geological Society of South Africa **2**, 13–31.
- KEYSER, A.W. & SMITH, R.M.H. 1979. Vertebrate biozonation of the Beaufort Group with special reference to the Western Karoo Basin. *Annals of the Geological Survey South Africa* **12**, 1–36.
- KITCHING, J.W. 1970. A short review of the Beaufort Zoning in South Africa. In: *Second Gondwana Symposium – Proceedings and Papers*, 309–313. Pretoria, Council for Scientific and Industrial Research.
- KITCHING, J.W. 1972. *On the distribution of the Karoo vertebrate fauna, with special reference to certain genera and the bearing of this distribution on the zoning of the Beaufort beds*. Unpublished Ph.D thesis, University of the Witwatersrand, Johannesburg.
- KITCHING, J.W. 1977. The distribution of the Karoo vertebrate fauna, 131. *Memoirs of the Bernard Price Institute for Palaeontological Research, University Witwatersrand* **1**.
- KITCHING, J.W. 1984. A reassessment of the biozonation of the Beaufort Group. *Palaeontology News* **4**(1), 12–13.
- LUCAS, S.G., 1990. Toward a vertebrate biochronology of the Triassic. *Albertiana* **8**, 36–41.
- LUCAS, S.G. 1993. The Shansiodon biochronology, Middle Triassic of Pangaea. *Albertiana* **11**, 40–42.
- LUCAS, S.G. 1995. Towards dicynodont biochronology. *Albertiana* **16**, 33–40.
- LUCAS, S.G. 1998. Global Triassic tetrapod biostratigraphy and biochronology. *Palaeogeography, Palaeoclimatology, Palaeoecology* **143**(4), 347–384.
- NICOLAS, M.V. 2007. *Tetrapod biodiversity through the Permo-Triassic Beaufort Group (Karoo Supergroup) of South Africa*. Unpublished Ph.D. thesis, University of the Witwatersrand, Johannesburg.
- NICOLAS, M.V. & RUBIDGE, B.S. 2009. Assessing content and bias in South African Permo-Triassic Karoo tetrapod fossil collections. *Palaeontologia africana* **44**, 13–20.
- OCHEV, V.G. & SHISHKIN, M.A. 1989. On the principles of global correlation of the continental Triassic on the tetrapods. *Acta Palaeontologica Polonica* **34**, 149–173.
- RUBIDGE, B.S. 1984. The cranial morphology and palaeoenvironment of *Eodicynodon* Barry (Therapsida: Dicynodontia). *Navorsing van die Nasionale Museum Bloemfontein, South Africa* **4**(14), 325–402.
- RUBIDGE, B.S. 1990. A new vertebrate biozone at the base of the Beaufort Group, Karoo sequence (South Africa). *Palaeontologia africana* **27**, 17–20.
- RUBIDGE, B.S. 1995a. Biostratigraphy of the *Eodicynodon* Assemblage Zone. In: Rubidge, B.S. (ed.), *Biostratigraphy of the Beaufort Group (Karoo Supergroup)*, 3–7. South African Committee for Stratigraphy. *Biostratigraphic Series 1*. Pretoria, Council for Geoscience.
- RUBIDGE, B.S. (ed.) 1995b. *Biostratigraphy of the Beaufort Group (Karoo Supergroup)*. South African Committee for Stratigraphy. *Biostratigraphic Series 1*. Pretoria, Council for Geoscience.
- RUBIDGE, B.S., KITCHING, J.W. & VAN DEN HEEVER, J.A. 1983. First record of a theropod (Therapsida: Pristerognathidae) from the Ecca of South Africa. *Navorsing van die Nasionale Museum Bloemfontein, South Africa* **4**, 229–235.
- RUBIDGE, B.S., KING, G. & HANCOX, P.J. 1994. The postcranial skeleton of the earliest dicynodont synapsid, *Eodicynodon* from the Upper Permian of South Africa. *Palaeontology* **37**(2), 397–408.
- RUBIDGE, B.S. 2005. Re-uniting lost continents – Fossil reptiles from the ancient Karoo and their wanderlust. *South African Journal of Geology* **108**(3), 135–172.
- SACS: South African Committee for Stratigraphy, 1980. *Stratigraphy of South Africa. Part 1: Lithostratigraphy of the Republic of South Africa, South West Africa/Namibia and the Republics of Boputhatswana, Transkei and Venda* (L.E. Kent, comp.), 690. Handbook of the Geological Survey, South Africa, 8.
- SEELEY, H.G. 1892. Researches on the structure, organization and classification of the fossil Reptilia. In: *Further Observations on Pareiasaurus*, 311–370. *Philosophical Transactions of the Royal Society, London*, B **183**.
- SHISHKIN, M.A. & OCHEV, V.G. 1992. On the problem of dating the Upper Beaufort vertebrate faunas. In: Abstracts, 24th Congress of the Geological Society South Africa, Bloemfontein, 366–368.
- SHISHKIN, M.A., RUBIDGE, B.S. & HANCOX, P.J. 1995. Vertebrate biozonation of the Upper Beaufort Series of South Africa – a new look on correlation of the Triassic biotic events in Euamerica and southern Gondwana. In: Sun, A. & Wang, Y. (eds), *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota*. Short papers, 39–41. Beijing: China Ocean Press.
- SMITH, R.M.H. 1995. Changing fluvial environments across the Permian-Triassic boundary in the Karoo Basin, South Africa and the possible causes of tetrapod extinctions. *Palaeogeography, Palaeoclimatology, Palaeoecology* **117**, 81–104.
- SMITH, R.M.H. & BOTHA, J. 2005. The recovery of terrestrial vertebrate diversity in the South African Karoo Basin after the end-Permian extinction. *Comptes Rendus Palévol* **4**, 555–568.
- VON HUENE, F. 1940. Die Saurier de Karroo-, Gondwana- und verwandten Ablagerungen in faunistischer, biologischer und phylogenetischer Hinsicht. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Abhandlungen* **83**, 275–303.
- WARD, P.D., MONTGOMERY, D.R. & SMITH, R.M.H. 2000. Altered river morphology in South Africa related to the Permian-Triassic extinction. *Science* **289**, 1740–1743.
- WATSON, D.M.S. 1914a. The zones of the Beaufort Beds of the Karoo System in South Africa. *Geological Magazine, New Series, Decade* **6**(1), 203–208.
- WATSON, D.M.S. 1914b. On the nomenclature of the South African pareiasaurians. *Annals of the Magazine of Natural History* **8**(14), 98–102.
- WELMAN, J., LOOCK, J.C. & RUBIDGE, B.S. 2001. New evidence for diachroneity of the Ecca-Beaufort contact (Karoo Supergroup, South Africa). *South African Journal of Science* **97**, 320–322.