

LIST OF FIGURES

Figure 2.1 Global climatic changes corresponding to the South African Middle and Later Stone Age (image reproduced with permission from C. Henshilwood). The years and corresponding marine oxygen isotope stages are taken from Martinson <i>et al.</i> 1987 and the ages for the Howiesons Poort and Still Bay from Jacobs <i>et al.</i> 2008	4
Figure 2.2 Dated Middle and Later Stone Age occupations at Blombos Cave. The LSA occupation phases L1 and L6 were dated by radiocarbon within the last 2 ka. The ages for the intermittent MSA occupations (denoted by the M1, M2 & M3 phases) were determined using OSL techniques and fall within the Last Glacial Period (Jacobs <i>et al.</i> 2006) (image reproduced with permission from C. Henshilwood; Jacobs <i>et al.</i> 2006)	5
Figure 2.3 Stratigraphy of the archaeological deposits at Blombos Cave including the ages of the Middle Stone Age levels (image reproduced with permission from C. Henshilwood).....	6
Figure 2.4 The southern Cape as defined in this study refers to the region between Cape Town and Mossel Bay and includes the De Hoop Nature Reserve and Blombos Cave	8
Figure 2.5 Simplified stratigraphy of the main geological formations at De Hoop and Blombos Cave	9
Figure 2.6 Key features of the southern Cape coastal platform	10
Figure 2.7 Main components of the atmospheric circulation which determine the weather and climate of South Africa	12
Figure 2.8 Main ocean currents affecting South Africa	13
Figure 2.9 Extent of South Africa's three rainfall zones. The abbreviations WRZ, SRZ and YRZ refer to the winter-rainfall zone, summer rainfall zone and year-round rainfall zone, respectively (From Chase & Meadows 2007)	14
Figure 2.10 Main soil types found in the De Hoop Nature Reserve and Blombos Cave.....	17
Figure 2.11 (a) 200 kyr sea level curve for South Africa (From Ramsay & Cooper 2002: 87)	19
Figure 2.11 (b) Relative position of the southern Cape coastline within the last 200 kyr (From Fisher <i>et al.</i> 2010)	20
Figure 2.12 Present position of the archaeological sites mentioned in the text	23

Figure 3.1 View of Blombos Cave (image reproduced with permission from C. Henshilwood)	34
Figure 3.2 Map showing the location of Blombos Cave and other sites mentioned in the text	35
Figure 3.3 Simplified stratigraphy of the main geological formations at Blombos Cave	36
Figure 3.4 Blombos Cave layout (image reproduced with permission from C. Henshilwood)	37
Figure 3.5 Stratigraphic levels based on excavations conducted from 1997 to 1999 (image reproduced with permission from C. Henshilwood)	38
Figure 3.6 Dated Middle Stone Age stratigraphic units at Blombos Cave and corresponding finds. The circles indicated the position of the OSL samples (Jacobs <i>et al.</i> 2006)	40
Figure 3.7 Location of the main southern African sites containing Still Bay and Howiesons Poort artefacts. Key: AA: Aar 1, AP: Apollo 11, BBC: Blombos Cave, BP: Boomplaas, BC: Border Cave, B1C: Bremen 1C, CH: Cave of Hearths, DRS: Diepkloof Rock Shelter, HAA: Haalenberg, HAS: Ha Soloja, HR: Highlands, HRS: Hollow Rock Shelter, HP: Howiesons Poort; KP: Kathu Pan, KRM: Klasies River Mouth, KKH: Klein Kliphuis, KLP: Klipfontein, MLK: Melikane, MON: Montagu Cave, MOS: Moshebi's Shelter, NBC: Nelson Bay Cave, NT: Ntloana Tsoana, OAK: Oakleigh, PAR: Paardeberg, PC: Peers Cave, PP: Pinnacle Point, POC: Pockenbanck, RCC: Rose Cottage Cave, SEH: Sehonghong, SB: Sibudu, UMH: Umhlatuzana, WK: Wonderwerk Cave (Modified from Jacobs & Roberts 2008: 21)	42
Figure 3.8 Bifacial points from Blombos Cave (image reproduced with permission from C. Henshilwood)	45
Figure 3.9 Blombos bone tools (image reproduced with permission from C. Henshilwood)	46
Figure 3.10 <i>Nassarius kraussianus</i> shell beads (image with permission from C. Henshilwood)	52
Figure 3.11 Engraved ochre from the c. 75 ka M1 level (image reproduced with permission from C. Henshilwood)	53
Figure 3.12 Incised ochre pieces from the c. 100 ka M3 level and the M1 and M2 levels (image reproduced with permission from C. Henshilwood)	54
Figure 4.1 (a) Extent of the De Hoop Nature Reserve	66

Figure 4.1 (b) Location of the caves surveyed in the reserve during April 2008	67
Figure 4.2 (a & b) Box Cave (34° 27.058'S, 20°42.011'E).....	68
Figure 4.3 (a-c) West Cave (34°27.049'S, 20°41.883'E).....	69
Figure 4.4 Morris's Cave (34°27.061'S, 20°42.016'E).....	70
Figure 4.5 (a & b) Bloukrantz Cave (34°27.557'S, 20°46.697'E).....	71
Figure 4.6 (a -c) Kaisers Gat II (34°28.248'S, 20°49.634'E).....	72
Figure 4.7 Longitudinal section through <i>Blou1</i>	74
Figure 4.8 Cross-section of the <i>KG2.1</i> stalagmite.....	75
Figure 4.9 The <i>KG2.2</i> flowstone sample from Kaisers Gat II	76
Figure 4.10 The <i>KG2.3</i> sample from Kaisers Gat II	77
Figure 4.11 Radioactive decay series for ^{238}U including the half-lives for the long-lived isotopes; ^{238}U , ^{234}U and ^{230}Th	80
Figure 4.12 An example of an isochron plot.....	83
Figure 4.13 (a-e) Eichron TRU resin column preparations for U and Th elution.....	86
Figure 4.14 Diagram illustrating the main components of the ICP MS.....	88
Figure 4.15 Transformation of the liquid sample as it moves from the nebulizer through the plasma, leaving the ICP torch via the interface region en route to the mass spectrometer	89
Figure 4.16 The three most abundant carbon isotopes are the stable ^{12}C and ^{13}C and the radioactive ^{14}C . These isotopes have the same number of electrons and protons within the atomic nucleus but different numbers of neutrons	94
Figure 4.17 Factors above the cave that influence the speleothem isotopic signal.....	97
Figure 4.18 Three stable isotopes of oxygen.....	98
Figure 4.19 The hydrological cycle.....	100
Figure 4.20 Rayleigh fractionation and the isotopic composition of precipitation during rain out events	101

Figure 4.21 Global meteoric water line (GMWL) representing the isotopic variations in water sampled from rivers, lakes, snow and rain in different regions of the world	102
Figure 4.22 Sub-sampling approach used for the stable isotope measurements	106
Figure 4.23 Diagram of the MAT 253 mass spectrometer	107
Figure 4.24 Schematic of the main components of the MAT 253 Kiel carbonate device	108
Figure 5.1 Age-depth profile for sample KG2.3 from the Kaisers Gat II Cave using the stratigraphic data and ages presented in tables 5.1 and 5.2, respectively. The U-series sample locations are represented by white blocks and the arrow indicates the assumed direction of growth	118
Figure 5.2 Inverted age-depth profile of the KG2.3 stalagmite and the corresponding sample numbers	119
Figure 5.3 Variation in the KG2.3 $^{230}\text{Th}/^{232}\text{Th}$ activity ratios	120
Figure 5.4 Plot of the $^{234}\text{U}/^{238}\text{U}$ and initial $^{234}\text{U}/^{238}\text{U}$ activity ratios for KG2.3	121
Figure 5.5 Activity ratio of $^{230}\text{Th}/^{234}\text{U}$ for KG2.3	122
Figure 5.6 An age-stratigraphic plot for sample <i>Blou1</i> from Bloukrantz Cave. The U-series sample locations are represented by red blocks	124
Figure 5.7 (a) HENDY Test applied to the Bloukrantz Cave (<i>Blou1</i>) and (b) Kaisers Gat II (KG2.3) speleothems. The <i>Blou1</i> data is represented by the closed (red) dots. The open (blue) dots represent the KG2.3 values.....	133
Figure 5.8 Comparison of the stable carbon and oxygen isotope signals preserved in the speleothems from the Bloukrantz Cave (<i>Blou1</i>) and Kaisers Gat.....	134
Figure 5.9 Variations in $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ sampled from the <i>Blou1</i> stalagmite	135
Figure 5.10 Variations in the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ isotopic profile of the KG2.1 stalagmite	136
Figure 5.11 $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values obtained from the KG2.2 flowstone	137
Figure 5.12 $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ data for the KG2.3 stalagmite	138
Figure 5.13 Trend line representing the main growth phase in the development of the <i>Blou1</i> stalagmite	139

Figure 5.14 Age estimations based on the <i>Blou1</i> U-series dates and constructed using a linear interpolation formula	140
Figure 5.15 Isotope time-series curve for <i>Blou1</i> corresponding to MIS 3	141
Figure 5.16 $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values and uncorrected U-series ages obtained for the KG2.3 sample. The ages are not interpolated because of the discrepancy between the stratigraphy and corresponding age determinations	147
Figure 5.17 $\delta^{13}\text{C}$ time-series for <i>Blou1</i>	149
Figure 5.18 $\delta^{18}\text{O}$ isotope record for <i>Blou1</i>	149
Figure 6.1 (a, b) Comparison between the Dome F $\delta^{18}\text{O}$ record and the De Hoop isotope data from <i>Blou1</i> and KG2.3. The KG2.3 isotope data has not been interpolated and the isotope values does not represent an age- interpolation The Dome F isotope data was obtained from ftp://ftp.ncdc.noaa.gov/pub/data/paleo/icecore/antarctica/domefuji/df-d18o-340ka-dfo2006.txt , Kawamura <i>et al.</i> 2007)	152
Figure 6.2(a, b) Palaeovegetation data obtained from the $\delta^{13}\text{C}$ signal in <i>Blou1</i> and KG2.3 compared with the Indian Ocean sea surface temperatures (SSTs) from the MD962077 sediment core. (The data corresponding to MD962077 was obtained from Bard and Rickaby (2009) at http://www.nature.com/nature/journal/v460/n7253/supinfo/nature08189.html)	153
Figure 6.3 $\delta^{18}\text{O}$ signal in the Dome F ice core compared with the Indian Ocean temperatures recorded in MD962077 (The data for the Dome F ice core was obtained from ftp://ftp.ncdc.noaa.gov/pub/data/paleo/icecore/antarctica/domefuji/df-d18o-340ka-dfo2006.txt , Kawamura <i>et al.</i> 2007 & the data for MD962077 was obtained from Bard and Rickaby (2009) at http://www.nature.com/nature/journal/v460/n7253/supinfo/nature08189.html)	154