

## 19. SUMMARY

1. Observations of the behaviour and ecology of Thick-tailed Bushbabies (Galago crassicaudatus) have been carried out over a period of eighteen months between February 1970 and August 1971. Detailed information was collected during fifteen months in a single study area while comparative observations were made in two other widely separated localities. Individual bushbabies, or groups, were followed for long periods at night using red light from a headband torch and a strong beam of white light from a hand lantern when necessary. A total of 663 animal observation hours were made, concentrating on the evenings and mornings when the animals were most active. Eleven bushbabies were caught in traps, of which four were housed under semi-natural conditions and the remainder marked for easy identification prior to their release. The data which were obtained are compared with those for other nocturnal prosimians, particularly the Lesser Bushbaby (G. senegalensis), the only other prosimian in southern Africa, which is sympatric with the study species in some areas.

2. The distribution of G. crassicaudatus is outlined in comparison with that of G. senegalensis. Effective isolation of breeding populations appears to have resulted in a number of sub-species, but the classification remains unclear. Three populations of G. crassicaudatus, which are generally considered as separate sub-species, were studied in different habitat types: G.c. umbrosus Thos. in riverine bush of the North Eastern Transvaal; G.c. garnetti (Ogilby) in forest habitats of Zululand and G.c. lönnbergi Schwarz in open woodland of Eastern Rhodesia. Each study area is described in terms of its vegetation, topography and climate, with a list of other local primate species. A few size and weight measurements are presented for individuals of the first sub-species from the main study area.

3. G. crassicaudatus are found at various altitudes and show tolerance to marked seasonal and diurnal variations in temperature. The species occurs in relatively open habitats but it is typically confined to dense evergreen vegetation. Estimations of population densities indicate that G. crassicaudatus are most abundant in high rainfall, humid

forest areas where there are many trees bearing fleshy fruits. This contrasts with G. senegalensis which are most common in semi-arid acacia thornveld. The relatively high population densities of both species when compared with related forms in tropical forest habitats can be explained as a result of the few nocturnal arboreal competitors.

4. G. crassicaudatus are strictly nocturnal. They leave cover after sunset and return before sunrise at times which are dependent on the level of light intensity beneath the trees. Activity at night follows a broadly bimodal pattern, with periods of rest tending to increase towards midnight. Long spells spent in a single tree are often followed by sudden movement through several trees. This pattern is basically similar to that of G. senegalensis, although this species shows many short bursts of movement during each hour. The difference between the two species in this respect may be related to divergence in their feeding habits. The similarity of the circadian activity cycles of these primates in the wild and under artificial conditions in captivity suggests an 'internal clock' which governs the general level of activity at any one time.

5. Fruits and resins (gums) represent the staple diet of G. crassicaudatus in the main study area but seeds, nectar and insects are also consumed. Moisture may occasionally be licked from leaves but otherwise bushbabies have not been seen to drink. Gum is extremely important in the winter when soft fruits are not available. Particular gum licks are well known to the animals, they are visited regularly and appear to be remembered. Gum is obtained by licking, or by first biting and then licking, but not by scraping with the lower anterior teeth. A remarkable number of gum licking postures may be assumed which enable this species to reach gum which is unavailable to food competitors. Continuous exploration ensures that small new sources of food are not missed but large sources may be smelled from a distance. The animals return repeatedly to certain food trees for as long as they remain productive. Predation on birds and other small vertebrates has often been reported for G. crassicaudatus but evidence is presented to show that it is not a universal characteristic. It appears to be a local habit which may spread within a population while being absent from another nearby. G. senegalensis also eat gum, particularly during the winter, but otherwise they are predominantly insectivorous. They catch insects by means of a ritualized prey catching strike, which G. crassicaudatus lack, and

they are able to survive in the absence of free water.

6. The locomotory postures of G. crassicaudatus are described and illustrated and compared with those of G. senegalensis and other prosimians. The overall pattern of locomotion differs from that of other Galaginae and is in many ways similar to that of the Lorisinae. Despite the morphological specializations which suit an active leaping mode of progression, this species is behaviourally adapted to a slower quadrupedal gait. This may be interpreted as a relatively recent adaptation for obtaining food by stealth in a forest environment, which has not been matched by skeletal changes. Undoubtedly the ability to move quietly also enables bushbabies to avoid detection by potential predators but, nevertheless, they are still capable of rapid escape. Similarities between the locomotion of G. crassicaudatus and the slower moving Perodicticus potto are considered to be a result of convergent adaptation.

7. G. crassicaudatus make regular use of a number of sleeping places during the day which provide concealment and direct protection from predators and extremes of temperature. Nest building is rare but it has been seen during the breeding season when young infants may require a suitable support. The offspring may be afforded additional protection from predators at this time by their mothers' habit of moving them from one sleeping place to another. This behaviour is similar to that of G. senegalensis which, however, continue to build nests after they are no longer essential for the support of the young. Members of this species will sleep in exposed places, relying to a greater extent on their colouration for camouflage, and they will move in the face of a threatening stimulus or a change in weather conditions.

8. G. crassicaudatus have a restricted breeding season at least in the southern part of their range. All births occur within a single period of approximately three weeks at the start of the rainy season when food and dense cover is readily available. Twins are most common but triplets and a singleton have also been recorded. Mating is correspondingly restricted to a short period in mid-winter and it is accompanied by a general increase in the level of excitability within a population. G. senegalensis have two birth and mating periods during the year but, like G. crassicaudatus, practically all infants are weaned by the end of the wet summer months.

9. G. crassicaudatus sleep either alone or in small family groups of two to six animals which may include an adult male, an adult female

and the offspring of one or two generations. The male and maturing offspring will often sleep alone, while the female and her youngsters usually sleep together. An examination of the social dynamics and mother/offspring relationship within a single group has been used to illustrate the development of the young and the nature of associations between individuals. In contrast to most other nocturnal prosimians, in which the sleeping groups split up for much of the night, G. crassicaudatus are commonly found in small groups whose members forage together. It is the female and the young which move together as a cohesive unit. Even when the infants are too small to follow the mother she will carry them from place to place, either in her mouth, one at a time, or with them clinging to the fur of her back, or both. This close association continues until the offspring are at least sub-adult. The range of the adult male may coincide to a large extent with that of the maternal group but he generally remains alone. He occasionally joins the group and moves with it without any signs of serious hostility. Even larger groupings may be formed through attraction to a particular food source.

10. The movements and use of space of G. crassicaudatus are discussed with reference to night range and annual home range. Rapid movements between feeding trees are made by way of regular pathways but when moving slowly the animals make a variety of deviations which allow exploitation of food sources at different heights over a large area. There is evidence that sight and hearing coupled with a kinaesthetic sense are important aids to movement rather than scent trails through the trees. In the main study group the adult male moved over a wider range than the female and offspring which utilized an annual home range of seven hectares. The maternal group spent most time in a central core area containing many sleeping sites, food trees and resting places yet covering only 4% of the total range. The male made use of the same fixed points but he frequented a different core area. The use of fixed points varied during the year and seems to depend on custom and convenience as much as on their suitability. Seasonal changes in the focus of activity can be related to the productivity of favourite food trees. G. crassicaudatus, like many other primate species, show little in the way of active territorial defence but groups may remain attached to a single range for long periods, while much of their behaviour maintains internal cohesion and ensures the dispersal of maturing individuals and separation between groups. Differences in the pattern of movement in a variety of

habitats indicate that the height or density of the vegetation has little direct effect on the distribution of this species, although it is somewhat less versatile in its movements than the smaller G. senegalensis.

11. Various aspects of social behaviour in G. crassicaudatus are described and discussed in relation to the establishment and maintenance of social ties and the spacing mechanisms between conspecifics. Bushbabies are able to groom almost all parts of the body by autogrooming using the tongue, ventral anterior dentition, or tooth-scraper, and the toilet claws on the second pedal digits. Yet allogrooming is performed at a high frequency which suggests that it serves important social functions. The amount of time spent allogrooming in this species is greater than for G. senegalensis in which group members spend less time together. It appears to enhance intra-group harmony and it may aid friendly contact between compatible strangers. Play is considered as a second cohesive force which serves a number of other functions as well. G. crassicaudatus perform six marking and/or rubbing actions in a variety of situations. Particular places are marked repeatedly and these may act as information posts of significance in a territorial or social context, or they may merely reassure the marker. It is suggested that the complex ritualized pattern of urine washing was originally derived as an adaptation to help maintain an effective grip in small active primates following the loss of claws. A detailed comparison of the vocalizations of G. crassicaudatus and G. senegalensis shows up remarkable structural and functional similarities in their vocal repertoires, while the differences correlate with divergencies in group structure and the rearing of the young. Finally, agonistic behaviour is outlined and compared with similar interactions by bushbabies in captivity where the types of encounters are often unnatural. Antagonism between individuals is mediated by mutual recognition of the sex and status of a social fellow or rival. Intra-group tension is mild and spacing mechanisms ensure that fighting between strangers is rare. G. crassicaudatus has a basically similar social structure to other nocturnal prosimians which have been studied but it shows trends towards a better developed social way of life through increased social contact between the mother and her offspring.

12. Some potential predators of G. crassicaudatus are listed and their importance discussed briefly. A number of the behavioural characteristics of the species enable it to avoid detection in most cases. It

is particularly vulnerable on the ground and it is often reluctant to descend unless this is unavoidable. Bushbabies may interact with a number of non-predatory species and the nature of some relationships is noted.

13. In conclusion the similarities and differences between the ecology and behaviour of G. crassicaudatus and G. senegalensis are discussed and the important factors which bring about their ecological separation are outlined. By making comparisons with other closely related species it is possible to speculate on the probable condition of the ancestral species and on the ways in which the recent forms have become specialized.

# LIST OF REFERENCES

- Altmann, S.A. 1962. A field study of the sociobiology of rhesus monkeys, Macaca mulatta. Ann. N.Y. Acad. Sci., 102 : 338 - 435.
- Altmann, S.A. 1965. Sociobiology of rhesus monkeys. II. Stochastics of social communication. J. Theoret. Biol., 8 : 490 - 522.
- Altmann, S.A. 1967. The structure of primate social communication. In S.A. Altmann (ed.), Social communication among primates. Chicago: Univ. Chicago Press. Pp. 325 - 62.
- Anderson, M. 1971. A watched Potto never grows : a cronicle of the pre-natal and first months of a Perodicticus potto. Discovery, 6 : 89 - 98.
- Andersson, A.B. 1969. Communication in the lesser bushbaby (Galago senegalensis moholi). Unpubl. M.Sc. Dissertation. Witwatersrand University.
- Andrew, R.J. 1962. The situations that evoke vocalization in primates. Ann. N.Y. Acad. Sci., 102 : 296 - 315.
- Andrew, R.J. 1963a. The origin and evolution of the calls and facial expressions of the primates. Behaviour, 20 : 1 - 109.
- Andrew, R.J. 1963b. Trends apparent in the evolution of vocalization in the Old World monkeys and apes. Symp. zool. Soc. Lond., 10 : 89 - 101.
- Andrew, R.J. 1964. The displays of the primates. In J. Buettner-Janusch (ed.), Evolutionary and genetic biology of primates. Vol. II. New York: Academic Press, Pp. 227 - 309.

- Andrew, R.J. & Klopman, R.B. 1974. Urine washing : comparative notes.  
In R.D. Martin, G.A. Doyle & A.C. Walker (eds.), Prosimian Biology.  
London : Duckworth.
- Astley-Maberly, C.T. 1967. The Game Animals of Southern Africa. Jo-  
hannesburg : Nelson.
- Bates, B.C. 1970. Territorial behavior in primates : a review of re-  
cent field studies. Primates, 11 : 271 - 4.
- Bearder, S.K. 1969. Territorial and intergroup behaviour of the lesser  
bushbaby, Galago senegalensis moholi (A. Smith), in semi-natural  
conditions and in the field. Unpubl. M.Sc. Dissertation. Witwaters-  
rand University.
- Bearder, S.K. & Doyle, G.A. 1974a. Ecology of bushbabies, Galago sene-  
galensis and Galago crassicaudatus, with some notes on their be-  
haviour in the field. In R.D. Martin, G.A. Doyle & A.C. Walker  
(eds.), Prosimian Biology. London : Duckworth.
- Bearder, S.K. & Doyle, G.A. 1974b. Field and laboratory studies of  
social organization in bushbabies (Galago senegalensis). Journal  
of Human Evolution, 3 : 37 - 50.
- Bishop, A. 1962. Control of the hand in lower primates. Ann. N.Y. Acad.  
Sci., 102 : 316 - 37.
- Bishop, A. 1964. Use of the hand in lower primates. In J. Buettner -  
Janusch (ed.), Evolutionary and genetic biology of the primates.  
Vol. 11. New York : Academic Press, Pp. 133 - 225.
- Blackwell, K. & Menzies, J.L. 1968. Observations on the biology of  
the potto (Perodicticus potto, Miller). Mammalia, 32 : 447 - 51.
- Bluntschli, H. 1933. In den Urwalden auf Madagascar. Umschau, 36 :  
769 - 1019 and 37 : 30 - 36.

- Buettner-Janusch, J. 1963. An introduction to the primates. In J. Buettner-Janusch (ed.), Evolutionary and genetic biology of the primates. Vol. I. New York : Academic Press, Pp. 1 - 64.
- Buettner-Janusch, J. 1964. The breeding of galagos in captivity and some notes on their behavior. Folia primat., 2 : 93 - 110.
- Buettner-Janusch, J. & Andrew, R.J. 1962. The use of the incisors by primates in grooming. Am. J. Phys. Anthropol., 20 : 127 - 9.
- Burt, W.H. 1943. Territoriality and home range concepts as applied to mammals. J. Mammal., 24 : 346 - 52.
- Butler, H. 1960. Some notes on the breeding cycle of the Senegal galago, Galago senegalensis senegalensis, in the Sudan. Proc. zool. Soc. Lond., 135 : 423 - 30.
- Cansdale, G.S. 1960. Bushbaby book. London : Phoenix House.
- Carpenter, C.R. 1934. A field study of the behavior and social relations of howling monkeys (Alouatta palliata). Comp. Psychol. Monogr., 10 : 1 - 168.
- Carpenter, C.R. 1965. The howlers of Barro Colorado Island. In I. DeVore (ed.), Primate Behavior : field studies of monkeys and apes. New York : Holt, Rinehart & Winston. Pp. 250 - 91.
- Charles-Dominique, P. & Martin, R.D. 1970. Evolution of lorises and lemurs. Nature, 227 : 257 - 60.
- Charles-Dominique, P. 1971. Eco-éthologie des prosimiens du Gabon. Biol. Gabonica, 7 : 121 - 228.
- Charles-Dominique, P. & Hladik, C.M. 1971. Le Lépilémur du Sud de Madagascar : écologie, alimentation et vie sociale. La Terre et la Vie, 25 : 3 - 66.

- Charles-Dominique, P. 1972. Ecologie et vie sociale de Galago demidovii (Fischer 1808; Prosimii). Z. Tierpsychol., Beiheft, 9 : 7 - 41.
- DeBoer, L.E.M. 1973a. Cytotaxonomy of the Lorisioidea (Primates : Prosimii). I. Chromosome studies and karyological relationships in the Galagidae. Genetica, 44 : 155 - 93.
- DeBoer, L.E.M. 1973b. Cytotaxonomy of the Lorisioidea (Primates : Prosimii). II. Chromosome studies in the Lorisidae and karyological relationships within the Superfamily. Genetica, 44 : 330 - 67.
- DeVore, I. 1963. Comparative ecology and behavior of monkeys and apes. In S.L. Washburn (ed.), Classification and human evolution : New York : Wenner-Gren Foundation. Pp. 301 - 19.
- DeVore, I. & Hall, K.R.L. 1965. Baboon ecology. In I. DeVore (ed.), Primate behavior : field studies of monkeys and apes. New York : Holt, Rinehart and Winston. Pp. 20 - 52.
- Dolhinow, P.J. & Bishop, N. 1970. The development of motor skills and social relationships among primates through play. In J.P. Hill (ed.), Minnesota symposia on child psychology. Vol. IV. Minneapolis : Univ. Minnesota Press.
- Doyle, G.A. 1974. Behavior of prosimians. In A.M. Schrier & F. Stollnitz (eds.), Behavior of non-human primates. Vol. V. New York : Academic Press.
- Doyle, G.A., Andersson, A. & Bearder, S.K. 1969. Maternal behaviour in the lesser bushbaby (Galago senegalensis moholi) under semi-natural conditions. Folia primat., 11 : 215 - 38.
- Doyle, G.A., Andersson, A. & Bearder, S.K. 1971. Reproduction in the lesser bushbaby (Galago senegalensis moholi) under semi-natural conditions. Folia primat., 14 : 15 - 22.

- Doyle, G.A. & Bearder, S.K. Naturalistic studies of the lesser bushbaby in the field. In Oehser (ed.), Research Reports - 1968 Projects. Washington : National Geographic Society. In press.
- Doyle, G.A. & Bekker, T. 1967. A facility for naturalistic studies of the lesser bushbaby (Galago senegalensis moholi). Folia primat., 7 : 161 - 8.
- Doyle, G.A., Pelletier, A. & Bekker, T. 1967. Courtship, mating and parturition in the lesser bushbaby (Galago senegalensis moholi) under semi-natural conditions. Folia primat., 7 : 169 - 97.
- Drews, D.R. 1971. Group formation in captive Galago crassicaudatus. Notes on the dominance concept. Research manuscript, Duke University.
- Eaton, G.G., Slob, A. & Resko, J.A. 1973. Cycles of mating behaviour in the Thick-tailed bushbaby (Galago crassicaudatus crassicaudatus) under laboratory conditions. Anim. Behav., 21 : 309 - 15.
- Ehrlich, A. 1968a. Food motivated behavior in prosimians. Folia primat., 8 : 66 - 71.
- Ehrlich, A. 1968b. Activity levels in prosimians. Folia primat., 8 : 72 - 6.
- Ehrlich, A. 1970. Response to novel objects in three lower primates : greater galago, slow loris and owl monkey. Behaviour, 37 : 54 - 63.
- Eibl-Eibesfeldt, I. 1953. Eine besondere Form des Duftmarkierens beim Riesen-Galago, Galago crassicaudatus (E. Geoffroy 1812). Z. Säugetierk., 1 : 171 - 73.
- Eisenberg, J.F., Muckenhirn, N.A. & Rudran, R. 1972. The relation between ecology and social structure in primates. Science, 176 : 863 - 74.
- Elliot-Smith, G. 1902. On the morphology of the brain in the mammalia, with special reference to the lemurs, recent and extinct. Trans. Linn. Soc. Lond., Ser. 2. zool., 8 : 319 - 432.

- Elliot-Smith, G. 1927. Essays on the evolution of man. 2nd Ed. Oxford : University Oxford Press.
- Fogden, M.P.L. 1974. A preliminary field-study of the Western Tarsier, Tarsioc bancanus Horsfield. In R.D. Martin, G.A. Doyle & A.C. Walker (eds.), Prosimian Biology. London : Duckworth.
- Frisch, J. 1959. Research on primate behavior in Japan. Am. Anthropol., 61 : 584 - 96.
- Gartlan, J.S. 1963. Structure and function in primate society. Folia primat., 8 : 89 - 120.
- Gérard, P. 1932. Etudes sur l'ovogénèse et l'ontogénèse chez les lémuriens du genre Galago. Archs. Biol., Paris, 43 : 93 - 151.
- Haagner, A.K. 1920. South African mammals. London : Witherby.
- Haddow, A.J. & Ellice, J.M. 1964. Studies on bushbabies (Galago spp.) with special reference to the epidemiology of yellow fever. Trans. roy. Soc. trop. Med. Hyg., 58 : 521 - 38.
- Hall-Craggs, E.C.B. 1965. An analysis of the jump of the lesser galago (Galago senegalensis). J. Zool., 147 : 20 - 9.
- Heinroth, O. 1911. Beiträge zur Biologie, namentlich Ethologie und Psychologie der Anatiden. Verh. V. Int. Ornith. Congr., Berlin : 589 - 702.
- Hill, W.C.O. 1938. A curious habit common to lorises and platyrrhine monkeys. Ceylon J. Sci. B. 21 : 65.
- Hill, W.C.O. 1953. Primates : comparative anatomy and taxonomy. I. Strepsirhini, Edinburgh : Univ. Edinburgh Press.
- Hockett, C.F. 1960. Logical considerations in the study of animal communication. In W.E. Lanyon & W.N. Tavalga (eds.) Animal sounds and communication. Washington : American Institute of Biological Sciences. Pp. 392 - 430.

- Hoof, J.A.R.A.M. van 1962. Facial expressions in higher primates. Symp. zool. Soc. Lond., 8 : 97 - 125.
- Hubel, D.H. & Wiesel, T.N. 1962. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. J. Physiol., 160 : 106 - 54.
- Huxley, J.S. 1914. The courtship habits of the Great Crested Grebe (Podiceps cristatus); with an addition to the theory of sexual selection.. Proc. zool. Soc. Lond., 1914 : 491 - 562.
- Ilse, D.R. 1955. Olfactory marking of territory in two young male loris, Loris tardigradus lydekkerianus, kept in captivity at Poona. Brit. J. Anim. Behav., 3 : 118 - 20.
- Immelman, W.F.E., Wicht, C.L. & Ackerman, D.P. 1973. Our green heritage. Cape Town : Tafelberg.
- Jay, P. 1965. Field studies. In A.M. Shrier, H.F. Harlow & F. Stollnitz (eds.), Behavior of non-human primates. Vol. II. New York : Academic Press, Pp. 525 - 91.
- Jay, P. 1963. Comments on studies of variabilities in species behavior. In P. Jay (ed.), Primates : studies in adaptation and variability. New York : Holt, Rinehart & Winston. Pp. 173 - 9.
- Jewell, P.A. 1963. The concept of home range in mammals. Symp. zool. Soc. Lond., 18 : 85 - 109.
- Jewell, P.A. & Oates, J.F. 1969. Ecological observations on the lorisoid primates of African lowland forest. Zoologica Africana, 4 : 231 - 48.
- Jolly, A. 1964. Prosimians manipulation of simple object problems. Anim. Behav., 12 : 560 - 70.
- Jolly, A. 1966a. Observations on Galago crassicaudatus. Unpubl. paper sent to Dr. Sauer.
- Jolly, A. 1966b. Lemur behavior : a Madagascar field study. Chicago : Univ. Chicago Press.

- Jolly, A. 1966c. Lemur social behavior and primate intelligence. Science, 153 : 501 - 6.
- Kaufmann, J.H. 1962. Ecology and social behavior of the coati, Nasua narica, on Barro Colorado Island, Panama. Univ. Calif. Publications Zoology, 60 : 95 - 222.
- Kingdon, J. 1971. East African Mammals. Vol. I. London - New York : Academic Press, Pp. 290 - 327.
- Lancaster, J.B. 1968. Primate communication systems and the emergence of human language. In P.C. Jay (ed.), Primates : studies in adaptation and variability. New York : Holt, Rinehart & Winston. Pp. 439 - 57.
- Lorenz, K. 1937. Über die Bildung des Instinkbegriffes. Naturwiss., 25 : 289 - 300, 307 - 18, 324 - 31.
- Lorenz, K. 1950. The comparative method in studying innate behaviour patterns. Symp. Soc. exp. Biol., 4 : 221 - 54.
- Lowther, F. de L. 1939. The feeding and grooming habits of the Galago. Zoologica, 25 : 477 - 80.
- Lowther, F. de L. 1940. A study of the activity of a pair of Galago senegalensis moholi in captivity, including the birth and post-natal development of twins. Zoologica, 25 : 433 - 62.
- Lumsden, W.H.R., Ellice, J.M., Mason, P.J. & Hewitt, L.E. 1955. Galago demidovii in the Entebbe area. E. Afr. Virus Res. Inst. Ann. Rep., No. 6 : 11 - 13.
- McKenna, M.C. 1966. Paleontology and the origin of the primates. Folia primat., 4 : 1 - 25.
- Manley, G.H. 1966. Reproduction in lorisoid primates. Symp. zool. Soc. Lond., 15 : 493 - 509.

- Marker, M.E. 1971. Karst landforms of the North-eastern Transvaal. Unpubl. Ph.D. Thesis. Witwatersrand University.
- Marler, P. 1961. The logical analysis of animal communication. J. Theoret. Biol., 1 : 295 - 317.
- Marler, P. 1965. Communication in monkeys and apes. In I. DeVore (ed.), Primate behavior : field studies of monkeys and apes. New York : Holt, Rinehart & Winston. Pp. 544 - 84.
- Marler, P. 1968. Aggregation and dispersal : two functions in primate communication. In P. Jay (ed.), Primates : studies in adaptation and variability. New York : Holt, Rinehart & Winston. Pp. 420 - 38.
- Martin, R.D. 1967. Behaviour and taxonomy of tree-shrews (Tupaiaidae). Unpubl. D. Phil. Thesis, University of Oxford.
- Martin, R.D. 1968. Reproduction and ontogeny in tree-shrews (Tupaia belangeri) with reference to their general behaviour and taxonomic relationships. Z. Tierpsychol., 25 : 409 - 532.
- Martin, R.D. 1972a. A preliminary field study of the Lesser Mouse Lemur (Microcebus murinus, J.F. Miller 1777). Z. Tierpsychol., Beiheft 9 : 43 - 89.
- Martin, R.D. 1972b. Adaptive radiation and behaviour of the Malagasy lemurs. Phil. Trans. R. Soc. Lond. B, 264 : 295 - 352.
- Masefield, G.B., Wallis, M., Harrison, S.G. & Nicholson, B.E. 1969. The Oxford book of food plants. Oxford : Oxford Univ. Press.
- Mason, W.A. 1966. Social organization of the South American monkey, Calli-  
cebus molloch : a preliminary report. Tul. Studies in Zool., 13 : 23 - 28.
- Mason, W.A. 1968. Use of space by Calli-  
cebus molloch. In P.C. Jay (ed.), Primates : studies in adaptation and variability. New York : Holt, Rinehart & Winston. Pp. 200 - 16.

- Matschie, P. 1905. Otolemur badius, eine neue Art afrikanischer Halbaffen. Sber. Ges. naturf. Freund Berl.; 8 : 277 - 9.
- Montagna, W. 1962. The skin of lemurs. Ann. N.Y. Acad. Sci., 102 : 190 - 209.
- Montagna, W. & Ellis, R.A. 1959. The skin of primates. I. The skin of the potto (Perodicticus potto). Am. J. phys. Anthrop., 17 : 137.- 62.
- Montagna, W. & Ellis, R.A. 1960. The skin of primates. II. The skin of the slender loris (Loris tardigradus). Am. J. phys. Anthrop., 18 : 19 - 44.
- Montagna, W. & Yun, J.S. 1962a. The skin of primates. VII. The skin of the greater bushbaby (Galago crassicaudatus). Am. J. phys. Anthrop., 20 : 149 - 66.
- Montagna, W. & Yun, J.S. 1962b. The skin of primates. X. The skin of the ring-tailed lemur (Lemur catta). Am. J. phys. Anthrop., 20 : 95 - 118.
- Moynihan, M. 1964. Some behavior patterns of Platyrrhine monkeys. I. The night monkey (Aotus trivigatus). Smithson. Misc. Coll., 146 : 1 - 84.
- Moynihan, M. 1967. Comparative aspects of communication in New World primates. In D. Morris (ed.), Primate ethology. London : Weidenfeld & Nicolson. Pp. 236 - 66.
- Napier, J.R. 1971. The roots of man. London : George Allen and Unwin Ltd.
- Napier, J.R. & Walker, A.C. 1967. Vertical clinging and leaping - a newly recognised category of locomotor behaviour in primates. Folia primat., 6 : 204 - 19.
- Pariente, G. 1974. Influence of light on the activity rhythms of two Malagasy lemurs : Phaner furcifer and Lepilemur mustelinus leuconus. In R.D. Martin, G.A. Doyle & A.C. Walker (eds.), Prosimian Biology. London : Duckworth.

- Petter, J-J. 1962a. Ecological and behavioral studies on Madagascar lemurs in the field. Ann. N.Y. Acad. Sci., 102(2) : 267 - 81.
- Petter, J-J. 1962b. Recherches sur l'écologie et l'éthologie des lémuriens, Malgaches. Mém. du Mus. Nat.d'Hist. Nat. Ser. A (Zoologie.), 27(1) : 1 - 146.
- Petter, J-J. 1962c. Remarques sur l'écologie et l'éthologie comparées des lémuriens Malgaches. La Terre et la Vie, 4 : 394 - 416.
- Petter, J-J. 1974. A study of population density and home range of Indri in Madagascar. In R.D. Martin, G.A. Doyle and A.C. Walker (eds.), Prosimian Biology. London : Duckworth.
- Petter, J-J. & Hladik, C.M. 1970. Observation sur le domaine vital et la densité de populations de Loris tardigradus dans les forêts de Ceylan. Mammalia, 3 : 394 - 409.
- Petter, J-J. & Petter-Rousseaux, A. 1959. Contribution à l'étude du Aye-aye. Naturaliste Malgache, 11 : 153 - 64.
- Petter, J-J. & Petter-Rousseaux, A. 1964. Première tentative d'estimation des densités de peuplement de lémuriens Malgaches. La Terre et la Vie, 4 : 427 - 35.
- Petter, J-J. & Petter-Rousseaux, A. 1967. The Aye-aye of Madagascar. In S.A. Altmann (ed.), Social communication among primates. Chicago : Univ. Chicago Press. Pp. 195 - 205.
- Petter, J-J. & Peyrieras, A. 1970. Observations éco-éthologiques sur les lémuriens Malgaches du genre Hapalemur, La Terre et la Vie, 24 : 356 - 82.
- Petter, J-J., Schilling, A. & Pariente, G. 1971. Observations éco-éthologiques sur deux lémuriens Malgaches nocturnes : Phaner furcifer et Microcebus coquereli. La Terre et la Vie, 25 : 287 - 327.

- Pinto, D. 1972. Patterns of activity in three nocturnal prosimians : Galago senegalensis moholi, Galago crassicaudatus umbrosus and Microcebus murinus murinus. Unpubl. M.Sc. Thesis. Witwatersrand University.
- Rahaman, H. & Parthasarathy, M.D. 1970. Observations on the habits and life span of the slender loris, Loris tardigradus lydekkerianus (L.), Myforest, 6(4): 23 - 6.
- Ralls, K. 1971. Mammalian scent marking. Science, 171 : 443 - 9.
- Randolph, M. 1971. Role of light and circadian rhythms in the nocturnal behavior of Galago crassicaudatus. J. Comp. Physiol. Psych., 74 (1) : 115 - 22.
- Roberts, A. 1951. The mammals of South Africa. South Africa : Central News Agency.
- Roberts, P. 1971. Social interactions of Galago crassicaudatus. Folia primat., 14 : 171 - 81.
- Rosenblum, L.A., Kaufmann, I.C. & Steynes, A.J. 1966. Some characteristics of adult social and autogrooming patterns in two species of macaque. Folia primat., 4 : 438 - 51.
- Rowell, T.E. 1962. Agonistic noises of the rhesus monkey (Macaca mulatta). Symp. zool. Soc. Lond., 8 : 91 - 6.
- Rowell, T.E. & Hinde, R.A. 1962. Vocal communication by the rhesus monkey (Macaca mulatta). Proc. zool. Soc. Lond., 138 : 279 - 94.
- Sanderson, I.T. 1940. The mammals of the North Cameroons forest area. Trans. zool. Soc. Lond., 24 : 623 - 725.
- Sanderson, I.T. 1957. The monkey kingdom. New York : Doubleday.
- Sauer, E.G.F. & Sauer, E.M. 1963. The South West African bushbaby of the Galgo senegalensis group. J. S.W. Africa Scient. Soc., Windhoek, 16 : 5 - 36.

- Schilling, A. 1974. A study of marking behaviour in !emur catta. In R.D. Martin, G.A. Doyle & A.C. Walker (eds.), Prosimian Biology. London : Duckworth.
- Schultz, A.H. 1969. The life of primates. London : Weidenfeld & Nicolson.
- Schwartz, E. 1931. On the African long-tailed lemurs or galagos. Ann. Mag. nat. Hist., 10 : 41 - 67.
- Seitz, E. 1967. Untersuchungen zur Aktivitätsrhythmik dunkelaktiver Halbaffen der Unterfamilie Lorissinae (Flower & Lydekker 1891). In : Starck, Schneider & Kuhn. (eds.), Proc. 1st. Congress of the int. Primat. Soc., Pp. 322 - 6.
- Shortridge, G.D. 1934. The mammals of South West Africa. Vol. 1. London : Heinemann.
- Simons, E.L. 1962. Fossil evidence relating to the early evolution of primate behavior. Ann. N.Y. Acad. Sci., 102 : 282 - 94.
- Simpson, G.G. 1945. The principles of classification and a classification of mammals. Bull. Am. Mus. nat. Hist., 85 : 1 - 350.
- Smithers, R.H.N. 1966. The mammals of Rhodesia, Zambia and Malawi. London : Collins.
- Smithers, R.H.N. 1968. A check list and atlas of the mammals of Botswana. Salisbury : The Trustees of the National Museums of Rhodesia.
- Southern, H.N. 1955. Nocturnal animals. Sci. Am. 193(4) : 88 - 98.
- Struhsaker, T.T. 1967a. Auditory communication among vervet monkeys (Cercopithecus aethiops). In S.A. Altmann (ed.), Social communication among primates. Chicago : Univ. Chicago Press. Pp. 281 - 324.
- Struhsaker, T.T. 1967b. Ecology of vervet monkeys (Cercopithecus aethiops) in the Masai-Amboseli Game reserve, Kenya. Ecology, 48(6): 891 - 904.

- Struhsaker, T.T. 1970. Notes on Galagoides demidovii in Cameroon. Mammalia, 34 : 207 - 11.
- Tansley, K. 1965. Vision in vertebrates. London : Chapman and Hall Ltd.
- Tenaza, R., Ross, B.A., Tanticharoenyos, P. & Berkson, G. 1969. Individual behaviour and activity rhythms of captive slow lorises (Nycticebus coucang). Anim. Behav., 17 : 664 - 9.
- Terry, R.L. 1970. Primate grooming as a tension reduction mechanism. J. Psychol., 76 : 129 - 36.
- Tinbergen, N. 1951. The study of instinct. Oxford : Oxford Univ. Press.
- Tinbergen, N. 1963. On aims and methods of ethology. Z. Tierpsychol., 20 : 410 - 29.
- Turnbull-Kemp, P. 1967. The leopard. Cape Town : Howard Timmins.
- Ullrich, W. 1961. Zur Biologie und Soziologie der Colobusaffen (Colobus guereza caudatus Thomas 1885). Der Zool. Garten, 25(6) : 305 - 68.
- Vincent, F. 1969. Contribution à l'étude des prosimiens Africains. Le Galago de Demidoff : reproduction (biology, anatomy, physiology) et comportement. Unpubl. Ph.D. Thesis. Paris University.
- Walker, A.C. 1967a. Locomotor adaptation in recent and fossil Madagascan lemurs. Unpubl. Ph.D. Thesis. London University.
- Walker, A.C. 1967b. Pattern of extinction among the subfossil Madagascan lemuroids. In P.S. Martin & H.E. Wright (eds.), Pleistocene extinctions : the search for a cause. Pp. 425 - 32.
- Walker, A.C. 1969. The locomotion of the lorises with special reference to the potto. E. Afr. wildl. J., 7 : 1 - 5.
- Walker, A.C. 1970. Post-cranial remains of the Miocene Lorisidae of East Africa. Am. J. phys. Anthrop., 33 : 249 - 61.

- Walker, A.C. 1974. A review of the Miocene Lorisidae of East Africa. In R.D. Martin, G.A. Doyle & A.C. Walker (eds.), Prosimian Biology. London : Duckworth.
- Walker, E.P. 1954. The monkey book. New York : Macmillan.
- Washburn, S.L. & Hamburg, D.A. 1965. The study of primate behavior. In I. DeVore (ed.), Primate behavior : field studies of monkeys and apes. New York : Holt, Rinehart & Winston. Pp. 1 - 13.
- Whitman, C.O. 1919. The behaviour of pigeons. Carnegie Inst. Wash. Publ., 257(3) : 1 - 161.
- Ying, K.L. & Butler, H. 1971. Chromosome polymorphism in the lesser bushbabies (Galago senegalensis). Can. J. Genet. Cytol., 13 : 793 - 800.

## APPENDICES

Appendix 1. Mean temperatures ( $t, ^\circ\text{C}$ ), precipitation ( $p, \text{mm}$ ) and relative humidity ( $h, \%$ ) for the study areas of a) G.c. umbrosus b) G.c. garnetti and c) G.c. lönnberqi

|      | Month | $t, ^\circ\text{C}$ |      | $p, \text{mm}$ | $h, \%$<br>0800 hrs. |
|------|-------|---------------------|------|----------------|----------------------|
|      |       | Max.                | Min. |                |                      |
| (a)  | J     | 29.1                | 17.7 | 352            | 82                   |
|      | F     | 28.6                | 17.7 | 307            | 85                   |
|      | M     | 27.7                | 16.4 | 280            | 88                   |
|      | A     | 26.8                | 13.4 | 95             | 87                   |
|      | M     | 25.1                | 8.4  | 44             | 86                   |
|      | J     | 23.2                | 5.1  | 20             | 85                   |
|      | -     |                     |      |                |                      |
|      | J     | 23.1                | 5.0  | 29             | 83                   |
|      | A     | 24.8                | 6.8  | 28             | 79                   |
|      | S     | 27.0                | 10.0 | 49             | 73                   |
|      | O     | 28.8                | 13.4 | 109            | 73                   |
|      | N     | 29.0                | 15.4 | 189            | 78                   |
|      | D     | 29.8                | 16.8 | 266            | 80                   |
| YEAR |       | 26.8                | 12.2 | 1768           | 80                   |

|     | Month | t, °C |      | p, mm | h, %<br>0800 hrs. |
|-----|-------|-------|------|-------|-------------------|
|     |       | Max.  | Min. |       |                   |
| (b) | J     | 27.7  | 17.6 | 127   | 79                |
|     | F     | 28.0  | 18.1 | 145   | 85                |
|     | M     | 26.6  | 17.3 | 203   | 88                |
|     | A     | 25.4  | 15.5 | 121   | 88                |
|     | M     | 23.9  | 13.3 | 103   | 85                |
|     | J     | 22.2  | 11.4 | 98    | 85                |
|     | -     |       |      |       |                   |
|     | J     | 22.0  | 11.0 | 75    | 84                |
|     | A     | 23.3  | 11.8 | 59    | 84                |
|     | S     | 24.2  | 13.2 | 63    | 85                |
|     | O     | 24.9  | 14.6 | 81    | 83                |
|     | N     | 25.9  | 15.9 | 106   | 81                |
|     | D     | 27.4  | 17.3 | 111   | 79                |
|     | YEAR  | 25.1  | 15.2 | 1292  | 81                |
| (c) | J     | 26.9  | 17.5 | 34    | 89                |
|     | F     | 26.1  | 16.1 | 23    | 89                |
|     | M     | 27.1  | 15.9 | 25    | 83                |
|     | A     | 26.6  | 14.5 | 19    | 83                |
|     | M     | 22.7  | 9.5  | 78    | 77                |
|     | J     | 20.6  | 8.0  | 05    | 79                |
|     | -     |       |      |       |                   |
|     | J     | 21.5  | 7.5  | 02    | 69                |
|     | A     | 24.3  | 10.0 | 11    | 68                |
|     | S     | 26.3  | 11.4 | 10    | 60                |
|     | O     | 26.6  | 14.5 | 05    | 70                |
|     | N     | 26.2  | 16.1 | 45    | 85                |
|     | D     | 27.5  | 16.6 | 41    | 83                |
|     | YEAR  | 25.2  | 13.1 | 298   | 78                |

Appendix 2. Scientific names for exotic trees and fruits referred to in the text. (For detailed descriptions and illustrations see Masfield et al., 1969 and Immelman et al., 1973.)

| <u>Common name(s)</u>              | <u>Species</u>                 |
|------------------------------------|--------------------------------|
| Anonaceous fruits - Custard apples | <u>Anona spp.</u>              |
| Avocado pear                       | <u>Persea americana</u>        |
| Banana                             | <u>Musa spp.</u>               |
| Bluegum                            | <u>Eucalyptus globulus</u>     |
| Cherry guava - Strawberry guava    | <u>Psidium cattleianum</u>     |
| Fig                                | <u>Ficus carica</u>            |
| Grapevines                         | <u>Vitis spp.</u>              |
| Guava                              | <u>Psidium guajava</u>         |
| Indian mahogany                    | <u>Cedrela toona</u>           |
| Jacaranda                          | <u>Jacaranda mimosaeifolia</u> |
| Litchi                             | <u>Litchi chinensis</u>        |
| Mango                              | <u>Mangifera indica</u>        |
| Mulberry                           | <u>Morus nigra</u>             |
| Papaya - Paw paw                   | <u>Carica papaya</u>           |
| Passion fruit - Purple granadilla  | <u>Passiflora edulis</u>       |
| Pines                              | <u>Pinus spp.</u>              |
| Wattle (Australian)                | <u>Acacia cyanophylla</u>      |

Appendix 3. Predominant indigenous trees and climbers of the riparian bush in the North Eastern Transvaal showing known food trees (+) and those bearing fleshy fruits (+).

|                                  |                                     |
|----------------------------------|-------------------------------------|
| <u>Acacia ataxacantha</u> +      | <u>Cussonia spicata</u> +           |
| <u>Acacia karoo</u> +            | <u>Dalbergia armata</u>             |
| <u>Adina microcephala</u>        | <u>Diospyros maspiliiformis</u> + + |
| <u>Albizia versicolor</u> +      | <u>Dombeya rotundifolia</u>         |
| <u>Andrachne ovalis</u>          | <u>Ekebergia capensis</u> + +       |
| <u>Annona senegalensis</u> +     | <u>Encephalartos transvenosus</u> + |
| <u>Antidesma venosum</u> +       | <u>Erythrina lysistemon</u>         |
| <u>Apodytes dimidiata</u>        | <u>Fagara davyi</u>                 |
| <u>Bauhinia galpini</u>          | <u>Ficus capensis</u> + +           |
| <u>Bequaertiodendron</u>         | <u>Ficus ingens</u> +               |
| <u>megalismontanum</u> +         | <u>Ficus petersii</u> +             |
| <u>Brachylaena discolor</u>      | <u>Ficus sonderi</u> +              |
| <u>Bridelia micrantha</u> + +    | <u>Ficus stuhlmanni</u> + +         |
| <u>Bridelia mollis</u> +         | <u>Ficus sycamorus</u> + +          |
| <u>Canthium ventosum</u> +       | <u>Gardenia amoena</u> +            |
| <u>Canthium mundianum</u> +      | <u>Gardenia jovis-tonantis</u> +    |
| <u>Carissa bispinosa</u> +       | <u>Gardenia spatulifolia</u> +      |
| <u>Cassia spp.</u>               | <u>Grewia occidentalis</u>          |
| <u>Catha edulis</u>              | <u>Halleria lucida</u> +            |
| <u>Celtis agricana</u> +         | <u>Harpephyllum caffrum</u> +       |
| <u>Cephalanthus natalensis</u> + | <u>Heteropyxis natalensis</u>       |
| <u>Chaetachme aristata</u>       | <u>Lannea discolor</u> +            |
| <u>Combretum kraussii</u>        | <u>Lannea edulis</u> +              |
| <u>Combretum collinum</u>        | <u>Lonchocarpus capassa</u>         |
| <u>Combretum molle</u>           | <u>Maesa lanceolata</u>             |
| <u>Combretum zeyheri</u>         | <u>Maytenus heterophylla</u>        |
| <u>Croton sylvestris</u>         | <u>Mimusops zeyheri</u> + +         |
| <u>Cryptocarya liebertiana</u> + | <u>Mimusops ovata</u> +             |
| <u>Curtisia dendata</u>          | <u>Ocina atropurpurea</u>           |

Ochna holstii  
Parinaria curatellifolia +  
Pavetta scumanniana  
Piliostigma thonningii  
Protorhus longifolia +  
Pterolobium exosum  
Pterocarpus angolensis  
Pterocarpus rotundifolius  
Rapanea melanophloeos +  
Rauvolfia caffra +  
Rawsonia lucida +  
Rhoicissus tomentosa +  
Rhus chirindensis +  
Rothmannia capensis +  
Sclerocarya caffra +  
Scolopia zeyheri +

Scotia brachypetala +  
Scheffiera umbellifera +  
Strychnos spinosa +  
Strychnos innocua  
Strychnos mitis +  
Strychnos usambarensis +  
Syzygium cordatum + +  
Syzygium gerrardii +  
Syzygium guineense +  
Terminalia sericea  
Tapiphyllum parvifolium +  
Trema orientalis +  
Trichilia emetica  
Vangueria infausta +  
Ximenia caffra +

|                                   |                                  |
|-----------------------------------|----------------------------------|
| <u>Ochna holstii</u>              | <u>Scotia brachypetala</u> +     |
| <u>Parinaria curatellifolia</u> + | <u>Scheffiera umbellifera</u> +  |
| <u>Pavetta scumanniana</u>        | <u>Strychnos spinosa</u> +       |
| <u>Piliostigma thonningii</u>     | <u>Strychnos innocua</u>         |
| <u>Protorhus longifolia</u> +     | <u>Strychnos mitis</u> +         |
| <u>Pterolobium oxosum</u>         | <u>Strychnos usambarensis</u> +  |
| <u>Pterocarpus angolensis</u>     | <u>Syzygium cordatum</u> + +     |
| <u>Pterocarpus rotundifolius</u>  | <u>Syzygium gerrardii</u> +      |
| <u>Rapanea melanophloeos</u> +    | <u>Syzygium guineense</u> +      |
| <u>Rauvolfia caffra</u> +         | <u>Terminalia sericea</u>        |
| <u>Rawsonia lucida</u> +          | <u>Tapiphyllum parvifolium</u> + |
| <u>Rhoicissus tomentosa</u> +     | <u>Trema orientalis</u> +        |
| <u>Rhus chirindensis</u> +        | <u>Trichilia emetica</u>         |
| <u>Rothmannia capensis</u> +      | <u>Vangueria infausta</u> +      |
| <u>Sclerocarya caffra</u> +       | <u>Ximenia caffra</u> +          |
| <u>Scolopia zeyheri</u> +         |                                  |

Appendix 4. Predominant indigenous trees and climbers of the coastal dune forest at Mtunzini in Zululand showing plants bearing fleshy fruits or gum (+).

|                                   |                                |
|-----------------------------------|--------------------------------|
| <u>Acacia sieberiana</u> +        | <u>Dalbergia obovata</u>       |
| <u>Acokanthera oblongifolia</u> + | <u>Deinbollia oblongifolia</u> |
| <u>Albizia adianthifolia</u>      | <u>Dichrostachys cinerea</u>   |
| <u>Albizia versicolor</u> +       | <u>Dovyalis rhamnoides</u>     |
| <u>Aloe bainesii</u>              | <u>Dovyalis zeyheri</u> +      |
| <u>Allophylus natalensis</u>      | <u>Drypetes natalensis</u>     |
| <u>Annona senegalensis</u> +      | <u>Ekebergia capensis</u> +    |
| <u>Andidesma venosum</u> +        | <u>Erythrina lysistemon</u>    |
| <u>Apodytes dimidiata</u> +       | <u>Euclea natalensis</u> +     |
| <u>Barringtonia racemosa</u>      | <u>Eugenia zuluensis</u> +     |
| <u>Bersama lucens</u>             | <u>Fagara capensis</u> +       |
| <u>Brachylena discolor</u>        | <u>Ficus capensis</u> +        |
| <u>Bridelia micrantha</u> +       | <u>Ficus burtt-davyi</u> +     |
| <u>Burchellia bubalina</u>        | <u>Ficus hippopotami</u> +     |
| <u>Calpurnia aurea</u>            | <u>Ficus natalensis</u> +      |
| <u>Canthium obovatum</u> +        | <u>Grewia occidentalis</u>     |
| <u>Canthium ventosum</u>          | <u>Halleria lucida</u> +       |
| <u>Carissa bispinosa</u>          | <u>Harpephyllum caffrum</u> +  |
| <u>Carissa macrocarpa</u>         | <u>Hibiscus tiliaceus</u>      |
| <u>Cassine aethiopica</u>         | <u>Hyphaene crinita</u>        |
| <u>Cassine papillosa</u>          | <u>Ilex mitis</u>              |
| <u>Clausena anisata</u>           | <u>Macaranga capensis</u>      |
| <u>Clerodendrum glabrum</u>       | <u>Maesa lanceolata</u>        |
| <u>Colpoon compressum</u>         | <u>Maytenus heterophylla</u>   |
| <u>Cordia caffra</u>              | <u>Maytenus undata</u>         |
| <u>Cussonia chartacea</u>         | <u>Mimusops caffra</u> +       |
| <u>Cussonia kraussii</u>          | <u>Myrica serrata</u>          |
| <u>Cussonia spicata</u>           | <u>Ochna arborea</u>           |
| <u>Cyathea dregei</u>             | <u>Ochna natalita</u>          |
| <u>Dalbergia armata</u>           | <u>Olea woodiana</u> +         |

Passerina falcifolia +  
Peddiea africana +  
Phoenix reclinata +  
Popowia obovata  
Protorhus longifolia +  
Psycotria capensis +  
Rapanea melanophloeos  
Raphia australis  
Rauvolfia caffra +  
Rhus macowanii  
Sapium ellipticum  
Sapium integerrimum  
Scheffiera umbellifera  
Sclerocarya caffra +  
Scolopia zeyheri +  
Scutia myrtina +

Sideroxylon inerme +  
Strelitzia nicolai  
Strychnos innocua +  
Strychnos spinosa +  
Syzygium cordatum +  
Trema orientalis +  
Tricalysia sonderiana  
Trichilia dregeana +  
Trimeria rotundifolia  
Turraea floribunda  
Vangueria infausta +  
Vepris undulata +  
Voacanga thouarsii  
Xeromphis obovata  
Ximenia caffra

Appendix 5. Predominant indigenous trees and shrubs of the woodlands near Umtali, Rhodesia.

|                              |                                 |
|------------------------------|---------------------------------|
| <u>Acacia abyssinica</u>     | <u>Encephalartos manikensis</u> |
| <u>Acacia campylacantha</u>  | <u>Ficus capensis</u>           |
| <u>Acacia karoo</u>          | <u>Ficus sycomorus</u>          |
| <u>Acacia sieberiana</u>     | <u>Ficus quibeba</u>            |
| <u>Albizzia gummifera</u>    | <u>Fluggea microcarpa</u>       |
| <u>Annona cuneata</u>        | <u>Gardenia imperialis</u>      |
| <u>Annona senegalensis</u>   | <u>Grewia caffra</u>            |
| <u>Antidesma venosum</u>     | <u>Maytenus spp.</u>            |
| <u>Azanza garckeana</u>      | <u>Mimusops zeyheri</u>         |
| <u>Bauhinia galpinii</u>     | <u>Olea africana</u>            |
| <u>Bauhinia petersiana</u>   | <u>Pavetta assimilis</u>        |
| <u>Bolusanthus speciosus</u> | <u>Peltophorum africanum</u>    |
| <u>Bridelia micrantha</u>    | <u>Phoenix reclinata</u>        |
| <u>Calodendrum capense</u>   | <u>Pittosporum viridiflorum</u> |
| <u>Carissa edulis</u>        | <u>Rhus longipes</u>            |
| <u>Cassia capensis</u>       | <u>Rothmannia manganjae</u>     |
| <u>Cassia floribunda</u>     | <u>Sapium ellipticum</u>        |
| <u>Celtis africana</u>       | <u>Scheffiera umbellifera</u>   |
| <u>Chaetacme aristata</u>    | <u>Spathodea campanulata</u>    |
| <u>Clerodendrum glabrum</u>  | <u>Steganotaenia araliacea</u>  |
| <u>Combretum mollii</u>      | <u>Synadenium sp.</u>           |
| <u>Craibia brevicaudata</u>  | <u>Syzigium cordatum</u>        |
| <u>Cussonia kirkii</u>       | <u>Teclea nobilis</u>           |
| <u>Cussonia natalensis</u>   | <u>Trichelea emetica</u>        |
| <u>Cussonia spicata</u>      | <u>Vernonia amygdalina</u>      |
| <u>Diospyros lycioides</u>   |                                 |
| <u>Diospyros sericea</u>     |                                 |

**Author** Bearder Simon Kenneth

**Name of thesis** Aspects Of The Ecology And Behaviour Of The Thick-tailed Bushbaby, Galago Crassicaudatus. 1974

***PUBLISHER:***

University of the Witwatersrand, Johannesburg

©2013

***LEGAL NOTICES:***

**Copyright Notice:** All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed, or otherwise published in any format, without the prior written permission of the copyright owner.

**Disclaimer and Terms of Use:** Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.