



Check kids' blood pressure to prevent hypertension in later life

Adapted Media Release

Published: Tuesday 18 August 2015

3 SHARE

The road to **hypertension** in adulthood might start as early as in childhood and adolescence in the urban black South African population, says a Wits-lead research team.

"Approximately one-third to a half of the children who were hypertensive at some time during childhood and adolescence were hypertensive at age 18 years. The risk of having elevated **blood pressure** at 18 years of age was lowest at age five years and highest at age 14 years," the researchers reported.

Researchers from the MRC/WITS Developmental Pathways for Health Research Unit (DPHRU) and the Department of Nutrition, University of North Carolina in the USA published a study, entitled Blood pressure tracking in urban black South African children: birth to twenty cohort in the *BMC Paediatrics*.

Juliana Kagura (PhD student at DPHRU), Professor Linda Adair (University of North Carolina and honorary Professor at Wits), Dr Mogi Musa (Postdoctoral fellow at DPHRU), Professor Shane Norris (DPHRU) and Wits Emeritus Professor John Pettifor collaborated on this paper

The researchers said that hypertension is a rapidly increasing global public health problem that disproportionately affects low and middle income countries. South Africa has the highest prevalence of hypertension in adults (78%), according to a recent global report from the World Health Organization - Strategic Advisory Group of Experts collaborative study.

The study is said to be the first in South Africa to show that elevated blood pressure in childhood and adolescence, classified according to blood pressure charts for age, sex and height, significantly persists into early adulthood.

Using birth to twenty cohort data of children born in Soweto, Johannesburg in 1990, investigators reported that approximately a third to half of the participants of black ethnicity with elevated blood pressure at one occasion between childhood and adolescence had sustained the elevated blood pressure status at 18 years of age.

"This work may suggest the importance of routine blood pressure measurement in children for early identification of at-risk children, which may inform timely interventions to prevent complications associated with elevated blood pressure," the researchers cautioned.

References

Blood pressure tracking in urban black South African children: birth to twenty cohort, Juliana Kagura, Linda S Adair, Mogi G Musa, John M Pettifor and Shane A Norris, *BMC Paediatrics*, doi:10.1186/s12887-015-0402-z, published 15 July 2015.

Source: Wits University

This page was printed from: <http://www.medicalnewstoday.com/releases/298299.php>

Visit www.medicalnewstoday.com for medical news and health news headlines posted throughout the day, every day.

© 2004-2016 All rights reserved. MNT is the registered trade mark of MedLexicon International Limited.