

Constructing evidence-based clinical intrapartum care algorithms for decision-support tools

M Bonet,^a  L Ciabati,^b LL De Oliveira,^b R Souza,^c  JL Browne,^d M Rijken,^d  S Fawcus,^e GJ Hofmeyr,^{f,g}  T Liabsuetrakul,^h  Ç Gülümser,ⁱ  A Blennerhassett,^{j,k} D Lissauer,^{l,m}  S Meher,ⁿ F Althabe,^a  OT Oladapo,^a  on behalf of the WHO Intrapartum Care Algorithms Working Group*

^a UNDP/UNFPA/UNICEF/WHO/World Bank Special Program of Research, Development and Research Training in Human Reproduction (HRP), Department of Sexual and Reproductive Health and Research, World Health Organization, Geneva, Switzerland ^b Department of Social Medicine, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil ^c Department of Obstetrics and Gynaecology, School of Medicine, University of Campinas (UNICAMP), Campinas, SP, Brazil ^d Julius Global Health, Julius Centre for Health Sciences and Primary Care, University Medical Centre, Utrecht University, Utrecht, The Netherlands ^e Department of Obstetrics and Gynaecology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa ^f Effective Care Research Unit, Walter Sisulu University and Eastern Cape Department of Health, University of the Witwatersrand, East London, South Africa ^g Department of Obstetrics and Gynaecology, University of Botswana, Gaborone, Botswana ^h Epidemiology Unit, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla, Thailand ⁱ Department of Obstetrics and Gynaecology, University of Health Science School of Medicine, Ankara, Turkey ^j Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK ^k World Health Organization Collaborating Centre for Global Women's Health Research, Birmingham, UK ^l Malawi-Liverpool-Wellcome Trust Research Institute, Queen Elizabeth Central Hospital, College of Medicine, Blantyre, Malawi ^m Institute of Life Course and Medical Sciences, William Henry Duncan Building, University of Liverpool, Liverpool, UK ⁿ Birmingham Women's and Children's NHS Foundation Trust, Birmingham, UK
Correspondence: M Bonet, Department of Sexual and Reproductive Health and Research, World Health Organization, Avenue Appia 20, Geneva 27, CH-1211 Switzerland. Email: bonetm@who.int

Accepted 17 September 2021. Published Online 11 April 2022.

Aim To describe standardised iterative methods used by a multidisciplinary group to develop evidence-based clinical intrapartum care algorithms for the management of uneventful and complicated labours.

Population Singleton, term pregnancies considered to be at low risk of developing complications at admission to the birthing facility.

Setting Health facilities in low- and middle-income countries.

Search strategy Literature reviews were conducted to identify standardised methods for algorithm development and examples from other fields, and evidence and guidelines for intrapartum care. Searches for different algorithm topics were last updated between January and October 2020 and included a combination of terms such as 'labour', 'intrapartum', 'algorithms' and specific topic terms, using Cochrane Library and MEDLINE/PubMED, CINAHL, National Guidelines Clearinghouse and Google.

Case scenarios Nine algorithm topics were identified for monitoring and management of uncomplicated labour and

childbirth, identification and management of abnormalities of fetal heart rate, liquor, uterine contractions, labour progress, maternal pulse and blood pressure, temperature, urine and complicated third stage of labour. Each topic included between two and four case scenarios covering most common deviations, severity of related complications or critical clinical outcomes.

Conclusions Intrapartum care algorithms provide a framework for monitoring women, and identifying and managing complications during labour and childbirth. These algorithms will support implementation of WHO recommendations and facilitate the development by stakeholders of evidence-based, up to date, paper-based or digital reminders and decision-support tools. The algorithms need to be field tested and may need to be adapted to specific contexts.

Keywords Algorithms, childbirth, first stage of labour, intrapartum care, labour complications, second stage of labour, third stage of labour.

Tweetable abstract Evidence-based intrapartum care clinical algorithms for a safe and positive childbirth experience.

Please cite this paper as: Bonet M, Ciabati L, De Oliveira LL, Souza R, Browne JL, Rijken M, Fawcus S, Hofmeyr GJ, Liabsuetrakul T, Gülümser Ç, Blennerhassett A, Lissauer D, Meher S, Althabe F, Oladapo OT; WHO Intrapartum Care Algorithms Working Group. Constructing evidence-based clinical intrapartum care algorithms for decision-support tools. BJOG 2024;131(S2):6–16.

*A list of the WHO Intrapartum Care Algorithms Working Group members can be found in the Appendix.

Introduction

The burden of maternal and perinatal deaths and severe morbidity is high around the time of childbirth, representing over a third of maternal deaths,^{1,2} half of all stillbirths and a quarter of neonatal deaths,³ particularly in low- and middle-income countries.^{2,3}

Investment in high-quality care around the time of childbirth has been identified as the most impactful strategy to reduce maternal and perinatal deaths and severe morbidities.^{4,5} The appropriate monitoring of women and babies throughout labour and childbirth, early identification of those at risk of labour complications, timely use of safe and effective interventions when needed and avoidance of unnecessary labour interventions would avert most of the preventable intrapartum-related adverse outcomes. High-quality intrapartum care is also key to optimising the wellbeing of women and their babies and their experience of care.⁶

In the last decade the World Health Organization (WHO) has released and continuously updated⁷ several evidence-based recommendations addressing the care that all women and their babies should receive during uneventful or complicated labours and childbirth. However, it is well known that passive dissemination of recommendations is insufficient for its optimal implementation.^{8,9} Different strategies have been used to ensure that high-quality, evidence-based practices are implemented in different contexts, such as the use of reminders and decision/job aids, in the form of posters, checklists, screening tools and flowcharts/algorithms to assist health workers in making diagnosis and clinical management decisions.^{10–12}

More recently, digital decision-support systems have emerged as potential platforms for reminders and decision/job aids based on decision rules (prompts and alerts) developed according to clinical protocols/recommendations or using machine learning techniques and predictive analysis to extract patterns from clinical data.¹³

To support the development of evidence-based reminders and decision/job aids, WHO commissioned the development of evidence-based clinical algorithms for the management of uneventful and complicated labours, by skilled health personnel providing care during labour and childbirth in any health facility, with a focus on low-resource settings. This manuscript presents the standardised, iterative methods used to develop these intrapartum care algorithms. It is expected that these algorithms will support implementation of WHO recommendations and facilitate the development by stakeholders of evidence-based, up-to-date, paper-based or digital reminders and decision-support tools.

Methods

The intrapartum care algorithms were developed by a multidisciplinary group of experts using an iterative,

standardised two-phased process including: (1) Evidence-based phase: to identify intrapartum care topics and case scenarios, identify evidence-based recommendations and practices in the literature, assess its hierarchy, and develop a first version of the algorithms; (2) Expert-based phase: to peer review the evidence identified and revise the algorithms, and validate the final versions of the algorithms.

A Steering Group, composed of WHO staff and selected content experts, developed the standard operating procedures, identified a first set of intrapartum care topics and case scenarios, convened the technical working group, oversaw the development and consolidation of the various algorithms and approved the final versions of the algorithms. The WHO Intrapartum Care Algorithms Working Group is a purposefully selected international, multidisciplinary group of individuals with expertise in clinical obstetrics and midwifery, evidence-based synthesis and guideline development. The experts were assigned to work in teams of two to four experts with dual function of developing topic-specific algorithms and peer-reviewing algorithms developed by other teams, so that at least one member of a team served as an external reviewer for algorithms that they had not been involved with. Teams could invite new members or consult external content experts as appropriate and in consultation with the steering group. No representatives of pregnant women were involved in the development of the algorithms.

The standard operating procedure was developed based on previous literature on methods for clinical algorithm development, previously proposed standards and examples of algorithm development from other fields. Based on the evidence, algorithms were defined as flowcharts providing evidence-based step-by-step visual interpretation of a logical sequence of information gathering, decision-making and/or associated action for optimal monitoring and management of women and babies during labour and childbirth.^{14–18}

The main components of the processes to develop clinical algorithms were also identified through literature review: description of principles, end users, target population and setting, and care period covered.

Evidence-based phase

Identification of topics and case scenarios

An initial list of intrapartum care topics and case scenarios was developed by the steering group, based on a rapid review of intrapartum care recommendations from WHO, international and national professional societies. This list was refined by the WHO Intrapartum Care Algorithms Working Group teams, based on the findings of the topic-specific literature searches conducted by each of the teams. For any given topic, the first step was to identify up to five

scenarios. References were to be provided to support the choice of the clinical case scenarios, based on epidemiological data of most common deviations, severity of related complications or clinical outcomes, if available.

Identification of evidence

A literature review was completed by each team to identify published clinical practice guidelines, meta-analyses, systematic reviews and relevant single studies for each topic. Specific search terms were developed, including combinations of terms such as 'labour', 'intrapartum', 'algorithms' and specific topic terms. Specific search strategies are described in each manuscript included in this supplement on intrapartum care algorithms.^{19–26} Briefly, searches included The Cochrane Library and MEDLINE/PubMED, CINAHL, National Guidelines Clearinghouse and Google. In addition, international and national guidance documents were identified by consulting web pages of professional societies. Searches were first completed in November 2018 and updated between January and October 2020. No language restrictions were applied when possible but this depended on the language proficiency of each of the teams. The teams screened titles and abstracts, extending manual searches through the reference lists of relevant articles and extracted the recommendations and supporting evidence into an EXCEL file (Appendix S1). Any inconsistencies were solved by consultation with another person in the teams or a member of the WHO working group, as appropriate.

The groups identified the highest level of evidence available for each scenario based on hierarchy of available evidence sources. The highest level of evidence found was used to support decisions along the algorithm using the template in Appendix S1. Priority was given to WHO recommendations and WHO guideline derivatives,^{27,28} given its internationally recognised methods and standards²⁹ for development of recommendations. These ensure that WHO guidelines are of the highest quality, ensure transparent decision-making processes, involve geographical and gender-balanced groups of content experts and stakeholders and ensure that recommendations are universally applicable and people-centred. Where no WHO recommendations or derivatives existed other international or national guidelines using Grading of Recommendations Assessment, Development and Evaluation (GRADE) methodology were prioritised, followed by recommendations developed using other methodologies. In the absence of guidelines, a combination of evidence from existing studies and expert opinion was used to determine key points for consideration in the algorithm. The highest level of evidence found was used to support the decisions along the algorithms, in the order of up-to-date systematic review with or without meta-analyses, other systematic review, validated decision rules, randomised controlled trials, non-randomised controlled trials, observational studies and consensus documents.

Prioritisation of recommendations and interventions in the algorithm

After collating the evidence, a selection process for inclusion of the evidence in the algorithm was conducted by each team. Selection was based on relevance of the evidence to the key decision points, strength of the evidence (if available from the source document), severity of the condition targeted by an intervention (interventions with the most impact on maternal and/or perinatal outcomes) and frequency (e.g. interventions that apply to a significant proportion of women). The selection also accounted for the applicability and feasibility in low-resource settings. Any lack of evidence or inconsistencies in the evidence were discussed with the working group, and decisions informed by the highest level of evidence and consensus for each of the algorithms.

All recommendations and supporting evidence that were applicable to each case scenario were extracted, using standardised tools (Appendix S1) by each of the teams, focusing on specific elements of care, as described in Box 1.

Construction of clinical algorithms

Algorithms were then developed for each case scenario, based on a common structure (Figure 1), starting with the identification of the case scenario and definition of the condition, followed by initial maternal and fetal assessments and management, identification of the probable cause, differential diagnosis and additional management until spontaneous vaginal birth, need for medical review to expedite birth, formulate an individualised care plan in complex cases or referral. Algorithms could also link to other relevant algorithms in the series for identification and/or management of complications where appropriate or to normal labour algorithms in case of resolution of the complication.

In the algorithms, content was broken up into boxes presented in a logical, sequential, succinct way, with unambiguous alternatives and clear stopping points.^{14,15} Each algorithm was composed of standardised, variably shaped boxes representing a decision/action point in terms of suspicion of a condition, diagnosis, monitoring or management. Boxes were colour-coded to improve readability, as follows: clinical state (grey rounded rectangle), decision point (red hexagon), action task (blue rectangle), or link to a different algorithm (pink oval) (Box 2). Different types of boxes could be linked using arrows that should not intersect depending on the clinical flow (e.g. a decision box could be linked to another decision box, an action box, a clinical state box or a link box). The preferred flow was from top to bottom, and left to right, where possible. Only arrows originating from decision boxes were labelled, using 'yes' or 'no'. Clinical state, decision and action boxes were numbered sequentially, left to right and top to bottom. Box numbers corresponded to the table of evidence, showing the evidence source for the action and decision points. Algorithms were made to fit on

Box 1 Elements of care considered in the intrapartum care algorithms

Suspicion of the condition/initial assessment (*what observations will warrant a clinical suspicion of the case scenario?*).

Definition (confirmation) (*what observations define or are characteristic of the case scenario?*).

Differential diagnosis (*what are the various diagnoses that the observations could indicate?*).

Monitoring (*what further observations or assessments are needed to monitor clinical progression of the scenario?*).

Management (*what treatment options can be used to slow the progression, correct the deviation, or avert adverse clinical outcomes?*).

Information and integration of women's views and preferences.

Box 2 Description of different algorithm boxes

Rounded rectangle: This box was used to indicate a starting or ending point, defining the clinical state, diagnosis or final outcome (birth). At the beginning of the algorithm it has only one exit path. At the end of the algorithm it has only an entry path and no exit path.

Rectangle. This box was used to indicate specific actions, either diagnostic or therapeutic, including re-evaluation, initial management, reassessment, maintenance therapy or non-clinical interventions.

Multiple actions that do not need to be sequential in time were listed in one box. When multiple actions are presented, each action was listed in a different line. Only first-option treatments or medications were included, as appropriate and other options were described in the accompanying algorithm manuscript.

Hexagon or diamond: This box required a branching decision (i.e. contains decision-relevant questions), with a response that led to one of two alternative paths. It always has one entry path and two exit paths. This box could refer to abnormal findings, suspicion or confirmation of diagnosis, identification of the cause, clinical laboratory test or differential diagnosis. The decision points are represented with yes/no, or conditional logic like if/then. Multiple questions could be asked in one box, with criteria specified for a 'yes' response (e.g. are two of three present, any present, all present?).

Small oval: This box was used to link different algorithms or proposed to link boxes for graphic clarity at page breaks. Text should read go to algorithm X.

one page. Annotations were allowed to provide clarifications or additional information as footnotes. DRAW.IO (<https://drawio-app.com>; SEIBERT/MEDIA GmbH), an open source diagramming software, was used to construct the algorithms and facilitated remote working.

Expert-based phase

For the purpose of peer review, algorithms were swapped across teams to validate case scenarios, all algorithm boxes and choice of recommendations and evidence based. All the algorithms were then reviewed by the algorithm developing teams in an iterative process that included one 2-day face-to-face meeting of the WHO working group in December 2018, several virtual meetings between the teams and/or the steering group, and written feedback, as deemed necessary. Finally, all algorithms were reviewed for consistency in case definitions, clinical management, formatting and language by the WHO steering group, during 6 days of face-to-face meetings.

Results**Principles**

The algorithms developed would, as far as possible, reflect most up-to-date evidence-based recommendations and therefore reflect interventions and clinical practices that comply with the

characteristics of good-quality maternity care: safe, effective, timely, efficient, equitable and people-centred,⁶ and promote a positive childbirth experience.³⁰ The use of interventions to accelerate, terminate, regulate or monitor labour will be limited to when clinically indicated. In this sense, the algorithms will maximise opportunities for spontaneous vaginal birth, or instrumental vaginal births where clinically appropriate and resources are available, and limit indications for caesarean section. The algorithms developed should never override the healthcare provider's and woman's best judgement.

End users

The target users are skilled health personnel providing care during labour and childbirth, working alone or as part of clinical teams, particularly midwives, non-specialised clinicians (i.e. clinicians without specialist training in obstetrics but who also provide care for women in labour), as well as obstetric specialists.

Target population and setting

The target population is pregnant women with singleton, term pregnancies considered to be at low risk of developing complications at admission to the birthing facility, in active first stage of labour or beyond. This in line with the definition of term 'healthy pregnant women' as per the WHO




Figure 1. Common structure of evidence-based, clinical intrapartum care algorithms.

recommendation on intrapartum care for a positive child-birth experience.³⁰ Management of complications first identified or arising after admission to the birthing facilities are covered in the algorithms.

Health facilities in low- and middle-income countries were the priority setting. The algorithms will mainly be used in the place where labour and childbirth are intended to take place, e.g. labour wards, delivery rooms and

Table 1. Clinical intrapartum care algorithm topics and corresponding case scenarios

Topic	Case scenarios	Differential diagnosis
Spontaneous labour resulting in normal vaginal birth	First stage of labour without complications Second stage of labour without complications Third stage of labour without complications	Latent first stage of labour
Fetal heart rate abnormalities ¹⁹	Fetal bradycardia or late decelerations	Uterine rupture Placental abruption Cord prolapse Maternal hypotension Uterine hyperstimulation Unexplained Maternal hyperthermia Maternal infection Maternal dehydration Unexplained
	Fetal tachycardia	Uteroplacental insufficiency Fetal growth restriction Preterm prelabour rupture of membranes Fetal distress Placental abruption Placenta or vasa praevia Uterine rupture Intra-amniotic infection Inadequate uterine activity Suspected uterine rupture Placental abruption Obstructed labour Uterine tachysystole during augmentation
Liquor abnormalities ²⁰	Oligohydramnios	Malpresentation Cephalopelvic disproportion/ Obstructed labour Delay in progress of labour Inadequate uterine activity Malpresentation Cephalopelvic disproportion Obstructed labour
	Meconium-stained amniotic fluid Bloody amniotic fluid or vaginal bleeding	Postpartum haemorrhage Intrapartum bleeding Maternal infection Suspected chorioamnionitis Suspected dehydration Supine hypotension Hypotension Eclampsia Severe pre-eclampsia Mild pre-eclampsia Postpartum haemorrhage Intrapartum bleeding Maternal infection Suspected chorioamnionitis Suspected amniotic fluid embolism Suspected dehydration
Uterine contractions abnormalities ²¹	Foul-smelling or purulent amniotic fluid Uterine hypoactivity	Chorioamnionitis Pyelonephritis Lower urinary tract infection Pneumonia
	Excessive uterine contractions	
Labour progress abnormalities ²²	Slow progress of active first stage	
	Slow progress of second stage of labour	
Maternal pulse and blood pressure abnormalities ²³	Maternal tachycardia	
	Maternal bradycardia	
	Hypertension	
	Hypotension	
Maternal temperature abnormalities ²⁴	Hyperthermia Hypothermia	

Table 1. (Continued)

Topic	Case scenarios	Differential diagnosis
Maternal urine abnormalities ²⁵	Glycosuria	Other suspected source of infection Cold exposure Previously undiagnosed diabetes Low renal glucose excretion threshold during pregnancy
	Oliguria	Urinary retention Maternal dehydration Pre-eclampsia Shock Underlying maternal cardiac or renal problems
	Proteinuria	Pre-eclampsia Maternal urinary tract infection Maternal dehydration
	Ketonuria	Obstructed labour Maternal diabetes Physiological ketonuria in labour
Third stage of labour abnormalities ²⁶	Postpartum haemorrhage after vaginal birth	Uterine atony Genital tract trauma Retained placenta/placental products Coagulopathy Suspected amniotic fluid embolism Inverted uterus
	Uterine atony	
	Genital tract trauma	1st or 2nd degree tear 3rd or 4th degree tear Cervical tear High vaginal tear Vulval or vaginal haematoma Uterine rupture
	Retained placenta/placental products	

The algorithms are based on best evidence available to date. Responsibility for the interpretation and use of the algorithms lies with the reader. In no event shall the authors be liable for damages arising from its use.

operating theatres. However, the algorithms are applicable to any healthcare setting or other places in the healthcare facilities (e.g. emergency rooms). Possible adaptations that may be required to different settings were acknowledged in each of the manuscripts in this supplement.

Care period covered

The algorithms cover assessments and management from the time of admission of pregnant women to the health facility with the diagnosis of active first stage of labour, or beyond, until immediate postnatal care (including the first hour after childbirth) for women and their babies.

Topic of algorithms and case scenarios

Nine topics were identified for development of clinical algorithms for uneventful and complicated labours. Each topic includes from two to four case scenarios, as shown in Table 1.

The iterative process to develop the algorithms yields around four to seven versions of the algorithms before final validation. In this process, statements were edited, redundant boxes and arrows were collapsed or removed, or additional boxes were added, ensuring that the algorithms used common definitions and homogeneous language. All events and decisions in the algorithm depend on the answers to the questions and observations that precede them. Linkages between algorithms were made in cases where multiple complications could have been identified and to avoid repetition across different algorithms. In the case of multiple complications, it was clearly highlighted where medical review was indicated to refer women with complicated labour for specialist care.

All algorithms include considerations or interventions to ensure people-centred, respectful care, including for example action points on effective communication (e.g. discussion with woman and companion about complication identified, request for consent for certain interventions) or relief of labour pain.

Discussion

The evidence-based, clinical intrapartum care algorithms developed using the methodology described in this manuscript will provide guidance on how to manage uneventful and complicated labours, particularly for less specialised birth attendants. These algorithms form the backbone upon which paper-based tools or digital decision-support systems could be built. They could also be employed to support training of healthcare providers.

To the best of our knowledge, this is the first series of intrapartum care algorithms designed using a robust, standardised approach for the management of uncomplicated and complicated labours, applicable in a variety of settings (including low-resource settings), independent of the models of intrapartum care (midwifery-led or other models). Our literature searches allowed us to identify other intrapartum care algorithms developed either for uncomplicated labours only and/or with a focus on high-resource settings.^{31,32} Although, existing algorithms were useful during the prioritisation of recommendations and interventions, our algorithms may differ given considerations to different models of care or resources available. Many pre-existing algorithms were often boxes and lines or arrows without guidance for when or whether to go from one box to the next; our algorithms incorporated a clear flow and decision boxes to aid in these decisions. Discrepancies between existing and current algorithms are described in each of the manuscripts in this supplement, where appropriate.

Strengths and limitations

The intrapartum clinical algorithms were developed using standardised procedures including identification of key case scenarios, review of the evidence and robust peer-review. They were derived from a comprehensive, up-to-date review of the literature and existing guidelines covering assessments, differential diagnosis and management of uncomplicated births or those presenting with a wide range of complications. Our approach facilitated linkage between relevant algorithms for the management of other conditions, or pointed to escalation for specialist input where needed.

Some of the challenges in developing the algorithms overlap across the different topics and case scenarios, and included the lack of or insufficient evidence for some topics or specific components of the algorithms, including how to prioritise use or sequence of use of certain interventions when multiple treatment choices are available, or when availability may vary according to the setting (e.g. for management of postpartum haemorrhage). Another common limitation was inconsistency among guidelines and recommendation on terminology and definitions used, thresholds to identify deviations from normality (e.g. fetal heart rate

abnormalities¹⁹), frequency of assessment (e.g. temperature²⁴) or management (e.g. recommended dosage for regimens for oxytocin augmentation²¹). These challenges where relatively easily overcome when WHO guideline derivative documents^{27,28} provided some guidance, but it was more difficult when this was not the case, and therefore necessitated reliance on expert opinion, including expert consensus of the WHO Intrapartum Care Algorithms Working Group, in consultation with external experts if deemed necessary. Justifications to those decisions are provided in each of the manuscripts included in this supplement, but in general, this included selection of lower thresholds to detect complications or reference to the more commonly used definition or practice among guidelines to allow wider acceptability and use. The algorithms focus on intrapartum care for women who are at low risk of complications at the time of admission to health facilities for childbirth, and therefore they do not cover management of labour in high-risk pregnant women, those who may present with multiple intrapartum complications where clinical decision-making becomes complex, or home births. Management may need to deviate from recommendations in cases where multiple conditions that impact on each other are present in the same woman, e.g. fluid management if pre-eclampsia and sepsis are both present. Although the algorithms point to timely referrals to higher-level facilities where appropriate, they do not provide details on how safe referrals should be organised. It was sometimes challenging to reflect all aspects of people-centred, respectful care in the algorithms, but additional considerations are provided in the manuscripts. Finally, no representatives of the women were involved in the development of these algorithms, but they are represented in WHO and some national guideline development panels.

Interpretation

Introducing intrapartum care clinical algorithms in facilities with less specialised health personnel may improve maternal and perinatal outcomes. Indeed, evidence-based, easy-to-use, decision-support labour tools could foster optimal labour management and optimise task shifting by supporting decision-making by less specialised health personnel.³³ These intrapartum care algorithms could serve as a prompt to consider a range of differential diagnoses, and as guidance towards appropriate monitoring, investigations and management, with timely referral for specialist care. These would also require functioning referral systems, and competent human resources in sufficient quantities with access to essential services and physical resources (infrastructure, medicines, supplies and equipment).

Implementation of standardised, evidence-based algorithms may improve adherence to guidelines and reduce unnecessary variation in practice.^{18,34} However, algorithms

by their static nature and focus on certain conditions cannot consider the whole complexity of labour and childbirth. The algorithms may provide general guidance on a logical sequence of decision-making, but in all cases evaluation of each individual situation, including women's preferences, should take precedence. The tensions between providing standardised pathway management and exercising clinical judgement to offer individualised care has been highlighted by others.³⁴ Our intrapartum care algorithms may need to be adapted to local context, taking into consideration the availability of skilled personnel and resources.

Over the last decade, WHO has issued and updated multiple guidelines with the overall aim of improving health outcomes and experience of care.⁷ These guidelines include recommendations on clinical and non-clinical interventions for the care of healthy women and babies, and for the management of maternal and neonatal complications around the time of childbirth. The close collaboration with WHO ensured that the algorithms are in line with current WHO recommendations. However, keeping these algorithms up to date will be challenging, and ways to allow them to become more responsive to recommendation updates need to be explored. Digital platforms that can connect guideline development, dissemination and implementation tools could facilitate this task, enhancing knowledge transfer and uptake by end users.^{7,13}

These intrapartum care algorithms, by translating narrative guideline recommendations and other evidence-based information into flowcharts, should facilitate understanding of the evidence by programmers and non-specialists for the development of evidence-based, accurate digital decision-support systems. In future, incorporation of these intrapartum algorithms into interactive software or mobile applications may facilitate end users' application and implementation for management in labour. Data-driven artificial intelligence, including machine learning and statistical predictive techniques, could be used to complement these digital tools.^{13,33}

The intrapartum care algorithms still need to be evaluated prospectively, especially in resource-limited settings, to identify any problems in their implementation, and whether they do in fact improve clinical care and maternal and perinatal outcomes. This will require field testing with feedback from users, to identify problems of interpretation, feasibility and other logistical barriers to implementation, such as skill and resource constraints, and revisions and updates based on field testing results. The implementation of algorithms in clinical practice merits an evaluation of its effects not only on clinical outcomes but unanticipated effects on communication and collaboration between healthcare providers, occupational divisions and satisfaction of women with care.

Conclusions

These intrapartum care algorithms provide a framework for monitoring women during labour and childbirth, identifying and managing complications. It is expected that these algorithms will support the implementation of WHO recommendations and facilitate the development by stakeholders of evidence-based, up-to-date, paper-based or digital reminders and decision-support tools. The algorithms need to be field tested and may need to be adapted to specific contexts.

Disclosure of interests

Dr Renato T Souza received funding from the HRP Alliance, part of the UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a co-sponsored programme executed by WHO, to complete his studies. Other co-authors have no conflicts of interest to declare. Full disclosure of interests available to view online as supporting information.

Contribution to authorship

MB and OTO conceived the methodology for the development of intrapartum care algorithms and developed the standard operating procedures. MB wrote the first draft of the paper. All authors implemented the methods described in this manuscript, contributed to revising the final version and approved the manuscript for publication.

Details of ethics approval

Ethical approval was not required for this work.

Funding

This work was funded by the Bill & Melinda Gates Foundation (Grant #OPP1084318); The United States Agency for International Development (USAID); and the UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a co-sponsored programme executed by the World Health Organization. The funders had no role in design, decision to publish or preparation of the manuscript.

Acknowledgements

We would like to acknowledge the WHO Intrapartum Care Algorithms Working Group for their valuable feedback and contribution to the algorithms.

Data availability

Data sharing not applicable to this article as no data sets were generated or analysed during the current study.

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Appendix S1. Template for reporting algorithms. ■

References

- Kassebaum NJ, Bertozzi-Villa A, Coggeshall MS, Shackelford KA, Steiner C, Heuton KR, et al. Global, regional, and national levels and causes of maternal mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:980–1004.
- Say L, Chou D, Gemmill A, Tunçalp Ö, Moller A-B, Daniels J, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health* 2014;2:e323–33.
- Lawn JE, Blencowe H, Waiswa P, Amouzou A, Mathers C, Hogan D, et al. Stillbirths: rates, risk factors, and acceleration towards 2030. *Lancet* 2016;387:587–603.
- Bhutta ZA, Das JK, Bahl R, Lawn JE, Salam RA, Paul VK, et al. Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *Lancet* 2014;384:347–70.
- Michalow J, Chola L, McGee S, Tugendhaft A, Pattinson R, Kerber K, et al. Triple return on investment: the cost and impact of 13 interventions that could prevent stillbirths and save the lives of mothers and babies in South Africa. *BMC Pregnancy Childbirth* 2015;15:39.
- Tunçalp Ö, Were WM, MacLennan C, Oladapo OT, Gülmezoglu AM, Bahl R, et al. Quality of care for pregnant women and newborns—the WHO vision. *BJOG* 2015;122:1045–9.
- Vogel JP, Dowswell T, Lewin S, Bonet M, Hampson L, Kellie F, et al. Developing and applying a ‘living guidelines’ approach to WHO recommendations on maternal and perinatal health. *BMJ Glob Health* 2019;4:e001683.
- Haddad SM, Souza RT, Cecatti JG, Barreix M, Tamrat T, Footitt C, et al. Building a digital tool for the adoption of the World Health Organization’s antenatal care recommendations: methodological intersection of evidence, clinical logic, and digital technology. *J Med Internet Res* 2020;22:e16355.
- Wang Z, Norris SL, Bero L. Implementation plans included in World Health Organisation guidelines. *Implement Sci* 2016;11:76.
- Allanson ER, Tunçalp Ö, Vogel JP, Khan DN, Oladapo OT, Long Q, et al. Implementation of effective practices in health facilities: a systematic review of cluster randomised trials. *BMJ Glob Health* 2017;2:e000266.
- Peven K, Bick D, Purssell E, Rotevatn TA, Nielsen JH, Taylor C. Evaluating implementation strategies for essential newborn care interventions in low- and low middle-income countries: a systematic review. *Health Policy Plan* 2020;35:ii47–65.
- Stokes T, Shaw EJ, Camosso-Steinovic J, Imamura M, Kanguru L, Hussein J. Barriers and enablers to guideline implementation strategies to improve obstetric care practice in low- and middle-income countries: a systematic review of qualitative evidence. *Implement Sci* 2016;11:144.
- Haddad SM, Souza RT, Cecatti JG, Barreix M, Tamrat T, Footitt C, et al. Correction: building a digital tool for the adoption of the World Health Organization’s antenatal care recommendations: methodological intersection of evidence, clinical logic, and digital technology. *J Med Internet Res* 2020;22:e24891.
- Rosenfeld RM, Shiffman RN, Robertson P. Clinical practice guideline development manual, third edition: a quality-driven approach for translating evidence into action. *Otolaryngol-Head Neck Surg* 2013;148:S1–55.
- Proposal for clinical algorithm standards. Society for Medical Decision Making Committee on Standardization of Clinical Algorithms. *Med Decis Making* 1992;12:149–54.
- Hanekom S, Berney S, Morrow B, Ntoumenopoulos G, Paratz J, Patman S, et al. The validation of a clinical algorithm for the prevention and management of pulmonary dysfunction in intubated adults—a synthesis of evidence and expert opinion. *J Eval Clin Pract* 2011;17:801–10.
- Beitz JM, van Rijswijk L. Developing evidence-based algorithms for negative pressure wound therapy in adults with acute and chronic wounds: literature and expert-based face validation results. *Ostomy Wound Manage* 2012;58:50–69.
- Hegarty J, Savage E, Cornally N, Byrne S, Henn P, Flynn M, et al. A Systematic Literature Review to Support a Framework for the Development of Standards for Clinical Practice Guidance. Dublin, Ireland: Department of Health; 2015.
- Cheung KW, Tan LN, Meher S, the WHO Intrapartum Care Algorithms Working Group. Clinical algorithm for the management of intrapartum maternal urine abnormalities. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16726>
- Liabsuetrakul T, Meher S, the WHO Intrapartum Care Algorithms Working Group. Intrapartum care algorithms for liquor abnormalities: oligohydramnios, meconium, blood and purulent discharge. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16728>
- Gülümser Ç, Yassa M, the WHO Intrapartum Care Algorithms Working Group. Clinical management of uterine contraction abnormalities; an evidence-based intrapartum care algorithm. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16727>
- Pasquale J, Chamillard M, Diaz V, Gialdini C, Bonet M, Oladapo OT, et al. Clinical algorithms for identification and management of delay in the progression of first and second stage of labour. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16775>
- Haddad SM, Souza RT, Cecatti JG, the WHO Intrapartum Care Algorithms Working Group. Management of maternal pulse and blood pressure abnormalities during labour and childbirth: an evidence-based algorithm for intrapartum care decision support. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16776>
- Blennerhassett A, Dunlop C, Lissauer D, the WHO Intrapartum Care Algorithms Working Group. Clinical management of deviations in maternal temperature during labour and childbirth: an evidence-based intrapartum care algorithm. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16730>
- Cheung KW, Tan LN, Meher S, the WHO Intrapartum Care Algorithms Working Group. Clinical algorithm for the management of intrapartum maternal urine abnormalities. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16726>
- Marcus JK, Fawcus S, the WHO Intrapartum Care Algorithms Working Group. Clinical algorithms for management of third stage abnormalities. *BJOG*. 2021. <https://doi.org/10.1111/1471-0528.16729>
- WHO, UNFPA, UNICEF. *Pregnancy, Childbirth, Postpartum and Newborn Care: A Guide for Essential Practice*. Geneva, Switzerland: World Health Organization; 2015.
- Managing Complications in Pregnancy and Childbirth: A Guide for Midwives and Doctors*. 2nd edn. Geneva, Switzerland: World Health Organization; 2017.
- WHO Handbook for Guideline Development*. 2nd edn. Geneva, Switzerland: World Health Organization; 2014.
- WHO Recommendations: Intrapartum Care for a Positive Childbirth Experience*. Geneva, Switzerland: World Health Organization; 2018.
- Intrapartum Care for Healthy Women and Babies*. NICE clinical guideline 190. Updated February 2017. London, UK: National Institute for Health and Care Excellence; 2014.

- 32 Creedon D, Akkerman D, Atwood L, Bates L, Harper C, Levin A, et al. *Management of Labor*. Bloomington (MN): Institute for Clinical Systems Improvement (ICSI); 2013. Mar. 66p.
- 33 Souza JP, Oladapo OT, Bohren MA, Mugerwa K, Fawole B, Moscovici L, et al. The development of a Simplified, Effective, Labour Monitoring-to-Action (SELMA) tool for Better Outcomes in Labour Difficulty (BOLD): study protocol. *Reprod Health* 2015;12: 49.
- 34 Hunter B, Segrott J. Re-mapping client journeys and professional identities: a review of the literature on clinical pathways. *Int J Nurs Stud* 2008;45:608–25.

Appendix

WHO Intrapartum Care Algorithms Working Group

WHO Intrapartum Care Algorithms Working Group: (in alphabetical order) Livia Ciabati, Lariza Laura De Oliveira

(University of São Paulo, Brazil); Dr Renato Souza (University of Campinas, Brazil); Joyce Browne, Marcus Rijken (UMC Utrecht, the Netherlands); Sue Fawcus (University of Cape Town, South Africa); Justus Hofmeyr (Frere Maternity Hospital East London, South Africa); Tip-pawan Liabsuetrakul (Prince of Songkla University, Thailand); Cagri Gülümser (University of Health Sciences, Turkey); Anna Blennerhassett (University of Birmingham, United Kingdom), David Lissauer (University of Liverpool, United Kingdom), Shireen Meher (Birmingham Women's and Children's NHS Foundation Trust). World Health Organization Secretariat (Switzerland): Fernando Althabe, Mercedes Bonet, A. Metin Gülmezoglu, Olufemi T Oladapo.

Disclaimer. The algorithms are based on best evidence available to date. Responsibility for the interpretation and use of these algorithms lies with the reader. In no event shall the World Health Organization or co-authors be liable for damages arising from its use.