

# **LARGE DISCS WITH LARGE CUPS – A DIAGNOSTIC CHALLENGE IN AFRICAN PATIENTS**

Darshana Soma

A research report submitted to the Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, in partial fulfillment of the requirements for the degree of Master of Medicine in Ophthalmology.

Johannesburg, 2009

## **DECLARATION**

I, Darshana Soma, hereby declare that this research report is my own work. It is being submitted for the degree of Master of Medicine (Ophthalmology) at the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at this or any other university.

Signed \_\_\_\_\_

On this \_\_\_\_\_ day of \_\_\_\_\_ 2009

The work reported in this research report was carried out at the Department of Ophthalmology, St John Eye Hospital (Chris Hani Baragwanath Hospital), Johannesburg, South Africa.

## **DEDICATION**

This research report is dedicated to my parents and my siblings in recognition of their continual support and encouragement throughout my life.

## **ETHICS APPROVAL**

The project was approved by the Human Research Ethics Committee (Medical) at the University of the Witwatersrand.

Clearance certificate number: M070435

## **PRESENTATIONS ARISING FROM THIS STUDY**

The results of this study were presented at the annual congress of the Ophthalmological Society of South Africa (OSSA) in March 2008.

Title: Large optic discs with large cups – a diagnostic challenge in African patients

Presenter: Darshana Soma

## **ABSTRACT**

### **Objectives**

To determine in a cohort of 69 African patients with large optic discs and large optic cups, that proportion of patients with physiologic cupping (normal eyes) misdiagnosed as glaucomatous. To evaluate the possible relationship between optic disc size and central corneal thickness.

### **Design and method**

A case series of 69 patients with large discs (vertical disc height measuring  $\geq 1.8\text{mm}$ ) and large cups (vertical cup to disc ratio  $\geq 0.6$ ) was evaluated to determine what proportion had glaucoma and what proportion was normal. Patients categorized as normal were further evaluated to determine what proportion were previously misdiagnosed and treated for glaucoma. Patients with a suspected diagnosis of glaucoma, normal tension glaucoma or primary open angle glaucoma were recruited from the glaucoma clinic at St John Eye Hospital.

Outcome measures included corrected vertical disc height (VDH), vertical cup to disc ratio (CDR), central corneal thickness (CCT), the relationship between VDH and vertical cup height, the relationship between VDH and CCT, adjusted intraocular pressure (A-IOP), retinal nerve fiber layer analysis and visual fields.

## **Results**

Sixty-nine African patients (138 eyes) with large discs and large cups were evaluated. 41 (59%) were females and 28 (41%) were males. The mean age was 56 years. Of the 69 patients, 51 (74%) had physiologic cupping (normal eyes) and 18 (26%) patients were glaucomatous. Of the group of 51 patients with physiologic cupping, there were 9 patients who were previously misdiagnosed with glaucoma and who had received treatment.

VDH ranged between 1.9 and 3.2mm (mean  $\pm$ SD,  $2.3 \pm 0.26$ mm). The distribution analysis of VDH measurements noted the largest cluster around 2.3mm. CCT ranged between 454 $\mu$ m and 618 $\mu$ m (mean $\pm$ SD,  $516 \pm 37$  $\mu$ m). 107 (77.5%) of the 138 eyes had CCT < 544 $\mu$ m.

## **Conclusion**

Large cup to disc ratio in relation to large disc size can be normal. It can be misdiagnosed as glaucomatous if objective retinal nerve fiber layer analysis is not carried out. In this study, 9 (18%) patients from a group of 51 patients with physiologic cupping were misdiagnosed as glaucomatous. There was no linear correlation between CCT and VDH in this study. Pearson's correlation coefficient was 0.13. The majority (77.5%) of eyes had thin corneas (CCT < 544 $\mu$ m).

## **ACKNOWLEDGEMENTS**

Many thanks to:

My supervisors, Prof. Grant McLaren and Prof. Trevor Carmichael for their never-ending assistance, encouragement, patience, guidance and wealth of knowledge that they have imparted to me. It is greatly appreciated.

Dr. Hassan Alli for his assistance, encouragement and good advice.

To fellow colleagues who referred patients to me.

Thomas Ramaphosa, for assisting me with the visual fields and the scanning laser polarimetry.

To all the patients and staff at St John Eye Hospital for making this study possible.



# CONTENTS

	Page
DECLARATION.....	ii
DEDICATION.....	iii
ETHICS APPROVAL.....	iv
PRESENTATIONS ARISING FROM THIS STUDY.....	v
ABSTRACT.....	vi
ACKNOWLEDGEMENTS.....	viii
CONTENTS.....	ix
LIST OF FIGURES.....	x
LIST OF TABLES.....	xi
NOMENCLATURE.....	xii
CHAPTER 1.0 INTRODUCTION.....	1
CHAPTER 2.0 EXPERIMENTAL PROCEDURE.....	13
CHAPTER 3.0 RESULTS.....	20
CHAPTER 4.0 DISCUSSION.....	27
CHAPTER 5.0 CONCLUSION.....	33
APPENDIX A PATIENT INFORMATION AND CONSENT FORM.....	34
APPENDIX B DATA CAPTURE SHEET.....	35
REFERENCES.....	36

## LIST OF FIGURES

Figure	Page
1.1: The Heidelberg engineering IOPac advanced pachymeter .....	6
1.2: Polarization by the scanning laser polarimeter (GDxVCC) – courtesy of Carl Zeiss Meditec Inc.....	8
1.3: Nerve fiber analysis with variable corneal compensation (GDxVCC) - courtesy of Carl Zeiss Meditec Inc.....	10
1.4: Advanced serial analysis (GDxVCC) - courtesy of Carl Zeiss Meditec Inc....	11
2.1: The Haag Streit biomicroscope.....	15
2.2: The 60 diopter lens.....	15
2.3: The calibrated Goldmann tonometer.....	16
2.4: The Volk 3-mirror lens.....	16
2.5: The Shaffer-Etienne classification system.....	17
2.6: The Nikon handheld autorefractor.....	18
2.7: The Oculus automotive perimeter – used to measure visual fields.....	18
2.8: The scanning laser polarimeter (GDxVCC).....	19
2.9: Excel database and Stata analysis and statistical software program.....	19
3.1: Subsections of patients in the study.....	20
3.2: Distribution of central corneal thickness.....	21
3.3: Distribution of corrected vertical disc height.....	22
3.4: The relationship between VDH and vertical cup height.....	23
3.5: The relationship between VDH and CCT.....	24
4.1: Large discs with large cups.....	27
4.2: Subsections of patients in the study.....	31

## LIST OF TABLES

Table	Page
1.1: Correction factors required for different lenses.....	3
3.1: Analysis of visual fields.....	26

## NOMENCLATURE

- |     |               |   |
|-----|---------------|---|
| 1.  | A-IOP         | adjusted intraocular pressure                               |
| 2.  | CCT           | central corneal thickness                                   |
| 3.  | CDR           | vertical cup to disc ratio                                  |
| 4.  | D             | diopter   |
| 5.  | eg.           | example   |
| 6.  | GDxVCC        | nerve fiber analyzer GDx with variable corneal compensation |
| 7.  | HDD           | horizontal disc diameter                                    |
| 8.  | ie.           | that is   |
| 9.  | IOP           | intraocular pressure  |
| 10. | mm            | millimeters   |
| 11. | mmHg          | millimeters of mercury                                      |
| 12. | $\mu\text{m}$ | micrometers   |
| 13. | NFI           | nerve fiber index   |
| 14. | nm            | nanometers  |
| 15. | OSSA          | Ophthalmological Society of South Africa                    |
| 16. | RNFL          | retinal nerve fiber layer                                   |
| 17. | SD            | standard deviation  |
| 18. | TSNIT         | temporal, superior, nasal, inferior, temporal               |
| 19. | VDH           | corrected vertical disc height                              |