

List of figures

Figure 1.1:	Location of the Nylsvlei floodplain in the Limpopo Drainage region (after Morgan, 1996)	4
Figure 1.2:	The main vegetation inundation zones (after Higgins and Rogers, 1993)	8
Figure 1.3:	Map showing the southern portion of the Nylsvlei floodplain and catchments, the location of the study area, locations of the gauging stations and rain gauges, and the then-proposed site of the Olifantspruit Dam. Location shown on 1: 250 000 map 2428 'Nylstroom', produced by the Surveyor General, Pretoria.	18
Figure 1.4:	Schematic diagram of the hydraulic modelling area showing its division into three smaller regions. Significant inflows (Nyl River inflow, tributaries and rainfall) and outflows (Nyl River outflow and losses), and the location of water level gauges along the floodplain are indicated (after Birkhead <i>et al</i> , 2004)	19
Figure 3.1:	Stage level plot for the gauge at Middelfontein (A6H039) for the period 02/1998 to 05/2001 (after Birkhead <i>et al</i> , 2004)	49
Figure 3.2:	Stage level plot for the gauge at Deelkraal (A6H002) for the period 11/1997 to 04/2002 (after Birkhead <i>et al</i> , 2004)	49
Figure 3.3:	Stage level plots for the seven gauges in the Nylsvley Reserve (including Vogelfontein located at its downstream boundary) for the period 01/1996 to 06/2001 (after Birkhead <i>et al</i> , 2004)	50
Figure 3.4:	Stage level plot for the gauge at Mosdene (GP8/A6H038) for the period 02/1996 to 06/2001 (after Birkhead <i>et al</i> , 2004)	50
Figure 3.5:	Rating data and function plot for the gauge at Middelfontein (A6H039) (after Birkhead <i>et al</i> , 2004)	51
Figure 3.6:	Rating data and function plot for the gauge at upstream boundary of the Nylsvley Reserve (GP2) (after Birkhead <i>et al</i> , 2004)	52

Figure 3.7:	Rating data and function plot for the gauge at Vogelfontein (A6H037) (after Birkhead <i>et al</i> , 2004)	52
Figure 3.8:	Rating data and function plot for the gauge at Mosdene (A6H038) (after Birkhead <i>et al</i> , 2004)	52
Figure 3.9:	Discharge hydrographs for the Nyl River at Middelfontein and GP2 (after Birkhead <i>et al</i> , 2004)	55
Figure 3.10:	Discharge hydrographs for the Nyl River at GP2 and Vogelfontein (after Birkhead <i>et al</i> , 2004)	55
Figure 3.11:	Discharge hydrographs for the Nyl River at Vogelfontein and Mosdene (after Birkhead <i>et al</i> , 2004)	56
Figure 3.12:	Total daily rainfall in mm for the Nylsvley weather station (0590486) for the period 1921 to 2001 (after Birkhead <i>et al</i> , 2004)	57
Figure 3.13:	Plot of LIDAR random survey points for the section of the Nylsvley Nature Reserve between GP2 (Deelkraal Road) and GP4 (after Birkhead <i>et al</i> , 2004)	61
Figure 3.14:	Plot of LiDAR points at the upstream end of the study area at Middelfontein, showing the Middelfonteinspruit, Nyl River and N1 freeway.	61
Figure 4.1:	Components of the radiation balance at a soil surface: (a) incoming and outgoing radiation; (b) radiation interchange during the day; (c) radiation interchange at night (after Blight, 1997a)	76
Figure 4.2:	Graph showing cumulative growth in R_n , G and Le over the course of the day 18 December 2000, after Blight (pers. comm.) in the grassy and the degrassed areas	79
Figure 4.3:	Evapotranspiration measured at the Nylsvley Reserve (Blight, 2002a) and from the S pan at the Donkerpoort Dam	86
Figure 4.4:	Comparison between pan data from Donkerpoort Dam and stations located near the study area of the Nylsvlei floodplain	89

Figure 4.5:	Symons pan factors derived from energy balance evapotranspiration data for grassy and bare areas and the Symons pan evaporation data for the same days at the Donkerpoort Dam Symons pan (A6E006), and the recommended Symons pan factor for lakes and wetlands in South Africa from Midgley <i>et al</i> (1994)	91
Figure 4.6:	Comparison of the evapotranspiration derived from the energy balance measurements with various empirical model predictions and data based on class A pan evaporation from Scholes and Walker (1993)	93
Figure 4.7:	Evapotranspiration rates predicted using various pan factors and the rates derived from the energy balance measurements applied to the Donkerpoort Dam station pan data	94
Figure 5.1:	Simplified soil map of the Nylsvley Nature Reserve (after Frost, 1987)	99
Figure 5.2:	Geology of the Nylsvlei floodplain region (after Morgan, 1996)	101
Figure 5.3:	The Guelph Permeameter on the floodplain alluvium in the Nylsvley Nature Reserve, note the Nyl River channel behind	105
Figure 5.4:	Method to determine infiltration into a soil from an open hole in uniform soil (after Blight, 1997b)	109
Figure 5.5:	Infiltration measurements along the traverse conducted in the Nylsvley Nature Reserve in 2001 (after Blight <i>et al</i> , 2001)	111
Figure 5.6:	Depth profile of infiltration measured at the traverse compared with measurements in the channel and at the bridge (transect 2 in Figure 5.7) at the Nylsvley Nature Reserve (after Blight <i>et al</i> , 2001)	112
Figure 5.7:	Soil map of the Nylsvley Nature Reserve (after Harmse, 1977), with numbered sites of infiltration measurements using the Guelph Permeameter and ring infiltrometers, and numbered transects	118

Figure 6.1:	View north to the Waterberg foothills from Vogelfontein in the Nylsvley Reserve, showing the typical limited aerial extent of convective storms in the Nylsvlei catchments	124
Figure 6.2:	Map of the Nylsvlei floodplain and catchments showing the position of all evaporation, rainfall, geohydrological and flow-gauging stations (after Pitman & Bailey, 2003)	125
Figure 6.3:	Contour map at 10cm intervals for the upper portion of the Nylsvley Nature Reserve between GP2 (Deelkraal Road) and GP4 (after Birkhead <i>et al</i> , 2004)	131
Figure 6.4:	Annotated contour map (20cm intervals) of the Nylsvley Nature Reserve upstream of the Vogelfontein Road, showing the artificial dikes (after Birkhead <i>et al</i> , 2004)	134
Figure 6.5:	Vogelfontein causeway, looking upstream into the Nylsvley Nature Reserve. The influence of artificial topographical features (dikes and road) on the hydraulic behaviour is noticeable. A bird-hide is located in the tree-grove upstream of the road (photo K. Rogers) (after Birkhead <i>et al</i> , 2004)	134
Figure 6.6:	Contour map at 20cm intervals for the upper portion of the Nylsvley Reserve, with every fifth contour shaded in black (i.e. at 1m intervals) (after Birkhead <i>et al</i> , 2004)	136
Figure 6.7:	Position of cross-sections (yellow transects) downstream of Gauge Plate (GP) 4 in the Nylsvley Nature Reserve, superimposed on a background image of the floodplain and 20cm contour map. The Nyl River flows from the bottom to the top of the figure (after Birkhead <i>et al</i> , 2004)	138
Figure 6.8:	Photograph of the same area as in Figure 6.7, looking upstream towards Gauge Plate (GP) 4. A bird hide is located in the reed beds through which the Nyl River flows. (photo K. Rogers) (after Birkhead <i>et al</i> , 2004)	138
Figure 6.9:	Aerial view of the Deelkraal Dam, looking downstream, a channel runs past the dam to the bottom right of the photograph (photo K. Rogers)	139

Figure 6.10:	Measured inflows from the DWAF stage plate A6H039 for the 1999/2000 hydrological year, plotted against inundated area in the Middelfontein reach together with the best fit regression line	145
Figure 6.11:	Plot of modelled stage and discharge hydrographs, and measured values at Middelfontein (A6H039 – cross-section 51) for the period 26/02/1998 to 09/05/2001 (after Birkhead <i>et al</i> , 2004).	155
Figure 6.12:	Plot of modelled stage hydrographs and measured values at Deelkraal (A6H002 – cross-section 20) for the period 26/02/1998 to 09/05/2001 (after Birkhead <i>et al</i> , 2004)	156
Figure 6.13:	Plot of modelled stage and discharge hydrographs, and measured values at the downstream boundary of the Middelfontein reach (GP2 – cross-section 1) for the period 26/02/1998 to 09/05/2001 (after Birkhead <i>et al</i> , 2004)	157
Figure 6.14:	Aerial view of the Deelkraal gauge (A6H002) (top right) and the narrow channel and floodplain at this point (K. Rogers)	158
Figure 6.15:	Plot of modelled stage and discharge hydrographs, and measured values at Middelfontein (A6H039) for the period 26/02/1997 to 09/05/2001 (after Birkhead <i>et al</i> , 2004)	160
Figure 6.16:	Plot of modelled stage hydrographs and measured values at Deelkraal (A6H002) for the period 1973 to 1979 (after Birkhead <i>et al</i> , 2004)	160
Figure 6.17:	Plot of modelled stage hydrographs and measured values at Deelkraal (A6H002) for the period 1980 to 1985 (after Birkhead <i>et al</i> , 2004)	161
Figure 6.18:	Plot of modelled stage hydrographs and measured values at Deelkraal (A6H002) for the period 1986 to 1991 (after Birkhead <i>et al</i> , 2004)	161
Figure 6.19:	Plot of modelled stage hydrographs and measured values at Deelkraal (A6H002) for the period 10/1997 to 04/2001 (after Birkhead <i>et al</i> , 2004)	162

Figure 6.20:	Plot of modelled stage and discharge hydrographs, and measured values at GP2 in the Nylsvley Reserve for the period 01/1996 to 04/2001 (Birkhead <i>et al</i> , 2004)	163
Figure 6.21:	Discharge hydrographs at GP2 derived from (i) measured stages and rating function, (ii) hydraulic routing through the upstream wetland using (a) measured flows at Middelfontein and extrapolated tributary flows and (b) extrapolated flows throughout (after Birkhead <i>et al</i> , 2004)	164
Figure 7.1:	Flood peaks in the Olifantspruit from the modelled historical time series used to time the IFR high flow releases in 1980/1981	176
Figure 7.2:	Flood peaks in the Olifantspruit from the modelled historical time series used to time the IFR high flow releases in 1986/1987	176
Figure 7.3:	Comparison of maximum inundated areas of 25 continuous days duration with various maintenance high flow release durations from the Olifantspruit Dam for 1980/1981	179
Figure 7.4:	Comparison of maximum inundated areas of 25 continuous days duration with various maintenance high flow release durations from the Olifantspruit Dam for 1986/1987	180
Figure 7.5:	Comparison of maximum inundated areas of 25 continuous days duration with various maintenance high flow release durations from the Olifantspruit Dam for February with a base flow release of 1.5 m ³ /s	181
Figure 8.1:	Conceptual model of the periodicity of the life history response of <i>Oryza longistaminata</i> (after Marneweck, 2003)	188
Figure 8.2:	Map showing Tarboton's inundation extent definitions (from Morgan, 1996)	190
Figure 8.3:	Daily time-series of inundated areas for the whole study area from the historical catchment scenario and Tarboton's qualitative flooding descriptions	192
Figure 8.4:	Temporal occurrences of inundation areas in the Nylsvley Reserve reach for each catchment development scenario	193

List of figures

Figure 8.5:	Temporal occurrence of inundation areas exceeded in the Nylsvley Reserve reach compared to the virgin scenario	195
Figure 8.6:	Temporal occurrence of inundation areas exceeded in the Nylsvley Reserve reach compared to the historical scenario	195
Figure 8.7:	The largest areas inundated for 25 days, for each year and scenario in the Middelfontein reach	196
Figure 8.8:	The largest areas inundated for 25 days, for each year and scenario in the Nylsvley Reserve reach	197
Figure 8.9:	The largest areas inundated for 25 days, for each year and scenario in the Mosdene reach	197
Figure 8.10:	The largest areas inundated for 25 days, for each year and scenario in the entire study area	198
Figure 8.11:	Number of years in which different inundation areas of at least 25 continuous days duration were exceeded for the different scenarios in the Middelfontein reach	201
Figure 8.12:	Number of years in which different inundation areas of at least 25 continuous days duration were exceeded for the different scenarios in the Nylsvley Reserve reach	201
Figure 8.13:	Number of years in which different inundation areas of at least 25 continuous days duration were exceeded for the different scenarios in the Mosdene reach	202
Figure 8.14:	Number of years in which different inundation areas of at least 25 continuous days duration were exceeded for the different scenarios in the entire study area	202
Figure 8.15:	Inflow, I , (m^3/s) versus inundated area with depth between 0.1m and 0.5m, A , (km^2) regression for range $0.1 \leq I \leq 1.005$	205
Figure 8.16:	Inflow, I , (m^3/s) versus inundated area with depth between 0.1m and 0.5m, A , (km^2) regression for range $I > 1.005$	205
Figure 8.17:	The largest areas inundated for 25 days at depths between 0.1m and 0.5m, for each year and scenario for the Nylsvley Reserve reach	206

- Figure 8.18: Number of years in which different inundation areas of depths between 0.1m and 0.5m and of at least 25 continuous days duration were exceeded for the different scenarios in the Nylsvley Reserve reach 208