

**Palaeoenvironments of the Middle Stone Age at Sibudu Cave,
KwaZulu-Natal, South Africa: An analysis of archaeological
charcoal**

Volume I

Lucy Fiona Allott

A thesis submitted to the Faculty of Science, University of the Witwatersrand, Johannesburg,
in fulfillment of the requirements for the degree of Doctor of Philosophy

Johannesburg, 2005

DECLARATION

I declare that this thesis is my own unaided work. It is being submitted for the Degree of Doctor of Philosophy in the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination in any other University.

_____ day of _____ 2005

ABSTRACT

Analysis of charcoal from Middle Stone Age layers at Sibudu Cave, KwaZulu-Natal, shows evidence of environmental change during the Last Glacial. Layers analysed encompass the end of the cold stadial, Oxygen Isotope Stage (OIS) 4, and the warmer interstadial, OIS 3. Layers are divided – on the basis of lithic industry associations and taxonomic content – into the Howiesons Poort (HP) (dated to ca. 61 000 years ago), and the early, middle and late post-HP assemblages (ca. 60-55 000, 55-50 000 and 50-33 000 years ago respectively). This project aims to identify evidence for environmental change, characterise this change using qualitative and quantitative analyses, and compare the results to evidence from other proxies at Sibudu Cave and elsewhere in South Africa.

HP layers (GS, GR, GR2) are dominated by evergreen forest taxa, including *Podocarpus* spp., *Buxus* sp., *Brachylaena* sp., *Sapium/Spirostachys* and *Ptaeroxylon obliquum*. *Kirkia* sp. suggests a warm, woodland savanna habitat grew beyond the forest vegetation. Early post-HP layers (Eb, SPCA, BSp) contain taxa from evergreen, riverine forest communities, including *Erica* spp., *Leucosidea sericea*, and *Rapanea melanophloeos*. Some of the taxa in these layers suggest a shift in vegetation, possibly related to the marine regression of the Last Glacial, bringing taxa currently found further inland towards the site. Fewer evergreen forest components, and more bushveld taxa, are present in the middle post-HP (RSp, OMOD, MOD) than in the previous layers. Some of the taxa are only found in northern South Africa in regions that are significantly drier than modern KwaZulu-Natal. These layers also contain more *Acacia* spp. and other Fabaceae taxa and fewer *Erica* spp. than the samples from the early post-HP. This may be a result of environmental change, a change in wood selection, charcoal fragmentation, or sampling bias. Layer Bu, within the late post-HP, contains evergreen and deciduous taxa many of which are found in KwaZulu-Natal today. *Kirkia* sp. again provides evidence for a dry habitat not

currently found in the region. Many Types were defined in Bu, which may indicate a vegetation community for which there is currently no reference material.

Temperature and moisture indices produced using the Factor Analysis suggest warm and moist conditions during the HP layers. During the early post-HP conditions became cooler with intermediate moisture levels. Subsequently, conditions were warm and dry (middle post-HP) and then warm with a little more moisture indicated (late post-HP).

Good fuelwoods were routinely collected during the post-HP. In contrast, the HP layers are dominated by *Podocarpus* spp. and many of the good fuelwoods, such as *Acacia* spp. and *Erica* spp., are absent. This pattern may be a result of changes in the environment, sample bias or a development of awareness of wood properties between the HP and post-HP occupations.

The charcoal results corroborate palaeoenvironmental interpretations provided by seeds and macrofauna from Sibudu Cave. When the Sibudu Cave data are combined with data from other sites it is apparent that, although conditions in the summer rainfall region during the Last Glacial were generally dry, there is evidence for localised variation in vegetation and climate.

ACKNOWLEDGEMENTS

Thanks are extended to Professor Bernie Moon, head of the School of Geography, Archaeology and Environmental Studies, for allowing and supporting this study. I am also indebted to Professor Lyn Wadley for providing access to the Sibudu charcoal assemblage as well as providing excellent supervision, discussion and encouragement throughout my research. Thanks to my thesis committee for comments on the project proposal, and particularly to Marion Bamford for providing access to obscure literature.

I am grateful to the staff (Drs. Chester Cain and Bonny Williamson) and students (Dr. Grant Cochrane, Christine Scott, Marelize Lombard and Grant Hall) of the Ancient Cognition and Culture in Africa (ACACIA) research programme for fruitful discussions on many aspects of this project from its conception to conclusion. Christine Scott provided invaluable field assistance and I am particularly appreciative of Christine and Chester for patiently discussing ideas concerning approaches to charcoal analysis. Zenobia Jacobs has provided the single grain Optically Stimulated Luminescence dates for the Middle Stone Age sequence at Sibudu Cave and has kindly permitted the use of all the available dates, some of which are provisional.

During this project, reference collection specimens were collected by Lyn Wadley, Christine Scott and Grant Cochrane. Geoff Nichols, Professor Kevin Balkwill, Lyn Wadley and Christine Scott assisted with identifying voucher specimens. The National Biodiversity Institute is thanked for the use of data from the National Herbarium, Pretoria (PRE) Computerised Information System.

The School of Animal, Plant and Environmental Sciences (APES) provided access to a muffle furnace. Caroline Lalkhan, Abe Seema and Professor Mike Witcomb of the Electron Microscope Unit in APES assisted by preparing specimens for the Scanning Electron Microscope, providing technical advice, producing negatives and allowing

me access to the facilities after normal working hours. Members of the central photography unit at the University of the Witwatersrand produced photographs for the database. Lyn Wadley, Solveig Schiegl, Ina Plug, Chester Cain and Ray Renaut allowed me access to the results of their seed, phytolith, macrofauna, taphonomic and pollen analyses, respectively. I am grateful to Wendy Voorvelt for allowing me access to electronic section drawings and to Grant Cochrane for Figures 3.6 and 3.7.

Dr. Katarina Neumann assisted by teaching me detailed wood anatomy, charcoal identification, and analytical procedures in the unit of African Archaeology and Archaeobotany at the J.W.Göethe University, in Frankfurt. Dr. Francis Thackeray (of the northern Flagship Institute) generously gave time to discuss data analysis and presentation, and Professor Louis Scott (of the University of the Free State) advised upon appropriate data analysis techniques. He also read and made comments on an early version of the quantitative analysis. Professors Paul Fatti and Jackie Galpin of the statistics department at the University of the Witwatersrand recommended appropriate statistical techniques.

Proof reading and editorial work was undertaken by Professor Lyn Wadley and Dr. Ross Damiani. Ross Damiani also assisted with graphics production, provided advice on thesis organisation, and his tireless patience throughout the research. Family and friends are also thanked for assistance with formatting and printing.

This project was funded by the University of the Witwatersrand through post-graduate merit awards (2001-2003) and University Council Scholarships (2002 and 2003). A National Research Foundation (NRF) “Women in Science” scholarship and a Karen Fredman Local Merit Scholarship were received in 2002 and 2003, respectively. An ACACIA bursary, awarded by Professor Lyn Wadley and funded by the NRF, was held between 2001 and 2003. The Palaeoanthropology Scientific Trust awarded a grant in 2004 to enable the completion of this project.

CONTENTS

	Page
DECLARATION	ii
ABSTRACT	iii
ACKNOWLEDGEMENTS	v
LIST OF FIGURES	x
LIST OF TABLES	xiv
LIST OF APPENDICES	xvi

VOLUME 1

CHAPTER 1. INTRODUCTION

1.1. Introduction	1
1.2. Aims	1
1.3. Chapter Summary	6

CHAPTER 2. EVIDENCE FOR PALAEOENVIRONMENTS IN SOUTHERN AFRICA DURING THE LAST GLACIAL

2.1. Introduction	8
2.2. Global Environmental changes between the Last Interglacial and the Last Glacial Maximum	10
2.3. Local Palaeoenvironmental evidence	12
2.3.1. Local Sea Levels	13
2.3.2. River Geomorphology and Palaeochannels	16
2.3.3. Sediment and Pollen Records	17
2.3.4. Oxygen Isotope Records	23
2.4. Palaeoenvironmental Indicators from Archaeological Contexts	23
2.4.1. Winter Rainfall Region	24
2.4.2. Summer Rainfall Region	36
2.5. Summary	40
2.5.1. Winter Rainfall Region	40
2.5.2. Summer Rainfall Region	42

CHAPTER 3. SIBUDU CAVE LOCATION, EXCAVATIONS AND ENVIRONMENTAL PROXIES

3.1. Site Location and Archaeology	45
3.2. Modern Vegetation and Local Climate	48
3.3. Local Geology and Lithic Raw Material Sources	49
3.4. Sibudu Cave Research	50
3.4.1. Fauna	51

3.4.2. Seeds	52
3.4.3. Phytoliths	54
3.4.4. Pollen	55
3.5. Summary	56
CHAPTER 4. SAMPLING STRATEGIES AND ANALYTICAL THEORY	
4.1. Introduction	58
4.2. Multiple Stages of Sampling	58
4.2.1. Sampling Agencies for Archaeological Charcoal Assemblages	60
4.3. Analytical Theory	69
4.3.1. Qualitative versus Quantitative	69
4.4. Summary	72
CHAPTER 5. SAMPLING, CHARCOAL IDENTIFICATION AND ANALYTICAL METHODS	
5.1. Introduction	74
5.2. Sibudu Charcoal Assemblage	74
5.2.1. Sampling Procedures	74
5.3. Identification	78
5.3.1. Southern African Comparative Collections	78
5.3.2. The University of the Witwatersrand Reference Collection and Database	81
5.4. Analysis of Sibudu Cave Charcoal	85
5.4.1. Sample Size Adequacy	85
5.4.2. Qualitative	87
5.4.3. Quantitative	89
5.5. Summary	92
CHAPTER 6. QUALITATIVE ANALYSIS RESULTS AND PRELIMINARY INTERPRETATIONS	
6.1. Introduction	93
6.2. Qualitative Analysis	94
6.2.1. Howiesons Poort (HP)	94
6.2.2. Post-Howiesons Poort (post-HP)	100
6.3. Summary	123
CHAPTER 7. QUANTITATIVE ANALYSIS RESULTS AND PRELIMINARY INTERPRETATIONS	
7.1. Introduction	127
7.2. Quantitative Analysis	127
7.2.1. Factor Analysis	127
7.2.2. Correspondence Analysis	138
7.3. Summary	140
CHAPTER 8. DISCUSSION	

8.1. Introduction	144
8.2. Sibudu Cave Charcoal Assemblage	144
8.2.1. Howiesons Poort (HP)	147
8.2.2. Post - Howiesons Poort (post-HP)	152
8.3. Palaeoenvironmental Proxies from the Middle Stone Age at Sibudu Cave	160
8.3.1. Charcoal and Botanical Remains	160
8.3.2. Botanical Remains and Fauna	164
8.4. The Contribution of Sibudu Cave data to Understanding Palaeoenvironments in southern Africa during the Last Glacial	165
8.4.1. Oxygen Isotope Stage 4 (OIS 3)	165
8.4.2. Oxygen Isotope Stage 3 (OIS 4)	167
8.4.3. Correlations between Temperature and Moisture Records and the Sibudu Cave Summary Statistics Indices	169
8.5. Summary	170

CHAPTER 9. CONCLUSION & FUTURE RESEARCH

9.1. Introduction	172
9.2. Global climates during the Last Interglacial/Glacial cycle	172
9.3. Evidence for changes in climate during the Last Interglacial/Glacial cycle in South Africa	173
9.4. Evidence for climate change using palaeoenvironmental proxies from Sibudu Cave	177
9.4.1. Macrofauna, seeds, phytoliths and pollen	177
9.4.2. Charcoal	178
9.5. Summary	181
9.6. Directions for Future Research	183

REFERENCES	186
------------	-----

VOLUME 2

FIGURES	1
TABLES	42
APPENDICES	66

LIST OF FIGURES

	Page
CHAPTER 2.	
Figure 2.1. Bathymetric Map of Tugela Canyon (after Zhiv 1975) with projected sea-level at 120 m below mean sea level_____	1
Figure 2.2. Offshore Palaeoshorelines 1-4 (after Ramsay 1996)_____	2
Figure 2.3. Plot of sea-level curve (after Williams <i>et al.</i> 1981) and the inferred age and depth relationships of Palaeocoastlines 1-4, on the Sodwana Bay shelf (after Ramsay 1996)_____	3
Figure 2.4. Current interpretation of the Kosi Bay and Port Durford Formations (adapted from Maud & Botha 2000)_____	4
Figure 2.5. Map showing locations of sites with palaeoenvironmental data_____	5
Figure 2.6. Tswaing Crater (Partridge <i>et al.</i> 1997) proxy rainfall record aligned with Voordrag colluvium and palaeosol formations (adapted from Clarke <i>et al.</i> 2003) _____	6
Figure 2.7. “Tentative” correlation between pollen zones at Tswaing Crater and Wonderkrater (redrawn from Scott 1999)_____	7
Figure 2.8. Correlation of Boomplaas stratigraphic units to Klein’s (1978), Earlier Upper Pleistocene Units_____	8
CHAPTER 3.	
Figure 3.1. Location of Sibudu Cave (redrawn from Wadley & Jacobs 2004)_____	9
Figure 3.2. Plan of Sibudu Cave and excavation (after Wadley & Jacobs 2004)_____	10
Figure 3.3. a) B5 and B6 North wall Stratigraphy_____	11
Figure 3.3. b) B5 East wall Stratigraphy_____	12

Figure 3.4. Stratigraphy of East Section wall showing layer Bu (after Wadley 2001)	13
Figure 3.5. Map from Acocks (1988) showing Veld Types in Savanna Biome (after Wadley 2004)	14
Figure 3.6. Map of Local Geology (supplied by G. Cochrane)	15
Figure 3.7. Dolerite Dyke formation dipping into Tongati River (supplied by G. Cochrane)	16
 CHAPTER 4.	
Figure 4.1. Flow chart showing stages of sampling	17
 CHAPTER 5.	
Figure 5.1. Photograph of Sibudu Cave Excavation (supplied by C. Scott)	18
Figure 5.2. View of Tongati River and vegetation from Sibudu Cave (supplied by L. Wadley)	19
Figure 5.3. Cumulative curve Howiesons Poort layers	20
Figure 5.4. Cumulative Curve Layer Eb	20
Figure 5.5. Cumulative Curve Layer SPCA	21
Figure 5.6. Cumulative Curve Layer BSp	21
Figure 5.7. Cumulative Curve Layer RSp	22
Figure 5.8. Cumulative Curve Layer OMOD	22
Figure 5.9. Cumulative Curve Layer MOD	23
Figure 5.10. Cumulative Curve Layer Bu	23
Figure 5.11. Diagram showing a hypothetical correlation between SSRF1 (temperature) and SSRF2 (moisture)	24

CHAPTER 6.

Figure 6.1. Distribution maps for taxa identified in the Howiesons Poort layers (GS, GR2 and GR)	25
Figure 6.2. Distribution maps for taxa identified in layer Eb	26
Figure 6.3. Distribution maps for taxa identified in the layer SPCA	27
Figure 6.4. Distribution maps for taxa identified in the layer BSp	28
Figure 6.5. Distribution maps for taxa identified in the layer RSp	29
Figure 6.6. Distribution maps for taxa identified in the layer OMOD	30
Figure 6.7. Distribution maps for taxa identified in the layer MOD	31
Figure 6.8. Distribution maps for taxa identified in the layer Bu	32
Figure 6.9. Location of PRECIS grid squares, 2931 AC and 2931 CA, and their proximity to Sibudu Cave	33
Figure 6.10. Graph showing identified taxa and fragment frequencies	34

CHAPTER 7.

Figure 7.1. SSRF1 temperature (warmer-cooler conditions). The indices could also be interpreted as an indication of moisture availability	35
Figure 7.2. SSRF2 moisture availability (wetter conditions-drier conditions). The graph could also be interpreted as showing the presence of frost resistant plants to more non-frost resistant taxa being present	36
Figure 7.3. SSRF3 habitat niches - forest vegetation to less forested (or different forest) and more open vegetation	37
Figure 7.4. Correspondence Analysis graph Axis 1 and 2	38

CHAPTER 8.

Figure 8.1. Vegetation Map (after van Wyk & van Wyk 1997) showing the modern vegetation around Sibudu Cave and the proximity of the Afromontane and inland forest vegetation to the site_____ **39**

Figure 8.2. Deuterium isotope ratios for the Vostok Ice Core (Jouzel *et al.* 1997) (adapted from Thackeray 1992) and Summary Statistics index on Rotated Factor 1 (SSRF1), based on archaeological charcoal samples, showing the placement of the Sibudu Cave sequence_____ **40**

Figure 8.3. Tswaing Crater, Voordrag colluvium accumulations (Clarke *et al.* 2003) and Summary Statistics index on Rotated Factor 2 (SSRF2) showing discrepancies between the Sibudu Cave record for available moisture and the proxy rainfall record_____ **41**

LIST OF TABLES

	Page
CHAPTER 2.	
Table 2.1. ^{14}C dates and relative sea levels (below mean sea level) for estuarine sediments (Maud 2000)	42
Table 2.2. Correlation of Klasies River Middle Stone Age Archaeological sequences with Avery's (1987) microfauna sample numbers, and Feathers's (2002) Luminescence dates	42
Table 2.3. Correlations between old and revised ESR dates by Grün <i>et al.</i> (1990) and Grün & Beaumont (2001), respectively, and Butzer's (1984) unit numbers used by Avery (1982) (Correlation based on Avery's updated interpretation of micromammal data (Avery 1992))	43
CHAPTER 3.	
Table 3.1. OSL dates and C^{14} dates (Wadley & Jacobs 2004), and provisional OSL dates Z. Jacobs (pers. comm.) for the Middle Stone Age deposits at Sibudu Cave	44
Table 3.2. Data from palaeoenvironmental proxies, including seeds, phytoliths and macrofauna (Wadley 2004, Schiegl <i>et al.</i> 2004, and Plug 2004), respectively	45
CHAPTER 6.	
Table 6.1. Environmental requirements and preferred habitats of taxa identified through charcoal and seed analyses from Sibudu Cave	48
Table 6.2. Fragment counts for Howiesons Poort (HP) charcoal identifications (from combined layers, GS, GR2 and GR)	54
Table 6.3. Fragment counts for Eb charcoal identifications	55
Table 6.4. Fragment counts for SPCA charcoal identifications	56
Table 6.5. Fragment counts for BSp charcoal identifications	57
Table 6.6. Fragment counts for RSp charcoal identifications	58

Table 6.7. Fragment counts for OMOD charcoal identifications_____	59
Table 6.8. Fragment counts for MOD charcoal identifications_____	60
Table 6.9. Fragment counts for Bu charcoal identifications_____	61

CHAPTER 7.

Table 7.1. Eigenvalues of the Correlation Matrix. Column 5 shows the cumulative percentage contribution of each factor to the data_____	62
Table 7.2. Factor Pattern with values less than 0.3 and greater than – 0.3 blanked out. Negative Values highlighted_____	62
Table 7.3. Factor Loadings for Factor 1 (F1) before varimax rotation was applied. (Taxa ordered from highest to lowest Factor Loadings)_____	63
Table 7.4. Factor Loadings for Factor 2 (F2) before varimax rotation was applied. (Taxa ordered from highest to lowest Factor Loadings)_____	63
Table 7.5. Factor Loadings for Factor 3 (F3) before varimax rotation was applied. (Taxa ordered from highest to lowest Factor Loadings)_____	64
Table 7.6. Factor Loadings for Rotated Factor 1 (RF1) (after varimax rotation was applied). (Taxa ordered from highest to lowest Factor Loadings)_____	64
Table 7.7. Factor Loadings for Rotated Factor 2 (RF2) (after varimax rotation was applied). (Taxa ordered from highest to lowest Factor Loadings)_____	65
Table 7.8. Factor Loadings for Rotated Factor 3 (RF2) (after varimax rotation was applied). (Taxa ordered from highest to lowest Factor Loadings)_____	65

LIST OF APPENDICES

	Page
APPENDIX A	
Reference Collection Database with no graphics_____	66
APPENDIX B	
Reference Database Images_____	92
APPENDIX C	
Wood anatomical characters used for charcoal identification_____	93
APPENDIX D	
Type Identification Database_____	103
APPENDIX E	
Summary Statistics Equations_____	124
APPENDIX F	
List of genera and species included in the factor analysis_____	130