

**A CROSS-SECTIONAL STUDY INVESTIGATING KNOWLEDGE, ATTITUDES AND
HEALTH CHOICES IN RELATION TO DIABETES MELLITUS AMONG NON-DIABETIC
PATIENTS ATTENDING CHIAWELO COMMUNITY PRACTICE, SOWETO, REPUBLIC OF
SOUTH AFRICA**

Dr M.R. TSHIBEYA

Student Number 1813673



A publishable article as research report submitted to the Faculty of Health Sciences, University of the Witwatersrand in partial fulfilment of the requirements for the degree of Master of Medicine in Family Medicine (MMed Fam Med)

Supervisor: Dr M. TORLUTTER

MBCHB, MMED (Family Medicine SU), DA, DMH, DipObst, Dip HIV Mx

Family Medicine University of the Witwatersrand

Johannesburg 2023

DECLARATION

I, MBUYI ROLAND TSHIBEYA, hereby declare that this research is my own unaided work, except where due acknowledgement for assistance received has been made. It is being submitted for the degree of Master of Family Medicine at the University of the Witwatersrand, Johannesburg. It has not been submitted previously for any other degree or examination at this or any other university.

Signed 

(Candidate)

Date: 28 February 2023

Signed _____

(Supervisor)

Date: 28 February 2023

ACKNOWLEDGEMENTS

I would like to thank my lecturers and supervisors in the Department of Family Medicine, Witwatersrand University. A special word of appreciation to my MMed supervisor, Dr Michele Torlutter, for all her help, not only to complete this research paper but also all other assignments, and for guiding my steps through this programme despite her heavy schedule. I appreciate all the guidance received by the statisticians of the University of the Witwatersrand for the data analysis process. My acknowledgement also goes to patients who agreed to participate in the study, as well as the clinical staff of Chiawelo Community Practice.

DEDICATION

I dedicate this research report firstly to my wife, Maddy, for always standing by me and encouraging me to fulfil my goals during the hopeless moments I encountered in this programme.

Secondly, to my children for all the years of deprivation and sacrifices they suffered to allow me to complete this work.

Thirdly, to my late mother and father who worked hard to offer me the comfort of achieving my dreams.

Table of Contents

INTRODUCTION	2
METHODS	4
Design and setting.....	4
Study population, sample size and sampling	4
Tools and data collection	4
Analysis.....	5
Ethical considerations	6
RESULTS.....	6
Socio-demographics	6
Knowledge.....	7
Attitudes and Health Choices.....	9
Univariate and Multivariate logistic regression	12
DISCUSSION	18
STUDY LIMITATIONS.....	22
CONCLUSIONS	23
REFERENCES	23

LIST OF TABLES

TABLE 1: Socio-demographic profiles of participants to evaluate knowledge, attitudes,health choices and related factors towards DM among members attending the CCP, Soweto 2020 (N = 165)	6
TABLE 2: Frequency distribution of participant responses to DM knowledge, 2020 (N=165).....	7
TABLE 3: Frequency distribution of patient responses to attitudes towards DM and self-reported health choices that prevent or mitigate DM, 2020 (N=165)	9
TABLE 4: Univariate and multivariate logistic regression predicting DM-associated knowledge among members attending the CCP, Soweto 2020 (N=165)	12
TABLE 5: Univariate and multivariate logistic regression predicting DM-associated attitudes among members attending the CCP, Soweto 2020 (N=165)	14
TABLE 6: Univariate and multivariate logistic regression predicting DM-associated health choices among members attending the CCP, Soweto, 2020 (N=165)	16

NOMENCLATURE

CCP: Chiawelo Community Practice

DM: Diabetes Mellitus

HIV: Human Immunodeficiency Virus

NCDs: Non-communicable Diseases

**A CROSS-SECTIONAL STUDY INVESTIGATING KNOWLEDGE, ATTITUDES AND
HEALTH CHOICES IN RELATION TO DIABETES MELLITUS AMONG NON-
DIABETIC PATIENTS ATTENDING CHIAWELO COMMUNITY PRACTICE,
SOWETO, REPUBLIC OF SOUTH AFRICA**

An article submitted according to the style guide and requirements of the **African Journal of Primary Health Care & Family Medicine** (PHCFM).

ABSTRACT

Background: The knowledge, attitudes and health choices of non-diabetic patients in resource-limited settings to prevent the onset of diabetes mellitus (DM) are poorly understood.

Aim: The aim of this study was to investigate and describe the knowledge, attitudes and health choices of non-diabetic patients in relation to DM at the Chiawelo Community Practice (CCP), Soweto, South Africa.

Methods: In this cross-sectional study, a self-reported questionnaire was administered to 165 adult participants attending the CCP from 2nd March to 17th April 2020. Descriptive analysis, Chi square and univariate logistic regression were included in the analysis. Multivariate analysis was done for variables with a p-value <0.25.

Results: In total, the study had 165 participants, of whom 112 (68%) were women, 35% were older than 45 years and 16% were younger than 25 years of age. Almost half (49%) of participants had good knowledge of DM, with 60% indicating good attitudes and 52% making good health choices to prevent DM. Participants who received education from clinicians were 4.31 times more likely to develop better attitudes ($p=0.003$) and 3.34 times more likely to adopt better health choices ($p=0.004$) towards DM compared to those who obtained information from media or other sources.

Conclusion: The study found that poor knowledge of DM does not necessarily translate into poor attitude towards the disease, which is noteworthy. The study also highlighted the important role of healthcare workers in influencing behaviour change.

Key words: Diabetes mellitus; non-diabetic patients; knowledge; attitudes; health choices; primary healthcare

INTRODUCTION

The burden of disease from diabetes mellitus (DM) is ever-increasing. Currently, one in 11 adults is affected by DM globally, with 90% of those having type 2 DM.¹ The prevalence of type 2 DM is projected to rise to 693 million by 2045, with half of the people affected remaining undiagnosed.² The world spent approximately \$850 billion on diabetic care in 2017.² The global mortality rate from DM in 2019 was 11.3%.³

On the African continent, it has been shown that DM often goes undiagnosed. A systematic review in 2020 in this region revealed a prevalence of 3.85% (95% CI. 3.10–4.60) of undiagnosed DM among adults.⁴ Compared to the rest of the world, Africa also has the highest percentage of deaths related to DM in individuals aged less than 60 years (73.1%).³

The prevalence of non-communicable diseases (NCDs) are reportedly increasing faster in poor and developing countries, and accounted for 57.4% of the total burden of disease in South Africa in 2016.^{5, 6} A study published in South Africa in 2017 showed that the prevalence of DM in individuals older than 15 years was 10.1%, which demonstrated a slightly higher rate compared to the global statistics as stated above.⁷ According to Statistics South Africa, DM was the fifth most common cause of death in South Africa in 2012.⁸ South Africa is experiencing an increase in NCDs, including DM, due to several risk factors that may be modifiable or non-modifiable. The effect of urbanisation on populations as well as rural-urban differences have had a major impact on the increase in the risk for DM. Another impact has been the effects and treatment of the human immunodeficiency virus (HIV) and AIDS. Modifiable risk factors may include diminished physical exercises, diet, obesity and psychosocial factors. Of particular concern is the number of obese children, which in sub-Saharan Africa has increased to 10 million in 2012 from four million in 1990.⁹ Non-modifiable risk factors include family history, age and genetics.¹⁰ The focus of this research is mainly on interventions that address the modifiable risk factors of type 2 DM.

A study of 6 958 patients in a Danish cohort, conducted in 2018, revealed that slightly more than one-third (35%) of type 2 DM patients had micro- and macrovascular complications related to DM at the time of diagnosis.¹¹ During the latent phase of the disease, micro- and macrovascular complications can progress significantly. The risk of contracting diabetic neuropathy, which is the most common microvascular complication, and other microvascular complications, depends on the

severity and duration of hyperglycaemia.¹² Diabetic neuropathy can develop seven years before type 2 DM is diagnosed.¹² Once DM is established, the long-term complications may include cardiovascular disease (stroke and ischemic heart disease), renal failure, blindness and limb amputation.¹³ DM also impairs the control of diseases such as HIV and tuberculosis.¹⁴ It is a condition that requires behaviour change in relation to the modifiable risk factors. If detected early, in the asymptomatic phases, the trajectory of the disease may be altered for better outcomes.¹⁵ This research was important to assess awareness of the general population to the risks of DM, and the importance of lifestyle changes during the latent phase of illness when people do not yet know they have the disease.

Studies done in Asian and other African countries amongst patients diagnosed with DM have shown a significant relationship between good knowledge and attitudes towards diabetes and better blood glucose control.^{16,17,18} Health promotion in clinics and the community may therefore improve knowledge, attitudes and health choices in guarding against DM. Increased knowledge, however, does not necessarily result in a change of behaviour, and behaviour change is difficult for most people.¹⁹

Healthcare workers require better understanding of the current knowledge and behaviours of the local community towards DM and DM prevention before appropriate interventions can be developed to address gaps. Furthermore, the way in which information is delivered to patients must be addressed. Clinical services are not always patient-centred, and health practitioners often have an authoritative style, which might not be conducive to conveying information to individuals with different background knowledge, beliefs, or specific ideas or needs.²⁰

There is currently a paucity of scientific papers on knowledge, attitudes and health choices among the non-diabetic population in a facility setting in South Africa. This study, which was conducted in the Chiawelo Community Practice (CCP), Soweto, South Africa, aimed to fill this gap. Approximately 1 500 patients attend the CCP monthly. Of the total population registered in Ward 11 by the CCP's community health workers, 1.17% reported being treated for DM at the clinic at the launch of this study, which may represent underdiagnosis of DM in this setting. Given the increasing prevalence, underdiagnosis and long pre-symptomatic phase in which prevention can be achieved, in addition to the morbidity and mortality and high costs associated with established disease, healthcare workers need to better understand the current lifestyle and health choices of

patients. This information will assist in improving the behaviour and lifestyle choices in the community and better inform primary and secondary prevention measures. Information from this study can also be used to strengthen health services through several clinical governance activities, including quality improvement, capacity building, health education and community-oriented promotion and prevention strategies.

METHODS

Design and setting

This was a descriptive, cross-sectional study conducted at the CCP in a peri-urban region of Soweto, Johannesburg. The CCP is situated within the Chiawelo Community Health Centre, which is a government facility that serves mainly socioeconomically disadvantaged patients. It is a community-oriented primary healthcare facility run by family physicians, and provides a variety of services, including acute and chronic care, prevention and health promotion services, HIV management and family planning. Patients are seen by different health workers (health promoters, community health workers, nurses or doctors) in the chronic and acute areas of the facility, based on their reason for seeking medical advice or care.

Study population, sample size and sampling

The study population comprised patients residing in Ward 11 who attended the CCP. While conducting this study, approximately 1 500 patients were attending the CCP monthly. The study took place between March and April 2020 and included adult patients undiagnosed with type 2 DM, aged 18 years or older. With the 'no' response rate of the questionnaire at 11.85%, a sample size of 161 was reached using a web-based statistical and epidemiologic calculator for public health. With an attrition of 10%, the sample size was 177.²¹ Twelve participants were excluded because they had left blank spaces on the questionnaires, therefore 165 participants were included in the analysis.

Tools and data collection

The study used a structured, self-reported questionnaire designed by Kassahun and Mekonen. It was validated in Ethiopia but adjusted to adapt it to the South African context, such as the currency from the Ethiopian birr to the South African rand.²² It was researcher-administered in the case of

illiterate patients. The questionnaire was piloted with five patients randomly selected from the CCP and found to be suitable in terms of the appropriateness of language use, content, clarity, sequence and flow of the data collection tool. A research assistant, trained in the use of the research tool, assisted the primary researcher with obtaining informed consent as well as interpretation where a language barrier existed.

Every third patient in the queue was selected for the study. This occurred during routine weekday mornings while patients were waiting for their consultation in the acute and chronic areas of the CCP. The primary investigator explained the purpose of the study to the patients. Selected patients who agreed to participate were taken to a private room where the research assistant obtained their informed consent to participate in the study in their preferred languages. Patients able to read English completed the questionnaire themselves. It took 20 minutes on average to complete the questionnaire.

The data collection sheets were coded, transcribed to an MS Excel spreadsheet under password protection, and stored on a flash disc kept safely in the researcher's office. Confidentiality was strictly maintained and only the primary researcher had access to the information.

Analysis

Data transcribed in MS Excel was uploaded to STATA version 15 for analysis. The normality of data distribution was determined using skewness, scatter plots and kurtosis. For continuous variables, the median and standard deviations and interquartile ranges were utilised for normal and skewed data. For proportions and to assess differences within groups, the researcher used the Chi square test or Fisher's exact test. To check for associations, univariate logistic regression was done, and odds ratios used to interpret the results. Multivariate analysis was done for variables with $p < 0.25$. An adjusted odds ratio (AOR) with $p < 0.05$ was regarded as statistically significant in this analysis. From the five-point Likert scale, the values were collapsed to two continuous data sets. The median was used to divide the categories.²³

Ethical considerations

This research respected all ethical norms for research in medical fields involving human participants, preserving the ethical value of the World Medical Association's Declaration of Helsinki. This study was not sponsored and there was no conflict of interest, with autonomy to publish research results. Before conducting this study, a clearance certificate with registration number M190347 was obtained from the Human Research Ethics Committee (HREC) (medical) of the University of the Witwatersrand. The National Human Research Database and the Johannesburg District Research Committee granted authorisation to conduct the study in the CCP, with registration numbers GP201905007 and 2019-05-004, respectively.

RESULTS

Socio-demographics

TABLE 1: Socio-demographic profiles of participants to evaluate knowledge, attitudes, health choices and related factors towards DM among members attending the CCP, Soweto 2020 (N = 165)

Gender	N	Proportion %
Male	53	32
Female	112	68
Age category	N	Proportion %
≤24	27	16
25-34	42	25
35-44	39	24
45+	57	35
Education	N	Proportion %
Illiterate	3	2
Literate	162	98
Marital status	N	Proportion %
Single	103	62
Married	34	21
Divorced/separated	5	3
Widowed	13	8
Cohabiting	10	6
Occupation	Total	Proportion %
Housewife	28	17
Student	25	15
Merchant	5	3
Farmer	2	1

Government/private	29	18
Daily labourer	23	14
Other	53	32
Average monthly income	N	Proportion %
0-4999	126	76
5000-9999	28	17
10000-14999	4	2
15000+	7	4
Exposure to DM education	N	Proportion %
Yes	139	84
No	26	16
Information sources	N	Proportion %
Media	60	36
Health practitioners	49	30
Friends/ relatives	28	17
Other	28	17
Family history of DM	N	Proportion %
Yes	99	60
No	66	40

Table 1 illustrates the socio-demographics of the study population. The study had a total of 165 participants, with 112 (68%) being female. There was a good mix of age ranges in the study, with an anticipated large number of patients aged over 45 (57, 35%) based on the clinic statistics showing that older people were more inclined to come for a consultation. The majority of participants were literate (161, 98%), single (103, 62%), and in a low income range (126, 76%). Most had previously been exposed to DM education (139, 84%), with the main sources being media (60, 36%) and health practitioners (49, 30%). Over half had a positive family history of DM (99, 60%).

Knowledge

TABLE 2: Frequency distribution of participant responses to DM knowledge, 2020 (N=165)

Definition of DM	Yes. N (%)	No. N (%)
DM is a disease of inadequate production of insulin	99 (60)	66(40)
DM is a disease of the body not reacting to insulin	80 (48)	85(52)
DM is not curable	115 (70)	50(30)
DM is a disease that can affect any part of the body	99 (60)	66(40)

Signs and symptoms of DM	Yes. N (%)	No. N (%)
Frequent urination	117 (71)	48(29)
Excessive thirst	122 (74)	43(26)
Excessive hunger	121 (73)	44(27)
Weight loss	112 (68)	53(32)
High blood sugar level	111 (67)	54(33)
Blurred vision	129 (78)	36(22)
Slow healing of wounds	123 (75)	42(25)
Feeling of weakness	111 (67)	54(33)
Risk factors of DM	Yes. N (%)	No. N (%)
Older age	64 (39)	101(61)
Hereditary or family history of DM	100 (61)	65(39)
Overweight or obesity	105 (64)	60(36)
Pregnancy	64 (39)	101(61)
Poor dietary habits	89 (54)	76(46)
Lack of exercise and Sedentary life	94 (57)	71(43)
Complications of DM	Yes. N (%)	No. N (%)
Eye problems or blindness	130 (79)	35(21)
Kidney failure	107 (65)	58(35)
Heart failure	93 (56)	72(44)
Stroke	113 (68)	52(32)
Amputation of limb	117 (71)	48(29)
Prevention of complications related to DM	Yes. N (%)	No. N (%)
Insulin injection	131 (79)	34(21)
Tablets and capsules	126 (77)	39(23)
Regular exercises	131 (79)	34(21)
Healthy diet practice	140 (85)	25(15)
Routine eye check and care	121 (73)	44(27)
Feet and toes medical check up	95 (57)	70(43)
Weight reduction	103 (62)	62(38)

In this study, a range of scores were developed for knowledge. The range for knowledge was 0—31, with a median of eight. Scores below eight were regarded as good, with scores eight and above being regarded as poor knowledge.

Table 2 illustrates the responses to questions on the definition, symptoms and signs, risk factors, complications and prevention of DM. Most participants were able to define DM correctly. Ninety-

nine (60%) knew DM was a disease of inadequate production of insulin, 115 (70%) knew DM was not curable and 99 (60%) knew DM could affect any part of the body. Only 80 (48%) participants knew about the more detailed relationship between insulin and DM.

In general, participants knew about the signs and symptoms of DM. One hundred and seventeen (71%) knew about frequent urination, 122 (74%) about excessive thirst, 121 (73%) about excessive hunger. One hundred and twelve (68%) participants were aware of weight loss, 111 (67%) knew about high blood sugar, 129 (78%) about blurred vision, and 123 (75%) about slow wound healing. Participants, however, did not perform well on some questions related to risk factors for DM. Sixty-four (39%) knew about older age and gestational diabetes (DM that occurs in pregnancy) as risk factors, with 89 (54%), slightly more than half, knowing that poor dietary habits and a sedentary life increased the risk. However, 100 participants (61%) were aware that family history was a risk factor, with 105 (64%) knowing that obesity or being overweight increased the risk.

The majority of participants were aware of DM complications, with 130 (79%) knowing about the eyes being affected, 107 (65%) about the kidneys, 93 (56%) about the heart, 113 (68%) about the brain and 117 (71%) about the limbs. Most participants also knew how to treat DM: 131 (79%) knew insulin injections were used to treat DM, and 126 (76%) knew about the use of tablets and capsules. In terms of lifestyle management, 131 (79%) knew that regular exercise improved DM, 140 (85%) knew that a healthy diet was conducive to the treatment of DM, and 103 (62%) knew that weight reduction assisted in the prevention and management of the disease. In terms of the monitoring and prevention of DM complications, 121 (73%) participants knew of the eye test requirement, and 95 (57%) were aware of the necessity for foot care.

Attitudes and Health Choices

TABLE 3: Frequency distribution of patient responses to attitudes towards DM and self-reported health choices that prevent or mitigate DM, 2020 (N=165)

Attitude towards DM Questions	Agree	Disagree
	N (%)	N (%)
Would you mind disclosing your DM status?	130 (79)	35 (21)
Do you think you should be examined for DM?	148 (90)	17 (10)

Do you think family members should be screened for DM?	128 (77)	37 (23)
Is support from friends and family advisable in addressing DM?	142 (86)	23 (14)
Should we avoid consumption of excessive sugar to control DM?	112 (68)	53 (32)
DM does not seriously affect marriage	60 (36)	105(64)
DM does not affect daily activities	54 (32)	111(68)
Physical exercises can prevent DM	79 (38)	86 (62)
Do you discuss smoking cessation with your clinician?	70 (42)	95 (58)
Maintaining a healthy weight helps in the management of DM	125 (76)	40 (24)
A well-controlled blood glucose level can prevent DM complications	123 (75)	42 (25)

Health choices Questions	Frequent	Less frequent
	N (%)	N (%)
How often do you consume fatty food	48 (29)	117(71)
How often do you exercise 30 - 60 min	90 (54)	75 (46)
How often do you try maintaining your weight	85 (51)	80 (49)
How often do you smoke and drink alcohol	26 (15)	139(85)
How often do you regularly screen for your blood sugar	60 (37)	105(63)

Kassahun and Mekonen defined health choices or practices as the usual participation of the community to prevent a chronic condition, and attitude as the manner in which a community thinks and acts towards a chronic condition.²² For attitude, the score range was 0—11 with a median of seven. Scores of seven and above were regarded as good, and below seven as poor. For health choices the range was 0—5, with median three. Scores of three and above were good and scores below three were poor.

Table 3 illustrates participant responses to the questions assessing their attitudes towards DM, as well as options related to health choices that mitigate DM. Most participants tended to agree with the statements on screening, examination, disclosure to family, supportive relationships and sugar and weight management. However, the responses related to the effect of DM on marriage, daily activities and the management of certain lifestyle factors such as smoking and exercising tended to be more ambiguous.

The participants agreed with the following questions and statements as follows:

- Should be examined for DM (148, 90%)
- Support from friends and family is advisable in addressing DM (142, 86%)
- Maintaining a healthy weight helps in the management of DM (125, 76%)
- A well-controlled blood glucose level may prevent complications (123, 75%)

Of note is that some participants disagreed with the following statements:

- To discuss stopping smoking with their healthcare practitioner (95, 58%)
- DM seriously affects daily activities (111, 68%)
- DM does not seriously affect marriage (105, 64%)

Regarding the questions on health choices, 139 (85%) participants reported that they smoked, or that they drank alcohol less frequently or not at all, whereas 26 (15%) participants reported harmful smoking and drinking habits in the frequent or very frequent ranges. Twenty-nine percent of participants reported that they consumed fatty foods frequently or very frequently, 83 (51%) were conscious of maintaining their weight in the frequent and very frequent ranges, and 75 (46%) participants exercised less frequently or not at all. In terms of screening for DM, presumably in the clinic setting as these participants were non-diabetic, 61 (37%) reported frequent or very frequent blood sugar checks.

Univariate and Multivariate logistic regression

TABLE 4: Univariate and multivariate logistic regression predicting DM-associated knowledge among members attending the CCP, Soweto 2020 (N=165)

Variable category	Knowledge level				# COR (95% CI)	P- value	## AOR (95%CI)	P- value
	Not knowledgeable		Knowledgeable					
	N	(%)	N	%				
Gender								
Male	25	30	28	35	0.8(0.42-1.54)	0.509		
Female (Ref)	59	70	53	65				
Age								
24 and below	14	17	13	16	1.28(0.51-3.21)	0.593		
25-34	22	26	20	25	1.31(0.59-2.92)	0.506		
35-44	22	26	17	21	1.54(0.68-3.5)	0.3		
45 and above	26	31	31	38				
Marital status								
Single (Ref)	59	70	44	54				
Married	11	13	23	28	0.36(0.16-0.81)	0.013	0.33(0.12-0.88)	0.027
Divorced/separated	5	6	0	0	-		-	
Widowed	6	7	7	9	0.64(0.2-2.04)	0.449	0.69(0.19-2.59)	0.587
Cohabiting	3	4	7	9	0.32(0.08-131)	0.112	0.23(0.04-1.27)	0.093
Level of education								
Unable to read & write	3	4	0	0				
Able to read &write	81	96	81	100				
Occupation								
Housewife	14	17	14	17	1.12(0.45-2.8)	0.808		
Student	14	17	11	14	1.43(0.55-3.71)	0.467		
Merchant	2	2	3	4	0.75(0.12-4.84)	0.759		
Farmer	2	2	0	0	-			
Government or Private employer	14	17	15	18	1.04(0.42-2.59)	0.924		

Daily labourer	13	15	10	12	1.46(0.54-3.90)	0.455		
Other(Ref)	25	30	28	35				
Average family income								
0-4999 (Ref)	69	82	57	70				
5000-9999	8	9	20	25	0.33(0.14-0.81)	0.015	0.52(0.18-1.50)	0.224
10000-14999	4	5	0	0	-		-	
15000 and above	3	4	4	5	0.62(0.13-2.88)	0.542	0.58(0.08-4.13)	0.589
Family history of DM								
Yes (Ref)	34	40	65	80				
No	50	60	16	20	5.97(2.97-12.02)	<0.001	6.95(3.06-15.8)	<0.001
Exposure to DM health education								
Yes(Ref)	70	83	69	85	1.15(0.50-2.66)	0.744		
No	14	17	12	15				
Information source								
Media	36	13	24	30				
Health practitioners	19	22	30	37	0.42(0.19-0.91)	0.029	0.79(0.31-1.99)	0.61
Friends or relatives	9	11	19	23	0.32(0.12-0.81)	0.017	0.55(0.18-1.67)	0.291
Other	20	24	8	10	1.67(0.63-4.39)	0.302	1.97(0.64-6.03)	0.236

COR: Crude odds ratio

AOR: Adjusted odds ratio

Table 4 reflects that 81 (49%) participants had knowledge of DM, with 84 (51%) indicating that they did not have knowledge about the disease. All 81 (100%) participants who were able to read and write were knowledgeable about DM. More than half of the participants (57, 71%) who came from the income category R0—4999 knew about DM. The main predictors of knowledge of DM in this study were marital status and family history of DM after adjustment. Those who were married were 0.33 times less likely to have poor knowledge of DM, compared to single people after adjustment ($p=0.027$). Participants with no family history of DM were 6.95 times more likely to have poor knowledge of DM compared to those who had a family history after adjustment ($p<0.001$).

TABLE 5: Univariate and multivariate logistic regression predicting DM-associated attitudes among members attending the CCP, Soweto 2020 (N=165)

Variable category	Attitudes level				COR (95% CI)	P-value	AOR (95%CI)	P-value
	Poor		Good					
	N	(%)	N	%				
Gender								
Male (Ref)	27	41	26	26	0.51(0.27-1.00)	0.05	0.66(0.32-1.35)	0.251
Female	39	59	73	74				
Age								
24 and below (Ref)	11	17	16	17	0.79(0.31-2.01)	0.616		
25-34	16	24	26	26	0.88(0.38-2.00)	0.759		
35-44	19	29	20	20	0.57(0.25-1.31)	0.183		
45 and above	20	30	37	37				
Marital status								
Single (Ref)	41	62	62	63				
Married	14	21	20	20	0.94(0.43-2.08)	0.888		
Divorced/separated	2	3	3	3	0.99(0.16-6.20)	0.993		
Widowed	5	8	8	8	1.06(0.32-3.46)	0.926		
Cohabiting	4	6	6	6	0.99(0.26-3.73)	0.99		
Level of education								
Unable to read & write	2	3	1	1	0.33(0.03-3.67-8)	0.365		
Able to read & write	64	97	98	99				
Occupation								
Housewife (Ref)	9	14	19	19	1.61(0.62-4.23)	0.326		
Student	9	14	16	16	1.36(0.51-3.63)	0.536		
Merchant	2	3	3	3	1.15(0.18-7.46)	0.884		

Farmer	1	1	1	1	0.77(0.05-12.92)	0.854		
Government/Private employer	10	15	19	19	1.46(0.57-3.72)	0.432		
Daily labourer	12	18	11	11	0.70(0.26-1.88)	0.481		
Other	23	35	30	31				
Average family income								
0-4999 (Ref)	48	72	78	79				
5000-9999	15	23	13	13	0.53(0.23-1.21)	0.135		
10000-14999	2	3	2	2	0.62(0.08-4.51)	0.633		
15000 and above	1	2	6	6	3.69(0.43-31.6)	0.233		
Family history of DM								
Yes (Ref)	31	47	68	69				
No	35	53	31	31	0.4(0.21-0.77)	0.006	0.46(0.23-0.94)	0.033
Exposure to DM health education								
Yes(Ref)	54	82	85	56				
No	12	18	14	14	0.74(0.32-1.72)	0.486		
Information source								
Media	29	44	31	31				
Health practitioners	7	11	42	43	5.61(2.18-14.5)	<0.001	4.31(1.62-11.46)	0.003
Friends or relatives	16	24	12	12	0.7(0.28-1.73)	0.442	0.6(0.24-1.53)	0.286
Other	14	21	14	14	0.94(0.38-2.29)	0.884	1(0.4-2.53)	0.983

Table 5 demonstrates that individuals with no family history of DM were 0.46 times less likely to have good attitudes towards DM compared to those with a family history of DM after adjustment ($p=0,033$). Those who obtained their information from health practitioners were 4.31 times more likely to have good attitudes towards DM in comparison to those whose information came from media ($p=0,003$).

TABLE 6: Univariate and multivariate logistic regression predicting DM-associated health choices among members attending the CCP, Soweto, 2020 (N=165)

Variable category	Health choice level				COR (95% CI)	P-value	AOR (95%CI)	P-value
	Poor choices		Good choices					
	N	%	N	%				
Gender								
Male (Ref)	27	34	26	30	0.83(0.43-1.61)	0.588		
Female	52	66	60	70				
Age								
24 and below	10	13	17	20	1.53(0.60-3.91)	0.374		
25-34	23	29	19	22	0.74(0.33-1.65)	0.468		
35-44	19	24	20	23	0.95(0.42-2.14)	0.897		
45 and above	27	34	30	35				
Marital status								
Single (Ref)	47	59	56	66				
Married	14	18	20	23	1.20(0.55-2.63)	0.651		
Divorced/separated	4	5	1	1	0.21(0.02-1.94)	0.169		
Widowed	7	9	6	7	0.72(0.23-2.29)	0.577		
Cohabiting	7	9	3	3	0.36(0.09-1.47)	0.154		
Level of education								
Unable to read & write	2	3	1	1	0.45(0.04-5.09)	0.521		
Able to read & write	77	97	85	99				
Occupation								

Housewife (Ref)	14	18	14	16	1.12(0.45-2.8)	0.808		
Student	9	11	16	19	1.99(0.75-5.30)	0.168		
Merchant	2	3	3	3	1.68(0.26-10.9)	0.586		
Farmer	2	3	0	0	-			
Government or private employer	12	15	17	20	1.59(0.64-3.96)	0.323		
Daily labourer	12	15	11	13	1.03(0.39-2.74)	0.958		
Other	28	35	25	29				
Average family income								
0-4999 (Ref)	64	81	62	72				
5000-9999	10	13	18	21	1.86(0.8-4.34)	0.152		
10000-14999	3	4	1	1	0.34(0.03-3.40)	0.361		
15000 and above	2	2	5	6	2.58(0.48-13.8)	0.268		
Family history of DM								
Yes (Ref)	39	49	60	70				
No	40	51	26	30	0.42(0.22-0.80)	0.008	0.53(0.27-1.05)	0.069
Exposure to DM health education								
Yes (Ref)	66	84	73	85				
No	13	16	13	15	0.90(0.39-2.09)	0.814		
Information source								

Media	38	48	22	25				
Health practitioners	15	19	34	40	3.92(1.75-8.74)	0.001	3.34(1.47-7.62)	0.004
Friends or relatives	11	14	17	20	2.67(1.06-6.71)	0.037	2.36(0.92-6.03)	0.074
Other	15	19	13	15	1.50(0.60-3.71)	0.385	1.57(0.62-3.94)	0.341

Table 6 reflects that participants who received their information from health practitioners were 3.34 times more likely to have good health choices compared to media as information source ($p=0,004$).

DISCUSSION

This study aimed to understand the knowledge, attitudes and behaviour of a peri-urban segment of the population, living under poor socio-economic circumstances. We wanted to understand the spread of ideas and lifestyle choices across different age groups amongst those undiagnosed with DM, thereby providing an opportunity for early intervention.

The first important consideration was the demographics of the study population. Sixty-eight percent of participants were women; consequently the study was skewed towards the knowledge, attitudes and behaviours of women. This may be important because it has been found that women generally have better help-seeking behaviour and possibly better health choices than men,²⁴ although this was not investigated in this study. Conversely, women are generally more socio-economically disadvantaged than men, which may affect their educational attainment and ability to carry out desired health behaviours.²⁴ Compared to the clinic population, the differences between the health behaviours of men and women might be even greater in the general community, which would need further research.

The gender distribution of those attending clinics is not unique to this study. A study on diabetes-related knowledge, attitudes and health choices among adult type 2 DM patients attending community healthcare centres in the Free State Province in South Africa showed that female patients (76.1%) attended clinics more than men (23.9%).²⁵ Men generally present late and have

different help-seeking behaviour.²⁴ A study conducted in Trinidad on the role of gender on attendance and compliance at an outpatient clinic for type 2 DM, found that more women (74.2%) than men (25.8%) attended the clinic and that they had better compliance to treatment than men.²⁶ Men therefore become a special group that may be difficult to access, requiring a different approach to health promotion, lifestyle intervention and prevention and early detection of DM. The same applies to younger people who are harder to access from the clinic base. Attention also needs to be paid to children and obesity, where a family-oriented approach becomes important.

A need for intervention with reference to improved knowledge and change in behaviour was identified in this study, while recognising the complex relationship between knowledge and behaviour change. Studies have shown differing results when considering this complexity. Abbasi *et al.* conducted a knowledge, attitudes and health choices study in 2018 on Type 2 DM patients in the Kuala Muda district of Malaysia, and revealed that good knowledge led to better attitudes and health choices.²⁷ This was also found by Kassahun and Mekonen in 2017.²²

Other studies on diabetic knowledge, attitudes and health choices have shown that poor knowledge correlated with poor attitudes and health choices.^{25, 28, 29, 30} Conversely, it was found that good knowledge correlated with poor attitudes and health choices.¹⁹ A South African study conducted in 2010 confirmed these findings, showing that even though participants were educated on hypertension, which is also a chronic condition, adequate knowledge did not necessarily result in behavioural change.³¹ In this study, however, when considering the relationship between DM knowledge, attitudes and health choices, it was found that poor knowledge did not translate into poor attitudes as even participants who were not knowledgeable about DM had good attitudes and were trying to lead a healthier lifestyle in the absence of diabetes. Although only 81 (49%) of participants amongst the CCP population had good DM knowledge, 99 (60%) had good attitudes towards DM prevention and 86 (52%) had good health choices.

Our study was different in that it looked at the non- or pre-diabetic population. It illuminated the gap in DM knowledge, with less than half (49%) of the participants being knowledgeable, which needs addressing. This study also showed that health workers were more effective than other sources of information at educating patients, as participants who received education from health workers were 4.31 times more likely to develop better attitudes to monitoring, prevention and behaviour change compared to those who obtained information from media.

This leads to a further discussion on the importance of different information sources to educate the community and how they compare, and if they actually result in behaviour change. This study highlighted the fact that participants obtained medical knowledge from different sources. Media was the most common source of information (36%), followed by health practitioners (30%) and friends or relatives (17%). Media is therefore an important source of information, which should be utilised to a greater extent to ensure that correct messaging is achieved. It should be kept in mind that inaccurate health news can cause harm to health and tarnish the reliability of healthcare providers and medicine.³²

This study also showed that healthcare workers were underperforming and needed to improve at providing health education, promotion and prevention. A similar finding was observed in a community in south Nigeria, where 27.9% of participants had health practitioners as their source of information.³³ Patient-friendly delivery of information that is facilitative rather than directive needs implementing and may take place as health messaging to groups of patients by medical staff, or one-on-one in the consulting room by clinicians. It appears that a personalised touch is more effective than media at resulting in behaviour change, which is another area for exploration. Improved interviewing skills and brief motivational counselling skills by clinicians should be harnessed during consultations, as well as community-based interventions that have a more personalised touch and are more likely to translate into behaviour change.

Sixty percent of participants in this study had a positive family history of DM, demonstrating the prevalence and burden of DM and NCDs in this community. Of those, 65 (80%) were knowledgeable about the disease, whereas participants without a family history of DM were 5.97 times less knowledgeable, which was statistically significant ($p < 0.001$). This demonstrates that for most participants a family history translated into better knowledge. It was also shown that those without a family history for DM performed 0.46 times lower on attitudes than those with a family history of DM ($p = 0.033$). Nevertheless, attitudes to lifestyle management can be improved in those with DM in the family. It is important for healthcare workers to take note of this, and it may be important to target and treat not only diabetic patients but also their family units. Due to the strong genetic links of the disease, it could be a case of finding an early lifestyle intervention in such families. A family history of DM is often a missed risk factor in consultations and may link with

undiagnosed DM and impaired fasting glucose information.^{34, 35, 36} This is an area where targeted prevention strategies and effective screening programmes may improve outcomes.

A study conducted in Ethiopia in 2017 confirmed our findings that participants without diabetes in the family had less knowledge than those who did.³⁷ In their study, knowledge was found to be 2.94 times (95% CI 1.87 - 4.86) higher for those who had DM in the family. They also found the probability of having good health choices to be 3.38 times (CI 2.05-5.58) higher compared to participants who did not have a positive family history. These findings underscore that health systems and delivery can be strengthened by focusing on families with diabetic members, where the risk is higher.

In this study, most participants were single (62%) and those who were married were 0.33 times less likely to have poor knowledge of DM compared to single people after adjustment ($p=0.027$). Marriage has been reported as a positive contributor to health-related behaviours and compliance with treatment.³⁸ Involvement of spouses in their partners' disease management improves stress management and the everyday quality of interactions between the couple.³⁹ This information should be carefully sought in consultations using a holistic technique, and patients outside marriage should especially be targeted for improving lifestyle habits, which should also be prioritised within the community. We could not demonstrate a statistically significant association between DM and knowledge, attitudes and health choices, possibly because this study may not have been sufficiently representative.

Other findings in this research indicated that a large proportion of patients experienced ambivalence towards managing important lifestyle risk factors. This was demonstrated in the attitudes section where attitudes towards the effect of chronic diseases on marriage and relationships, effects on daily activities, and choices around exercise and smoking had a larger range of responses. This is an area where targeted prevention strategies may assist in the non-diabetic population to avert disease. In this study, 38% of participants admitted smoking and drinking alcohol. A study involving 732 participants in Limpopo, South Africa, however, revealed that only 13.7% were smokers and only 16.3% consumed alcohol.⁴⁰ These differing findings were confirmed by the results of a study conducted in urban Pretoria and rural Bela Bela in South Africa which showed that the prevalence of both smoking and alcohol was twice as frequent among urban participants compared to rural participants.⁴¹ They also found that tobacco use was highly

associated with the use of alcohol. Based on the above findings, targeting these lifestyle risk factors is therefore an important component in the prevention and management of most NCDs, especially in urban areas.

The current study found that only 52% of patients undiagnosed with DM were practising good health choices. This is of public health concern, given the increasing incidence of NCDs and the consequent implications. This study revealed that 15% of participants were in the range of harmful drinking and smoking, and 29% reported high fat diets, which may be an indicator of poor dietary choices. Not only should media be better utilised to spread information and address behaviour change, but health practitioners should also be trained in the facilitatory techniques of brief behaviour change counselling.⁴² This should be done in health facilities and also extend into communities. Changing behaviour requires a paradigm shift in the way we live and needs careful consideration as we see the impact urbanisation has on health. It is an important target in primary care facilities to promote health and assist people in making healthy choices. This might be challenging, as individuals might not feel in control of their personal circumstances and their environment. Healthcare providers, however, have to be able to assist patients in moving towards empowerment.⁴³

STUDY LIMITATIONS

The study sample was selected from a facility-based population, which may not be representative of the community at large as it did not include those with pre-diabetes who were not utilising the health facility and were therefore presumed healthy. The sample size was small, even though a web-based statistical and epidemiologic calculator for public health was used. Also, the study was conducted in a brief timeframe as a cross-sectional survey and may not describe the whole picture of this peri-urban community that has more than 3 000 households. Women were over-represented in the study, with men generally having poorer help-seeking behaviour and possibly different knowledge, attitudes and health choices. The younger population, who may have riskier lifestyle attitudes and behaviours, were also underrepresented. We could also not demonstrate in this study the relationship between food insecurity and attitudes to health or lifestyle choices. The fact that the median was used may constitute a weakness or limitation in terms of measuring good and poor knowledge, attitudes or health choices.

The study may be biased due to the nature of the study tool that gave participants a tick sheet of options on diabetic knowledge, which may not be a true representation of knowledge. This may have given a false impression of the participants' knowledge, where lack of understanding of questions, fear, embarrassment or dishonesty in reporting accurately may have played a role. Culture and tradition may also apply to the aforementioned considerations. Furthermore, it is always advisable to run the Cronbach's Alpha test after the data collection to support a good internal consistency of the study instrument in the local population, which we did not do. The study had a power of 80 in the sample size, strengthening the analysis adjustment, thereby catering for other biases such as type 1 and type 2 errors.⁴⁴

CONCLUSIONS

From the findings of this study, we concluded that the type of information source and a family history of DM influenced knowledge, attitudes and health choices. The findings also highlighted the important role of healthcare workers in influencing behaviour change. Contrary to previous studies, this study illustrated that poor knowledge of DM did not necessarily translate into poor attitudes in preventing chronic conditions.

The study was useful because it highlighted the gaps where targeted prevention strategies are required to prevent DM and non-communicable diseases (NCDs). This should be tailored to the different groups in the population studied by using media optimally, through health promotion by healthcare workers, and by community-based interventions.

REFERENCES

1. Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat Rev Endocrinol*. 2018; 14(2):88-98. doi.org/10.1038/nrendo.2017.151
2. Cho NH, Shaw JE, Karuranga S, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract*. 2018; 138:271-81. doi: 10.1016/j.diabres.2018.02.023
3. Saeedi P, Salpea P, Karuranga S, et al. Mortality attributable to diabetes in 20-79 years old adults, 2019 estimates: Results from the International Diabetes Federation Diabetes Atlas, 9th ed. *Diabetes Res Clin Pract*. 2020; 162:108086. doi: 10.1016/j.diabres.2020.108086
4. Dessie G, Mulugeta H, Amare D, et al. A systematic analysis on prevalence and sub-regional distribution of undiagnosed diabetes mellitus among adults in African countries. *J Diabetes Metab Disord*. 2020; 19(2):1931-41. doi: 10.1007/s40200-020-00635-9

5. Lalkhen H, Mash R. Multi-morbidity in non-communicable diseases in South African primary healthcare. *S Afr Med J*. 2015; 105(2):134-8. doi: 10.7196/samj.8696
6. Weimann A, Dai D, Oni T. A cross-sectional and spatial analysis of the prevalence of multi-morbidity and its association with socioeconomic disadvantage in South Africa: A comparison between 2008 and 2012. *Soc Sci Med*. 2016; 163:144-56. doi: 10.1016/j.socscimed.2016.06.055
7. Stokes A, Berry KM, Mchiza Z, et al. Prevalence and unmet need for diabetes care across the care continuum in a national sample of South African adults: Evidence from the SANHANES-1, 2011-2012. *PLoS One*. 2017; 12(10): e0184264. doi.org/10.1371/journal.pone.0184264
8. Mortality and causes of death in South Africa, 2015: Findings from death notification. *Afr SS*; 2015. Report No: P0309.3
9. Chan M. Obesity and diabetes: The slow-motion disaster. *Milbank Q* 2017; 95(1):11-14. doi: 10.1111/1468-0009.12238.
10. Damsgaard CT, Michaelsen KF, Molbo D, Mortensen EL, Sørensen TI. Trends in adult body-mass index in 200 countries from 1975 to 2014: A pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016;387(10026):1377-96. doi.org/10.1016/S0140-6736(16)30054-X
11. Gedebjerg A, Almdal TP, Berencsi K, et al. Prevalence of micro- and macrovascular diabetes complications at time of type 2 diabetes diagnosis and associated clinical characteristics: A cross-sectional baseline study of 6958 patients in the Danish DD2 cohort. *J Diabetes Complic*. 2018; 32(1):34-40. doi: 10.1016/j.jdiacomp.2017.09.010
12. Fowler MJ. Microvascular and macrovascular complications of diabetes. *Clin Diabetes*. 2008; 26(2):77-82. <https://doi.org/10.2337/diaclin.26.2.77>
13. Harding JL, Pavkov ME, Magliano DJ, Shaw JE, Gregg EW. Global trends in diabetes complications: A review of current evidence. *Diabetol*. 2019 Jan; 62(1):3-16. doi.org/10.1007/s00125-018-4711-2
14. Young F, Critchley JA, Johnstone LK, Unwin NC. A review of co-morbidity between infectious and chronic disease in Sub-Saharan Africa: TB and diabetes mellitus, HIV and metabolic syndrome, and the impact of globalization. *Global Health*. 2009 Dec; 5(1):1-9. doi: 10.1186/1744-8603-5-9
15. Paulweber B, Valensi P, Lindström J, et al. A European evidence-based guideline for the prevention of type 2 diabetes. *Horm Metabol Res*. 2010 Apr; 42(S 01):S3-6. doi.org/10.1055/s-0029-1240928
16. Ku GM, Kegels G. Effects of the First Line Diabetes Care (FiLDCare) self-management education and support project on knowledge, attitudes, perceptions, self-management practices and glycaemic control: A quasi-experimental study conducted in the Northern Philippines. *BMJ Open Access*. 2014; 4(8). doi: 10.1136/bmjopen-2014-005317

17. Asmelash D, Abdu N, Tefera S, Baynes HW, Derbew C. Knowledge, attitude, and practice towards glycemic control and its associated factors among diabetes mellitus patients. *J Diabetes Res.* 2019; 2019 (Article ID 2593684). doi: 10.1155/2019/2593684
18. Niroomand M, Ghasemi SN, Karimi-Sari H, Kazempour-Ardebili S, Amiri P, Khosravi MH. Diabetes knowledge, attitude and practice (KAP) study among Iranian in-patients with type-2 diabetes: A cross-sectional study. *Diabetes Metab Syndr.* 2016; 10(Suppl 1):S114-S9. doi: 10.1016/j.dsx.2015.10.006.
19. Rathod GB, Rathod S, Parmar P, Parikh A. Study of knowledge, attitude and practice of general population of Waghodia towards diabetes mellitus. *IJCRR.* 2014 Jan 1; 6(1):63-68.
20. DiClemente RJ, Crosby RA, Kegler MC (Eds). *Emerging theories in health promotion practice and research: Strategies for improving public health.* Book Review. San Francisco, CA. Jossey-Bass, John Wiley and Sons. 2002; 335-337.
21. Sullivan KM, Dean A, Soe MM. On academics: OpenEpi: A web-based epidemiologic and statistical calculator for public health. *Public health reports.* 2009 May;124(3):471-4. doi: 10.1177/003335490912400320
22. Kassahun CW, Mekonen AG. Knowledge, attitude, practices and their associated factors towards diabetes mellitus among non-diabetes community members of Bale Zone administrative towns, South East Ethiopia. A cross-sectional study. *PLoS One* 2017; 12(2). doi: 10.1371/journal.pone.0170040
23. Harpe SE. How to analyze Likert and other rating scale data. *Currents in pharmacy teaching and learning.* 2015 Nov 1;7(6):836-50. doi.org/10.1016/j.cptl.2015.08.001
24. Sagar-Ouriaighli I, Godfrey E, Bridge L, Meade L, Brown JS. Improving mental health service utilization among men: A systematic review and synthesis of behavior change techniques within interventions targeting help-seeking. *AJMH.* 2019 Jun; 13(3). doi: 10.1177/1557988319857009
25. Le Roux M, Walsh C, Reid M, Raubenheimer J. Diabetes-related knowledge, attitude and practices (KAP) of adult patients with type 2 diabetes mellitus in the Free State Province, South Africa. *SAJCN* 2019; 32(4). doi.org/10.1080/16070658.2018.1468536
26. Babwah F, Baksh S, Blake L, et al. The role of gender in compliance and attendance at an outpatient clinic for type 2 diabetes mellitus in Trinidad. *Rev Panam Salud Publica.* 2006; 19:79-84. doi:10.1590/S1020-49892006000200002
27. Abbasi YF, See OG, Ping NY, Balasubramanian GP, Hoon YC, Paruchuri S. Diabetes knowledge, attitude, and practice among type 2 diabetes mellitus patients in Kuala Muda District, Malaysia - A cross-sectional study. *Diabetes Metab Syndr.* 2018; 12(6):1057-63. doi: 10.1016/j.dsx.2018.06.025
28. Kiberenge MW, Ndegwa ZM, Njenga EW, Muchemi EW. Knowledge, attitude and practices related to diabetes among community members in four provinces in Kenya: A cross-sectional study. *Pan Afr Med J.* 2010; 7:2.

29. Alaofè H, Hounkpatin WA, Djrolo F, Ehiri J, Rosales C. Knowledge, attitude, practice and associated factors among patients with type 2 diabetes in Cotonou, Southern Benin. *BMC Public Health* 2021; 21(1): 339. doi:10.1186/s12889-021-10289-8
30. Gul N. Knowledge, attitudes and practices of type 2 diabetic patients. *J Ayub Med Col, Abbottabad*. 2010 Sep 1; 22(3):128-31.
31. Nkosi NG, Wright SCD. Knowledge related to nutrition and hypertension management practices of adults in Ga-Rankuwa day clinics. *Curationis*. 2010; 33(2):33-40. doi:10.4102/curationis.v33i2.1083
32. Dhoju S, Rony MMU, Kabir MA, Hassan N. A large-scale analysis of health journalism by reliable and unreliable media. *Stud Health Technol Inform*. 2019; 264:93-7. doi:10.3233/SHTI190190
33. Alele FO, Ilesanmi OS. Knowledge and attitude of a semi-urban community in the south-south region of Nigeria towards diabetes mellitus. *Am J Pub Health Res*. 2014 Apr 15; 2(3):81-5. doi: 10.12691/ajphr-2-3-3
34. Worede A, Alemu S, Gelaw YA, Abebe M. The prevalence of impaired fasting glucose and undiagnosed diabetes mellitus and associated risk factors among adults living in a rural Koladiba town, northwest Ethiopia. *BMC Res Notes*. 2017; 10(1):251. doi.org/10.1186/s13104-017-2571-3
35. Megerssa YC, Gebre MW, Birru SK, Goshu AR, Tesfaye D. Prevalence of undiagnosed diabetes mellitus and it's risk factors in selected institutions at Bishoftu Town, East Shoa, Ethiopia. *J Diabetes Metab*. 2013; 2013:1-7. doi:10.4172/2155-6156.S12-008
36. Omar SM, Musa IR, ElSouli A, Adam I. Prevalence, risk factors, and glycaemic control of type 2 diabetes mellitus in eastern Sudan: a community-based study. *Ther Adv Endocrinol Metab*. 2019; 10: 2042018819860071. doi: 10.1177/2042018819860071
37. Wolde M, Berhe N, van Die I, Medhin G, Tsegaye A. Knowledge and practice on prevention of diabetes mellitus among diabetes mellitus family members, in suburban cities in Ethiopia. *BMC Res Notes*. 2017; 10(1):551. doi:10.1186/s13104-017-2871-7
38. August KJ, Sorkin DH. Marital status and gender differences in managing a chronic illness: The function of health-related social control. *Soc Sci Med*. 2010; 71(10):1831-8. doi:10.1016/j.socscimed.2010.08.022
39. August KJ, Rook KS, Franks MM, Parris Stephens MA. Spouses' involvement in their partners' diabetes management: Associations with spouse stress and perceived marital quality. *J Fam Psychol* 2013; 27(5):712-21. doi:10.1037/a0034181
40. Maimela E, Alberts M, Modjadji SE, et al. The prevalence and determinants of chronic non-communicable disease risk factors amongst adults in the Dikgale health demographic and surveillance system (HDSS) site, Limpopo Province of South Africa. *PLoS One*. 2016 Feb 16; 11(2):e0147926. doi:10.1371/journal.pone.0147926
41. World Health Organization. Substance use in Southern Africa: knowledge, attitudes, practices and opportunities for intervention: summary of baseline assessments in South Africa, the

United Republic of Tanzania and Zambia. World Health Organization; 2003. Available at: <https://apps.who.int/iris/handle/10665/42668> (Accessed 22 June 2022).

42. Morton K, Beauchamp M, Prothero A, et al. The effectiveness of motivational interviewing for health behaviour change in primary care settings: A systematic review. *Health Psychol Rev.* 2015 Jan 1; 9(2):205-23. doi:10.1080/17437199.2014.882006
43. Koelen MA, Lindström B. Making healthy choices easy choices: The role of empowerment. *Eur J Clin Nutr* 2005; 59(Suppl 1):S10-S23. doi:10.1038/sj.ejcn.1602168
44. Jones S, Carley S, Harrison M. An introduction to power and sample size estimation. *EMJ.* 2003 Sep; 20(5):453. Doi: 10.1136/emj.20.5.453