



Understanding the female athlete: molecular mechanisms underpinning menstrual phase differences in exercise metabolism

Tanja Oosthuysen^{1,2} · Juliette A. Strauss³ · Anthony C. Hackney⁴

Received: 12 September 2022 / Accepted: 7 November 2022 / Published online: 19 November 2022
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract

Research should equitably reflect responses in men and women. Including women in research, however, necessitates an understanding of the ovarian hormones and menstrual phase variations in both cellular and systems physiology. This review outlines recent advances in the multiplicity of ovarian hormone molecular signaling that elucidates the mechanisms for menstrual phase variability in exercise metabolism. The prominent endogenous estrogen, 17- β -estradiol (E2), molecular structure is bioactive in stabilizing plasma membranes and quenching free radicals and both E2 and progesterone (P4) promote the expression of antioxidant enzymes attenuating exercise-induced muscle damage in the late follicular (LF) and mid-luteal (ML) phases. E2 and P4 bind nuclear hormone receptors and membrane-bound receptors to regulate gene expression directly or indirectly, which importantly includes cross-regulated expression of their own receptors. Activation of membrane-bound receptors also regulates kinases causing rapid cellular responses. Careful analysis of these signaling pathways explains menstrual phase-specific differences. Namely, E2-promoted plasma glucose uptake during exercise, via GLUT4 expression and kinases, is nullified by E2-dominant suppression of gluconeogenic gene expression in LF and ML phases, ameliorated by carbohydrate ingestion. E2 signaling maximizes fat oxidation capacity in LF and ML phases, pending low-moderate exercise intensities, restricted nutrient availability, and high E2:P4 ratios. P4 increases protein catabolism during the luteal phase by indeterminate mechanisms. Satellite cell function supported by E2-targeted gene expression is countered by P4, explaining greater muscle strengthening from follicular phase-based training. In totality, this integrative review provides causative effects, supported by meta-analyses for quantitative actuality, highlighting research opportunities and evidence-based relevance for female athletes.

Keywords Premenopausal women · Eumenorrhea · Estrogen and progesterone signaling · Exercise metabolism · Exercise-induced muscle damage

Communicated by Michael I Lindinger.

✉ Tanja Oosthuysen
oosthuysen@polka.co.za

¹ School of Physiology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

² Division of Physiological Sciences, Department of Human Biology, University of Cape Town, Sports Science Institute of South Africa, Boundary Road, Newlands, Cape Town 7700, South Africa

³ Research Institute for Sport & Exercise Sciences, Liverpool John Moores University, Liverpool, UK

⁴ Department of Exercise and Sport Science, Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Abbreviations

ACC	Acetyl-CoA carboxylase
Akt	Protein kinase B
AMPK	5'AMP-activated protein kinase
AP-1	Activator protein 1
BCOAD	Branched-chain 2-oxo-acid dehydrogenase
cAMP	Cyclic-AMP
CD36	Fatty acid translocase
CEBPA	CCAAT/enhancer-binding protein alpha
CK	Creatine kinase
CPT-1	Carnitine palmitoyl transferase-1
CREB	CAMP-response element-binding protein
DOMS	Delayed onset muscle soreness
EIMD	Exercise-induced muscle damage
E2	17- β -Estradiol
EF	Early follicular
ER	Estrogen receptor

ERE	Estrogen response elements
ERK	Extracellular signal-related kinases
FABPc	Fatty acid binding protein
FAS	Fatty acid synthase
FATmax	Exercise intensity at maximal fat oxidation
FATP	Fatty acid transport protein
FOXO1	Forkhead box protein O1
FSH	Follicle-stimulating hormone
FSR	Fractional protein synthetic rate
G6Pc	Glucose-6-phosphatase
GLUT4	Glucose transporter 4
GnRH	Gonadotropin-releasing hormone
GPOR	G-protein estrogen receptor
β -HAD	β -Hydroxyacyl-CoA dehydrogenase
IMTG	Intramuscular triglyceride
IRS	Insulin receptor substrate
LCFA	Long chain fatty acid
LF	Late follicular
LH	Luteinizing hormone
LPL	Lipoprotein lipase
MCAD	Medium-chain fatty acyl-CoA dehydrogenase
MF	Mid-follicular
ML	Mid-luteal
mPR	Membrane progesterin receptors
mtGPAT	Mitochondrial glycerol-3-phosphate acyl transferase
mtTFA	Mitochondrial transcription factor A
Myo-D	Myogenic differentiation factor D
NF κ B	Nuclear factor kappa-B
NRF-1	Nuclear respiratory factor-1
OC	Oral contraceptive
OVX	Ovariectomized
p38MAPK	P38 mitogen activated protein kinase
P4	Progesterone
Pax7	Paired box homeotic gene 7
PDK-1	Phosphoinositide-dependent protein kinase-1
PDK-4	Pyruvate dehydrogenase kinase 4
PEPCK	Phosphoenolpyruvate-carboxykinase
PGRMC	Progesterone G-protein receptor membrane complex
PI3K	Phosphoinositide-3-kinase
PPAR	Peroxisome proliferator-activated receptor
PR	Progesterone receptor
PRE	Progesterone response elements
Ra	Rate of appearance
Rac1	Ras-related C3 botulinum toxin substrate 1
Rd	Rate of disappearance
RDI	Recommended daily intake
Slc2a4	Solute carrier family 2 member 4
SOD	Superoxide dismutase
SP-1	Specific protein 1
SREBP1c	Sterol regulatory element-binding protein 1c

T3	Triiodothyronine
T4	Thyroxine
TBC1D	TBC1Domain family member
TCA	Tricarboxylic acid
TFP α	Trifunctional protein alpha

Introduction

Since the early 1980s researchers have observed menstrual phase differences in the metabolic response to exercise that can influence exercise capacity or performance (for example, Jurkowski et al. 1981; Lavoie et al. 1987; Nicklas et al. 1989). Subsequent research evaluated the influence of the menstrual phases or purposeful manipulation of the ovarian hormones on various aspects of carbohydrate, fat, and protein metabolism during exercise to isolate the underlying modulations and the metabolic state that is required for these changes to be notable (for reviews see, Oosthuysen and Bosch 2010; Hackney 2021; Boisseau and Isacco 2022). More recently, a clearer understanding of the molecular pathways and cellular interactions of the ovarian hormones has helped to elucidate the mechanisms that bring about the menstrual phase-specific differences. Understanding the ovarian hormone metabolic signaling mechanisms will improve future research design striving to optimize training and performance in female athletes. This review aims to describe the menstrual phase ovarian hormonal effects on exercise metabolism while highlighting the molecular (signaling) mechanisms. With that objective, this review: (i) defines the menstrual phases; (ii) presents the molecular structure of estrogen as a mechanistic factor instrumental in regulating exercise-induced muscle damage (EIMD); (iii) introduces the receptor-activated signaling pathways of estrogen and progesterone; (iv) reviews the effects of the menstrual cycle on carbohydrate, fat and protein metabolism during exercise and the regulatory signaling mechanisms for each (including meta-analyses where appropriate to describe the magnitude of effect on the oxidation or turnover of the various endogenous energy sources); and (v) introduces the effect of the menstrual cycle on muscle regeneration and strength. Additionally, practical relevance for female athletes and areas warranting further research exploration are highlighted.

Defining the menstrual phases

The ovarian status of women is multiplexed and requires methodological considerations when conducting research in women (Elliott-Sale et al. 2021). The current review focuses on premenopausal, eumenorrheic women who experience natural, ovulatory menstrual cycles and

excludes women on hormonal contraceptives; with reference to studies on women using hormonal contraceptives for isolated comparative purposes only. The reproductive physiology of a eumenorrheic (normal, regular length, ovulatory) menstrual cycle is comprehensively described by Mihm et al. (2011), where the predictable rise and fall in the circulating ovarian hormones, estrogen and progesterone, are explained in the context of the feedback loops of the anterior pituitary hormones, follicle-stimulating hormone (FSH) and luteinizing hormone (LH), which in-turn respond to hypothalamic gonadotropin-releasing hormone (GnRH), to co-ordinate follicle development and regression. This is a worthwhile read for researchers who aim to include female participants in studies. In this current review, we only consider the predictable flux of estrogen and progesterone across a typical eumenorrheic cycle, which is partitioned into the following phases: early follicular (EF), mid-follicular (MF), late follicular (LF), ovulation, early luteal, mid-luteal (ML) and late luteal phases (Fig. 1). It is noteworthy that while a large proportion of premenopausal women experience menstrual

cycles of ~28 days, the length of a eumenorrheic cycle can vary between 21 to 35 days (Elliott-Sale et al. 2021). Mihm et al. (2011) explains that women with the more typical shorter cycle length follow a 2-wave follicular cycle while women with longer cycle lengths follow a 3-wave follicular cycle, with respect to follicle development that culminates in the selection of a dominant follicle. Thus, the longer menstrual cycle in some eumenorrheic women is owing to a longer follicular phase whereas the luteal phase is more consistently ~14 days (Mihm et al. 2011). This means that the time to ovulation is variable but after ovulation has occurred, menses can be expected ~2 weeks later. Janse de Jonge et al. (2019) has outlined what are deemed acceptable methods for predicting and verifying the menstrual phases and ovulation. Accordingly, menstrual phase comparative studies are required to verify the menstrual cycle phases by reporting measured estrogen and progesterone concentrations. To this end, only menstrual phase comparative studies that reported estrogen and/or progesterone concentrations are included in this review. Stricker et al. (2006) measured serum ovarian hormone concentrations in daily samples taken from 20 eumenorrheic women and these median values can be used to guide researchers in defining the acceptable concentration ranges when verifying menstrual phases in female participants (Table 1). Most comparative studies compare EF (or MF) and ML phases because these phases represent the extremes of low and high ovarian hormones (combined estrogen and progesterone), respectively. The LF phase is occasionally included in studies, presenting an ideal time to evaluate the effect of estrogen without progesterone.




Fig. 1 Diagram defining the menstrual cycle phases showing typical serum 17-β-estradiol and progesterone concentrations

Table 1 Reference ranges* for serum 17-β-estradiol (E2) and progesterone (P4) concentrations that define the menstrual phases

Menstrual phase	Recommended timing	Serum E2 (pmol/L)		Serum P4 (nmol/L)		Evidence of ovulation
		Range of median values	Range of 95% CI values	Range of median values	Range of 95% CI values	
EF	Days 1–7 from onset of menses	129–154	58–225	0.6–1.3	0.3–3.8	
MF	8–4 days before the day of ovulation	163–363	113–491	0.3–0.6	0.3–1.6	
LF	0–2 days before the day of ovulation (or LH surge)	651–939	337–1518	0.6–2.5	0.3–4.1	
ML	5–10 days after ovulation	451–552	267–807	32.4–42.5	22.4–60.4	Requires a minimum P4 > 16 nmol/L

EF early follicular, LF late follicular, MF mid-follicular, ML mid-luteal, LH luteinizing hormone

*Reference ranges are provided as the highest and lowest median values and highest and lowest 95% confidence interval (95% CI) values over the specified days according to findings by Stricker et al. (2006) from $n=20$ eumenorrheic women (aged 20–36 years) who supplied daily blood samples for a complete menstrual cycle

Estrogen modulates exercise-induced muscle damage by virtue of its molecular structure

Estrogen and progesterone (P4) are synthesized primarily in the ovaries by canonical steroidogenesis with cholesterol as the primary precursor, and to a lesser extent in other tissues that express the enzyme, aromatase and on availability of a C19 steroid precursor (Barakat et al. 2016). The pathway outlining human steroidogenesis has been schematically constructed (see Häggström and Richfield 2014). This useful reference shows the key enzymes and intersecting paths to produce firstly the progestogens, which progresses to produce either the corticoids or androgens and finally the estrogens. The three main endogenous estrogens include: estrone (E1), 17- β -estradiol (E2), and estriol (E3), with a fourth estrogen, estetrol (E4) only present during pregnancy, where the assigned numeral defines the number of hydroxyl groups. E2 is the most potent and prominent form that is measured in most menstrual phase comparative studies and is used as the default when referring to endogenous estrogen in this review. E2 is metabolized by hydroxylation to form less potent catechol estrogen metabolites. The hydroxylated E2 is then deactivated by methylation via catechol-O-methyltransferase. In this

way, E2 competes with catecholamines for deactivation and can thereby increase the half-life of catecholamine action. Or alternatively, E2 is oxidized to estrone (E1) before being deactivated by conjugation with sulfate or glucuronide groups (Fuentes and Silveyra 2019). All active estrogens, including some estrogen metabolites and estrogen mimetics or dietary phytoestrogens are biologically active by binding to various estrogen receptors (Gregorio et al. 2021).

The molecular structure of E2 (Fig. 2a) also accounts for certain bioactive effects independent of ligand–receptor-activated signaling. The phenolic structure and hydroxyl groups of E2 affords it strong lipophilic and antioxidant potency that are key features in E2 protection of phospholipid membrane stability and integrity against lipid peroxidation (Kendall and Eston 2002). E2 is proposed to act like tocopherol in free radical quenching where the hydroxyl groups function as proton donors to reduce and quench free radicals. As with cholesterol, the phenolic structure of E2 allows it to intercalate in plasma membrane bilayers to improve membrane stability while halting lipid peroxidation chain reactions (Kendall and Eston 2002) (Fig. 2a). These features are particularly relevant for endurance exercise metabolism, where prolonged high rates of oxidative energy production are associated with free radical production that compromises the function and integrity of the sarcolemma resulting in




Fig. 2 Diagram showing the molecular structure of 17- β -estradiol (E2) and a schematic of how it protects plasma membrane integrity (a), that is contrasted to the molecular structure of progesterone (P4) (b) which is not a structural antioxidant

skeletal muscle inflammation and damage. To this end, the post exercise leakage of muscle proteins, such as creatine kinase (CK), into circulation is largely explained by compromised sarcolemma function because of exercise-associated oxidative damage and membrane instability (Tiidus 2003; Enns and Tiidus 2010). For this reason, the change in serum CK activity following exercise is a marker of sarcolemma stability. In a series of early studies, Tiidus and colleagues found that E2 treatment versus placebo in ovariectomized (OVX) rodents attenuated the increase in serum CK activity after exercise and this protection increased markedly with the duration of estrogen therapy (reviewed by Tiidus 2003; Enns and Tiidus 2010). Similarly, estrogen treatment reduced the influx of neutrophils into muscle and the activity of the protease, calpain in muscle following exercise. Tiidus (2003) described the sequence for estrogen's protection as follows: the sarcolemma is stabilized by estrogen, which results in less calcium leaking into the cytosol and thus less calpain activation. Reduced calpain activity results in less expression of chemoattractant peptides that explains the decrease in neutrophil infiltration and overall reduced muscle damage; encompassing the primary pro-inflammatory phase of the muscle damage/repair process (Peake et al. 2017). Therefore, it is not surprising that sex-differences in resting serum CK activity and post exercise CK response have been reported with lower resting CK levels in women and reduced post exercise CK response (Sewright et al. 2008; Luk et al. 2021; Kumagai et al. 2022) or quicker return to resting levels in women than men (Oosthuyse and Bosch 2017). Furthermore, given that population variability in the genotypes of alleles within genes can translate into some differences in the functionality of the expressed protein, it is interesting that men having a TT genotype polymorphism in the gene for the E2 producing enzyme, aromatase have higher E2 concentrations and reduced CK response after an ultramarathon than men with CT/CC genotype (Kumagai et al. 2022). Moreover, recent evidence showed women to have a muted intramuscular cytokine response to muscle damaging exercise compared with men (Luk et al. 2021) with previous evidence of reduced infiltration of granulocytes into the muscle in women versus men (Stupka et al. 2000). Interestingly, while the serum CK and delayed onset muscle soreness (DOMS) recovery times are associated in men, they are unrelated in women (Sewright et al. 2008; Oosthuyse and Bosch, 2017). Nociceptive sensitivity may be influenced by menstrual phase, causing DOMS to poorly reflect restoration of sarcolemma function and muscle repair in women (Oosthuyse and Bosch 2017).

Considering that E2's plasma membrane protection depends partly on the molecular structure of E2 and thus only appreciated when E2 is present and possibly more so with elevated concentrations, it is likely menstrual phase specific. Accordingly, the increase in serum CK activity 24 h

and 48 h after 20 min of downhill running was negatively correlated with serum E2 concentrations in a combined sample of eumenorrheic women who participated once in either their EF, LF or ML phase (Oosthuyse and Bosch 2017). Likewise, serum CK activity was consistently lower 24 h and 72 h after 60–90 min of level running at 70% VO_2 peak in the ML compared with MF phase (Williams et al. 2015; Hackney et al. 2019); with a negative correlation between the increase in serum CK activity and serum E2 concentrations (Williams et al. 2015) and a decreased serum interleukin-6 response at all time points after exercise in the ML versus MF phase (Hackney et al. 2019). A recent meta-analysis attempted to quantify the effect of the menstrual phases on markers of EIMD by including all prior studies in females and found no clear menstrual phase effect on the serum CK response (Romero-Parra et al. 2021). However, the inclusion of studies that only considered a single menstrual phase or did not provide adequate verification of menstrual phases by means of ovarian hormone concentrations suggests the outcome of that meta-analysis should be graded as low certainty and interpreted with caution. Such a between-study comparison of effect lacks sensitivity owing to differing severities of exercise protocols that would undoubtedly influence the CK response variably independent of menstrual phase. Furthermore, when an unaccustomed mechanical stress is extremely high, such as during maximal force eccentric exercise, the severity of structural damage to muscle fibers might not be mitigated by the effect of E2 to preserve membrane function. In that case, a similar large serum CK response may occur irrespective of menstrual phase (for example see, Funaki et al. 2022). Therefore, menstrual phase differences in EIMD due to E2 ability to stabilize membranes and protect against lipid peroxidation may be expected following endurance-type exercise (for example, Williams et al. 2015; Oosthuyse and Bosch, 2017; Hackney et al. 2019), which imposes lower muscle forces over prolonged periods requiring a high flux of oxidative phosphorylation.

The cellular protective effect of E2 persists even in the presence of the estrogen receptor inhibitor, tamoxifen (Paroo et al. 2002), supporting the functional role of the molecular structure of E2 to be instrumental in conferring these benefits independent of certain typical ligand–receptor signaling. However, E2 signaling is complex and multifaceted with several direct and indirect genomic and non-genomic paths of action that are introduced herein. Indeed, E2 signaling does increase the expression of the antioxidant defense enzymes, superoxide dismutase (SOD) (Strehlow et al. 2003) and glutathione peroxidase (Viña et al. 2008). And therefore, the antioxidant role of E2 appears to be a sum of its role as both a structural antioxidant and genomic activator. Agreeably, a recent study quantified the presence of reactive oxygen metabolites and tested the antioxidant potential in blood samples

taken before and after high-intensity intermittent exercise in women during their EF and ML phases (Matsuda et al. 2020). They reported a 10% increase in circulating reactive oxygen metabolites in the EF, but not ML phase after exercise, together with a tendency for a greater increase in antioxidant potential after exercise in ML than EF phase. Albeit the ML phase is associated with an increase in both E2 and P4.

P4, unlike E2, is not a structural antioxidant and is susceptible to reduction at the ketone groups or carbon double bond (positioned between C4 and C5; from where the abbreviation, P4 derives its numeral) (Fig. 2b). P4, however, has been shown to reduce free radical damage by increasing the expression and activity of the antioxidant defense enzymes, SOD, glutathione peroxidase, and glutathione reductase via P4 receptor signaling (Hernández-Rabaza et al. 2019). For this reason, the synergistic effect of E2 and P4 in the ML phase to reduce oxidative stress supports evidence for the ML phase to be better than the EF phase for exercise that risks EIMD.

Although beyond the intended scope of this review, it must be noted that the condition of oral contraceptive (OC) use may alter these responses as discussed previously (Oosthuyse and Bosch 2017). Furthermore, recent accumulation of evidence supports a higher level of oxidative stress and inflammatory markers such as C-reactive protein in female athletes taking combined OCs compared with non-OC users (Cauci et al. 2017, 2021; Larsen et al. 2020; Quinn et al. 2021).

E2 and P4 receptor-activated signaling

Ligand–receptor-activated signaling pathways of E2 and P4 are mechanistically similar and have been reviewed in detail (for E2 signaling see: Foryst-Ludwig and Kintscher 2010; Yaşar et al. 2016; Fuentes and Silveyra 2019; Puglisi et al. 2019) (for P4 signaling see: Garg et al. 2017; Gonzalez 2020; Medina-Laver et al. 2021). Signaling occurs either via the classical path of binding nuclear hormone receptors




Fig. 3 Diagram showing the receptor-activated signaling pathways of estrogen (E2) and progesterone (P4). The classical direct genomic effects (a) of E2 and P4 occur via cytosolic binding of respective nuclear receptor transcription factors, estrogen receptors (ER) and progesterone receptors (PR) in dimer complexes that translocate to the nucleus to regulate gene expression on binding their respective response elements (ERE or PRE, respectively) in the promoter of target genes or by binding certain DNA-bound transcription factors (TF) to transactivate or trans-repress target gene expression. In the same way, E2 regulates mitochondrial DNA expression. The indirect genomic and rapid non-genomic effects (b) occur via binding plasma membrane-bound receptors complexed with G proteins, namely, ER and PR or alternate membrane receptors, G-protein coupled estrogen

receptor (GPER) and progesterone receptor membrane complex 1 (PGRMC1), membrane progestin receptors (mPRs), respectively. On E2 and P4 binding, these membrane-bound receptors activate various signaling kinases that phosphorylate DNA-bound ER or PR or other TF to regulate target gene expression indirectly. In addition, the activated kinases bring about other rapid non-genomic cellular responses and membrane-bound ER signaling regulates endoplasmic/sarcoplasmic reticular calcium (Ca^{2+}) ion channel activity to increase cytosolic Ca^{2+} content and related signaling. *p38MAPK* p38 mitogen activated protein kinase, *ERK1/2* extracellular signal-related kinases1/2, *PI3K* phosphoinositide-3-kinase, *Akt* RAC (Rho family)-alpha serine/threonine-protein kinase (also known as protein kinase B), *AMPK* AMP-activated protein kinase

or the more recently discovered alternate route of binding membrane receptors (Fig. 3), as detailed below.

Nuclear hormone receptors

Both E2 and P4 influence metabolism by binding estrogen and progesterone nuclear hormone receptors that occur as E2 receptor-alpha ($ER\alpha$) and E2 receptor-beta ($ER\beta$) or P4 receptor-A (PR-A) and P4 receptor-B (PR-B) isoforms, respectively. Receptor isoform specificity is relevant because the various receptor isoforms are linked to differing metabolic outcomes (Foryst-Ludwig and Kintscher, 2010; Mauvais-Jarvis et al. 2013; Gregorio et al. 2021). E2 and P4 bind to the respective isoforms with equal affinity and the net outcome is, therefore, dependent on the tissue-specific isoform abundance (Yaşar et al. 2016). E2 and P4 diffuse through plasma membranes to bind their respective nuclear hormone receptors in the cytosol. On binding, they translocate to the nucleus where the ligand bound receptors assemble typically as dimer complexes to act as ligand-activated transcription factors and bind estrogen response elements (ERE) or progesterone response elements (PRE), respectively, in the promoter regions of target genes to activate or repress gene transcription (Yaşar et al. 2016; Puglisi et al. 2019). E2 and P4 bound to their cognate nuclear hormone receptors can also regulate expression of genes that do not have ERE or PRE in the promoter regions by binding to other DNA-bound transcription factors of those respective target genes (Foryst-Ludwig and Kintscher, 2010; Yaşar et al. 2016; Puglisi et al. 2019; Gonzalez 2020). The relevant nuclear gene targets (or proteins of gene targets) regulated by E2 and P4 are made known in the sections to follow while reviewing each of the metabolic pathways.

Mitochondrial DNA also includes EREs and E2-ER binding on mitochondrial DNA has been shown to enhance the expression of a protein subunit of ATP synthase and mitochondrial manganese SOD expression (Yaşar et al. 2016). Furthermore, nuclear E2-ER binding activates nuclear respiratory factor-1 (NRF-1) expression, which subsequently activates nuclear expression of mitochondrial DNA transcription factors such as, mitochondrial transcription factor A (mtTFA) (Yaşar et al. 2016; Ventura-Clapier et al. 2019). In this way, E2 promotes mitochondrial biogenesis and capacity for oxidative phosphorylation. Both $ER\alpha$ and $ER\beta$ isoforms have been identified in mitochondria as well as membrane-bound E2 receptors (Ventura-Clapier et al. 2019).

Membrane-bound receptors

E2 and P4 induce rapid signaling in seconds or minutes by binding membrane-bound receptors coupled to G proteins that activate various kinases, namely: phosphoinositide-3-kinase (PI3K)/Akt, extracellular signal-related kinases

(ERK), p38 mitogen activated protein kinase (p38MAPK), and others, and thereby E2 and P4 act in a non-genomic manner to phosphorylate and regulate the activity of other cellular proteins (Puglisi et al. 2019; Gonzalez 2020). The activated kinases can also phosphorylate nuclear bound E2 or P4 nuclear hormone receptors that suffices to activate or repress expression of target genes having ERE or PRE in their promoter regions without the need for direct E2 or P4 ligand binding (Yaşar et al. 2016; Puglisi et al. 2019). Similarly, the activated kinases can phosphorylate various other nuclear bound transcription factors to indirectly regulate gene expression of other target genes lacking ERE or PRE and without any interaction of E2 and P4 nuclear hormone receptors (Yaşar et al. 2016; Puglisi et al. 2019). In this way, E2 and P4 can impose indirect genomic regulation when binding their membrane-bound receptors. Furthermore, E2 regulates calcium ion channels and sarcoplasmic/endoplasmic reticular calcium transporters to modulate cytosolic and mitochondrial calcium content and associated signaling cascades (Mahmoodzadeh and Dworatzek 2019; Puglisi et al. 2019). E2 is also functional in the phosphorylation of 5'AMP-activated protein kinase (AMPK) and rapidly increases its activity within less than 10 min of administering E2 in vitro or in situ (D'Eon et al. 2005, 2008; Rogers et al. 2009; Gorres et al. 2011), which in skeletal muscle occurs selectively via $ER\alpha$ but not $ER\beta$ (Gorres et al. 2011). Conversely, in vascular endothelial cells, E2 binding $ER\beta$ (but not $ER\alpha$) increased intracellular calcium content and activation of calcium calmodulin-dependent protein kinase kinase- β to phosphorylate AMPK, independent of the more typical upstream AMPK kinase, liver kinase B1 (Yang and Wang, 2015); thus, emphasizing tissue-specific signaling.

It is proposed that the membrane-bound receptors can include: (1) the classical E2 and P4 nuclear hormone receptors that are in this instance membrane-bound and (2) further receptor family classes referred to as G-protein estrogen receptors (GPER) (Puglisi et al. 2019) or progesterone G-protein receptor membrane complex (PGRMC) and membrane progesterin receptors (mPR) (Garg et al. 2017; Gonzalez 2020; Medina-Laver et al. 2021). Recent evidence in adipocytes suggests that of the E2 classical nuclear hormone receptor isoforms, only $ER\alpha$ translocate to the plasma membrane in the presence of E2, whereas $ER\beta$ remain nuclear bound (Campello et al. 2017; Gregorio et al. 2021). For this reason, it is likely that E2 membrane-receptor signaling and rapid non-genomic effects occur mostly via specifically E2- $ER\alpha$ or E2-GPER interactions and not E2- $ER\beta$; albeit possibly tissue specific. Studies in mice report expression of $ER\alpha > GPER > ER\beta$ in both soleus (type 1 fibers) and extensor digitorum longus (type 2 fibers) muscles (Baltgalvis et al. 2010) and expression of $ER\alpha > ER\beta$ in not only skeletal muscle but also liver and adipose (Ribas et al. 2010).

P4 represses ER expression

Evidence exists for E2 and P4 cross-interactions that are often antagonistic (Campbell and Febbraio 2001, 2002) with the net outcome in the luteal phase dependent on the relative serum concentrations. We now know that P4 antagonism of E2 can in part or in whole be explained by P4 inhibition or repression of ER expression and protein content (Jayaraman and Pike 2009; Ekenros et al. 2017). In an in vitro study in neurons, E2 promoted ER α and ER β expression and P4 downregulated this effect rapidly within 15 min and the level of repression increased with greater incubation time (Jayaraman and Pike 2009). P4 repression was maximal at typical ML concentrations of 30 nmol/L (Jayaraman and Pike 2009). Furthermore, skeletal muscle ER and PR expression and protein content have been compared between EF, ovulation, and ML phases in sedentary eumenorrheic women, where ER mRNA and protein content was found to be lowest in the ML phase (Ekenros et al. 2017). The order of skeletal muscle ER α mRNA content occurred as: EF > ovulation > ML that coincided with typical serum P4 concentrations that were lowest in EF, modestly higher at ovulation, and substantially higher in ML. Accordingly, ER α protein content was greater in the EF phase than ML phase. In that study, the modest serum E2 concentrations reported during the ovulation sampling period (200 pmol/L) might suggest that sampling may have occurred in the recent post ovulation phase when E2 concentrations fall off rapidly and it would have been insightful had sampling been conducted in the LF phase coincident to peak E2 concentrations. Moreover, PR protein content was found to be greater in the ML than both at ovulation and EF phase suggesting that P4 promotes PR abundance, which conversely does not appear to be antagonized by the concomitant increase in E2 in the ML phase (Ekenros et al. 2017) and PR expression is activated by E2-ER signaling (Medina-Laver et al. 2021). These mechanistic uncovering of P4's modulation of E2 signaling supports early suggestions to consider the relative concentrations of E2 and P4 in the ML phase when interpreting outcomes (D'Eon et al. 2002; Horton et al. 2002).

Mechanisms behind menstrual cycle effects on carbohydrate metabolism

Overview of exercise-stimulated plasma glucose uptake

Our understanding of insulin signaling, and exercise-stimulated plasma glucose uptake has advanced. The basic framework of the insulin signaling pathway includes insulin binding to membrane-bound insulin receptor tyrosine kinase that phosphorylates insulin receptor substrate (IRS)

that then binds phosphoinositide-3-kinase (PI3K) resulting in the generation of PI-3,4,5-triphosphate to activate phosphoinositide-dependent protein kinase-1 (PDK-1) and initiates two signaling pathways. Firstly, the PI3K pathway phosphorylates and activates serine/threonine-protein kinase B also known as Akt, which phosphorylates the Rab-GTPase activating proteins, TBC1Domain family member1 (TBC1D1) or TBC1D4 (also known as AS160), and thereby relieves inhibition on Rab-GTPase that then facilitates translocation of GLUT4 storage vesicles to the plasma membrane (Sakamoto and Holman 2008). Secondly, PI3K activates Ras-related C3 botulinum toxin substrate 1 (Rac1), a Rho-family GTPase, that regulates scaffold proteins and remodels the actin cytoskeleton to facilitate GLUT4 translocation and therefore plasma glucose uptake (Yue et al. 2020). Exercise-stimulated plasma glucose uptake acts independently of insulin and the current viewpoint is that it is partly reliant on AMPK to directly phosphorylate TBC1D1/4 (de Wendt et al. 2021); although evidence suggests that while AMPK is essential for increasing glucose uptake after exercise, it is not essential during exercise (Kjøbsted et al. 2019; McConnell, 2020). Instead, exercise-stimulated glucose uptake is promoted by mechanical stress or stretch that activates integrin receptors and focal adhesions, or dystrophin-glycoprotein complexes, which then promote Rac1 activity and GLUT4 translocation (Sylov et al. 2017; Flores-Opazo et al. 2020) (Fig. 4). However, it is also possible that the AMPK and Rac1 pathways driving exercise-stimulated glucose uptake are inter-related (Yue et al. 2020; de Wendt et al. 2021). A recent viewpoint suggests that the increase in glucose uptake with exercise is not only dependent on GLUT4 translocation but also possibly an increase in GLUT4 activity (Richter, 2021). Furthermore, during exercise it can be assumed that all the plasma glucose taken up is shuttled into glycolysis and oxidation (Sylov et al. 2017) and therefore the measured rate of plasma glucose uptake is often used as a surrogate for the rate of plasma glucose utilization during exercise.

E2 and P4 influence of plasma glucose uptake

E2 and P4 signaling feed into insulin signaling pathways (Wada et al. 2010; Mauvais-Jarvis et al. 2013) and in addition E2 influences GLUT4 expression (reviewed by Gregorio et al. 2021) to affect glucose uptake (Fig. 4), as outlined in the sections that follow.

E2 regulation of GLUT4 expression

While there is no ERE in the promoter of the gene (solute carrier family 2 member 4, *Slc2a4*) encoding GLUT4, E2-ER exerts genomic effects by binding various *Slc2a4* transcription factors (Gregorio et al. 2021). E2 influence is dependent on the ER isoform where E2-ER β trans-represses




Fig. 4 Diagram showing E2 and P4 signaling to influence exercise carbohydrate metabolism. In skeletal muscle, E2 transactivates or trans-represses GLUT4 gene expression via ER α or ER β , respectively, that bind specific DNA-bound transcription factors (1). GLUT4 translocation to the plasma membrane is promoted by E2 independent of insulin via membrane-bound ER α that directly stimulates phosphoinositide-3-kinase (PI3K), which then signals via protein kinase B (Akt) (2a) or Ras-related C3 botulinum toxin substrate 1 (Rac1) (2b) and via independent phosphorylation of AMP-activated protein kinase (AMPK) (3). P4 antagonizes these E2 effects mainly by inhibition of ER expression and protein content. In hepatocytes, E2 via PI3k/Akt signaling phosphorylates forkhead box protein O1 (FOXO1) to cause its nuclear export and degradation and thereby

trans-represses expression of gluconeogenic target genes (4). Conversely, in a glucose-deprived state typical of fasted exercise, P4 transactivates expression of these gluconeogenic genes via binding membrane-bound PGRMC1 that activates adenylate cyclase and cyclic-AMP production that activates protein kinase A (PK-A) to phosphorylate and activate the transcription factor, cAMP-response element-binding protein (CREB) (5). *CEBPA* CCAAT/enhancer-binding protein alpha, *G6Pc* glucose-6-phosphatase, *IRS* insulin receptor substrate, *NF κ B* nuclear factor kappa-B, *PDK1* phosphoinositide-dependent protein kinase-1, *PEPCK* phosphoenolpyruvate-carboxykinase, *PIP3* PI-3,4,5-triphosphate, *Rab* Rab-GTPase, *SP-1* specific protein-1, *TBC1D1/4* TBC1Domain family member1 or 4 (also known as AS160)

and E2-ER α transactivates GLUT4 expression (Campello et al. 2012; Barreto-Andrade et al. 2018). For example, E2-ER α binds the inflammation-stimulated transcription factor, nuclear factor kappa-B (NF κ B), and thereby blocks its repression of GLUT4 expression (Campello et al. 2012); E2-ER α binds specific protein 1 (SP-1) on the promoter of *Slc2a4* causing transactivation of GLUT4 expression (Barreto-Andrade et al. 2018); E2-ER α binds CCAAT/enhancer-binding protein alpha (CEBPA) to transactivate GLUT4 expression and also directly increases the expression and content of CEBPA and promotes its translocation to the nucleus (Fatima et al. 2019). The mechanism explaining ER β trans-repression of GLUT4 expression is not clear but may be via inhibition of SP-1 (Gregorio et al. 2021).

P4 alone did not change adipocyte GLUT4 expression in vitro (Wada et al. 2010) but did reduce GLUT4 protein content in adipose tissue and skeletal muscle of OVX rodents compared with sham-operated rodents

(Campbell and Febbraio 2002). GLUT4 content was suppressed even when P4 was administered with E2 and P4 inhibition was only overcome with a supraphysiological E2 dose (Campbell and Febbraio 2002). Thus, P4 effect on GLUT4 expression or protein content may be indirect possibly in part due to P4 repression of ER α expression. Accordingly, skeletal muscle GLUT4 mRNA content was greater in women in their MF phase compared with men and was increased twofold in men after 8 days of oral E2 supplementation that mimicked LF phase serum E2 concentrations (946 pmol/L) (Fu et al. 2009). Furthermore, a comparison between menstrual phases revealed a trend for 2.2-fold greater skeletal muscle GLUT4 mRNA content during the MF phase than luteal phase (Fu et al. 2009), where the luteal phase was characterized by modest serum E2 (203 pmol/L) and P4 (6 nmol/L) concentrations more typical of the early or late luteal phase and possibly represented a P4-dominant effect.

E2 and P4 regulation of GLUT4 translocation

E2 binding specifically ER α at the plasma membrane induces PI3K/Akt phosphorylation and promotes GLUT4 translocation independent of insulin (Barros et al. 2006; Rogers et al. 2009; Gorres et al. 2011; Campello et al. 2017). Thus, E2-ER α can initiate both the classical TBC1D1/4 pathway and Rac1 pathway. In addition, E2-ER α phosphorylation of AMPK initiates AMPK phosphorylation of TBC1D1/4 (Rogers et al. 2009; Gorres et al. 2011).

In this way, E2 via ERA increases GLUT4 expression and GLUT4 translocation to the plasma membrane and is thus expected to enhance exercise-stimulated plasma glucose uptake. Accordingly, exercise-stimulated glucose uptake is 1.4-fold greater in the skeletal muscle of female compared with male mice (Kang et al. 2021). Furthermore, endurance training increases ER α abundance in skeletal muscle (Lemoine et al. 2002; Wiik et al. 2005). Therefore, E2's effect on plasma glucose uptake might be expected to be most notable in endurance trained women.

However, P4 decreases insulin-stimulated glucose uptake by decreasing the abundance of IRS and interfering with signaling distal to IRS and thereby reduces GLUT4 translocation in adipocytes (Wada et al. 2010); albeit possibly tissue specific considering a P4 positive effect on glucose uptake and GLUT4 expression in neurons (Wu et al. 2019). Accordingly, exercise-stimulated plasma glucose uptake in skeletal muscle was reduced by OVX in rodents and restored after 15 days of treatment with E2 alone, but not with P4 alone (Campbell and Febbraio 2002). P4 when combined with E2, prevented the increase in exercise-stimulated glucose uptake that could be overcome with a supraphysiological dose of E2 (Campbell and Febbraio 2002). For these reasons, the relative increase in E2 and P4 in the ML phase will determine the net outcome for potential menstrual phase changes specifically to skeletal muscle exercise-stimulated glucose uptake and GLUT4 content, with the LF phase likely presenting the most favorable time for this E2-dominant effect to be realized and certainly warrants investigation in eumenorrheic athletes.

However, despite convincing evidence for the tissue-specific effects of E2 to enhance exercise-stimulated glucose uptake, this can be constrained by ovarian hormone influences on whole-body energy homeostasis via multi-tissue governance, as uncovered in the sections that follow.

Whole-body plasma glucose uptake during exercise

A substantial number of studies have measured whole-body plasma glucose uptake during exercise either during different menstrual phases in fed (Campbell et al. 2001; Suh et al. 2002) or fasted (Campbell et al. 2001; Zderic et al. 2001; Horton et al. 2002; Devries et al. 2006) eumenorrheic

women, or after purposeful controlled manipulation of ovarian hormone concentrations in non-OC users (D'Eon et al. 2002), or in amenorrheic women (Ruby et al. 1997) and men (Carter et al. 2001; Devries et al. 2005) on short-term E2 treatment. We have combined the findings from these studies in a summative meta-analysis; where the mean (and SD) rate of plasma glucose uptake measured in $\mu\text{mol/kg}\cdot\text{min}$ in the EF phase/placebo trial is compared with the ML phase/hormone supplemented trial in each study to derive a mean difference and effect size as the standardized mean difference using Review Manager (RevMan version 5.4.1) (Fig. 5a).

While most studies included in the meta-analysis did apply a within-participant crossover design, results were only presented as between-individual means (and SD) for each phase or condition instead of the mean within-individual change. For this reason, the findings presented reflect independent between-group and not dependent within-group outcomes and the meta-analysis outcome likely underestimates the true effect, adopting a conservative approach. In all studies, stable isotopic-tracer methodology was applied by constant intravenous infusion of [6,6- ^2H]glucose to attain steady-state enrichment before plasma glucose rate of appearance (Ra) and rate of disappearance (Rd) could be derived from validated equations based on the principles of tracer dilution (Kim et al. 2016), where glucose Rd represents the rate of whole-body plasma glucose uptake. The meta-analysis outcome shows, first, in agreement with the strong antagonistic effect of P4 as discussed above, the ML phase is associated with a decrease in the rate of plasma glucose uptake compared with the EF phase, with a moderate effect during fasted exercise and a small non-significant effect when exercising fed (Fig. 5a). Therefore, the ovarian hormone effects are more notable during periods of metabolic stress or a high demand for endogenous substrates. Second, while we may expect conditions of short-term E2 treatment alone in men or amenorrheic women to increase the rate of plasma glucose uptake during exercise, instead studies also report a significant decrease with E2 treatment compared with placebo (Fig. 5a). This seemingly discrepant finding may be a result of men and amenorrheic women, who experience chronic low E2 concentrations, also having low ER α content in skeletal muscle and greater relative ER β content as is typical in E2 deficient conditions, such as menopause (Foryst-Luwig and Kintscher, 2010; Park et al. 2017). However, in the study by D'Eon et al. (2002), where ovarian hormones in eumenorrheic women were suppressed with a GnRH antagonist followed by treatment with E2 alone or combined E2 and P4, the rate of plasma glucose uptake during exercise also tended to be reduced with E2 treatment (or E2 plus P4) compared with ovarian suppression. This suggests that despite prior evidence for tissue-specific effects in skeletal muscle for E2 to increase exercise-stimulated glucose uptake, on a whole-body level E2 reduces the rate of



a**Plasma glucose Rd ($\mu\text{mol/kg}\cdot\text{min}$)****b****Plasma glucose Ra ($\mu\text{mol/kg}\cdot\text{min}$)**

Fig. 5 Forest plot showing the mean difference (95% confidence interval CI) of menstrual phase or estrogen (E2) supplement treatment on rate of plasma glucose uptake (Rd) (**a**) and rate of plasma glucose appearance (Ra) (**b**) during exercise when fasted or when fed;

showing the heterogeneity indices, z-scores, p-values, and standardized mean difference as effect size (ES) for each outcome. Effect size scores are interpreted as: 0.0–0.2 is trivial; 0.2–0.6 is small; 0.6–1.2 is moderate; > 1.2 is large effects, respectively

plasma glucose uptake during exercise. The current meta-analysis demonstrates an overall outcome for the ML phase/E2 treatment compared with the EF phase/placebo treatment to produce a small negative effect of $-4.18 \mu\text{mol/kg}\cdot\text{min}$ on the rate of plasma glucose uptake during exercise lasting 50–120 min at 50–70% $\text{VO}_{2\text{max}}$ (Fig. 5a), that equates to -2.7 g/h of glucose taken up for 60 kg women. Such a magnitude of effect may appear to have little physiological relevance over the duration of exercise that has been tested, but it remains to be determined whether longer duration exercise produces larger and more meaningful menstrual phase effects.

Moreover, oral contraceptive therapy for 4 months has been found to reduce the rate of plasma glucose uptake during exercise by up to 20% ($-8.44 \mu\text{mol/kg}\cdot\text{min}$) even when exercising fed (3 h postprandial) compared with before-OC use in eumenorrheic women (Suh et al. 2003). It may be that OC potency is greater than the naturally cycling endogenous hormones resulting in a larger effect that becomes notable even with less metabolic stress. Such a decrease in the rate of plasma glucose uptake with OC use versus non-OC use equates to -5.5 g/h of glucose taken up for 60 kg women, an amount that represents 5% of total carbohydrate oxidation (reported as 1.75 g/min or 105 g/h) during exercise at 65% $\text{VO}_{2\text{max}}$ (Suh et al. 2003). Some suggest that the oral route of delivery of hormonal contraceptives imposes high hepatic exposure via the portal vein, which could magnify ovarian hormone receptor binding and downstream signaling responses in hepatocytes (Magkos et al. 2022).

As is typically expected, the rate of plasma glucose uptake (R_d) approximates the rate of glucose appearance in plasma (R_a) under the rigorous homeostatic regulation of blood glucose concentration (Kim et al. 2016). Accordingly, menstrual phase or hormonal conditional outcomes on the rate of plasma glucose appearance during exercise mimics the rate of plasma glucose uptake and is similarly presented in a summative meta-analysis (Fig. 5b). During fasted exercise, plasma glucose R_a largely reflects hepatic glucose production derived from hepatic glycogenolysis and gluconeogenesis. Considering evidence for E2 to increase skeletal muscle-specific glucose R_d capacity during exercise, and yet whole-body plasma glucose R_d is reduced in the presence of elevated E2 (and P4) specifically during fasted exercise, it is likely that plasma glucose R_d in ML (and possibly LF) phase is limited by plasma glucose R_a and the influence of E2 (and P4) on hepatic glucose production.

Ovarian hormone influence on hepatic glucose production

In the 1970s, a study on rodents reported that 21-days of E2 and P4 treatment suppressed gluconeogenesis measured during the intravenous delivery of isotopic-glucogenic

precursors at rest (Matute and Kalkhoff 1973). More recently, the respective molecular signaling mechanisms have been uncovered. A study using OVX mice with E2 treatment alone or in combination with an Akt inhibitor or ER α inhibitor found that E2 suppresses gluconeogenesis and thus hepatic glucose production (Yan et al. 2019). E2 acts by binding ER α at the plasma membrane to induce PI3K and Akt phosphorylation, where Akt, in turn, phosphorylates forkhead box protein O1 (FOXO1) resulting in its nuclear export and degradation and thereby prevents FOXO1 binding to the promoters of the key gluconeogenic enzymes, glucose-6-phosphatase (G6Pc) and phosphoenolpyruvate-carboxykinase (PEPCK) (Yan et al. 2019). In this way, E2 like insulin, represses the expression of these gluconeogenic enzymes (Fig. 4).

Conversely, P4's effect on gluconeogenesis is variable depending on carbohydrate availability (Lee et al. 2020). With carbohydrate provision, P4 enhances insulin secretion to suppress gluconeogenesis in healthy insulin-sensitive states, as seen in male mice with P4 treatment (Lee et al. 2020). Alternatively, in a glucose-depleted condition, as may occur during fasted exercise, P4 in hepatocytes increases glucose production by increasing the expression of the key gluconeogenic enzymes (G6Pc and PEPCK). An in vitro model using hepatocytes shows, P4 binds P4 receptor membrane complex 1 (PGRMC1) to activate adenylate cyclase and induce cyclic-AMP (cAMP) that in turn activates protein kinase A, which phosphorylates the transcription factor, cAMP-response element-binding protein (CREB) to promote expression of G6P and PEPCK (Lee et al. 2020) (Fig. 4). However, considering plasma glucose R_a during fasted exercise is reduced in the ML phase, we can assume that in eumenorrheic women, E2 suppression of hepatic glucose production predominates.

Overriding hepatic glucose production annuls the decrease in plasma glucose uptake during exercise in the ML phase

Only one previous study compared plasma glucose R_d when a carbohydrate supplement (60 g/h of glucose) was ingested during 2 h of exercise at 70% $\text{VO}_{2\text{max}}$ during the EF and ML phases (Campbell et al. 2001). Ingesting carbohydrate compared with placebo increased glucose R_a similarly in the EF and ML phase (Fig. 5b), sourced mainly from absorption of the ingested supplement with hepatic glucose production contributing minimally. With a surplus plasma glucose supply, the rate of plasma glucose uptake increased similarly in both EF and ML phase (Fig. 5a), implying that the combined increase of E2 and P4 in the ML phase failed to produce an expected E2-augmented exercise-stimulated glucose R_d .

Furthermore, a recent study overcame the limitation of E2 or P4 influence on gluconeogenesis and hepatic glucose

production by implementing an intravenous 10 mM hyperglycemic clamp during 90 min of exercise at 60% VO_2max in the follicular phase and luteal phase and the rate of glucose uptake was inferred from the rate of glucose infusion required to maintain the clamp (Hulton et al. 2021). The menstrual phase timing of the luteal phase trials did not coincide with the ML phase in all participants and possibly rather represents the early luteal or late luteal phase in some participants considering serum E2 concentration increased from a mean ~ 100 pmol/L in follicular phase to ~ 200 pmol/L in luteal phase and serum P4 concentration attained only a mean of 14 nmol/L in the luteal phase with P4 increasing above the 16 nmol/L threshold to confirm an ovulatory cycle in only 2 of the 7 participants. Notwithstanding, this study also found that the rate of plasma glucose uptake during exercise remains unchanged between follicular and luteal phases when plasma glucose availability is not limiting (attaining a high peak glucose Rd of 131 $\mu\text{mol/kg}\cdot\text{min}$ that equates to 1.4 g/min for a 60 kg women) (Hulton et al. 2021). These findings remain to be confirmed by further studies that encompass LF and ML phases with the provision of oral multiple-transportable carbohydrates during exercise at maximal tolerable doses (Trommelen et al. 2017).

It is well established that the availability of carbohydrate as an energy source is important for promoting best performances in moderate-high-intensity exercise (Burke and Hawley 2018). Accordingly, overnight-fasted high-intensity time trial performance, performed after 2 h of steady-state exercise, was compromised in ML phase compared with the EF phase, coincident to reduced plasma glucose Ra (or hepatic glucose production) and glucose Rd (Campbell et al. 2001). Interestingly, when overnight-fasted premenopausal women ingested a glucogenic substrate in the form of rapidly absorbed casein protein hydrolysate versus placebo during 80 min of moderate-hard exercise in a crossover design matched for menstrual phase, metabolism shifted in favor of greater carbohydrate oxidation that coincided with a small increase in distance covered and total work done during the final 20 min time trial (Oosthuysen et al. 2021). Thus, the restriction of E2 (and P4) on gluconeogenesis during the ML and possibly LF phases can have negative implications for exercise performance under certain conditions of prolonged exercise requiring high-intensity efforts without adequate carbohydrate supplementation. For this reason, the ability to store muscle and liver glycogen is also important.

Influence of the ovarian hormones on glycogen storage and use

Studies have found a greater ability to replete muscle glycogen stores after glycogen depleting exercise in the ML phase compared with MF phase (Nicklas et al. 1989; Hackney 1990) and when following a normal mixed diet

including ~ 5 g/kg carbohydrate (MF: 575 mmol/kg d.w. vs. ML: 728 mmol/kg d.w.) (McLay et al. 2007). However, this menstrual phase difference disappears when following a high carbohydrate-loading diet for 3 days (~ 8 g/kg carbohydrate) (McLay et al. 2007). Similarly, when ingesting a high carbohydrate dose (1.2 g/kg per hour) during the immediate 5 h recovery period after glycogen depleting exercise, a similar muscle glycogen repletion is achieved in EF, LF and ML phases (Matsuda et al. 2022b). E2 increases glycogen synthase activity (Beckett et al. 2002) possibly by acting via membrane-bound $\text{ER}\alpha$ to stimulate PI3K/Akt signaling to phosphorylate glycogen synthase kinase-3 α thereby releasing its inhibition on glycogen synthase activity (Yan et al. 2019) and by increasing glycogen synthase expression as seen in men after 8 days of oral E2 supplementation (Fu et al. 2009). However, it may be less relevant when following a high carbohydrate-rich diet and during acute recovery when ingesting a high-dose carbohydrate supplement that results in an already maximal insulin-stimulated signaling response.

Muscle glycogen use during moderate-intensity endurance exercise (60–70% VO_2max) based on analysis of muscle biopsies has been reported to be 24 and 25% less in the ML versus EF phase in two independent studies, respectively, coincident to greater fat oxidation (Hackney 1999; Devries et al. 2006) where glycogen use was inversely correlated with E2 concentration in the ML phase (Hackney 1999). Likewise, the rate of glycogen use at 60% VO_2max , estimated from the difference between the rate of total carbohydrate oxidation and tracer measured plasma glucose Rd, was 24 and 28% less in eumenorrheic women on controlled E2 supplements mimicking the LF phase also coincident to greater fat oxidation compared with ovarian hormone suppression and compared with combined E2 and P4 supplements, respectively (D'Eon et al. 2002). Interestingly, the combined E2 and P4 supplements caused a supraphysiological increase in P4 possibly reflecting a P4-dominant condition (D'Eon et al. 2002). Thus, the glycogen sparing effect of E2 may occur secondary to greater fat oxidation during moderate-intensity endurance exercise and can be countered by P4 and hence is dependent on the relative increase in E2 and P4 (Hackney et al. 2022). Conversely, high-intensity exercise necessitates carbohydrate-sourced energy production, and consequently, E2 suppression of hepatic gluconeogenesis might explain recent findings for greater muscle glycogen use, measured by ^{13}C -magnetic resonance spectroscopy, during overnight-fasted high-intensity intermittent exercise to exhaustion in the LF than EF phase (Matsuda et al. 2022a). However, the greater muscle glycogen use during exercise in the LF phase in this study was found to be significant despite no menstrual phase differences in the absolute starting or ending muscle glycogen concentration (Matsuda et al. 2022a).

Mechanisms behind menstrual cycle effects on fat metabolism

E2 signaling increases fat oxidation capacity

CAMPBELL and Febbraio (2001) reported that 15-days of E2 treatment in OVX rodents increased the maximal activity of enzymes regulating long chain fatty acid (LCFA) oxidation in skeletal muscle, namely carnitine palmitoyl transferase-1 (CPT-1) that governs LCFA transport across the mitochondrial membrane and β -hydroxyacyl-CoA

dehydrogenase (β -HAD), the third enzyme in beta oxidation. Coincident treatment with P4 blocked E2's effect on these enzymes, where P4 antagonism could be prevented by a supraphysiological dose of E2 (Campbell and Febbraio 2001).

Subsequently, a collection of studies largely from three different laboratories have uncovered the molecular signaling pathway whereby E2 promotes fat oxidation (Fig. 6). The cumulative findings from these studies that employed either an OVX-rodent model, in vitro cell incubation, men on E2 supplements or sex-differences have been reviewed in detail elsewhere (Tarnopolsky 2008; Oosthuysen and Bosch,




Fig. 6 Diagram showing E2 signaling to regulate fat availability away from adipose tissue and towards skeletal muscle lipid storage and oxidation. In skeletal muscle, depicted on the bottom right, E2 bound ER α activates gene expression of the nuclear hormone receptors, peroxisome proliferator-activated receptor- α (PPAR α and PPAR δ) directly or indirectly via phosphorylation of AMP-activated protein kinase (AMPK). PPARs enhance expression of plasma membrane (fatty acid transport protein, FATP and fatty acid translocase, CD36), cytosolic (fatty acid binding protein cytosolic, FABPc) and mitochondrial (carnitine palmitoyltransferase 1, CPT1) long chain fatty acid (LCFA) transporters, beta oxidation enzymes (medium-chain acyl-CoA dehydrogenase, MCAD; trifunctional protein alpha, TFP α ; β -hydroxy-acyl-CoA dehydrogenase, β -HAD) and kinases (pyruvate dehydrogenase kinase 4, PDK4) that direct metabolism away from carbohydrate and in favor of fat oxidation. E2 also increases lipid

availability in skeletal muscle first by increasing skeletal muscle localized lipoprotein lipase (LPL) to support plasma very low density lipoprotein triglyceride (VLDL-TG) hydrolysis for LCFA uptake, second by increasing the expression of the transcription factor, sterol regulatory element-binding protein 1c (SREBP1c) and downstream targets for skeletal muscle IMTG storage (mitochondrial glycerol-3-phosphate acyl transferase, mtGPAT; acetyl-CoA carboxylase, ACC; fatty acid synthase, FAS) and third E2 promotes a redistribution of LCFA away from adipose tissue and toward skeletal muscle, as depicted on the top left, by decreasing adipose-localized LPL activity, decreasing lipid storage by repressing SREBP1c expression and promoting adipose lipolysis by increasing β -adrenergic receptor (β -AR) sensitivity to epinephrine, decreasing α_2 -AR sensitivity and directly or indirectly increasing perilipin activity. P4 antagonizes E2 effects possibly by repressing ER expression

2012). Briefly, in skeletal muscle, E2 bound ER α activates gene transcription of the nuclear hormone receptors, peroxisome proliferator-activated receptor- α (PPAR α) and PPAR δ either directly by binding ERE in the gene promoter regions or indirectly via membrane-bound ER α to induce phosphorylation of AMPK (Campbell et al. 2003; D'Eon et al. 2005, 2008; Fu et al. 2009; Salehzadeh et al. 2011). PPAR α and PPAR β activate the expression of downstream targets (some of which may also be under direct genomic regulation of E2) to increase the protein content of plasma membrane (fatty acid transport protein (FATP), fatty acid translocase (CD36)), cytosolic (fatty acid binding protein (FABPc)) or mitochondrial (CPT-1) LCFA transporters and mitochondrial beta oxidation enzymes (medium-chain fatty acyl-CoA dehydrogenase (MCAD); β -HAD; beta oxidation enzyme complex, trifunctional protein alpha (TFP α)) and the glycolytic down-regulator, pyruvate dehydrogenase kinase 4 (PDK4) to direct substrate utilization away from carbohydrate and in favor of fat oxidation (Campbell et al. 2003; D'Eon et al. 2005; Fu et al. 2009; Maher et al. 2010a, 2010b). Recent evidence suggests that the Rab-GTPase activating proteins, TBC1D1/4 that regulate GLUT4 translocation are also involved in regulating LCFA transporter content and trafficking to the plasma membrane and thus LCFA oxidation (Benninghoff et al. 2020; Mikłosz et al. 2021). TBC1D1/4 maintain the Rab-GTPase proteins in an inactive GDP-bound state and this inhibition is released upon phosphorylation of TBC1D1/4 by its upstream effectors (Akt and AMPK), which then allows Rab-GTP binding and hydrolysis to facilitate transporter translocation. In this way, investigations are needed to establish whether E2 via membrane-bound ER α phosphorylation of PI3K/Akt and AMPK may possibly also contribute to LCFA transporter translocation and increased LCFA uptake in skeletal muscle.

In addition, E2 increases lipid availability specifically in skeletal muscle by firstly, increasing the expression of the transcription factor, sterol regulatory element-binding protein 1c (SREBP1c), which activates gene expression of downstream targets of skeletal muscle intramuscular triglyceride (IMTG) storage, namely, mitochondrial glycerol-3-phosphate acyl transferase (mtGPAT) for re-esterification and acetyl-CoA carboxylase (ACC) and fatty acid synthase (FAS) for lipogenesis (Fu et al. 2009). Secondly, E2 via increased PPAR δ increases expression and content of skeletal muscle localized lipoprotein lipase (LPL) to support hydrolysis of plasma triglyceride for skeletal muscle LCFA uptake (D'Eon et al. 2005). Thirdly, E2 promotes a redistribution of LCFA away from adipose tissue and toward skeletal muscle by reducing adipose tissue localized LPL activity to reduce triglyceride uptake in adipose (Ellis et al. 1994; Homma et al. 2000) and by reducing adipose-specific lipid storage via tissue-specific inhibition of SREBP1c expression in adipose and liver (D'Eon et al. 2005) and by promoting

adipose triglyceride lipolysis by enhancing beta-adrenergic sensitivity to epinephrine (Benoit et al. 1982), reducing antilipolytic alpha₂-adrenergic responsiveness (Schmidt et al. 2014) and increasing perilipin protein content (albeit without identifying subtype specificity) (D'Eon et al. 2005). The role of different perilipin protein isoforms in the regulation of triglyceride storage and lipolysis in various tissues and sex-differences in perilipin content have been reviewed (MacPherson and Peters 2015). The actions of perilipins are important for preventing the accumulation of lipid oxidation intermediates and accordingly, perilipin actions differ from the basal state to the hormone-stimulated lipolytic state (MacPherson and Peters 2015).

P4 when administered in combination with E2 in OVX rodents modulates E2 effect to increase PPAR α and PDK4 (Campbell et al. 2003). Without further studies investigating the direct mechanistic effect of P4 on lipid transport or metabolic pathways, currently P4 modulation of E2 actions can possibly only be attributed to P4's repression of ER expression and protein content (Jayaraman and Pike 2009; Ekenros et al. 2017).

Menstrual cycle effect on whole-body fat oxidation

Taking the prior points into account, it is not surprising that sex-differences in the rate of whole-body fat oxidation are frequently reported particularly during fasted exercise (for Reviews see, Devries, 2016; Boisseau and Isacco 2022; Cano et al. 2022). Surprisingly, however, menstrual phase differences are less consistent, with only some reporting a greater rate of fat oxidation in the LF (Hackney et al. 1991) and ML phases compared with EF phase during fasted exercise (for example, Hackney et al. 1994, 1999; Campbell et al. 2001; Zderic et al. 2001; Willett et al. 2021). However, E2 promotion of fat oxidation during exercise is confirmed with men on E2 supplements (Hamadeh et al. 2005) and, as already mentioned, in premenopausal women on controlled E2 treatment versus ovarian suppression or combined E2 and supraphysiological P4 treatment (D'Eon et al. 2002). The change in the rate of fat oxidation during exercise from EF to ML phase has been found to correlate positively with the change in the E2/P4 ratio (Hackney et al. 2022). Moreover, in one study, a greater rate of fat and lower carbohydrate oxidation in the ML versus MF phase was significant at low (35% VO₂max) and moderate (60% VO₂max) exercise intensities but not significant at a higher intensity (75% VO₂max) (Hackney et al. 1994) possibly where a higher rate of carbohydrate utilization becomes obligatory. Therefore, it seems that a notable shift in whole-body substrate utilization with menstrual phase is only evident: (i) during fasted exercise; or (ii) mainly when exercise duration places a sufficiently high demand on endogenous energy stores; and (iii) possibly only at or below the exercise intensity marking the metabolic

threshold after which carbohydrate utilization predominates; and (iv) as such is dependent on a high relative concentration of E2 to P4 in the ML phase.

For example, the exercise intensity eliciting maximal fat oxidation (FATmax) and the peak fat oxidation rate is not different between MF, LF and ML phases when measured in a standard 3-min step incremental test to exhaustion (Frandsen et al. 2020). This is not surprising because the 3-min steady-state steps are too short to pressurize energy metabolism and fuel reserves sufficiently to reveal potential menstrual phase differences that may possibly become evident should exercise duration at each workload be substantially longer. Conversely, with a similar test protocol, sex-differences in peak fat oxidation rate expressed relative to fat-free mass were evident, being greater in women than men (Chrzanowski-Smith et al. 2021) with this sex-difference evident only in women < 45 years but not > 45 years (Frandsen et al. 2021) that likely differentiates premenopausal versus peri- or post-menopausal life stages and thus cyclical E2 availability versus chronic E2-deficiency as the underlier.

Fat oxidation source: Intramuscular triglyceride or plasma long chain fatty acid-derived

Sex-differences in fat oxidation may be partly owing to greater IMTG content in women (because of a greater number of lipid droplets and not greater lipid droplet size) and greater localization of lipid droplets near mitochondria in women (Devries et al. 2007; Beaudry and Devries 2019), together with reports for superior mitochondrial functional capacity in women compared with men matched for training status (Cardinale et al. 2018; Montero et al. 2018), that are E2-attributable perturbations according to signaling mechanisms already discussed. Albeit early variations in methods of IMTG quantification (i.e. biochemical extraction, immunofluorescence microscopy or proton magnetic resonance spectroscopy) have resulted in an unclear consensus for sex-differences in IMTG use during exercise (Steffensen et al. 2002; Roepstorff et al. 2002, 2006; White et al. 2003; Devries et al. 2007). Furthermore, whether the cyclical fall and rise in E2 with the respective menstrual phases notably alters IMTG content, sub-cellular location and use during exercise in eumenorrheic women has not been evaluated and certainly warrants investigation. Improvements in histological staining techniques has advanced our understanding of IMTG myofibrillar-localization and association with perilipin proteins, IMTG utilization during exercise and resynthesis in recovery (Jevons et al. 2020; Strauss et al. 2020) but are yet to be applied in menstrual phase comparative studies.

Most interestingly, IMTG use during exercise is not reduced by ingestion of carbohydrate supplements in men (Fell et al. 2021) with confirmation needed for the same response in women. However, IMTG use is dependent on

exercise duration and intensity, and varies inversely with plasma fatty acid availability and use (Lundsgaard et al. 2018). At high exercise intensities, fat oxidation is limited at the point of LCFA entry into the mitochondria by free carnitine availability where the excess acetyl-CoA produced by the glycolytic enzyme, pyruvate dehydrogenase is buffered by carnitine acetyltransferase resulting in the entrapment of free carnitine as acetylcarnitine to support the maintenance of a high-demand glycolytic rate (Lundsgaard et al. 2018). This limits the activity of CPT1 where free carnitine is required for the conversion of long chain fatty acyl-CoA to fatty-acyl-carnitine for mitochondrial uptake. Thus, unlike the assumption of plasma glucose uptake, not all plasma LCFA taken up into skeletal muscle during exercise is necessarily oxidized. Excess LCFA uptake is shuttled into re-esterification and IMTG storage. During low-moderate intensity exercise, with lower glycolytic flux, IMTG roughly accounts for 30% of fat oxidation (Lundsgaard et al. 2018) with plasma-derived LCFA largely from adipose tissue lipolysis accounting for 60% of fat oxidation, in men (Lundsgaard et al. 2018) and women (Jacobs et al. 2005).

Only two menstrual phase comparative studies have measured the rate of lipolysis and the rate of plasma LCFA uptake during exercise by intravenous infusion of the stable-isotopic tracers, [1,1,2,3,3-²H]glycerol and [1-¹³C]palmitate, respectively. The first study reports no change between EF and ML phases in glycerol kinetics (Casazza et al. 2004) and no change in the rate of plasma LCFA appearance, uptake, or oxidation (Jacobs et al. 2005) at either 45% $\text{VO}_{2\text{max}}$ or 65% $\text{VO}_{2\text{max}}$. However, the small sample size ($n=5$) and high variability in especially plasma LCFA kinetics and oxidation in the ML phase encourages further follow up studies. Furthermore, in this former study the women participated fed (3 h postprandial) and exercise lasted only 60 min, which might also partly explain the lack of differences. The second study included a larger sample of women ($n=11$) who were overnight-fasted and exercised for 90 min at 50% $\text{VO}_{2\text{max}}$, but some menstrual phase timing errors resulted in a few missing data points and data being evaluated using a less sensitive unpaired groups design (Horton et al. 2006). Nonetheless, this latter study also reports no differences between EF, MF and ML phases in the rate of plasma glycerol or LCFA appearance and uptake during exercise but was without measurement of plasma LCFA oxidation. The findings from these two studies have been combined in a summative meta-analysis that may encourage further studies to investigate menstrual cycle effects on particularly plasma LCFA oxidation and flux during exercise (Fig. 7a). This meta-analysis, however, does present a clear outcome for no effect on plasma glycerol Ra and Rd, reflecting an unchanged whole-body lipolytic rate during exercise between EF and ML phases in these studies (Fig. 7b). In addition, E2 treatment in amenorrheic women for 6 days (Ruby et al. 1997)



a**b**

Fig. 7 Forest plot showing the mean difference (95% confidence interval CI) of menstrual phase on rate of plasma long chain fatty acid (LCFA) appearance (Ra), uptake (Rd), and oxidation (Rox) during exercise (**a**) and rate plasma glycerol appearance (Ra), and uptake (Rd) during exercise as an indication of whole-body lipolytic flux (**b**);

showing the heterogeneity indices, z -scores, p -values, and standardized mean difference as effect size (ES) for each outcome. Effect size scores are interpreted as: 0.0–0.2 is trivial; 0.2–0.6 is small; 0.6–1.2 is moderate; > 1.2 is large effects, respectively

or men for 8 days (Carter et al. 2001) failed to alter plasma glycerol kinetics during exercise further supporting the likelihood of no notable ovarian hormone effect on whole-body lipolysis between menstrual phases or with short-term

E2 treatment. Of note, the above studies also all failed to find a difference in total whole-body fat oxidation during exercise between menstrual phases or ovarian conditions. It could be that the serum E2 concentration recorded in the

ML phase in these studies (E2: 311 pmol/L, P4: 34.7 nmol/L and E2: 393 pmol/L, P4 36.6 nmol/L, respectively) was not high enough for ovarian or menstrual phase effects on fat metabolism to become apparent; compared with studies that have reported a shift in substrate utilization towards greater fat oxidation in the ML phase (for example, E2: 519 pmol/L, P4 47.9 nmol/L (Willett et al. 2021) where the change in fat oxidation from EF phase correlated with the change in the E/P ratio (Hackney et al. 2022). Others have suggested that the latter studies where E2 treatment was administered (Ruby et al. 1997; Carter et al. 2001) were likely underpowered for the given variability considering that a clear shift towards greater whole-body fat oxidation during exercise was noted in a larger sample of 12 males receiving similar dose oral E2 supplements (Hamadeh et al. 2005). It would be insightful if, in future studies, plasma glycerol and LCFA kinetics and plasma LCFA oxidation were evaluated during exercise where a coincident shift in whole-body metabolism towards greater fat oxidation in the LF or ML compared with EF phase is evident.

Conversely, OC use for 4 months increased plasma cortisol concentration and whole-body lipolysis during exercise, as indicated by stable-tracer measured plasma glycerol Ra that increased by more than 20%, compared to before-OC use in eumenorrheic women (Casazza et al. 2004). However, the greater rate of lipolysis was coupled with a greater rate of FFA re-esterification and less plasma FFA being oxidized during exercise, measured by plasma [1-¹³C]palmitate tracer kinetics, after 4 months of OC use compared with before-OC use and hence indicates a futile cycle of substrate mobilization (Jacobs et al. 2005). Yet again, this demonstrates that OC use influences exercise metabolism differentially compared with naturally cycling endogenous ovarian hormones experienced by eumenorrheic women, where differences also extend to other physiological parameters, such as basal inflammatory status (Cauci et al. 2017, 2021; Larsen et al. 2020; Quinn et al. 2021) and bone turnover (Allaway et al. 2020).

Notably, the technical difficulties of administering LCFA tracers by intravenous infusion could be a deterring factor to explain the shortage of such menstrual phase comparative studies. The hydrophobic nature of LCFA tracers necessitates constitution with 5% human serum albumin as a carrier protein that could pose ethical risk limitations. Furthermore, use of LCFA carbon-13 tracers to measure plasma LCFA oxidation from expired [¹³C]-carbon dioxide enrichment requires correction for possible retention of the carbon-13 label in the exchange or transamination reactions that occur with tricarboxylic acid (TCA) cycle intermediates (Sidossis et al. 1995a, 1995b). To this end, the acetate correction factor was proposed that must be derived concurrently with plasma LCFA-tracer measured oxidation or in an additional identical exercise trial with the constant infusion of

carbon-13 labeled sodium acetate (Sidossis et al. 1995b). When the acetate correction factor was derived during 90 min of cycling at 60% $\dot{V}O_{2\max}$ in a small sample ($n=5$) of eumenorrheic women, 2–3-h post prandial, during the EF, LF, and ML phases it was found to be modestly lower in the ML (0.57) than EF (0.59) phase despite no difference in whole-body substrate utilization (Oosthuyse et al. 2003). While application of the correction factor would increase absolute plasma LCFA oxidation rate notably, the small difference between menstrual phases would only equate to a 6% error in the measured change in plasma LCFA oxidation between menstrual phases. Nevertheless, the greater carbon-label fixation in TCA cycle exchange reactions in the ML phase does suggest an increased flux in transamination reactions in the ML phase, which may be suggestive of greater protein catabolism during exercise.

Mechanisms behind menstrual cycle effects on protein metabolism

Increased protein catabolism in the ML phase

LAMONT et al. (1987) were first to report greater total urea nitrogen excretion (a sum of urine and sweat urea nitrogen) in the ML versus EF phase in recreationally active women measured over 3 days that included a 60 min cycling exercise on day 1, indicating greater protein catabolism in the ML phase. According to these findings, the protein requirement per day in the EF phase were notably less than in the ML phase (1.05 g/kg vs. 1.36 g/kg per day, respectively). In agreement, amino acid isotopic-tracer studies in eumenorrheic women have consistently reported a greater rate of leucine or phenylalanine oxidation during the ML versus EF phase when fasted (Lariviere et al. 1994; Toth et al. 2006) and fed (Kriengsinyos et al. 2004). These studies, however, were measured at rest and did not include exercise. The summative findings from these studies are presented in a meta-analysis showing a moderate effect of menstrual cycle phase on plasma amino acid oxidation rate (Fig. 8a) but no clear effect on plasma amino acid flux (where at rest $Ra=Rd$) (Fig. 8b). Moreover, lysine dietary requirements based on the indicator amino acid oxidation method was found to be higher in the ML than EF phase (37.7 mg/kg versus 35.0 mg/kg per day, respectively) (Kriengsinyos et al. 2004). In this study carbon-labeled lysine was ingested in increasing doses on separate occasions over several consecutive menstrual cycles in each participant and the rate of lysine oxidation measured from the carbon-label enrichment in expired carbon dioxide that increases disproportionately on reaching an ingestion dose in excess of the requirement to support protein synthesis. Considering that lysine oxidation was maintained at a higher rate at all ingestion doses in ML versus EF





a**Plasma Leucine Rox at rest ($\mu\text{mol/kg}\cdot\text{h}$)****Plasma Phenylalanine Rox at rest ($\mu\text{mol/kg}\cdot\text{h}$)****b****Plasma Leucine flux at rest ($\text{Ra} = \text{Rd}$) ($\mu\text{mol/kg}\cdot\text{h}$)****Plasma Phenylalanine flux at rest ($\text{Ra} = \text{Rd}$) ($\mu\text{mol/kg}\cdot\text{h}$)**

Fig. 8 Forest plot showing the mean difference (95% confidence interval CI) of menstrual phase on rate of plasma leucine oxidation (Rox) and plasma phenylalanine oxidation (Rox) at rest (a) and plasma leucine flux and plasma phenylalanine flux at rest (b) where rate of appearance (Ra) equals the rate of uptake (Rd); showing the hetero-

genicity indices, z-scores, p-values, and standardized mean difference as effect size (ES) for each outcome. Effect size scores are interpreted as: 0.0–0.2 is trivial; 0.2–0.6 is small; 0.6–1.2 is moderate; > 1.2 is large effects, respectively

phase, suggests the higher lysine requirement in ML phase is due to higher amino acid catabolism (Kriengsinyos et al. 2004).

Greater protein catabolism during the ML phase appears to be a P4-dominant effect because when men received oral E2 supplements (2 mg/day) for 8 days to mimic plasma E2

concentration during the LF phase (876 pmol/L), leucine oxidation was notably decreased at rest (PL: 28 ± 2 vs. E2: 23 ± 2 $\mu\text{mol/kg}\cdot\text{h}$) and during 90 min of exercise at 65% VO_2max (PL: 62 ± 5 vs. E2: 53 ± 4 $\mu\text{mol/kg}\cdot\text{h}$) (Hamadeh et al. 2005). This resulted in a less negative protein balance after E2 supplements compared with placebo. However,

leucine flux and the rate of protein synthesis were not altered by E2 supplements. As such, menstrual phase comparative studies during exercise are required to verify the same decrease in protein oxidation in the LF phase and the proven increase in protein oxidation in the ML phase (tested to-date only at rest, or without isolating the exercise period per se) compared with the EF phase.

The mechanism whereby P4 upregulates and E2 downregulates protein catabolism remains undefined. E2 does not appear to alter the expression of certain enzymes involved in amino acid catabolism. For example, in the same men just noted receiving 8 days of oral E2 supplementation to mimic the LF phase (Hamadeh et al. 2005), E2 supplementation did not alter skeletal muscle expression of the amino acid transamination enzyme, aspartate aminotransferase, or the rate-limiting enzyme for branched-chain amino acid oxidation, branched-chain 2-oxo-acid dehydrogenase (BCOAD), or its negative regulator, BCOAD kinase (Fu et al. 2009). However, skeletal muscle BCOAD kinase mRNA content was 1.2–1.3-fold greater in women than men at rest (Fu et al. 2009), which in its role as a negative regulator of BCOAD could reduce branch-chain amino acid oxidation and thereby partly explain the frequently reported lower rate of protein catabolism in women than men (Tarnopolsky 2008). In that same study, skeletal muscle mRNA content of BCOAD kinase in women was not different between the follicular (E2, 125 pmol/L; P4, 2.0 nmol/L) and luteal (E2, 203 pmol/L; P4, 6.0 nmol/L) phase, but interpretation of this finding is limited by the modest differences in ovarian hormone concentrations between phases suggesting that the measurement in the luteal phase may have corresponded with the late luteal phase (Fu et al. 2009). Future studies should investigate the effect of P4 on these gene targets. Furthermore, whether the ovarian hormones regulate skeletal muscle protein content or activity of transamination and branched-chain amino acid oxidation enzymes remains to be determined. Alternatively, some suggest P4 upregulation and E2 downregulation of thyroid function and availability of free triiodothyronine (T3) (Ben-Rafael et al. 1987; Torre et al. 2020) is causative of the greater protein catabolism in the luteal phase (Lariviere et al. 1994). Resting plasma free T3 concentration was found to be higher coincident to a greater leucine oxidation rate in the ML versus EF phase in one study (Lariviere et al. 1994). P4 binds PR within the thyroid gland to promote T3/thyroxine (T4) secretion and reduce its binding to the plasma protein, thyroid-binding globulin to maximize free T3 availability (Torre et al. 2020; Kaminski et al. 2021) thereby accounting for the higher resting metabolic rate (Benton et al. 2020) and thus possibly protein catabolism reported in the luteal phase compared with follicular phase.

The estimated average daily requirement of protein for female endurance athletes derived from 3-day nitrogen

balance measurements in two studies conducted in the MF phase in female cyclists training 13 and 11 h/week is between 1.28 and 1.63 g/kg per day with high interindividual variability (95% CI: 0.7–1.85 and 1.1–3.8 g/kg per day, respectively) (Rowlands and Wadsworth, 2011; Houltham and Rowlands et al. 2014). Based on these findings, recommended daily intake (RDI) of protein for female endurance athletes has been suggested as 1.6–2.0 g/kg per day, to allow for 12% interindividual variability (Mercer et al. 2020) but does not include menstrual phase specificity, which warrants further validation.

No menstrual phase effect on muscle protein synthesis

Only one study in eumenorrheic women has quantified the rate of muscle protein synthesis between menstrual phases. In this study, unilateral leg exercise was performed with 1-legged kicks and 1-leg acting as a resting control in the EF and ML phases (Miller et al. 2006). Twenty-four hours later, fractional protein synthetic rate (FSR) was measured from a 4 h constant infusion of [1 – ¹³C]leucine followed by a muscle biopsy to measure the percentage uptake of the label into muscle myofibrillar protein in each leg. While FSR was higher in the exercised leg 24-h post exercise compared with the resting leg, there was no menstrual phase differences in resting (EF: $0.053 \pm 0.009\%/h$ vs. ML: $0.055 \pm 0.013\%/h$) or 24-h post exercise (EF: $0.131 \pm 0.018\%/h$ vs. ML: $0.134 \pm 0.018\%/h$) FSR (Miller et al. 2006).

Nevertheless E2-deficiency, as occurs in menopause, is known to promote muscle atrophy by increased autophagic signaling through FOXO3 that is suppressed by E2 replacement, where E2 replacement increases muscle mass and strength further when combined with exercise training possibly by regulation of satellite cell activity (Hansen 2018). It may be that while menstrual phase does not influence resting or recovery muscle protein synthesis in eumenorrheic women, E2 could enhance exercise-stimulated anabolism by the activation and proliferation of muscle satellite cells following exercise in a menstrual phase-specific manner, as presented below.

Menstrual phase influences muscle regeneration and strengthening

ENNS and Tiidus (2008) have clearly demonstrated these effects in OVX rodents receiving E2 replacement by showing a greater increase in paired box homeotic gene 7 (Pax7)-positive cells, a marker of satellite cell number, a greater increase in myogenic differentiation factor D (Myo-D)-positive fibers indicating satellite cell activation, and a greater increase in DNA uptake of bromo-deoxyuridine suggesting

satellite cell proliferation after downhill running with E2 treatment versus placebo. This group went on to show that these E2 effects on satellite cells were ER α dependent (Thomas et al. 2010). E2-ER increases Myo-D expression indirectly by binding the transcription factor, activator protein 1 (AP-1) and thereby prevents AP-1 repressive action on Myo-D expression (Pedraza-Alva et al. 2009).

Interestingly, E2 influence over the satellite cell response to EIMD is more noticeable during the MF compared with ML phase as shown by Haines et al. (2018) after unilateral eccentric knee extensor exercise. These authors found ER α muscle content and the increase in ER-DNA binding after exercise to be greater in the MF than ML phase (Haines et al. 2018). Accordingly, the increase in expression of Myo-D, reflecting satellite cell activation, appeared greater and cyclin D1 reflecting mitotic cycling was significantly greater in the MF than ML phase after eccentric exercise (Haines et al. 2018). In this study, serum E2 concentration, although significantly higher in the ML phase (410 pmol/L) was also substantially elevated in the MF phase (325 pmol/L) and consequently muscle E2 concentration was not different between MF and ML phases. Thus, the noted difference in ER and satellite cell activity occurred independent of a difference in muscle E2 concentration and suggests that P4 in the ML phase suppressed E2 promotion of satellite cell differentiation and muscle regeneration after EIMD, possibly by downregulation of ER expression and protein content as already discussed (Jayaraman and Pike 2009; Ekenros et al. 2017). Although it is also true that EIMD is consistently less in ML than EF phase as detailed earlier and as such decreased plasma membrane disruption and EIMD in ML may also contribute to a reduced need for satellite cell activation and muscle repair in ML than EF phase.

Consequently, a study that performed unilateral leg resistance training with 1-leg trained only in the follicular phase and the other leg trained only in the luteal phase for 3 menstrual cycles found greater muscle strength gain and muscle diameter in the follicular phase trained-leg than the luteal phase trained-leg (Sung et al. 2014). This occurred coincident to an increase in the nuclei-to-fiber ratio in muscle from the follicular phase leg only, that is said to suggest notable satellite cell recruitment in the muscle trained in the follicular phase (Sung et al. 2014). Similar findings support greater muscle strength gain from high frequency (4–5 d/week) leg resistance training when performed specifically during the follicular phase rather than luteal phase (Reis et al. 1995; Wikström-Frisén et al. 2017), with only a single report of no difference in outcomes after follicular phase-based versus luteal phase-based arm resistance training at a lower frequency (1–3 d/week) (Sakamaki-Sunaga et al. 2016).

Therefore, muscle regeneration and strengthening are mostly greater during the follicular phase when E2 is present alone (even in low-moderate concentrations) without P4.

Conclusion and perspectives

The molecular mechanisms of E2 and P4 are diverse and underpin menstrual phase differences in exercise metabolism (summarized in Fig. 9). At this stage research has uncovered firstly, that E2 phenolic structure stabilizes membranes and quenches free radicals, while genomic effects of E2 and P4 increase antioxidant defense causing the ML phase to be the best time for unaccustomed endurance exercise or low intensity resistance loads owing to a modulated EIMD response. Secondly, E2 promotion of skeletal muscle glucose uptake is not realized in the LF or ML phase owing to E2-dominant repression of hepatic gluconeogenic enzyme expression and plasma glucose availability that can compromise high-intensity exercise efforts in carbohydrate-depleted states. For this reason, carbohydrate supplements during exercise are imperative in these menstrual phases and future studies should explore menstrual phase effects when ingesting multiple-transportable carbohydrates at recommended doses during exercise, as well as variability in efficacy of ingesting exogenous glucogenic sources, such as rapidly absorbed peptides. Thirdly, E2 signaling to increase glycogen synthase activity and thereby promote increased glycogen storage in the ML (or possibly LF) phase at normal daily carbohydrate intakes should be considered for recovery or when planning pre-event nutrition that will necessitate higher carbohydrate-loading intakes to equalize glycogen stores in the EF (or MF) phase. Fourthly, E2 signaling favors skeletal muscle fat uptake, storage, and utilization but exercise intensity and carbohydrate availability are the dominant regulators of substrate partitioning, such that a menstrual phase increase in whole-body fat oxidation is only realized in the ML (or LF) phase under carbohydrate-restricted exercise and with a sufficiently high E2:P4 ratio. Currently, insufficient studies have evaluated for menstrual phase differences in the lipid-derived source, namely plasma LCFA or IMTG, during exercise. Fifthly, protein catabolism has consistently been shown to be greater in the luteal phase at rest and the possibility for RDI of protein to be menstrual phase specific should be explored. In particular, assessment of menstrual phase differences in protein catabolism during exercise should be investigated and the possibility of recommending ingestion of rapidly absorbed peptides during exercise or more rigorous protein recovery supplements in the ML should be considered. Moreover, the molecular signaling mechanism explaining the P4-dominant effect promoting protein catabolism should be investigated. Lastly, athletes including adjunct strength training should periodize these training sessions in the follicular phase to benefit from greater muscle strength gains owing to E2 activation of satellite




Fig. 9 Summary of E2 and P4 cellular signaling to direct metabolic outcomes for maintenance of energy homeostasis during exercise and in recovery in a menstrual phase-specific manner, where: (1) the molecular structure of E2 stabilizes plasma membranes and quenches free radicals, and E2 and P4 promotes the expression of antioxidant enzymes that reduces exercise-induced muscle damage (EIMD) in the LF and ML phases; (2) E2 promotion of plasma glucose uptake during exercise via GLUT4 expression and kinases is nullified by E2-dominant suppression of gluconeogenic gene expression resulting in reduced plasma glucose uptake during exercise in the LF and ML phases; (3) E2 promotes glycogen synthase (GS) expression and activity by downregulating glycogen synthase kinase-3α (GSK3α), a negative regulator of glycogen synthase, and thereby explains greater glycogen storage in the LF and ML phases, although a similar outcome can be achieved with a high carbohydrate (CHO) diet and associated insulin signaling; (4) E2 signaling maximizes fat oxidation capacity and long chain fatty acid (LCFA) availability in skeletal muscle by preferential lipid storage in muscle and lipolysis in adipose resulting in greater fat oxidation in the LF and ML phases, albeit pending low-moderate exercise intensities, restricted nutrient availability and a high E2:P4 ratio; (5) P4 promotes and E2 antagonizes

protein oxidation by indeterminate mechanisms, resulting in greater protein catabolism in the luteal phase; (6) E2 targets gene expression to promote satellite cell recruitment, activation and proliferation following exercise that is antagonized by P4 in the ML phase resulting in frequent evidence for greater gains in muscle strength following follicular phase-based training; P4 antagonizes many E2 effects by downregulating the expression of E2 receptors, where inhibiting effects are denoted by a flat-ended line (⊥), while stimulating effects are denoted by an arrow line (→); AMPK AMP-activated protein kinase, AP-1 activator protein 1, E2 17-β-estradiol, EF early follicular phase, ER estrogen receptor, ERE estrogen response element, FOXO1 forkhead box protein O1, FP follicular phase, G6Pc glucose-6-phosphatase, GLUT4 glucose transporter 4, IMTG intramuscular triglyceride, LF late follicular phase, LP luteal phase, MyoD myogenic differentiation factor D, ML mid-luteal phase, P4 progesterone, PEPCK phosphoenolpyruvate-carboxykinase, PI3K phosphoinositide-3-kinase, PPAR peroxisome proliferator-activated receptor, PR progesterone receptor, PRE progesterone response element, Ra rate of appearance, Rd rate of disappearance, ROS reactive oxygen species, SREBP1c sterol regulatory element-binding protein 1c

cells that is suppressed by P4 in the luteal phase. Following the evidence presented herein, this review serves to guide researchers in conceptualizing future studies where research is lacking, and to test progressive interventions for female athletes to circumvent or gain advantage from the menstrual phase-specific responses.

Declarations

Author contribution TO, JAS and ACH, conceived the review design; TO drafted the manuscript; TO, JAS and ACH edited, revised, and approved the final version of the manuscript.

Funding Preparation of this manuscript required no financial support.

Conflict of interest The authors have no conflict of interests, financial or otherwise to declare.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

References

- Allaway HC, Misra M, Southmayd EA, Stone MS, Weaver CM, Petkus DL, De Souza MJ (2020) Are the effects of oral and vaginal contraceptives on bone formation in young women mediated via the growth hormone-IGF-I axis? *Front Endocrinol (Lausanne)* 11:334. <https://doi.org/10.3389/fendo.2020.00334>
- Baltgalvis KA, Greising SM, Warren GL, Lowe DA (2010) Estrogen regulates estrogen receptors and antioxidant gene expression in mouse skeletal muscle. *PLoS ONE* 5:e10164. <https://doi.org/10.1371/journal.pone.0010164>
- Barakat R, Oakley O, Kim H, Jin J, Ko CJ (2016) Extra-gonadal sites of estrogen biosynthesis and function. *BMB Rep* 49:488–496. <https://doi.org/10.5483/bmbrep.2016.49.9.141>
- Barreto-Andrade JN, de Fátima LA, Campello RS, Guedes JA, de Freitas HS, Machado MM (2018) Estrogen receptor 1 (ESR1) enhances Slc2a4/GLUT4 expression by a SP1 cooperative mechanism. *Int J Med Sci* 15:1320–1328. <https://doi.org/10.7150/ijms.26774>
- Barros RP, Machado UF, Warner M, Gustafsson JA (2006) Muscle GLUT4 regulation by estrogen receptors ERbeta and ERalpha. *Proc Natl Acad Sci U S A* 103:1605–1608. <https://doi.org/10.1073/pnas.0510391103>
- Beaudry KM, Devries MC (2019) Sex-based differences in hepatic and skeletal muscle triglyceride storage and metabolism. *Appl Physiol Nutr Metab* 44:805–813. <https://doi.org/10.1139/apnm-2018-0635>
- Beckett T, Tchernof A, Toth MJ (2002) Effect of ovariectomy and estradiol replacement on skeletal muscle enzyme activity in female rats. *Metabolism* 51:1397–1401. <https://doi.org/10.1053/meta.2002.35592>
- Benninghoff T, Espelage L, Eickelschulte S, Zeinert I, Sinowenka I, Müller F, Schöndeling C, Batchelor H, Cames S, Zhou Z, Kotzka J, Chadt A, Al-Hasani H (2020) The RabGAPs TBC1D1 and TBC1D4 control uptake of long-chain fatty acids into skeletal muscle via fatty acid transporter SLC27A4/FATP4. *Diabetes* 69:2281–2293. <https://doi.org/10.2337/db20-0180>
- Benoit V, Valette A, Mercier L, Meignen JM, Boyer J (1982) Potentiation of epinephrine-induced lipolysis in fat cells from estrogen-treated rats. *Biochem Biophys Res Commun* 109:1186–1191. [https://doi.org/10.1016/0006-291x\(82\)91902-7](https://doi.org/10.1016/0006-291x(82)91902-7)
- Ben-Rafael Z, Struass JF 3rd, Arendash-Durand B, Mastroianni L Jr, Flickinger GL (1987) Changes in thyroid function tests and sex hormone binding globulin associated with treatment by gonadotropin. *Fertil Steril* 48:318–320. [https://doi.org/10.1016/s0015-0282\(16\)59363-7](https://doi.org/10.1016/s0015-0282(16)59363-7)
- Benton MJ, Hutchins AM, Dawes JJ (2020) Effect of menstrual cycle on resting metabolism: a systematic review and meta-analysis. *PLoS ONE* 15:e0236025. <https://doi.org/10.1371/journal.pone.0236025>
- Boisseau N, Isacco L (2022) Substrate metabolism during exercise: sexual dimorphism and women's specificities. *Eur J Sport Sci* 22:672–683. <https://doi.org/10.1080/17461391.2021.1943713>
- Burke LM, Hawley JA (2018) Swifter, higher, stronger: what's on the menu? *Science* 362:781–787. <https://doi.org/10.1126/science.aau2093>
- Campbell SE, Angus DJ, Febbraio MA (2001) Glucose kinetics and exercise performance during phases of the menstrual cycle: effect of glucose ingestion. *Am J Physiol Endocrinol Metab* 281:E817–E825. <https://doi.org/10.1152/ajpendo.2001.281.4.E817>
- Campbell SE, Febbraio MA (2002) Effect of the ovarian hormones on GLUT4 expression and contraction-stimulated glucose uptake. *Am J Physiol Endocrinol Metab* 282:E1139–E1146. <https://doi.org/10.1152/ajpendo.00184.2001>
- Campbell SE, Febbraio MA (2001) Effect of ovarian hormones on mitochondrial enzyme activity in the fat oxidation pathway of skeletal muscle. *Am J Physiol Endocrinol Metab* 281:E803–E808. <https://doi.org/10.1152/ajpendo.2001.281.4.E803>
- Campbell SE, Mehan KA, Tunstall RJ, Febbraio MA, Cameron-Smith D (2003) 17β-Estradiol upregulates the expression of peroxisome proliferator-activated receptor α and lipid oxidative genes in skeletal muscle. *J Mol Endocrinol* 31:37–45. <https://doi.org/10.1677/jme.0.0310037>
- Campello RS, Alves-Wagner AB, Lucas TF, Mori RC, Furuya DT, Porto CS, Machado UF (2012) Estrogen receptor 1 agonist PPT stimulates Slc2a4 gene expression and improves insulin-induced glucose uptake in adipocytes. *Curr Top Med Chem* 12:2059–2069. <https://doi.org/10.2174/156802612804910197>
- Campello RS, Fátima LA, Barreto-Andrade JN, Lucas TF, Mori RC, Porto CS, Machado UF (2017) Estradiol-induced regulation of GLUT4 in 3T3-L1 cells: involvement of ESR1 and AKT activation. *J Mol Endocrinol* 59:257–268. <https://doi.org/10.1530/JME-17-0041>
- Cano A, Ventura L, Martinez G, Cugusi L, Caria M, Deriu F, Manca A (2022) Analysis of sex-based differences in energy substrate utilization during moderate-intensity aerobic exercise. *Eur J Appl Physiol* 122:29–70. <https://doi.org/10.1007/s00421-021-04802-5>
- Cardinale DA, Larsen FJ, Schiffer TA, Morales-Alamo D, Ekblom B, Calbet JAL, Holmberg HC, Boushel R (2018) Superior intrinsic mitochondrial respiration in women than in men. *Front Physiol* 9:1133. <https://doi.org/10.3389/fphys.2018.01133>
- Carter S, McKenzie S, Mourtzakis M, Mahoney DJ, Tarnopolsky MA (2001) Short-term 17beta-estradiol decreases glucose R(a) but not whole body metabolism during endurance exercise. *J Appl Physiol* 90:139–146. <https://doi.org/10.1152/jappl.2001.90.1.139>
- Casazza GA, Jacobs KA, Suh SH, Miller BF, Horning MA, Brooks GA (2004) Menstrual cycle phase and oral contraceptive effects on triglyceride mobilization during exercise. *J Appl Physiol* 97:302–309. <https://doi.org/10.1152/japplphysiol.00050.2004>
- Cauci S, Francescato MP, Curcio F (2017) Combined oral contraceptives increase high-sensitivity c-reactive protein but not

- haptoglobin in female athletes. *Sports Med* 47:175–185. <https://doi.org/10.1007/s40279-016-0534-9>
- Cauci S, Xodo S, Buligan C, Colaninno C, Barbina M, Barbina G, Francescato MP (2021) Oxidative stress is increased in combined oral contraceptives users and is positively associated with high-sensitivity c-reactive protein. *Molecules* 26:1070. <https://doi.org/10.3390/molecules26041070>
- Chrzanowski-Smith OJ, Edinburgh RM, Smith E, Thomas MP, Walhin JP, Koumanov F, Williams S, Betts JA, Gonzalez JT (2021) Resting skeletal muscle PNPLA2 (ATGL) and CPT1B are associated with peak fat oxidation rates in men and women but do not explain observed sex differences. *Exp Physiol* 106:1208–1223. <https://doi.org/10.1113/EP089431>
- de Wendt C, Espelage L, Eickelschulte S, Springer C, Toska L, Scheel A, Bedou AD, Benninghoff T, Cames S, Stermann T, Chadt A, Al-Hasani H (2021) Contraction-mediated glucose transport in skeletal muscle is regulated by a framework of AMPK, TBC1D1/4, and Rac1. *Diabetes* 70:2796–2809. <https://doi.org/10.2337/db21-0587>
- D'Eon TM, Rogers NH, Stancheva ZS, Greenberg AS (2008) Estradiol and the estradiol metabolite, 2-hydroxyestradiol, activate AMP-activated protein kinase in C2C12 myotubes. *Obesity (silver Spring)* 16:1284–1288. <https://doi.org/10.1038/oby.2008.50>
- D'Eon TM, Sharoff C, Chipkin SR, Grow D, Ruby BC, Braun B (2002) Regulation of exercise carbohydrate metabolism by estrogen and progesterone in women. *Am J Physiol Endocrinol Metab* 283:E1046–E1055. <https://doi.org/10.1152/ajpendo.00271.2002>
- D'Eon TM, Souza SC, Aronovitz M, Obin MS, Fried SK, Greenberg AS (2005) Estrogen regulation of adiposity and fuel partitioning. Evidence of genomic and non-genomic regulation of lipogenic and oxidative pathways. *J Biol Chem* 280:35983–35991. <https://doi.org/10.1074/jbc.M507339200>
- Devries MC (2016) Sex-based differences in endurance exercise muscle metabolism: impact on exercise and nutritional strategies to optimize health and performance in women. *Exp Physiol* 101:243–249. <https://doi.org/10.1113/EP085369>
- Devries MC, Hamadeh MJ, Graham TE, Tarnopolsky MA (2005) 17beta-estradiol supplementation decreases glucose rate of appearance and disappearance with no effect on glycogen utilization during moderate intensity exercise in men. *J Clin Endocrinol Metab* 90:6218–6225. <https://doi.org/10.1210/jc.2005-0926>
- Devries MC, Hamadeh MJ, Phillips SM, Tarnopolsky MA (2006) Menstrual cycle phase and sex influence muscle glycogen utilization and glucose turnover during moderate-intensity endurance exercise. *Am J Physiol Regul Integr Comp Physiol* 291:R1120–R1128. <https://doi.org/10.1152/ajpregu.00700.2005>
- Devries MC, Lowther SA, Glover AW, Hamadeh MJ, Tarnopolsky MA (2007) IMCL area density, but not IMCL utilization, is higher in women during moderate-intensity endurance exercise, compared with men. *Am J Physiol Regul Integr Comp Physiol* 293:R2336–R2342. <https://doi.org/10.1152/ajpregu.00510.2007>
- Ekenros L, Papoutsis Z, Fridén C, Dahlman Wright K, Lindén Hirschberg A (2017) Expression of sex steroid hormone receptors in human skeletal muscle during the menstrual cycle. *Acta Physiol (oxf)* 219:486–493. <https://doi.org/10.1111/apha.12757>
- Elliott-Sale KJ, Minahan CL, Janse de Jonge XAK, Ackerman KE, Sipilä S, Constantini NW, Lebrun CM, Hackney AC (2021) Methodological considerations for studies in sport and exercise science with women as participants: a working guide for standards of practice for research on women. *Sports Med* 51:843–861. <https://doi.org/10.1007/s40279-021-01435-8>
- Ellis GS, Lanza-Jacoby S, Gow A, Kendrick ZV (1994) Effects of estradiol on lipoprotein lipase activity and lipid availability in exercised male rats. *J Appl Physiol* 77:209–215. <https://doi.org/10.1152/jappl.1994.77.1.209>
- Enns DL, Tiidus PM (2008) Estrogen influences satellite cell activation and proliferation following downhill running in rats. *J Appl Physiol* 104:347–353. <https://doi.org/10.1152/japplphysiol.00128.2007>
- Enns DL, Tiidus PM (2010) The influence of estrogen on skeletal muscle: sex matters. *Sports Med* 40:41–58. <https://doi.org/10.2165/11319760-000000000-00000>
- Fatima LA, Campello RS, Barreto-Andrade JN, Passarelli M, Santos RS, Clegg DJ, Machado UF (2019) Estradiol stimulates adipogenesis and Slc2a4/GLUT4 expression via ESR1-mediated activation of CEBPA. *Mol Cell Endocrinol* 498:110447. <https://doi.org/10.1016/j.mce.2019.05.006>
- Fell JM, Hearris MA, Ellis DG, Moran JE, Jevons EF, Owens DJ, Strauss JA, Cocks M, Louis JB, Shepherd SO, Morton JP (2021) Carbohydrate improves exercise capacity but does not affect subcellular lipid droplet morphology, AMPK and p53 signalling in human skeletal muscle. *J Physiol* 599:2823–2849. <https://doi.org/10.1113/JP281127>
- Flores-Opazo M, McGee SL, Hargreaves M (2020) Exercise and GLUT4. *Exerc Sport Sci Rev* 48:110–118. <https://doi.org/10.1249/JES.0000000000000224>
- Forst-Ludwig A, Kintscher U (2010) Metabolic impact of estrogen signalling through ERalpha and ERbeta. *J Steroid Biochem Mol Biol* 122:74–81. <https://doi.org/10.1016/j.jsbmb.2010.06.012>
- Frandsen J, Amaro-Gahete FJ, Landgrebe A, Dela F, Ruiz JR, Helge JW, Larsen S (2021) The influence of age, sex and cardiorespiratory fitness on maximal fat oxidation rate. *Appl Physiol Nutr Metab* 46:1241–1247. <https://doi.org/10.1139/apnm-2021-0080>
- Frandsen J, Pistoljevic N, Quesada JP, Amaro-Gahete FJ, Ritz C, Larsen S, Dela F, Helge JW (2020) Menstrual cycle phase does not affect whole body peak fat oxidation rate during a graded exercise test. *J Appl Physiol* 128:681–687. <https://doi.org/10.1152/japplphysiol.00774.2019>
- Fu MH, Maher AC, Hamadeh MJ, Ye C, Tarnopolsky MA (2009) Exercise, sex, menstrual cycle phase and 17beta-estradiol influence metabolism-related genes in human skeletal muscle. *Physiol Genom* 40:34–47. <https://doi.org/10.1152/physiolgenomics.00115.2009>
- Fuentes N, Silveyra P (2019) Estrogen receptor signaling mechanisms. *Adv Protein Chem Struct Biol* 116:135–170. <https://doi.org/10.1016/bs.apcsb.2019.01.001>
- Funaki A, Gam H, Matsuda T, Ishikawa A, Yamada M, Ikegami N, Nishikawa Y, Sakamaki-Sunaga M (2022) Influence of menstrual cycle on leukocyte response following exercise-induced muscle damage. *Int J Environ Res Public Health* 19:9201. <https://doi.org/10.3390/ijerph19159201>
- Garg D, Ng SSM, Baig KM, Driggers P, Segars J (2017) Progesterone-mediated non-classical signaling. *Trends Endocrinol Metab* 28:656–668. <https://doi.org/10.1016/j.tem.2017.05.006>
- Gonzalez SL (2020) Progesterone for the treatment of central nervous system disorders: the many signaling roads for a single molecule. *Neural Regen Res* 15:1846–1847. <https://doi.org/10.4103/1673-5374.280314>
- Gorres BK, Bomhoff GL, Morris JK, Geiger PC (2011) In vivo stimulation of oestrogen receptor α increases insulin-stimulated skeletal muscle glucose uptake. *J Physiol* 589:2041–2054. <https://doi.org/10.1113/jphysiol.2010.199018>
- Gregorio KC, Laurindo CP, Machado UF (2021) Estrogen and glycemic homeostasis: the fundamental role of nuclear estrogen receptors ESR1/ESR2 in glucose transporter GLUT4 regulation. *Cells* 10:99. <https://doi.org/10.3390/cells10010099>
- Hackney AC (1990) Effects of the menstrual cycle on resting muscle glycogen content. *Horm Metab Res* 22:647. <https://doi.org/10.1055/s-2007-1004994>

- Hackney AC (1999) Influence of oestrogen on muscle glycogen utilization during exercise. *Acta Physiol Scand* 167:273–274. <https://doi.org/10.1046/j.1365-201x.1999.00605.x>
- Hackney AC (2021) Menstrual cycle hormonal changes and energy substrate metabolism in exercising women: a perspective. *Int J Environ Res Public Health* 18:10024. <https://doi.org/10.3390/ijerph181910024>
- Hackney AC, Curley CS, Nicklas BJ (1991) Physiological responses to submaximal exercise at the mid-follicular, ovulatory and mid-luteal phases of the menstrual cycle. *Scand J Med Sci Sports* 1:94–98. <https://doi.org/10.1111/j.1600-0838.1991.tb00277.x>
- Hackney AC, Kallman AL, Åggön E (2019) Female sex hormones and the recovery from exercise: menstrual cycle phase affects responses. *Biomed Hum Kinet* 11:87–89. <https://doi.org/10.2478/bhk-2019-0011>
- Hackney AC, Koltun KJ, Williett HN (2022) Menstrual cycle hormonal changes: estradiol- β -17 and progesterone interactions on exercise fat oxidation. *Endocrine* 76:240–242. <https://doi.org/10.1007/s12020-022-02998-w>
- Hackney AC, McCracken-Compton MA, Ainsworth B (1994) Substrate responses to submaximal exercise in the midfollicular and mid-luteal phases of the menstrual cycle. *Int J Sport Nutr* 4:299–308. <https://doi.org/10.1123/ijns.4.3.299>
- Häggström M, Richfield D (2014) Diagram of the pathways of human steroidogenesis. *WikiJ Med*. <https://doi.org/10.15347/2FWJM/2F2014.005>
- Haines M, McKinley-Barnard SK, Andre TL, Gann JJ, Hwang PS, Willoughby DS (2018) Skeletal muscle estrogen receptor activation in response to eccentric exercise up-regulates myogenic-related gene expression independent of differing serum estradiol levels occurring during the human menstrual cycle. *J Sports Sci Med* 17:31–39
- Hamadeh MJ, Devries MC, Tarnopolsky MA (2005) Estrogen supplementation reduces whole body leucine and carbohydrate oxidation and increases lipid oxidation in men during endurance exercise. *J Clin Endocrinol Metab* 90:3592–3599. <https://doi.org/10.1210/jc.2004-1743>
- Hansen M (2018) Female hormones: do they influence muscle and tendon protein metabolism? *Proc Nutr Soc* 77:32–41. <https://doi.org/10.1017/S0029665117001951>
- Hernández-Rabaza V, López-Pedrajas R, Almansa I (2019) Progesterone, lipoic acid and sulforaphane as promising antioxidants for retinal diseases: a review. *Antioxidants (basel)* 8:53. <https://doi.org/10.3390/antiox8030053>
- Homma H, Kurachi H, Nishio Y, Takeda T, Yamamoto T, Adachi K, Morishige K, Ohmichi M, Matsuzawa Y, Murata Y (2000) Estrogen suppresses transcription of lipoprotein lipase gene. Existence of a unique estrogen response element on the lipoprotein lipase promoter. *J Biol Chem* 275:11404–11411. <https://doi.org/10.1074/jbc.275.15.11404>
- Horton TJ, Miller EK, Bourret K (2006) No effect of menstrual cycle phase on glycerol or palmitate kinetics during 90 min of moderate exercise. *J Appl Physiol* 100:917–925. <https://doi.org/10.1152/jappphysiol.00491.2005>
- Horton TJ, Miller EK, Glueck D, Tench K (2002) No effect of menstrual cycle phase on glucose kinetics and fuel oxidation during moderate-intensity exercise. *Am J Physiol Endocrinol Metab* 282:E752–E762. <https://doi.org/10.1152/ajpendo.00238.2001>
- Houltham SD, Rowlands DS (2014) A snapshot of nitrogen balance in endurance-trained women. *Appl Physiol Nutr Metab* 39:219–225. <https://doi.org/10.1139/apnm-2013-0182>
- Hulton AT, Malone JJ, Campbell IT, MacLaren DP (2021) The effect of the menstrual cycle and hyperglycaemia on hormonal and metabolic responses during exercise. *Eur J Appl Physiol* 121:2993–3003. <https://doi.org/10.1007/s00421-021-04754-w>
- Jacobs KA, Casazza GA, Suh SH, Horning MA, Brooks GA (2005) Fatty acid reesterification but not oxidation is increased by oral contraceptive use in women. *J Appl Physiol* 98:1720–1731. <https://doi.org/10.1152/jappphysiol.00685.2004>
- Janse de Jonge X, Thompson B, Han A (2019) Methodological recommendations for menstrual cycle research in sports and exercise. *Med Sci Sports Exerc* 51:2610–2617. <https://doi.org/10.1249/MSS.0000000000002073>
- Jayaraman A, Pike CJ (2009) Progesterone attenuates oestrogen neuroprotection via downregulation of oestrogen receptor expression in cultured neurones. *J Neuroendocrinol* 21:77–81. <https://doi.org/10.1111/j.1365-2826.2008.01801.x>
- Jevons EFP, Gejl KD, Strauss JA, Ørtenblad N, Shepherd SO (2020) Skeletal muscle lipid droplets are resynthesized before being coated with perilipin proteins following prolonged exercise in elite male triathletes. *Am J Physiol Endocrinol Metab* 318:E357–E370. <https://doi.org/10.1152/ajpendo.00399.2019>
- Jurkowski JE, Jones NL, Toews CJ, Sutton JR (1981) Effects of menstrual cycle on blood lactate, O₂ delivery and performance during exercise. *J Appl Physiol Respir Environ Exerc Physiol* 51:1493–1499. <https://doi.org/10.1152/jappphysiol.1981.51.6.1493>
- Kaminski J, Junior CM, Pavesi H, Drobrzenski B, Amaral GMD (2021) Effects of oral versus transdermal estradiol plus micronized progesterone on thyroid hormones, hepatic proteins, lipids and quality of life in menopausal women with hypothyroidism: a clinical trial. *Menopause* 28:1044–1052. <https://doi.org/10.1097/GME.0000000000001811>
- Kang JH, Park JE, Dagoon J, Masson SWC, Merry TL, Bremner SN, Dent JR, Schenk S (2021) Sirtuin 1 is not required for contraction-stimulated glucose uptake in mouse skeletal muscle. *J Appl Physiol* 130:1893–1902. <https://doi.org/10.1152/jappphysiol.00065.2021>
- Kendall B, Eston R (2002) Exercise-induced muscle damage and the potential protective role of estrogen. *Sports Med* 32:103–123. <https://doi.org/10.2165/00007256-200232020-00003>
- Kim IY, Suh SH, Lee IK, Wolfe RR (2016) Applications of stable, non-radioactive isotope tracers in in vivo human metabolic research. *Exp Mol Med* 48:e203. <https://doi.org/10.1038/emmm.2015.97>
- Kjøbsted R, Roll JL, Jørgensen NO, Birk JB, Foretz M, Viollet B, Chadt A, Al-Hasani H, Wojtaszewski JF (2019) AMPK and TBC1D1 regulate muscle glucose uptake after, but not during, exercise and contraction. *Diabetes* 68:1427–1440. <https://doi.org/10.2337/db19-0050>
- Kriengsinyos W, Wykes LJ, Goonewardene LA, Ball RO, Pencharz PB (2004) Phase of menstrual cycle affects lysine requirement in healthy women. *Am J Physiol Endocrinol Metab* 287:E489–E496. <https://doi.org/10.1152/ajpendo.00262.2003>
- Kumagai H, Miyamoto-Mikami E, Takaragawa M, Kuriki K, Goto C, Shibata K, Yamada N, Hosono A, Fuku M, Suzuki S, Fuku N (2022) Genetic polymorphisms in CYP19A1 and ESR1 are associated with serum CK activity after prolonged running in men. *J Appl Physiol* 132:966–973. <https://doi.org/10.1152/jappphysiol.00374.2021>
- Lamont LS, Lemon PW, Bruot BC (1987) Menstrual cycle and exercise effects on protein catabolism. *Med Sci Sports Exerc* 19:106–110
- Lariviere F, Moussalli R, Garrel DR (1994) Increased leucine flux and leucine oxidation during the luteal phase of the menstrual cycle in women. *Am J Physiol* 267:E422–E428. <https://doi.org/10.1152/ajpendo.1994.267.3.E422>
- Larsen B, Cox A, Colbey C, Drew M, McGuire H, de St F, Groth B, Hughes D, Vlahovich N, Waddington G, Burke L, Lundy B, West N, Minahan C (2020) Inflammation and oral contraceptive use in female athletes before the Rio Olympic Games. *Front Physiol* 11:497. <https://doi.org/10.3389/fphys.2020.00497>
- Lavoie JM, Dionne N, Helie R, Brisson GR (1987) Menstrual cycle phase dissociation of blood glucose homeostasis during exercise.

- J Appl Physiol (1985) 62:1084–1089. <https://doi.org/10.1152/jappl.1987.62.3.1084>
- Lee SR, Choi WY, Heo JH, Huh J, Kim G, Lee KP, Kwun HJ, Shin HJ, Baek IJ, Hong EJ (2020) Progesterone increases blood glucose via hepatic progesterone receptor membrane component 1 under limited or impaired action of insulin. *Sci Rep* 10:16316. <https://doi.org/10.1038/s41598-020-73330-7>
- Lemoine S, Granier P, Tiffocche C, Berthon PM, Thieulant ML, Carré F, Delamarche P (2002) Effect of endurance training on oestrogen receptor alpha expression in different rat skeletal muscle type. *Acta Physiol Scand* 175:211–217. <https://doi.org/10.1046/j.1365-201X.2002.00992.x>
- Luk HY, Levitt DE, Appell C, Vingren JL (2021) Sex dimorphism in muscle damage-induced inflammation. *Med Sci Sports Exerc* 53:1595–1605. <https://doi.org/10.1249/MSS.0000000000002628>
- Lundsgaard AM, Fritzen AM, Kiens B (2018) Molecular regulation of fatty acid oxidation in skeletal muscle during aerobic exercise. *Trends Endocrinol Metab* 29:18–30. <https://doi.org/10.1016/j.tem.2017.10.011>
- MacPherson RE, Peters SJ (2015) Piecing together the puzzle of perilipin proteins and skeletal muscle lipolysis. *Appl Physiol Nutr Metab* 40:641–651. <https://doi.org/10.1139/apnm-2014-0485>
- Magkos F, Fabbri E, Patterson BW, Mittendorfer B, Klein S (2022) Physiological interindividual variability in endogenous estradiol concentration does not influence adipose tissue and hepatic lipid kinetics in women. *Eur J Endocrinol* 187:391–398. <https://doi.org/10.1530/EJE-22-0410>
- Maher AC, Akhtar M, Tarnopolsky MA (2010a) Men supplemented with 17beta-estradiol have increased beta-oxidation capacity in skeletal muscle. *Physiol Genom* 42:342–347. <https://doi.org/10.1152/physiolgenomics.00016.2010>
- Maher AC, Akhtar M, Vockley J, Tarnopolsky MA (2010b) Women have higher protein content of beta-oxidation enzymes in skeletal muscle than men. *PLoS ONE* 5:e12025. <https://doi.org/10.1371/journal.pone.0012025>
- Mahmoodzadeh S, Dworatzek E (2019) The role of 17β-estradiol and estrogen receptors in regulation of Ca²⁺ channels and mitochondrial function in cardiomyocytes. *Front Endocrinol (Lausanne)* 10(310):2019. <https://doi.org/10.3389/fendo.2019.00310>
- Matsuda T, Ogata H, Kanno M, Ishikawa A, Yamada M, Sakamaki-Sunaga M (2020) Effects of the menstrual cycle on oxidative stress and antioxidant response to high-intensity intermittent exercise until exhaustion in healthy women. *J Sports Med Phys Fit* 60:1335–1341. <https://doi.org/10.23736/S0022-4707.20.10868-5>
- Matsuda T, Takahashi H, Nakamura M, Kanno M, Ogata H, Ishikawa A, Yamada M, Kamemoto K, Sakamaki-Sunaga M (2022a) Influence of menstrual cycle on muscle glycogen utilization during high-intensity intermittent exercise until exhaustion in healthy women. *Appl Physiol Nutr Metab* 47:671–680. <https://doi.org/10.1139/apnm-2021-0532>
- Matsuda T, Takahashi H, Nakamura M, Ogata H, Kanno M, Ishikawa A, Sakamaki-Sunaga M (2022b) Influence of the menstrual cycle on muscle glycogen repletion after exhaustive exercise in eumenorrheic women. *J Strength Cond Res*. <https://doi.org/10.1519/JSC.0000000000004306>
- Matute ML, Kalkhoff RK (1973) Sex steroid influence on hepatic gluconeogenesis and glycogen formation. *Endocrinology* 92:762–768. <https://doi.org/10.1210/endo-92-3-762>
- Mauvais-Jarvis F, Clegg DJ, Hevener AL (2013) The role of estrogens in control of energy balance and glucose homeostasis. *Endocr Rev* 34:309–338. <https://doi.org/10.1210/er.2012-1055>
- McConell GK (2020) It's well and truly time to stop stating that AMPK regulates glucose uptake and fat oxidation during exercise. *Am J Physiol Endocrinol Metab* 318:E564–E567. <https://doi.org/10.1152/ajpendo.00511.2019>
- McLay RT, Thomson CD, Williams SM, Rehner NJ (2007) Carbohydrate loading and female endurance athletes: effect of menstrual cycle phase. *Int J Sport Nutr Exerc Metab* 17:189–205. <https://doi.org/10.1123/ijnsnem.17.2.189>
- Medina-Laver Y, Rodríguez-Varela C, Salsano S, Labarta E, Domínguez F (2021) What do we know about classical and non-classical progesterone receptors in the human female reproductive tract? A Review. *Int J Mol Sci* 22:11278. <https://doi.org/10.3390/ijms222011278>
- Mercer D, Convit L, Condo D, Carr AJ, Hamilton DL, Slater G, Snipe RM (2020) Protein requirements of pre-menopausal female athletes: systematic literature review. *Nutrients* 12:3527. <https://doi.org/10.3390/nu12113527>
- Mihm M, Gangooly S, Muttukrishna S (2011) The normal menstrual cycle in women. *Anim Reprod Sci* 124:229–236. <https://doi.org/10.1016/j.anireprosci.2010.08.030>
- Miklosz A, Łukaszuk B, Supruniuk E, Grubczak K, Moniuszko M, Choromańska B, Myśliwiec P, Chabowski A (2021) Does TBC1D4 (AS160) or TBC1D1 deficiency affect the expression of fatty acid handling proteins in the adipocytes differentiated from human adipose-derived mesenchymal stem cells (ADM-SCs) obtained from subcutaneous and visceral fat depots? *Cells* 10:1515. <https://doi.org/10.3390/cells10061515>
- Miller BF, Hansen M, Olesen JL, Flyvbjerg A, Schwarz P, Babraj JA, Smith K, Rennie MJ, Kjaer M (2006) No effect of menstrual cycle on myofibrillar and connective tissue protein synthesis in contracting skeletal muscle. *Am J Physiol Endocrinol Metab* 290:E163–E168. <https://doi.org/10.1152/ajpendo.00300.2005>
- Montero D, Madsen K, Meinild-Lundby AK, Edin F, Lundby C (2018) Sexual dimorphism of substrate utilization: differences in skeletal muscle mitochondrial volume density and function. *Exp Physiol* 103:851–859. <https://doi.org/10.1113/EP087007>
- Nicklas BJ, Hackney AC, Sharp RL (1989) The menstrual cycle and exercise: performance, muscle glycogen and substrate responses. *Int J Sports Med* 10:264–269. <https://doi.org/10.1055/s-2007-1024913>
- Oosthuyse T, Bosch AN (2010) The effect of the menstrual cycle on exercise metabolism: implications for exercise performance in eumenorrhoeic women. *Sports Med* 40:207–227. <https://doi.org/10.2165/11317090-000000000-00000>
- Oosthuyse T, Bosch AN (2012) Oestrogen's regulation of fat metabolism during exercise and gender specific effects. *Curr Opin Pharmacol* 12:363–371. <https://doi.org/10.1016/j.coph.2012.02.008>
- Oosthuyse T, Bosch AN (2017) The effect of gender and menstrual phase on serum creatine kinase activity and muscle soreness following downhill running. *Antioxidants (basel)* 6:16. <https://doi.org/10.3390/antiox6010016>
- Oosthuyse T, Bosch AN, Jackson S (2003) Effect of menstrual phase on the acetate correction factor used in metabolic tracer studies. *Can J Appl Physiol* 28:818–830. <https://doi.org/10.1139/h03-061>
- Oosthuyse T, Florence GE, Correia A, Smyth C, Bosch AN (2021) Carbohydrate-restricted exercise with protein increases self-selected training intensity in female cyclists but not male runners and cyclists. *J Strength Cond Res* 35:1547–1558. <https://doi.org/10.1519/JSC.0000000000004046>
- Park YM, Pereira RI, Erickson CB, Swibas TA, Kang C, Van Pelt RE (2017) Time since menopause and skeletal muscle estrogen receptors, PGC-1α and AMPK. *Menopause* 24:815–823. <https://doi.org/10.1097/GME.0000000000000829>
- Paroo Z, Dipchand ES, Noble EG (2002) Estrogen attenuates postexercise HSP70 expression in skeletal muscle. *Am J Physiol Cell Physiol* 282:C245–C251. <https://doi.org/10.1152/ajpcell.00336.2001>
- Peake JM, Neubauer O, Della Gatta PA, Nosaka K (2017) Muscle damage and inflammation during recovery from exercise. *J Appl*

- Physiol (1985) 122:559–570. <https://doi.org/10.1152/jappphysiol.00971.2016>
- Pedraza-Alva G, Zingg JM, Donda A, Pérez-Martínez L (2009) Estrogen receptor regulates MyoD gene expression by preventing AP-1-mediated repression. *Biochem Biophys Res Commun* 389:360–365. <https://doi.org/10.1016/j.bbrc.2009.08.153>
- Puglisi R, Mattia G, Carè A, Marano G, Malorni W, Matarrese P (2019) Non-genomic effects of estrogen on cell homeostasis and remodeling with special focus on cardiac ischemia/reperfusion injury. *Front Endocrinol (lausanne)* 10:733. <https://doi.org/10.3389/fendo.2019.00733>
- Quinn KM, Cox AJ, Roberts L, Pennell EN, McKeating DR, Fisher JJ, Perkins AV, Minahan C (2021) Temporal changes in blood oxidative stress biomarkers across the menstrual cycle and with oral contraceptive use in active women. *Eur J Appl Physiol* 121:2607–2620. <https://doi.org/10.1007/s00421-021-04734-0>
- Reis E, Frick U, Schmidtleicher D (1995) Frequency variations of strength training sessions triggered by the phases of the menstrual cycle. *Int J Sports Med* 16:545–550. <https://doi.org/10.1055/s-2007-973052>
- Ribas V, Nguyen MT, Henstridge DC, Nguyen AK, Beaven SW, Watt MJ, Hevener AL (2010) Impaired oxidative metabolism and inflammation are associated with insulin resistance in ER α -deficient mice. *Am J Physiol Endocrinol Metab* 298:E304–E319. <https://doi.org/10.1152/ajpendo.00504.2009>
- Richter EA (2021) Is GLUT4 translocation the answer to exercise-stimulated muscle glucose uptake? *Am J Physiol Endocrinol Metab* 320:E240–E243. <https://doi.org/10.1152/ajpendo.00503.2020>
- Roepstorff C, Steffensen CH, Madsen M, Stallknecht B, Kanstrup IL, Richter EA, Kiens B (2002) Gender differences in substrate utilization during submaximal exercise in endurance-trained subjects. *Am J Physiol Endocrinol Metab* 282:E435–E447. <https://doi.org/10.1152/ajpendo.00266.2001>
- Roepstorff C, Donsmark M, Thiele M, Vistisen B, Stewart G, Vissing K, Schjerling P, Hardie DG, Galbo H, Kiens B (2006) Sex differences in hormone-sensitive lipase expression, activity and phosphorylation in skeletal muscle at rest and during exercise. *Am J Physiol Endocrinol Metab* 291:E1106–E1114. <https://doi.org/10.1152/ajpendo.00097.2006>
- Rogers NH, Witczak CA, Hirshman MF, Goodyear LJ, Greenberg AS (2009) Estradiol stimulates Akt, AMP-activated protein kinase (AMPK) and TBC1D1/4, but not glucose uptake in rat soleus. *Biochem Biophys Res Commun* 382:646–650. <https://doi.org/10.1016/j.bbrc.2009.02.154>
- Romero-Parra N, Cupeiro N, Alfaro-Magallanes VM, Rael B, Rubio-Arias JA, Peinado AB, Benito PJ, IronFEMME Study Group (2021) Exercise-induced muscle damage during the menstrual cycle: a systematic review and meta-analysis. *J Strength Cond Res* 35:549–561. <https://doi.org/10.1519/JSC.0000000000.003878>
- Rowlands DS, Wadsworth DP (2011) Effect of high-protein feeding on performance and nitrogen balance in female cyclists. *Med Sci Sports Exerc* 43:44–53. <https://doi.org/10.1249/MSS.0b013e3181e93316>
- Ruby BC, Robergs RA, Waters DL, Burge M, Mermier C, Stolarczyk L (1997) Effects of estradiol on substrate turnover during exercise in amenorrheic females. *Med Sci Sports Exerc* 29:1160–1169. <https://doi.org/10.1097/00005768-199709000-00007>
- Sakamaki-Sunaga M, Min S, Kamemoto K, Okamoto T (2016) Effects of menstrual phase-dependent resistance training frequency on muscular hypertrophy and strength. *J Strength Cond Res* 30:1727–1734. <https://doi.org/10.1519/JSC.0000000000.001250>
- Sakamoto K, Holman GD (2008) Emerging role for AS160/TBC1D4 and TBC1D1 in the regulation of GLUT4 traffic. *Am J Physiol Endocrinol Metab* 295:E29–E37. <https://doi.org/10.1152/ajpendo.90331.2008>
- Salehzadeh F, Rune A, Osler M, Al-Khalili L (2011) Testosterone or 17 β -estradiol exposure reveals sex-specific effects on glucose and lipid metabolism in human myotubes. *J Endocrinol* 210:219–229. <https://doi.org/10.1530/JOE-10-0497>
- Schmidt SL, Bessesen DH, Stotz S, Peelor FF 3rd, Miller BF, Horton TJ (2014) Adrenergic control of lipolysis in women compared with men. *J Appl Physiol* (1985) 117:1008–1019. <https://doi.org/10.1152/jappphysiol.00003.2014>
- Sewright KA, Hubal MJ, Kearns A, Holbrook MT, Clarkson PM (2008) Sex differences in response to maximal eccentric exercise. *Med Sci Sports Exerc* 40:242–251. <https://doi.org/10.1249/mss.0b013e31815aedd4>
- Sidossis LS, Coggan AR, Gastaldelli A, Wolfe RR (1995a) Pathway of free fatty acid oxidation in human subjects. Implications for tracer studies. *J Clin Invest* 95:278–284. <https://doi.org/10.1172/JCI117652>
- Sidossis LS, Coggan AR, Gastaldelli A, Wolfe RR (1995b) A new correction factor for use in tracer estimations of plasma fatty acid oxidation. *Am J Physiol* 269:E649–E656. <https://doi.org/10.1152/ajpendo.1995.269.4.E649>
- Steffensen CH, Roepstorff C, Madsen M, Kiens B (2002) Myocellular triacylglycerol breakdown in females but not in males during exercise. *Am J Physiol Endocrinol Metab* 282:E634–E642. <https://doi.org/10.1152/ajpendo.00078.2001>
- Strauss JA, Shepherd DA, Macey M, Jevons EF, Shepherd SO (2020) Divergence exists in the subcellular distribution of intramuscular triglyceride in human skeletal muscle dependent on the choice of lipid dye. *Histochem Cell Biol* 154:369–382. <https://doi.org/10.1007/s00418-020-01898-2>
- Strehlow K, Rotter S, Wassmann S, Adam O, Grohé C, Laufs K, Böhm M, Nickenig G (2003) Modulation of antioxidant enzyme expression and function by estrogen. *Circ Res* 93:170–177. <https://doi.org/10.1161/01.RES.0000082334.17947.11>
- Stricker R, Eberhart R, Chevailler MC, Quinn FA, Bischof P, Stricker R (2006) Establishment of detailed reference values for luteinizing hormone, follicle stimulating hormone, estradiol and progesterone during different phases of the menstrual cycle on the Abbott ARCHITECT analyzer. *Clin Chem Lab Med* 44:883–887. <https://doi.org/10.1515/CCLM.2006.160>
- Stupka N, Lowther S, Chorneyko K, Bourgeois JM, Hogben C, Tarnopolsky MA (2000) Gender differences in muscle inflammation after eccentric exercise. *J Appl Physiol* (1985) 89:2325–2332. <https://doi.org/10.1152/jappl.2000.89.6.2325>
- Suh SH, Casazza GA, Horning MA, Miller BF, Brooks GA (2002) Luteal and follicular glucose fluxes during rest and exercise in 3-h postabsorptive women. *J Appl Physiol* (1985) 93:42–50. <https://doi.org/10.1152/jappphysiol.01080.2001>
- Suh SH, Casazza GA, Horning MA, Miller BF, Brooks GA (2003) Effects of oral contraceptives on glucose flux and substrate oxidation rates during rest and exercise. *J Appl Physiol* (1985) 94:285–294. <https://doi.org/10.1152/jappphysiol.00693.2002>
- Sung E, Han A, Hinrichs T, Vorgerd M, Manchado C, Platen P (2014) Effects of follicular versus luteal phase-based strength training in young women. *Springerplus* 3:668. <https://doi.org/10.1186/2193-1801-3-668>
- Sylov L, Kleinert M, Richter EA, Jensen TE (2017) Exercise-stimulated glucose uptake - regulation and implications for glycaemic control. *Nat Rev Endocrinol* 13:133–148. <https://doi.org/10.1038/nrendo.2016.162>
- Tarnopolsky MA (2008) Sex differences in exercise metabolism and the role of 17 β -estradiol. *Med Sci Sports Exerc* 40:648–654. <https://doi.org/10.1249/MSS.0b013e31816212ff>
- Thomas A, Bunyan K, Tiidus PM (2010) Oestrogen receptor- α activation augments post-exercise myoblast proliferation. *Acta*

- Physiol (oxf) 198:81–89. <https://doi.org/10.1111/j.1748-1716.2009.02033.x>
- Tiidus PM (2003) Influence of estrogen on skeletal muscle damage, inflammation and repair. *Exerc Sport Sci Rev* 31:40–44. <https://doi.org/10.1097/00003677-200301000-00008>
- Torre F, Calogero AE, Condorelli RA, Cannarella R, Aversa A, La Vignera S (2020) Effects of oral contraceptives on thyroid function and vice versa. *J Endocrinol Investig* 43:1181–1188. <https://doi.org/10.1007/s40618-020-01230-8>
- Toth MJ, Sites CK, Matthews DE, Casson PR (2006) Ovarian suppression with gonadotropin-releasing hormone agonist reduces whole body protein turnover in women. *Am J Physiol Endocrinol Metab* 291:E483–E490. <https://doi.org/10.1152/ajpendo.00600.2005>
- Trommelen J, Fuchs CJ, Beelen M, Lenaerts K, Jeukendrup AE, Cermak NM, van Loon LJ (2017) Fructose and sucrose intake increase exogenous carbohydrate oxidation during exercise. *Nutrients* 20:167. <https://doi.org/10.3390/nu9020167>
- Ventura-Clapier R, Piquereau J, Veksler V, Garnier A (2019) Estrogens, estrogen receptors effects on cardiac and skeletal muscle mitochondria. *Front Endocrinol (lausanne)* 10:557. <https://doi.org/10.3389/fendo.2019.00557>
- Viña J, Sastre J, Pallardó FV, Gambini J, Borrás C (2008) Modulation of longevity-associated genes by estrogens or phytoestrogens. *Biol Chem* 389:273–277. <https://doi.org/10.1515/BC.2008.027>
- Wada T, Hori S, Sugiyama M, Fujisawa E, Nakano T, Tsuneki H, Nagira K, Saito S, Sasaoka T (2010) Progesterone inhibits glucose uptake by affecting diverse steps of insulin signaling in 3T3-L1 adipocytes. *Am J Physiol Endocrinol Metab* 298:E881–E888. <https://doi.org/10.1152/ajpendo.00649.2009>
- White LJ, Ferguson MA, McCoy SC, Kim H (2003) Intramyocellular lipid changes in men and women during aerobic exercise: a (1) H-magnetic resonance spectroscopy study. *J Clin Endocrinol Metab* 88:5638–5643. <https://doi.org/10.1210/jc.2003-031006>
- Wiik A, Gustafsson T, Esbjörnsson M, Johansson O, Ekman M, Sundberg CJ, Jansson E (2005) Expression of oestrogen receptor alpha and beta is higher in skeletal muscle of highly endurance-trained than of moderately active men. *Acta Physiol Scand* 184:105–112. <https://doi.org/10.1111/j.1365-201X.2005.01433.x>
- Wikström-Frisén L, Boraxbekk CJ, Henriksson-Larsén K (2017) Effects on power, strength and lean body mass of menstrual/oral contraceptive cycle based resistance training. *J Sports Med Phys Fit* 57:43–52. <https://doi.org/10.23736/S0022-4707.16.05848-5>
- Willett HN, Koltun KJ, Hackney AC (2021) Influence of menstrual cycle estradiol- β -17 fluctuations on energy substrate utilization-oxidation during aerobic, endurance exercise. *Int J Environ Res Public Health* 18:7209. <https://doi.org/10.3390/ijerph18137209>
- Williams T, Walz E, Lane AR, Pebole M, Hackney AC (2015) The effect of estrogen on muscle damage biomarkers following prolonged aerobic exercise in eumenorrheic women. *Biol Sport* 32:193–198. <https://doi.org/10.5604/20831862.1150300>
- Wu H, Wu ZG, Shi WJ, Gao H, Wu HH, Bian F, Jia PP, Hou YN (2019) Effects of progesterone on glucose uptake in neurons of Alzheimer's disease animals and cell models. *Life Sci* 238:116979. <https://doi.org/10.1016/j.lfs.2019.116979>
- Yan H, Yang W, Zhou F, Li X, Pan Q, Shen Z, Han G, Newell-Fugate A, Tian Y, Majeti R, Liu W, Xu Y, Wu C, Allred K, Allred C, Sun Y, Guo S (2019) Estrogen improves insulin sensitivity and suppresses gluconeogenesis via the transcription factor Foxo1. *Diabetes* 68:291–304. <https://doi.org/10.2337/db18-0638>
- Yang S, Wang J (2015) Estrogen activates AMP-activated protein kinase in human endothelial cells via ER β /Ca(2+)/calmodulin-dependent protein kinase kinase β pathway. *Cell Biochem Biophys* 72:701–707. <https://doi.org/10.1007/s12013-015-0521-z>
- Yaşar P, Ayaz G, User SD, Güpür G, Muyan M (2016) Molecular mechanism of estrogen-estrogen receptor signaling. *Reprod Med Biol* 16:4–20. <https://doi.org/10.1002/rmb2.12006>
- Yue Y, Zhang C, Zhang X, Zhang S, Liu Q, Hu F, Lv X, Li H, Yang J, Wang X, Chen L, Yao Z, Duan H, Niu W (2020) An AMPK/Axin1-Rac1 signaling pathway mediates contraction-regulated glucose uptake in skeletal muscle cells. *Am J Physiol Endocrinol Metab* 318:E330–E342. <https://doi.org/10.1152/ajpendo.00272.2019>
- Zderic TW, Coggan AR, Ruby BC (2001) Glucose kinetics and substrate oxidation during exercise in the follicular and luteal phases. *J Appl Physiol* (1985) 90:447–453. <https://doi.org/10.1152/jappl.2001.90.2.447>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.