## References

Anderson CS, Linto J, Stewart-Wynne ED 1995 A Population-Based Assessment of the Impact and Burden of Caregiving for Long-term Stroke Survivors. Stroke 26: 843 - 849 Aprile I, Piazzini DB, Bertolini C, Caliandro P, Pazzaglia C, Tonali P, Padua L 2006 Predictive variables on disability and quality of life in stroke outpatients undergoing rehabilitation. Neuro Sci 27: 40-46 Ashburn A 1999 *Physical recovery following stroke*. Physiotherapy 83 (9): 480 - 490Bamford J, Sandercock P, Dennis M, Burn J, Warlow C 1991 Classification and natural history of clinically identifiable subtypes of cerebral infarction. Lancet 337: 1521 - 1526 Barer DH 1989 Continence after stroke: useful predictor or goal of therapy? Age Ageing 18 (3): 183 - 191 Belanger L, Bolduc M, Noel M 1988 Relative importance of after-effects, environmental and socio-economic factors on the social integration of stroke victims. Int J Rehabil Res 11 (3): 251 - 260 Bernspang B, Fisher AG 1995 Differences between persons with right or left cerebral vascular accident on the Assessment of Motor and Process skills. Arch Phys Med Rehabil 76 (12): 1144 - 1151 Brown D, Ransom J, Hass S, Petty G, O'Fallon M, Whisnant P, Leisbon L 1999 Use of Nursing Home After Stroke and Dependence on Stroke Severity: A Population-Based Analysis. Stroke 30: 924 - 929 Boden-Albala B, Litwak E, Elkind MSV, Rundek T, Sacco RL 2005 Social isolation and outcomes post stroke. Neurology 64: 1888 – 1892. Brittain KR, Perry SI, Peet SM, Shaw C, Dallosso H, Assassa RP, Williams K, Jagger C, Potter JF, Castleden CM 2000 Prevalence and Impact of Urinary Symptoms Among Community-Dwelling Stroke Survivors. Stroke 31:886 Collin C, Wade DT, Davies S, Horne V 1988 The Barthel ADL Index: A reliability study. Int Disabil Stud 10 (2): 61 - 6.3 Connor MD, Thorogood M, Casserly B, Dobson C, Warlow MD 2004 Prevalence of stroke survivors in rural South Africa: Results from the Southern Africa stroke prevention initiative (SASPI) Agincourt field site. Stroke 35: 627 – 632. D'Olhaberriague L, Litvan I, Mitsias P, Mansbach H 1996 A Reappraisal of Reliability and Validity Studies in Stroke. Stroke 27: 2331 – 2336. Dromerick AW, Edwards DF, Diringer MN 2003 Sensitivity to changes in disability after stroke: A comparison of four scales useful in clinical trials. Journal of Rehabilitation Research and Development 40 (1): 1 - 8. Eales CJ, Noakes TD, Stewart AV, Becker P 2004 Self-responsibility predicts the successful outcome of coronary artery bypass surgery. The South African Journal of Physiotherapy 60 (2): 11 - 19 Fritz VU 1997 Stroke incidence in South Africa. SAMJ 87 (5): 584 - 585 Gainotti G, Antonucci G, Marra C, Paolucci S 2001 Relation between depression after stroke, antidepressant therapy, and functional recovery. J. Neurol. Neurosurg. Psychiatry 71: 258 - 261

Glader EL, Stegmayr B, Johansson L, Hulter-Asberg K, Wester PO 2001 Differences in Long-Term Outcome Between Patients Treated in Stroke Units and in General Wards: A 2-Year Follow-Up of Stroke Patients in Sweden. Stroke 32: 2124.

Glass TA, Matchar DB, Belyea M, Feussner JR 1993 *Impact of social support on outcome in first stroke*. Stroke 24 (1): 64 – 70

Granger C, Dewis L, Peters N, Sherwood C, Barrett J 1979 *Stroke rehabilitation: an analysis of repeated Barthel Index measures.* Arch Phys Med Rehabil 60 (1): 14-17

Green J, Forster A, Young J 2001 A test-retest reliability of the Barthel Index, the Rivermead Mobility Index, the Nottingham Extended Activities of Daily Living Scale and the Frenchay Activities Index in stroke patients. Disability and Rehabilitation. 23: 670-676

Green J, Valvanne-Tommila H, Loubser H, Gordon R, Paunio P, Marosszeky JE, Richards B 2005 *Australia, Finland and South Africa – Delving into Data to Investigate Differences in Stroke Rehabilitation.* International Society of Physical and Rehabilitation Medicine – ISPRM:37 - 40

Greenberg E, Treger J, Ring H 2004 *Post-stroke follow-up in a rehabilitation center outpatient clinic*. Isr Med Assoc J 6 (10): 603 - 606 Guccionne AA, Felson DT, Anderson JJ, Anthony JM, Zhang Y, Wilson PW, Kelly-Hayes M, Wolf PA, Kreger BE, Kannel Wb 1994 *The effects of specific medical conditions on the functional limitations of elders in the Framingham Study*. AM J Public Health 84 (3): 351 - 358

Hale LA, Eales CJ 1998 *Recovery of walking function in stroke patients after minimal rehabilitation*. Physiotherapy Research International 3 (3): 194 – 205.

Hale LA, Eales CJ, Stewart A, Fritz VU 1999 *The problems experienced by black stroke patients in SOWETO, South Africa.* The South African Journal of physiotherapy 55 (2): 12 - 17.

Herrmann N, Black S, Lawrence J, Szekely C, Szalai J 1998 *The Sunnybrook Stroke Study: A Prospective Study of Depressive Symptoms and Functional Outcome.* Stroke 29: 618 - 624

Hobart JC, Lamping DL, Freeman JA, Langdon DW, McLellan DL, Greenwood RJ, Thompson AJ 2001 *Evidence-based measurement:Which disability scale for neurologic rehabilitation?* Neurology 57: 639 - 644 Horner RD, Swanson JW, Boworth HB, Matchar DB 2003 *Effects of Race and Poverty on the Process and Outcome of Inpatient Rehabilitation Services among Stroke Patients.* Stroke 34: 1027.

Hsueh IP, Lin JH, Jeng JS, Hsieh CL 2002 Comparison of the psychometric characteristics of the functional independence measure, 5 item Barthel index, and 10 item Barthel index in patients with stroke. J. Neurol. Neurosurg. Psychiatry 73: 188 - 190

Hughes J, Jelsma J, Maclean E, Darder M, Tinise X 2004 *The healthrelated quality of life of people living with HIV/AIDS*. Disabil Rehabil 18 (6): 371 – 6

Jakobsson UIF, Klevsgard R, Westergren A, Hallberg I 2003 *Old people in Pain: A comparative study.* Journal of pain and Symptom Management 26 (1): 625 - 636

Jonsson A, Lindgren I, Hallstrom B, Norrving B, Lindgren A 2005 Prevalence and Intensity of Pain after Stroke: A population-based study *focusing on patients' perspectives.* J. Neurol. Neurosurg. Psychiatry published online; doi: 10.1136/jnnp.2005.079145

Jorgensen HS, Reith J, Nakayama H, Kammersgaard LP, Raaschou HO, Olsen TS, 1999 What determines good recovery in patients with the most severe strokes? The Copenhagen stroke study. Stroke 30: 2008 - 2012 Kalra L 1994 The influence of stroke unit rehabilitation on functional recovery from stroke. Stroke 25 (4): 821 – 825

Kauhanen M, Korpelainen J, Hiltunen P, Nieminen P, Sotaniemi K, Myllyla V 2000 *Domains and Determinants of Quality of Life After Stroke Caused by Brain Infarction*. Arch Phys Med Rehabil 81: 1541 - 1546

Kelly-Hayes M, Beiser A, Kase CS, Scaramucci A, D'Agostino RB, Wolf PA 2003 *The Influence of Gender and Age on Disability following Ischemic Stroke: the Framingham study*. Journal of Stroke and Cerebrovascular Diseases 12: 119 – 126.

Kelly PJ, Furie KL, Shafqat S, Rallis N, Chang Y, Stein J 2003 *Functional recovery following rehabilitation after hemorrhagic and ischaemic stroke.* Arch Phys Med Rehabil 84: 968 - 972

Knapp P, Hewison J 1999 *Disagreement in Patient and Carer Assessment* of Functional Abilities after Stroke. Stroke 30: 934 – 938.

Lam GW, Foldspang A, Elving LB, Mommsen S 1992 Social context, social abstention, and problem recognition correlated with adult female urinary incontinence. Dan Med Bull 39 (6): 565 - 570

Louisette M, Therese A, Rejean H, Annie R, Marie-France D 2001 *Impact* of Motor, Cognitive and Perceptual Disorders on Ability to Perform Activities of Daily Living after Stroke. Stroke 32: 2602.

Mars M 2004 *HIV* – *Implications for exercise in treatment and rehabilitation.* The South African Journal of Physiotherapy 60 (4): 9 - 14 Mayo NE, Wood-Dauphinee S, Cote R, Gayton D, Carlton J, Buttery J, Tamblyn R 2000 *There's No Place Like Home.* Stroke 31: 1016 – 1023 Nakayama H, Jorgensen HS, Raaschou HO, Olsen TS 1994 *The influence of age on stroke outcome. The Copenhagen Stroke Study.* Stroke 25 (4): 808 - 813

Nunnaly JC 1978 *Psychometric theory*, 2<sup>nd</sup> edn.pp 230 – 245. McGraw-Hill, New York.

Paolucci S, Antonucci G, Grasso MG, Bragoni M, Coiro P, De Angelis D, Fusco FR, Morelli D, Venturiero V, Troisi E, Pratesi L 2003 *Functional Outcome of Ischemic and Haemorrhagic Stroke Patients After Inpatient Rehabilitation: A matched comparison.* Stroke 34: 2861.

Paolucci S, Antonucci G, Pratesi L, Traballesi M, Grasso M, Lubich S 1999 *Poststroke Depression and Its Role in Rehabilitation of Inpatients*. Arch Phys Med Rehabil 80: 985 – 989

Pinsky JL, Branch LG, Jette AM, Haynes SG, Feinleib M, Cornoni-Huntley JC, Bailey KR 1985 *Framingham Disability Study: relationship of diability to cardiovascular risk factors among persons free of diagnosed cardiovascular diseases.* Am J Epidemiol 122 (4): 644 – 56

Pohjahsvaara T, Erkinjuntti T, Vataja R, Kaste M 1997 Comparison of stroke Features and Disability in Daily Life in Patients With Ischaemic Stroke Aged 55 to 70 and 71 to 85 Years. Stroke 28: 729 - 735 Rexroth P, Fisher AG, Merritt BK, Gliner J 2005 *ADL differences in individuals with unilateral hemispheric stroke*. Can J Occup Ther 72 (4): 212 - 221

Sabine P, Aurelie D, Benoit N, Jean FP, Valerie K, Phillipe G 2002 *Prognostic Factors in the Recovery of the Ability to Walk After Stroke.* Journal of Stroke and Cerebrovascular Diseases 11 (6): 330 -335. Sangha H, Lipson D, Foley N, Salter K, Bhogal S, Ponahi G, Teasell RW 2005 *A comparison of the Barthel Index and the Functional Independece Measure as outcome measures in stroke rehabilitation: patterns of disability scale usage in clinical trials.* Int J Rehabil Res 28 (2): 135 – 139. Schepers V, Visser-Meily A, Ketelaar M, Lindeman E 2005 *Prediction of Social Activity 1 Year Poststroke.* Arch Phys Med Rehabil 86: 1472 - 1476 Schlote A, Kruger J, Topp H, Wallesch CW 2004 *Inter-rater of the Barthel Index, the Activity Index, and the Nottingham Extended Activities of Daily Living: The use of ADL instruments in stroke rehabilitation by medical and non medical personnel.* Rehabilitation (Stuttg) 43 (2): 75 – 82 Schmidt S, Herman L, Koenig P, Leuze P, Monahan M 1986 *Status of* 

stroke patients: a community assessment. Arch Phys Med Rehabil 67 (2): 99 - 102

Schwartz J, Speed N, Brunberg J, Brewer T, Brown M, Greden J 1993 Depression in stroke rehabilitation. Biol Psychiatry 33 (10): 694 - 699 Sharrack B, Hughes RAC, Soudain S, Dunn G 1999 The psychometric properties of clinical rating scales used in multiple sclerosis. Barin 122 (1): 141 – 159.

Stewart AV, Eales CJ, De Charmoy S 2000 *The knowledge profile of patients with hypertension*. The South African Journal of Physiotherapy 56 (4): 17 - 21

Steyn K, Fourie J, Bradshaw D 1992 *The impact of chronic diseases of lifestyle and their major risk factors on mortality in South Africa.* S Afr Med J 82: 227 - 231

Stineman M, Maislin G, Fiedler RC, Granger CV 1997 A Prediction Model for Functional Recovery in Stroke. Stroke 28: 550 - 556

Sturm JW, Donnan GA, Dewey HM, Macdonnell RAL, Gilligan KA, Thrift AG 2004 *Determinants of Handicap After Stroke*: The North East Melbourne Stroke Incidence Study (NEMESIS). Stroke 35: 715 - 720 The World Health Organization MONICA Project 1988 *(monitoring trends and determinants in cardiovascular disease: a major international collaboration.* J Clin Epidemiol 41 (2): 105 - 114

Tilling K, Sterne J, Rudd A, Glass T, Wityk R, Wolfe C 2001 A new method for predicting recovery after stroke. Stroke 32: 2867 – 2873 Van de Weg F, Kuik D, Lankhorst G 1999 Post-stroke depression and functional outcome: a cohort study investigating the influence of depression on functional recovery from stroke. Clin Rehabil 13 (3): 268 -272

Van Rooijen AJ, Rheeder P, Eales CJ, Becker P 2001 *Black female patients with type 2 diabetes mellitus: Knowledge, Attitudes and Physical activity.* The South African Journal of Physiotherapy 57 (3): 20 - 27