



Unpacking the issues on child stunting in Nairobi slums: a time for action

WHY IS THIS STUDY IMPORTANT?

Growth trajectories are central to monitoring child health. They provide essential indicators of childhood development and can predict determinants of adult health outcomes.

In African countries, approximately 40% of under-five children are affected by growth impairment including stunting (36%) and overweight (7%). Child stunting is more pronounced in Sub-Saharan Africa, in particular Eastern and Southern Africa, where the prevalence reached 40% among under-five children according to UNICEF 2007-2011 data. In Kenya, the prevalence of stunting among under-five children decreased from 35% in 2008-09 to 26% in 2014. This figure however hides intra-urban disparities as child stunting was estimated at 46% in 2010 in two Nairobi slums, Korogocho and Viwandani, covering a total population of an estimated 70,000 residents.

HOW WAS THE STUDY DONE?

This study used longitudinal data from the Nairobi Urban Health and Demographic Surveillance System (NUHDSS) to identify factors associated with linear growth among under-five children in two informal settlements in Nairobi.

We looked at 1,917 children born in 2007 and 2008 and followed them up till they were five years old. All children were less than two years at the time of recruitment.

The main outcome variable was the height for age z-scores (HAZ), computed at each child visit based on the World Health Organization (WHO) 2006 standards.

We fitted a generalized linear model on the HAZ to analyse linear growth trajectories among under-five children.

Highlights

- Findings from this study point to the need to prioritize the urban poor population in child health programs in Nairobi. They also call for changes in current programs addressing child undernutrition in Nairobi informal settlements slums.
- The study identifies important child and maternal factors that are closely related to linear growth among under-five children in two Nairobi's informal settlements. It is worth focusing on those factors that can be changed, particularly: child weight at birth, immunization status, child feeding and mother's education and socio-economic status. These factors should be considered when designing interventions aiming at addressing child health inequities in these settings.
- The interventions must also take into account the enabling role of women's education (secondary or higher) in ensuring good child health outcomes, including normal linear growth. To improve child health, young women need to be in school.
- Targeting single mothers in child health and economic interventions in these areas would contribute to lowering the prevalence of stunting among under-five children.
- In a context of poor access to formal housing, uncertain daily access to food and to basic social needs, policy-makers and program implementers must prioritize economic reinforcement interventions such as cash transfers and access to decent employment and housing.

KEY FINDINGS

- There is a decline in linear growth from after birth up to about 24 months, followed by regular catch-up to about 5 years, as shown in Figure 1. This is something that is peculiar to developing countries.
- Children living in Korogocho had better linear growth in their first months of life than those in Viwandani; then from about 8 months, the linear growth became greater for children living in Viwandani.
- Linear growth patterns for girls remained above that for boys throughout the period 0 to 5 years.
- Children with low birth weight remained below those with normal birth weight and below those whose birth weight was missing.
- Few differences in linear growth patterns were observed between children who had more than one illness symptoms over the last two weeks preceding the survey round and those who did not.
- Interestingly, children in the study areas, both those who were stunted and those who were not, brushed the lower boundary of the WHO height references relative to their age, reflecting persistent failure in achieving expected height for age.





Figure 1. Variation in child HAZ for specific characteristics



- More than half the children (58%) were stunted at some point before the age of five years, with a significantly greater proportion found in Korogocho compared to Viwandani. Children in Viwandani were more likely to grow faster than those in Korogocho.
- Children from least poor households were more likely to grow faster than those from the poorest households.
- Low birth weight was associated with slower linear growth while full immunization status predicted faster linear growth.
- Children who were exclusively breastfed in the first six months of life, presented better linear growth outcomes.
- Children of older mothers (25-34 years) were more likely to grow faster than those from younger mothers (age below 18). Children of currently married mothers were more likely to grow faster than those of formerly married mothers.



REFERENCES

1. Haymond M, Kappelgaard A-M, Czernichow P, Biller BMK, Takano K, Kiess W, et al. Early recognition of growth abnormalities permitting early intervention. *Acta Paediatrica* (Oslo, Norway : 1992). 2013;102(8):787-96.
2. Oddy WH, Kendall GE, Blair E, De Klerk NH, Stanley FJ, Landau LI, et al. Breast feeding and cognitive development in childhood: a prospective birth cohort study. *Paediatr Perinat Epidemiol*. 2003;17(1):81-90.
3. Victora C, Adair L, Fall C, Hallal P, Martorell R, Richter L, et al. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet*. 2008;371(9609):302.
4. Legler JD, Rose LC. Assessment of abnormal growth curves. *American family physician*. 1998;58(1):153-8.
5. WHO Multicentre Growth Reference Study Group. WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age. Geneva: World Health Organization; 2006. Report No.: 924154693X.
6. Adair LS, Cole TJ. Rapid child growth raises blood pressure in adolescent boys who were thin at birth. *Hypertension*. 2003;41(3):451-6.
7. Ong KKL, Ahmed ML, Emmett PM, Preece MA, Dunger DB, the Avon Longitudinal Study of P, et al. Association between postnatal catch-up growth and obesity in childhood: prospective cohort study. *BMJ : British Medical Journal*. 2000;320(7240):967-71.
8. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, De Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*. 2013;382(9890):427-51.
9. Mendez MA, Adair LS. Severity and timing of stunting in the first two years of life affect performance on cognitive tests in late childhood. *The Journal of nutrition*. 1999;129(8):1555-62.
10. De Onis M. Child growth and development. *Nutrition and health in developing countries*: Springer; 2008. p. 113-37.
11. Caulfield LE, de Onis M, Blössner M, Black RE. Under-nutrition as an underlying cause of child deaths associated with diarrhea, pneumonia, malaria, and measles. *The American journal of clinical nutrition*. 2004;80(1):193-8.
12. Graber E, Rapaport R. Growth and growth disorders in children and adolescents. *Pediatric annals*. 2012;41(4):e1-9.
13. Rapaport R, Bowlby D. Clinical aspects of growth and growth disorders. *Pescovitz OH & Eugster EA Pediatric endocrinology: mechanisms, manifestations, and management Philadelphia: Lippincott Williams&Willkins*. 2004:172-90.
14. UNICEF. Improving child nutrition: the achievable imperative for global progress: United Nations Children's Fund; 2013.
15. Kenya National Bureau of Statistics (KNBS) and ICF Macro. Kenya Demographic and Health Survey 2008-09. Calverton, Maryland; 2010.
16. Kenya National Bureau of Statistics (KNBS). Kenya Demographic and Health Survey 2014. 2015.
17. Kimani-Murage EW, Muthuri SK, Oti SO, Mutua MK, van de Vijver S, Kyobutungi C. Evidence of a Double Burden of Malnutrition in Urban Poor Settings in Nairobi, Kenya. *PloS one*. 2015;10(6):e0129943.
18. Black R, Allen L, Bhutta Z, Caulfield L, de Onis M, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*. 2008;371:243 - 60.
19. UNICEF/WHO. UNICEF/WHO Joint Committee on Health Policy: Report of the 27th Session Held at the Headquarters of the World Health Organization, Geneva, 23-25 January 1989. UN; 1989.



SOURCE: Cheikh M. Faye, Sharon Fonn, Johnathan Levin, Elisabeth Kimani-Murage. 2018. Analysing child linear growth trajectories among under-5 children in two Nairobi informal settlements, *Public Health Nutrition* doi:10.1017/S1368980019000491

AUTHORS: Cheikh Mbacké Faye¹², Sharon Fonn², Jonathan Levin², Elisabeth Kimani-Murage¹

AFFILIATIONS: ¹ African Population and Health Research Center, Tel. +254 20 400 1000. P.O Box 10787 – 00100, Nairobi, Kenya. ² University of the Witwatersrand, School of Public Health, 27 St Andrews Road, Johannesburg Parktown 2193, South Africa

FUNDING: This research was supported by the Consortium for Advanced Research Training in Africa (CARTA). CARTA is jointly led by the African Population and Health Research Center and the University of the Witwatersrand and funded by the Carnegie Corporation of New York (Grant No--B 8606.R02), Sida (Grant No:54100029), the DELTAS Africa Initiative (Grant No: 107768/Z/15/Z). The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS)'s Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (UK) (Grant No: 107768/Z/15/Z) and the UK government.

ACKNOWLEDGEMENTS: We sincerely thank the study communities in Korogocho and Viwandani for agreeing to participate in this study.



The Consortium for Advanced Research Training in Africa CARTA is an Africa-based, Africa-led initiative to rebuild and strengthen the capacity of African universities to locally produce well-trained and skilled researchers and scholars. The program addresses an important challenge facing the African academy i.e. the scarcity of a robust research and training infrastructure capable of offering the type of vibrant and sustained doctoral training necessary to attract, train and retain the continent's brightest minds. CARTA is structured to fast-track the career development of the next generation of academics; build communities of fellows and mentors; reduce their isolation and; provide them a nurturing environment for research. CARTA's broad development goal is to increase use of cross-institutional collaboration to build research capacity of doctoral students and strengthen university systems to lead and conduct research on critical areas around public and population health in the region. For more information on CARTA, visit: <http://cartafrica.org/>

Disclaimer

This brief represents the opinions of the authors, and is the product of professional research. It is not meant to represent the position or opinions of the CARTA consortium or its funders. Any errors are the fault of the authors.