

https://dspace.org/

The antimicrobial activity and essential oil composition of medicinal aromatic plants used in African traditional healing

Van Vuuren, Sandra Freda

2008-02-29T12:14:13Z
http://hdl.handle.net/10539/4505

Downloaded from a DSpace repository. https://dspace.org/

The Antimicrobial Activity and Essential Oil Composition of Medicinal Aromatic Plants Used in African Traditional Healing

Sandy van Vuuren



A thesis submitted to the Faculty of Health Sciences, University of the Witwatersrand, in fulfilment of the requirements for the Degree of Doctor of Philosophy

Johannesburg, 2007

Declaration

I, Sandy van Vuuren, declare that this thesis is my own work. It is being submitted for the Degree of Doctor of Philosophy, at the Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at this or any other University.

…….
Signature

……..
Date

Abstract

A study on the essential oil chemistry and microbiological activity of South African indigenous medicinal plants with the aim of establishing a scientific rationale for their anti-infective properties was undertaken. For the purpose of this study, nine medicinal aromatic plants were selected. The hydrodistilled essential oil was analyzed by gas chromatography combined with mass spectrometry (GC-MS) and various techniques were used to document the antimicrobial activity.

Disc diffusion studies on *Myrothamnus flabellifolius* indicated highest activities against the fungal test organisms *Cryptococcus neoformans* and *Aspergillus niger* with radial inhibition zones of 8 mm and 10 mm respectively. The highest antimicrobial activity noted in the minimum inhibitory concentration (MIC) assay was for *C. neoformans* (2 mg/mL). Time-kill studies demonstrated the death kinetic progression on *M. flabellifolius* essential oils where the killing rate was greatest for *Candida albicans*. *Osmitopsis asteriscoides*, a plant used traditionally for cuts and swellings showed highest disc diffusion antimicrobial efficacy against *Staphylococcus aureus* (3 mm) and *C. neoformans* (3 mm). The MIC study indicated highest susceptibilities (4 mg/mL) for *Escherichia coli* and *Staphylococcus epidermidis*. Death kinetics for the three organisms studied demonstrated that the killing rate was greatest for *C. albicans*. The role of the two major constituents (1,8-cineole and (-)-camphor) act synergistically to enhance antimicrobial activity. Disc diffusion assays undertaken on *Artemisia afra* showed highest activity against *Candida tropicalis* (5 mm). In the MIC assay the highest susceptibility was against *Serratia odorifera* (4 mg/mL). Time-kill assays on *Artemisia afra* showed a concentration dependent bactericidal activity, with evidence that the major constituents independently and in combination were not responsible for the overall activity of the plant. *Lippia javanica*, a plant used to treat coughs, colds and bronchitis, indicated highest susceptibility against the respiratory pathogen *Klebsiella pneumoniae* (5 mm) with the disc diffusion assay. The MIC assay indicated highest susceptibilities (4 mg/mL) against *C. neoformans* and *E. coli*. Death kinetic assays for three test organisms showed that the killing rate was the greatest for *K. pneumoniae*. The time-kill study for *L. javanica* in combination with *A. afra* demonstrated that the oils in combination act synergistically against *K. pneumoniae*. The antimicrobial activity of the essential oils and extracts were determined for *Helichrysum cymosum* subsp. *cymosum* where the extracts demonstrated at least a six times greater MIC efficacy than the essential oils. Using column

chromatography, the antimicrobially active compound was isolated from *H. cymosum* subsp. *cymosum* and identified as helihumulone. The traditional use of plants as a treatment for infectious diseases is not always restricted to a single part of the plant as was noted in the study on *Croton gratissimus* var. *subgratissimus*, where the leaf, bark and root extracts were investigated singularly and combined in various ratios to establish possible interaction. The MIC and fractional inhibitory concentration (FIC) results indicated variable efficacies for the plant combinations. The greatest synergistic profile was noted for *C. neoformans* in the leaf and root combination (MIC 0.4 mg/mL and FIC of 0.4). Further isobologram combination studies were thereafter conducted on varying ratios of leaf and root extracts, indicating greatest synergy for *Bacillus cereus, Enterococcus faecalis, C. albicans* and *C. neoformans*. While seasonal variation had very little impact on the MIC results obtained from *Heteropyxis natalensis*, the ratio of the two major compounds (1,8-cineol and limonene) fluctuated on a monthly basis. Moderate antimicrobial activity (3.0-16.0 mg/mL) was found for most pathogens with higher sensitivities for *C. neoformans*. The geographical variation of *H. natalensis* essential oil indicated similar profiles for Gauteng, Nelspruit and Waterberg samples. The Lagalametse sample, however, showed distinct variation both chemically and microbiologically where efficacy was higher than in all other samples. The impact of the enantiomeric configuration was investigated for limonene in combination with 1,8-cineole with (+/-)-limonene in combination with 1,8-cineole having the most significant synergistic ratios against *Pseudomonas aeruginosa.* The antimicrobial activities of the non-volatile and volatile fractions of *Tarchonanthus camphoratus* and *Plectranthus grandidentatus*, singularly and in combination demonstrated that the volatile constituents contribute to the total efficacy of the plant. Isobologram representation of the combination of various ratios of *T. camphoratus* and *P. grandidentatus* essential oil and non-volatile extracts devoid of essential oils present a predominant synergistic profile for all pathogens studied. A comparative study on five indigenous oils (*M. flabellifolius, O. asteriscoides, H. natalensis, A. afra* and *L. javanica*) was undertaken with five popular commercial oils (*Lavendula angustifolia, Thymus vulgaris, Melaleuca alternifolia*, *Mentha piperita* and *Rosmarinus officinalis*). The highest antimicrobial activity was noted for *Thymus vulgaris* in the MIC assay, followed by *M. flabellifolius, O. asteriscoides* and *M. alternifolia*. With the time-kill assay, *M. flabellifolius* showed the most rapid cidal effect against all three pathogens tested. The comparative evaluation of commercial essential oils with indigenous oils validated the use of South African aromatic plants for their anti-infective properties.

Publications arising from this study*

- AM Viljoen, MJ Klepser, E Ernst, D Keele, E Roling, **SF van Vuuren**, B Demirci, KHC Başer and B-E van Wyk. The composition and antimicrobial activity of the essential oil of the resurrection bush, *Myrothamnus flabellifolius*. South African Journal of Botany, 2002, 68, 100-105 **(abstract, pg 266)**.
- AM Viljoen, **SF van Vuuren,** B Demirci, KHC Başer and B-E van Wyk. *Osmitopsis asteriscoides* (Asteraceae) - The antimicrobial activity and composition of the essential oil of a Cape-Dutch remedy. Journal of Ethnopharmacology, 2003, 88, 137-143 **(abstract, pg 267)**.
- AM Viljoen, S Subramoney, **SF van Vuuren,** B-E van Wyk, KHC Başer and B Demirci. The composition, geographical variation and antimicrobial activity of *Lippia javanica* (Verbenaceae) leaf essential oils. Journal of Ethnopharmacology, 2005, 96, 271-277 **[recipient of best publication award, ICPPS conference, 2006]** **(abstract, pg 268)**.
- **SF van Vuuren**, AM Viljoen, RL van Zyl, FR van Heerden and KHC Başer. The antimicrobial, antimalarial and toxicity profiles of helihumulone, leaf essential oil and extracts of *Helichrysum cymosum* (L.) D. Don subsp. *cymosum*. South African Journal of Botany, 2006, 72, 287-290 **(abstract, pg 269)**.
- AM Viljoen, **SF van Vuuren**, T Gwebu, B Demirci and KHC Başer. The geographical variation and antimicrobial activity of African Wormwood (*Artemisia afra* Jacq.) essential oil. Journal of Essential Oil Research, Special Edition, 2006, 18, 19-25 **(abstract, pg 270)**.
- **SF van Vuuren** and AM Viljoen. A comparative investigation of the antimicrobial properties of indigenous South African aromatic plants with popular commercially available essential oils. Journal of Essential Oil Research, Special Edition, 2006, 18, 66-71 **(abstract, pg 271)**.
- **SF van Vuuren**, AM Viljoen, T Őzek, B Demirci and KHC Başer. Seasonal and geographical variation of *Heteropyxis natalensis* essential oil and the effect thereof on the antimicrobial activity. South African Journal of Botany, 2007, in press **(abstract, pg 272)**.

***Appendix B**

Conference proceedings relating to this thesis

- AM Viljoen, B-E van Wyk, KHC Başer, B Demirci and **SF van Vuuren**. The essential oil composition, medicinal uses and antimicrobial activity of the resurrection bush, *Myrothamnus flabellifolius* 31st International Symposium on Essential Oils, Hamburg, Germany, 2000. [poster]
- T Pelele, AM Viljoen, **SF van Vuuren,** B Demirci and KHC Başer. African wormwood – essential oil composition, geographical variation and antimicrobial properties of a coveted traditional herbal remedy. The 33rd International Symposium on Essential Oils, Lisbon, Portugal, 4-7 September, 2002. [poster]
- **SF van Vuuren**, AM Viljoen and ME Klepser. The antimicrobial activity of medicinal aromatic plants used in African traditional healing with special reference to method variability and essential oil composition. RAU Inter-University Post Graduate Symposium, 30 October, 2002. [**awarded a medal for best PhD podium presentation**]
- S Subramoney, **SF van Vuuren**, AM Viljoen, B Demirci and KHC Başer. Antimicrobial properties and geographical variation in essential oil composition of fever tea - *Lippia javanica* (Verbenceae). The 7th Conference of the International Society of Ethnopharmacology jointly with the South African Association of Botanists, University of Pretoria, 8-11 January, 2003.
- **SF van Vuuren,** AM Viljoen, E Ernst, B Demirci, T Özek and KHC Başer. The antimicrobial activity and death kinetics of the essential oil and chemical components of the South African endemic, *Osmitopsis asteriscoides*. The 7th Conference of the International Society of Ethnopharmacology jointly with the South African Association of Botanists, University of Pretoria, 8-11 January, 2003. **[awarded first prize for best ethnobotany poster]**
- LT Gwebu, **SF van Vuuren**, AM Viljoen, B Demirci and KHC Başer. The antimicrobial activity and essential oil chemistry of African wormwood (*Artemisia afra*). The South African Association of Botanists 30th Conference, Durban, 18-22 January, 2004. [podium presentation]
- **SF van Vuuren**, AM Viljoen, RL van Zyl and FR van Heerden. Biological activities of leaf essential oils and extracts of *Helichrysum cymosum* subsp. *cymosum*. The South African

Association of Botanists 30th Conference, Durban, 18-22 January, 2004. [**awarded first prize for best Ethnobotany poster**]

- **SF van Vuuren** and AM Viljoen. A comparative investigation of the antimicrobial properties of indigenous South African and commercially available essential oils. The 34th International Symposium on Essential Oils, Messina, Italy, 29 September – 2 October, 2004. [poster]
- **SF van Vuuren**, AM Viljoen, RL van Zyl, E Njenga and M Braithwaite. Some variables to be considered in pharmacognostic studies using South African medicinal plants as a demonstration model. Indigenous Plant Use Forum, Grahamstown, 27-30 June, 2005. [podium presentation]
- **SF van Vuuren** and AM Viljoen. Phyto-synergy: examples from indigenous aromatic medicinal plants used in antimicrobial therapy. University of Johannesburg Postgraduate Symposium, 2 November, 2005. [**awarded a medal for best PhD podium presentation**]
- **SF van Vuuren** and AM Viljoen. Exploring the Antimicrobial Activity of Indigenous Aromatic Plants. Agribusiness in Sustainable Natural African Plant Products. International Aromatic Plants and Essential Oil Mini-Symposium. Stellenbosch, 27 February - 1 March 2006. [podium presentation]
- **SF van Vuuren** and AM Viljoen. The biological activity of medicinal aromatic plants and their constituents. Indigenous Plant Use Forum, Botswana, 3-6 July, 2006. [podium presentation]
- **SF van Vuuren** and AM Viljoen. Aromatic plants and their constituents as a model to study phyto-synergy. The 37th International Symposium on Essential Oils, Grasse-Opio, France, 10-13 September, 2006. [podium presentation]

Publications submitted for review

- **SF van Vuuren** and AM Viljoen. *In vitro* evidence of phyto-synergy between two plants (*Lippia javanica* and *Artemisia afra*) and for plant part combinations (*Croton gratissimus* Burch. var. *subgratissimus*) in corroboration with their ethnobotanical use. Submitted to Journal of Ethnopharmacology.
- **SF van Vuuren** and AM Viljoen. The antimicrobial efficacy of limonene enantiomers and 1,8-cineole. Submitted to Flavour and Fragrance Journal.
- **SF van Vuuren,** AM Viljoen and KHC Başer *Tarchonanthus camphoratus and Plectranthus grandidentatus*; Antimicrobial activity and pharmacological interaction of the non-volatile and volatile fractions singularly and in combination. Submitted to South African Journal of Botany.

Dedication

To my husband Vernon for his continual support and encouragement,

and

my children Kim and Darrian. May this inspire them to continue learning.

Acknowledgements

First and foremost I would like to acknowledge and thank my supervisor Professor Alvaro Viljoen, who has played a pivotal role in my career as both a supervisor and mentor. As supervisor, his incessant enthusiasm, generous knowledge, guidance and support imparted to me have been invaluable. As mentor, he has been instrumental in establishing collaborative ties and creating a research niche area for me within the phytomedicinal field. For this I am extremely grateful.

Furthermore, I would like to express my gratitude and appreciation to the following people who have assisted me in various ways in the completion of this study:

- My sincerest thanks go to Prof. Başer, Dr Demirci and Dr Özek from the Department of Pharmacognosy, Faculty of Pharmacy, Anadolu University in Turkey, for assisting with the GC-MS analysis of the essential oils.
- Grants: University Research Committee, Medical Faculty Research Endowment Fund (WITS) and the National Research Foundation (Thuthuka) for financial assistance. Without this, the study would not have been possible.
- Plant collectors: Antonio De Castro for the collection of *O. asteriscoides* and selected *H. natalensis* samples and Jan Vlok for the collection of *H. cymosum* subsp. *cymosum* samples.
- The staff at the Johannesburg Botanical Garden is thanked for their generosity in permitting the collection of *H. natalensis* seasonal samples, and my special thanks to Andrew Hankey at the Walter Sisulu Botanical Garden whose consistent helpfulness in hosting my many collecting visits is greatly appreciated.
- The Department of Pharmacy and Pharmacology, University of Witwatersrand, for allowing me the time and capacity to follow my dream.
- Dr. Robyn van Zyl for her theoretical advice in the construction of the isobolograms. Furthermore, her valuable collaboration on the determination of toxicity profiles for all oils is greatly appreciated.
- Prof. Fanie van Heerden, for her valuable expertise in the NMR and MS analysis of *H. cymosum* subsp. *cymosum* samples.

- Robertet (France) for the generous provision of high quality commercial essential oils.
- Paul Steenkamp is gratefully acknowledged for assistance with the HPLC analyses for the bark, leaf and root extracts of C. *gratissimus* var. *subgratissimus*.
- Dr. Denise Lindsey for her guidance in confocal microscopy.
- The staff at the Witwatersrand Health Sciences Library for their assistance in literature acquisition.
- Photographic contribution of all laboratory procedures and plants were very patiently provided by Prof. Viljoen.
- To all my academic peers and fellow students within the botany field, for their inspiration, encouragement, motivation and hospitality in welcoming me into their world of plants.
- Last, but not least, my husband Vernon, for all his steady, unwavering support throughout this study.

Table of contents

	Page
Declaration……	ii
Abstract…….	iii
Publications arising from this study……	v
Conference proceedings relating to this thesis……..	vi
Publications submitted for review……..	viii
Dedication……..	ix
Acknowledgements……	x
Table of contents……..	xii
List of figures……	xix
List of tables……..	xxvii
Abbreviations……	xxxii

Chapter 1: Overview of the use of aromatic plants and their essential oils to treat microbial infections.

1.1	Introduction…….	1
1.2	A historical perspective …….	2
1.3	Present use of antimicrobials and the impact on disease……	4
1.4	Ethnomedicinal plants as antimicrobials……	5
1.5	Essential oils as antimicrobials …….	6
1.6	The South African perspective ……..	8
1.7	Thesis structure……...	10
1.7.1	Study objectives…….	12

Chapter 2: *Myrothamnus flabellifolius* Welw., antimicrobial efficacy determined by disc diffusion, minimum inhibitory concentration and time-kill methodology.

2.1 Introduction…… 13
2.2 Botanical description…….. 14
2.3 Distribution……. 14
2.4 Medicinal uses…… 14
2.5 Methods. …… 15
2.5.1 Chemical aspects…… 15
2.5.2 Antimicrobial aspects……. 17
2.6 Results and discussion…… 22
2.6.1 Essential oil chemistry…… 22
2.6.2 Antimicrobial activity……. 25
2.7 General conclusions…… 31

Chapter 3: *Osmitopsis asteriscoides* (P.J. Bergius) Less., the antimicrobial efficacy (disc diffusion, minimum inhibitory concentration, time-kill) and role of the major essential oil constituents.

3.1 Introduction……. 32
3.2 Botanical description…….. 32
3.3 Distribution……. 33
3.4 Medicinal uses…… 33
3.5 Methods…….. 34
3.5.1 Chemical aspects…… 34
3.5.2 Antimicrobial aspects …… 35
3.6 Results and discussion…… 37
3.6.1 Essential oil chemistry…… 37
3.6.2 Antimicrobial activity ……..... 39
3.7 General conclusions…… 47

Chapter 4: The antimicrobial activity of *Artemisia afra* Jacq. ex Willd essential oil and the role of the major constituents, singularly and in combination.

4.1 Introduction……. 48
4.2 Botanical description…….. 49
4.3 Distribution……. 49
4.4 Medicinal uses…… 50
4.5 Methods…….. 50
4.5.1 Chemical aspects…… 50
4.5.2 Antimicrobial aspects……. 51
4.6 Results and discussion…… 53
4.6.1 Essential oil chemistry…… 53
4.6.2 Antimicrobial activity …… 55
4.7 General conclusions…… 63

Chapter 5: *Lippia javanica* (Burm.f.) Spreng, the antimicrobial activity and an *in vitro* synergy study when combined with *Artemisia afra* to treat *K. pneumoniae* infections.

5.1 Introduction…… 65
5.2 Botanical description……. 66
5.3 Distribution…… 66
5.4 Medicinal uses…… 66
5.5 Methods…….. 67
5.5.1 Chemical aspects…… 67
5.5.2 Antimicrobial aspects……. 68
5.6 Results and discussion…… 68
5.6.1 Essential oil chemistry…… 68
5.6.2 Antimicrobial activity…… 71
5.7 General conclusions…… 78

Chapter 6: *Helichrysum cymosum* (L.) D. Don subsp. *cymosum*, antimicrobial activity of the essential oil, extract and isolated bioactive compound helihumulone.

6.1 Introduction……. 80
6.2 Botanical description…….. 80
6.3 Distribution……. 81
6.4 Medicinal uses…… 81
6.5 Methods…….. 82
6.5.1 Chemical aspects…… 82
6.5.1.1 Isolation of antimicrobial compound……. 87
6.5.2 Antimicrobial aspects……. 87
6.6 Results and discussion…… 87
6.6.1 Essential oil chemistry…… 87
6.6.2 Elucidation and identification of the isolated compound…… 90
6.6.3 Antimicrobial activity …… 91
6.7 General conclusions…… 93
Appendix: *Helichrysum cymosum* subsp. *cymosum* …….. 95

Chapter 7: *Croton gratissimus* Burch. var. *subgratissimus*, the antimicrobial activity of the essential oil and extracts, with evidence of synergy for plant part combinations.

7.1 Introduction…… 98
7.2 Botanical description…….. 98
7.3 Distribution……. 99
7.4 Medicinal uses…… 99
7.5 Methods…….. 100
7.5.1 Chemical aspects…… 100
7.5.2 Antimicrobial aspects……. 101
7.6 Results and discussion…… 103
7.6.1 Essential oil chemistry…… 103

7.6.2 The HPLC analysis of the leaf, bark and root extract……. 106
7.6.3 Antimicrobial activity..…… 109
7.4 General conclusions…… 119

Chapter 8: A seasonal and geographical study of *Heteropyxis natalensis* Harv. essential oil and effect of stereochemistry on antimicrobial activity.

8.1 Introduction……. 121
8.2 Botanical description…….. 122
8.3 Distribution …… 123
8.4 Medicinal uses…… 123
8.5 Methods…….. 123
8.5.1 Chemical aspects…… 123
8.5.2 Antimicrobial aspects…….. 126
8.6 Results and discussion…… 128
8.6.1 Essential oil chemistry…… 128
8.6.2 Antimicrobial activity …… 140
8.6.3 The enantiomeric influence on microbiological efficacy……. 145
8.7 General conclusions…… 152

Chapter 9: *Tarchonanthus camphoratus* L. and *Plectranthus grandidentatus* Gürke, antimicrobial activity and pharmacological interaction of the non-volatile and volatile fractions.

9.1 Introduction…… 154
9.2 Botanical description……. 156
9.3 Distribution…… 156
9.4 Medicinal uses…… 157
9.5 Methods…….. 158
9.5.1 Chemical aspects…… 158
9.5.2 Antimicrobial aspects……. 160
9.6 Results and discussion…… 162

9.6.1 Essential oil chemistry…… 162
9.6.2 Antimicrobial activity..…….. 168
9.7 General conclusions…… 180

Chapter 10: A comparative investigation of the antimicrobial properties of indigenous South African aromatic plants with popular commercially available essential oils.

10.1 Introduction…… 182
10.2 Botanical description and distribution…… 183
10.3 Medicinal uses…… 185
10.4 Methods……. 186
10.4.1 Chemical aspects…… 186
10.4.2 Antimicrobial aspects…….. 187
10.5 Results and discussion…… 187
10.5.1 Essential oil chemistry…… 187
10.5.2 Antimicrobial activity…… 190
10.6 General conclusions…… 195

Chapter 11: A review of the methods used to study the antimicrobial activity of aromatic plants.

11.1 Introduction…… 197
11.2 Plant collection and essential oil isolation……. 198
11.3 Equipment…….. 199
11.4 Chemical evaluation of essential oils……. 199
11.5 Effect of solvents, media selection and additions…….. 199
11.6 Reference standards…… 200
11.7 Pathogen selection…….. 200
11.7.1 The Gram-positive test organisms……. 200
11.7.2 The Gram-negative test organisms…… 202
11.7.3 The fungal test organisms……. 205

11.8 Inoculum……. 206
11.9 Disc diffusion assay…… 208
11.10 Minimum inhibitory concentration (MIC)……. 210
11.11 Bio-directed fractionation (bio-autographic assays)…….. 214
11.12 Time-kill analysis…… 214
11.13 Method variation……. 217
11.14 Validation of various methodologies…….. 218
11.15 Interactive combination studies…….. 219
11.16 Summary……..... 223

Chapter 12: General discussion, conclusion and further recommendations.

12.1 General discussion…… 225
12.2 Thesis summary…… 225
12.3 Future trends……. 234
12.4 Conclusion…… 238

References……. 240

Appendix A.

Raw MIC data (mg/mL) for isobolograms …… 262

Appendix B.

Publications arising from this study…… 265

List of figures

Page

Figure 1.1 and Figure 1.2 Egyptian hieroglyphic depictions of essential oils used for anointing and medicinal purposes. Figure 1.1 depicts an Egyptian with an incense burner preparing oils for inhalation. Figure 1.2 illustrates the administration of oils for the enhancement of body processes…….. 2

Figure 1.3 A schematic outlay of the study protocol for the antimicrobial investigation of aromatic plants…… 11

Figure 2.1 Aromatic foliage of *Myrothamnus flabellifolius*……. 14

Figure 2.2 *Myrothamnus flabellifolius* in habitat……. 14

Figure 2.3 The geographical distribution of *M. flabellifolius* in South Africa (SAN BI)…… 15

Figure 2.4 Clevenger apparatus used for distillation of plant material……. 16

Figure 2.5 A diagrammatic representation of the steps undertaken for each oil concentration carried out at all time intervals……... 21

Figure 2.6 Death kinetic method as detailed clockwise: aseptic addition of *M. flabellifolius* into culture media; incubation in shaking waterbath; layout of plates for dilution; plating out of successive dilutions……..... 22

Figure 2.7 Chemical structures for major compounds identified in the essential oil of *M. flabellifolius*……... 25

Figure 2.8 Time-kill expressed in Log_{10} reduction of *S. aureus* exposed to *M. flabellifolius* essential oil within the first six hours……..... 29

Figure 2.9 Time-kill expressed in Log_{10} reduction of *P. aeruginosa* exposed to *M. flabellifolius* essential oil over 24 hr……. 30

Figure 2.10 Time-kill expressed in Log_{10} reduction units of *C. albicans* exposed to *M. flabellifolius* essential oil within the first six hours……. 30

Figure 3.1 The leaves and flowers of *Osmitopsis asteriscoides*…….. 33

Figure 3.2 The geographical distribution of *Osmitopsis asteriscoides* in South Africa (SANBI)…… 33

Figure 3.3 Chemical structures for major compounds identified in the essential oil of *O. asteriscoides*……... 38

Figure 3.4 Death kinetic studies of *S. aureus* with exposure to the essential oil of *O. asteriscoides* as seen over a 24 hour period…… 41

Figure 3.5 Death kinetic studies of *P. aeruginosa* with exposure to the essential oil of *O. asteriscoides* as seen over a 24 hour period……... 42

Figure 3.6 Death kinetic studies of *C. albicans* with exposure to the essential oil of *O. asteriscoides* as seen over a 24 hour period…… 42

Figure 3.7 The confocal scanner laser images (2 µm) of the control (having no essential oil) and 0.5% *O. asteriscoides* over 24 hr……. 43

Figure 3.8 Death kinetic studies of *S. aureus* with exposure to the essential oil (0.5%) of *O. asteriscoides* over five hours…….. 44

Figure 3.9 Time-kill plot for *O. asteriscoides* essential oil (1%), 1,8-cineole, (+)-camphor, (-)-camphor, 1,8-cineole and (-)-camphor tested on *Candida albicans* (ATCC 10231)……..... 45

Figure 4.1 The leaves of *Artemisia afra*……... 49

Figure 4.2 The geographical distribution of *Artemisia afra* in southern Africa (SA NBI)……. 49

Figure 4.3 Chemical structures for major compounds identified in the essential oil of *A. afra*…….. 55

Figure 4.4 The death kinetics of *K. pneumoniae* on exposure to *A. afra* essential oil at concentrations 0.063%-0.75% over a 24 hr period…….. 58

Figure 4.5 The death kinetic study of *K. pneumoniae* on exposure to *A. afra* essential oil showing bactericidal activity within one hour…….. 58

Figure 4.6 The death kinetics of *C. neoformans* on exposure to *A. afra* essential oil at concentrations 0.063%-0.75% over a six hour period……. 58

Figure 4.7 The death kinetic representation for the various major compound combinations together with *A. afra* (0.5%) essential oil tested against *K. pneumoniae* ……. 61

Figure 4.8 The CFU's observed when performing the compound combination time-kill assay tested against *K. pneumoniae* …….. 62

Figure 5.1 The aerial leaves and white flower heads of *Lippia javanica*…… 66

Figure 5.2 The geographical distribution of *L. javanica* in South Africa (SANBI). 66

Figure 5.3 Chemical structures for the major compounds identified in the essential oil of *L. javanica*……... 70

Figure 5.4 Time-kill plot of *L. javanica* essential oil showing death kinetics of *K. pneumoniae* (NCTC 9633) represented over the first four hours of a 24 hr test period……. 74

Figure 5.5 Time-kill plot of *L. javanica* essential oil showing death kinetics of *C. neoformans* (ATCC 90112) represented over the first eight hours of a 24 hr test period…….. 74

Figure 5.6 Time-kill plot of *L. javanica* essential oil showing death kinetics of *B. cereus* (ATCC 11778) represented over four hours of a 24 hr test period…... 75

Figure 5.7 Time-kill plot of *L. javanica* and *A. afra* independently and in combination for 0.25% essential oil against *K. pneumoniae* (NCTC 9633) represented over 48 hr…….. 77

Figure 6.1 *Helichrysum cymosum* subsp. *cymosum* in habitat……. 81

Figure 6.2 Geographical distribution of *H. cymosum* subsp. *cymosum* in South Africa (SANBI)…… 81

Figure 6.3 Bio-autographic assay of the crude acetone extract of *H. cymosum* subsp. *cymosum* against *S. aureus*…… 84

Figure 6.4 Follow-up isolation of the biologically active compound by assay-guided fractionation using *S. aureus* as a test pathogen……... 84

Figure 6.5 Schematic representation of the bioactivity guided fractionation procedure…….. 86

Figure 6.6 Chemical structures for major compounds identified in the essential oil of *H. cymosum* subsp. *cymosum* 89

Figure 6.7 The HPLC profile of *H. cymosum* subsp. *cymosum* with the major peak identified as helihumulone……..... 90

Figure 6.8 The NMR carbon spectra (^{13}C) for helihumulone..... 96

Figure 6.9 The NMR carbon spectra (^{1}H) for helihumulone 97

Figure 7.1 The leaves and fruit of *Croton gratissimus* var. *subgratissimus*..... 99

Figure 7.2 The geographical distribution of *Croton gratissimus* var. *sub gratissimus* in southern Africa (SANBI)……... 99

Figure 7.3 The leaves, bark and root of *C. gratissimus* var. *subgratissimus*..... 100

Figure 7.4 Isobologram depicting antagonism, synergy and an additive effect where the relative ratio of Y is plotted to the relative ratio of X……. 103

Figure 7.5 Chemical structures for major compounds identified in the essential oil of *C. gratissimus* var. *subgratissimus* ……. 106

Figure 7.6 The *C. gratissimus* var. *subgratissimus* HPLC chromatogram for root (red), bark (green) and leaf (blue)……. 107

Figure 7.7 The MIC (mg/mL) of *C. gratissimus* var. *subgratissimus* essential oil (EO), and leaf, bark and root extracts……... 110

Figure 7.8 - Figure 7.17 The antimicrobial efficacy of the Gram-positive test organisms (*B. cereus, S. aureus, S. epidermidis, B. subtilis* and *E. faecalis* respectively) against the leaf, bark and root extract independently and in a 1:1 combination with the adjacent isobologram of leaf and root combination…….. 115

Figure 7.18 - Figure 7.23 The antimicrobial efficacy of the Gram-negative test organisms (*K. pneumoniae, P. aeruginosa* and *E. coli*) against the leaf, bark and root extract independently and in a 1:1 combination with the adjacent isobologram of leaf and root combination…… 116

Figure 7.24 - Figure 7.27 The antimicrobial efficacy of the yeasts (*C. albicans* and *C. neoformans*) against the leaf, bark and root extract independently and in a 1:1 combination with the adjacent isobologram of leaf and root combination…….. 118

Figure 8.1 and Figure 8.2 *Heteropyxis natalensis* (Figure 8.1) and leaves with small inconspicuous flowers (Figure 8.2)…… 122

Figure 8.3 The geographical distribution of *H. natalensis* in southern Africa (SANBI)…… 123

Figure 8.4 Chemical structures for major compounds identified in the essential oil of various *H. natalensis* samples……. 135

Figure 8.5 and Figure 8.6 The TLC chromatograms indicating similar profiles for plant A, sampled monthly (Figure 8.5) and Figure 8.6 representing similar profiles for three individual plants within a population……... 136

Figure 8.7 The monthly variation of limonene and 1,8-cineole recorded for three individual *H. natalensis* plants at the same locality……. 136

Figure 8.8 Dendrogram constructed from the essential oil data matrix (Table 8.4) for *H. natalensis*…… 139

Figure 8.9 The MIC plate for *E. faecalis* showing uniform microbial inhibition for selected *H. natalensis* samples…… 141

Figure 8.10 Dendrogram constructed from the microbial MIC data in Table 8.6 for *H. natalensis*…… 144

Figure 8.11 Isobologram plots for *S. aureus* (ATCC 12600) when exposed to the combination of (+)-limonene with 1,8-cineole, (-)-limonene with 1,8-cineole and a racemic mixture of limonene with 1,8-cineole. The 1:1 ratio as determined from the raw data is presented as a square…….. 149

Figure 8.12 Isobologram plots for *P. aeruginosa* (ATCC 9027) when exposed to the combination of (+)-limonene with 1,8-cineole, (-)-limonene with 1,8-cineole and a racemic mixture of limonene with 1,8-cineole……... 150

Figure 8.13 Isobologram plots for *C. neoformans* (ATCC 90112) when exposed to the combination of (+)-limonene with 1,8-cineole, (-)-limonene with 1,8-cineole and a racemic mixture of limonene with 1,8-cineole…….. 151

Figure 9.1 The leaves of *Tarchonanthus camphoratus* with woolly fruits……..... 156

Figure 9.2 The geographical distribution of *Tarchonanthus camphoratus* in Africa (SANBI)…… 156

Figure 9.3 The inflorescence of *Plectranthus grandidentatus*……..... 157

Figure 9.4 The geographical distribution of *Plectranthus grandidentatus* in South Africa (SANBI)…… 157

Figure 9.5 The soxhlet apparatus……. 160

Figure 9.6 Chemical structures for major compounds identified in the essential oil of *T. camphoratus*…… 165

Figure 9.7 Chemical structures for major compounds identified in the essential oil of *P. grandidentatus*……... 167

Figure 9.8 The comparative MIC (mg/L) for the volatile constituents (EO), non-volatile constituents (NV) and dried combined (DC) constituents for *T. camphoratus*……. 170

Figure 9.9 The antimicrobial efficacy of the Gram-positive test organisms (*S. aureus, B. cereus* and *E. faecalis* respectively) against varying concentrations of the non-volatile (NV) and volatile (EO) constituents of *T. camphoratus*…… 172

Figure 9.10 The antimicrobial efficacy of the Gram-negative test organisms (*E. coli, P. aeruginosa* and *K. pneumoniae* respectively) against varying concentrations of the non-volatile (NV) and volatile (EO) constituents of *T. camphoratus*……. 173

Figure 9.11 The antimicrobial efficacy of the yeasts (*C. albicans* and *C. neoformans* respectively) against varying concentrations of the non-volatile (NV) and volatile (EO) constituents of *T. camphoratus*……... 174

Figure 9.12 The comparative MIC (mg/L) for the volatile constituents (EO), non-volatile constituents (NV) and dried combined (DC) constituents for *P. grandidentatus*..... 177

Figure 9.13 The antimicrobial efficacy of the test organisms *S. aureus, K. pneumoniae* and *E. faecalis* against varying concentrations of the non-volatile (NV) and volatile (EO) constituents of *P. grandidentatus*…….. 178

Figure 10.1 *Lavendula angustifolia* (lavender)..... 184

Figure 10.2 *Thymus vulgaris* (thyme)..... 184

Figure 10.3 *Melaleuca alternifolia* (tea tree)..... 184

Figure 10.4 *Rosmarinus officinalis* (rosemary)……... 184

Figure 10.5 *Mentha piperita* (peppermint)…….. 184

Figure 10.6 Chemical structures for the major compounds identified in the commercial oils of *L. angustifolia, T. vulgaris, M. alternifolia, M. piperita* and *R. officinalis*……... 189

Figure 10.7 A comparative death kinetic study of *S. aureus,* exposed to 0.5% essential oil from five commercial and five indigenous plants over a 24 hr period…… 192

Figure 10.8 A comparative death kinetic study of *K. pneumoniae,* exposed to 0.5% essential oil from five commercial and five indigenous plants over a 24 hr period……... 193

Figure 10.9 A comparative death kinetic study of *C. albicans,* exposed to 0.5% essential oil from five commercial and five indigenous plants over a 24 hr period…… 194

Figure 11.1 and 11.2 Agar with seeded test organism where the diameter of the zone of inhibition is measured (Figure 11.1) and where the radius of the zone of inhibition is measured (Figure 11.2)…… 209

Figure 11.3 Death kinetic studies of *K. pneumoniae* exposed to the essential oil of *C. gratissimus* var. *subgratissimus* over 24 hr……. 215

Figure 11.4 Death kinetic studies of *B. cereus* exposed to the essential oil of *P. grandidentatus* over 24 hr…… 216

Figure 11.5 Death kinetic studies of *K. pneumoniae* exposed to the essential oil of *H. natalensis* over 24 hr…… 216

Figure 11.6 The isobole of the non-volatile constituents and volatile constituents combined in different ratios demonstrating the different profiles (antagonism, synergy and additive) dependent on the ratio studied (Table 11.6)…….. 223

Figure 12.1 Outline of thesis: The antimicrobial activity and essential oil composition of medicinal aromatic plants used in African traditional healing……... 225

List of tables

Page

Table 2.1 Plant collection data for *M. flabellifolius*……. 16

Table 2.2 Microbial organisms with corresponding reference numbers and acceptable MIC ranges for controls (ciprofloxacin for bacteria and amphotericin B for yeasts)…… 19

Table 2.3 The essential oil composition and integration percentage of *M. flabellifolius* as determined by GC-MS..... 23

Table 2.4 Zones of inhibition (measured in mm from disc edge to margin of culture growth) and MIC (mg/mL) of *M. flabellifolius* essential oil…… 26

Table 3.1 Plant collection data for *O. asteriscoides*……. 34

Table 3.2 Acceptable MIC ranges for ciprofloxacin antibiotic control…… 35

Table 3.3 Essential oil composition of *O. asteriscoides*..... 37

Table 3.4 Disc diffusion and MIC assays of *O. asteriscoides* essential oil……. 40

Table 3.5 The confocal cell viability (%) of *S. aureus* cells when exposed to 0.5% *O. asteriscoides* over 4 hr and 20 min……. 44

Table 3.6 The MIC of *O. asteriscoides* essential oil and major constituents against *C. albicans*……. 45

Table 4.1 Plant collection data for *A. afra*…… 50

Table 4.2 Acceptable MIC ranges for controls (ciprofloxacin for bacteria and amphotericin B for the yeast)……... 51

Table 4.3 Essential oil composition of *A. afra*……. 53

Table 4.4 Disc diffusion (mm from edge of disc) and MIC (mg/mL) of *A. afra* essential oil……... 56

Table 4.5 MIC (mg/mL) determination for the major compounds and essential oil of *A. afra*……... 60

Table 4.6 The CFU/agar plate obtained for the various major compound combinations together with *A. afra* essential oil tested against *K. pneumoniae*. 60

Table 5.1 Plant collection data for *L. javanica*..... 67
Table 5.2 Essential oil composition of *L. javanica*..... 68
Table 5.3 Disc diffusion (mm from edge of disc) and MIC (mg/mL) of *L. javanica* essential oil……. 72
Table 5.4 The CFU on agar plate obtained for 0.25% essential oils against *K. pneumoniae* of *L. javanica* and *A. afra* independently and in combination…… 76

Table 6.1 Plant collection data for *H. cymosum* subsp. *cymosum*……. 82
Table 6.2 Acceptable MIC ranges for ciprofloxacin control…… 87
Table 6.3 Essential oil composition of *H. cymosum* subsp. *cymosum*..... 88
Table 6.4 Zones of inhibition (measured in mm from edge of disc) of the acetone extract and essential oil of *H. cymosum* subsp. *cymosum* …..... 91
Table 6.5 MIC (mg/mL) for *H. cymosum* subsp. *cymosum* essential oil, acetone extract and helihumulone against ten pathogens…….. 92
Table 6.6 Correlation of NMR chemical shift data ^{13}C and ^{1}H for helihumulone with literature values obtained by Bohlmann *et al*. (1979)…….. 95

Table 7.1 Plant collection data for *C. gratissimus* var. *subgratissimus* ……... 101
Table 7.2 Essential oil composition of *C. gratissimus* var. *subgratissimus*……. 104
Table 7.3 The HPLC quantification of leaf, bark and root compounds of *C. gratissimus* var. *subgratissimus* ……..... 107
Table 7.4 The MIC values (mg/mL) for the leaf, root and bark *C. gratissimus* var. *subgratissimus* extracts, singularly and in combination with the FIC[*1]…… 111
Table 7. 5 Raw data for the isobologram construction of *B. cereus*…… 112

Table 8.1 Collection data for *H. natalensis* harvested in the Johannesburg Botanical Garden…….. 124

Table 8.2 Collection data for *H. natalensis* sourced from various localities…….. 125

Table 8.3 Raw data for the isobologram construction of *S. aureus*……. 127

Table 8.4 Essential oil composition as determined by GC-MS for *H. natalensis* samples (seasonal and geographical variation study)……... 129

Table 8.5 The mean MIC (mg/mL) for selected *H. natalensis* essential oils for January – December, 2004……... 142

Table 8.6 The mean MIC (mg/mL) for *H. natalensis* essential oils selected from different localities……. 143

Table 8.7 The mean MIC (mg/mL) for the major constituents limonene and 1,8-cineole independently, and in combination with FIC (in brackets), determined for 1:1 combinations……. 147

Table 9.1 Plant collection data for *T. camphoratus* and *P. grandidentatus*……. 158

Table 9.2 Raw data for the isobologram construction for the volatile (EO) constituent interaction with non-volatile constituents of *T. camphoratus* against *E. faecalis*……. 161

Table 9.3 Raw data for the isobologram construction for volatile (EO) constituent interaction with non-volatile constituents of *P. grandidentatus* against *S. aureus*……... 161

Table 9.4 Essential oil composition of *T. camphoratus*……... 163

Table 9.5 The essential oil composition of *P. grandidentatus* (Maistry, 2003)…… 166

Table 9.6 The mean MIC (mg/mL) for the essential oil (EO) and plants that have undergone soxhlet extraction (NV, FC, DC) for *T. camphoratus*[1] and *P. grandidentatus*[2]…… 169

Table 10.1 A summary of the major compounds and therapeutic use for the five indigenous and five commercial essential oils……. 189

Table 10.2 The MIC (mg/mL) of the five indigenous and five commercial oils……. 191

Table 11.1 The MIC (mg/mL) for oil samples when tested with a varying inoculum density ranging from 0.1-30 mL: 100 mL (24 hr culture: broth ratio)…….. 207

Table 11.2 The MIC (µg/mL) for controls when tested with a varying inoculum density ranging from 0.1-30 mL: 100 mL (24 hr culture: broth ratio)……. 208

Table 11.3 The mean MIC (mg/mL) for plant samples exposed to *K. pneumoniae* in microtitre plates that have been sealed in comparison to the uncovered microtitre plates…… 211

Table 11.4 The classification of the FIC index in accordance with the corresponding authors…….. 221

Table 11.5 The relative ratios of the volatile and non-volatile constituents of *T. camphoratus*……. 222

Table 11.6 The raw data for the volatile and non-volatile constituents of *T. camphoratus*……. 222

Table 12.1 The highest antimicrobial activities found for all plants in this study…….. 228

Table 12.2 Summary of the major oil constituents obtained from all plants in this study……..... 230

Table 12.3 The antimicrobial properties of 1,8-cineole as determined by corresponding authors..... 231

Table 12.4 The toxicity profiles for the eight essential oils studied (van Zyl, publication in preparation)…… 235

Table A1 Raw MIC data (mg/mL) for isobolograms (Chapter 7) where C. *gratissimus* roots and leaves were combined…… 262

Table A2 Raw MIC data (mg/mL) for isobolograms (Chapter 8) where (+), (-) and the racemic form of limonene was combined with 1,8-cineol……. 263

Table A3 Raw MIC data (mg/mL) for isobolograms (Chapter 9) where the volatile and non-volatile constituents for *T. camphoratus* were combined….. 264

Table A4 Raw MIC data (mg/mL) for isobolograms (Chapter 9) where the volatile and non-volatile constituents for *P. grandidentatus* were combined.. 264

Abbreviations

AACHRD- African Advisory Committee for Health Research and Development

ADCAV- Plant collection by Antonio De Castro and Prof. Alvaro Viljoen

ATCC- American Type Culture Collection

CONC- Concentration

CFU- Colony forming units

DC- Combined volatile and non-volatile constituents that have been prepared from dried plant material

DD- Disc diffusion

DMSO- Dimethyl sulfoxide

EO- Essential oil

EMRSA- Epidemic methicillin resistant *Staphylococcus aureus*

ESCMID- European Society of Clinical Microbiology and Infectious Diseases

EX- Extract

FC- Combined volatile and non-volatile constituents that have been prepared with fresh plant material

FIC- Fractional inhibitory concentration

FID- Flame ionization detector

g- Gram

GC- Gas chromatography

GC-MS- Gas chromatography combined with mass spectrometry

HPLC- High performance liquid chromatography

hr- Hour

IC_{50}- The half maximum efficacy

ICU- Intensive care unit

INT- *p*-Iodonitrotetrazolium violet

IU- International units

JHB BG- Johannesburg Botanical Garden

JV- Plant collection by Jan Vlok

MIC- Minimum inhibitory concentration

min- Minutes

mg- Milligram

mL- Millilitre

mm- Millimetre

MRSA- Methicillin resistant *Staphylococcus aureus*

NCCLS- National Committee for Clinical Laboratory Standards

NaCl- Sodium chloride

NCTC- National Culture Type Collection

NHLS- National Health Laboratory Services

NMR- Nuclear magnetic resonance spectroscopy

NV- Non-volatile constituents

PB- Plant collection by Priscilla Burgoyne

PDA- Photodiode array detector

$\mathbf{R_f}$- Distance travelled by spot from base line with respect to solvent front.

RRI- Relative retention index

SABS- South African Bureau of Standards

SANBI- South African National Biodiversity Institute

Spp- Species

STI- Sexually transmitted infections

SVV- Plant collection by Sandy van Vuuren

TMD- Thermabeam mass selective detector

TIC- Total ion chromatogram

TLC- Thin layer chromatography

tr- Trace

µL- Microlitre

UV-Ultra violet

VRE- Vancomycin-resistant *Enterococcus faecalis*

WSBG- Walter Sisulu Botanical Garden

WHO- World Health Organization.