

Tanaka DT, Grunstein MM. Substance P augments electrically-induced constriction of isolated rabbit tracheal smooth muscle. Am Rev Respir Dis 1985; 131 (Suppl): A281.

Tannenbaum GS, Martin JB. Evidence for an endogenous ultradian rhythm governing growth hormone secretion in the rat. Endocrinology 1976; 98: 562-570.

Tashkin DP, Conolly ME, Deutsch RI, Hui KK, Littner M, Scarpace P, Abrass I. Subsensitization of beta-adrenoceptors in airways and lymphocytes of healthy and asthmatic subjects. Am Rev Respir Dis 1982; 125: 185-193.

Tattersfield AE, Leaver DG, Pride NB. Effects of  $\beta$ -adrenergic blockade and stimulation on normal human airways. J Appl Physiol 1973; 35: 613-619.

Tharp GD. The role of glucocorticoids in exercise. Med Sci Sports 1975; 7: 6-11.

Tharp GD, Buuck RJ. Adrenal adaptation to chronic exercise. J Appl Physiol 1974; 37: 720-722.

Thomson NC, Daniel EE, Hargreave FE. Role of smooth muscle  $\alpha$ 1-receptors in nonspecific bronchial responsiveness in asthma. Am Rev Respir Dis 1982; 126: 521-525.

Tickner TR, Cramp DG, Foo AY, Johnson AJ, Bateman SM, Pidgeon J, Spiro SG, Clarke SW, Wills MR. Metabolic response to intravenous salbutamol therapy in acute asthma. Thorax 1977; 32: 182-184.

Tinkelman DG, Cavanaugh MJ, Cooper DM. Inhibition of exercise-induced bronchospasm by atropine. Am Rev Respir Dis 1976; 114: 87-94.

Titinchi S, Al Shammai M, Patel KR, Kerr JW, Clark B. Circadian variation in number and affinity of  $\beta_2$ -adrenoceptors in lymphocytes of asthmatic patients. Clin Sci 1984; 66: 323-328.

Tolone G, Bonasera L, Sajeve R. Hydrocortisone increases the responsiveness of mast cells to  $\beta$ -adrenergic agonists by an action distal to the  $\beta$ -adrenoreceptors. Br J Exp Pathol 1979; 60: 269-275.

Tomiyama A, Takayanagi I, Takagi K. Relaxation of intestinal smooth muscle and calcium movements. J Pharm Pharmacol 1973; 25: 65-68.

Tomori Z, Widdicombe JG. Muscular, bronchomotor and cardiovascular reflexes elicited by mechanical stimulation of the respiratory tract. J Physiol (Lond) 1969; 200: 25-49.

Townley RG, McGeady S, Bewtra A. The effect of beta adrenergic blockade on bronchial sensitivity to acetyl-beta-methacholine in normal and allergic rhinitis subjects. J Allergy Clin Immunol 1976; 57: 358-366.

Townley RG, Reeb R, Fitzgibbons T, Adolphson RL. The effect of corticosteroids on the beta-adrenergic receptors in bronchial smooth muscle. J Allergy 1970; 45: 118.

Undem BJ, Dick EC, Buckner CK. Inhibition by vasoactive intestinal peptide of antigen-induced histamine release from guinea-pig minced lung. Eur J Pharmacol 1983; 88: 247-250.

Vaccarezza JR. The suprarenal function in allergic asthma. Determination of the plasma and urinary 17-OH-Corticoids and of the urinary 17-Ketosteroids before and after ACTH Zn. Dis Chest 1961; 40: 121-127.

Van de Werve G, Hue L, Hers H-G. Hormonal and ionic control of the glycogenolytic cascade in rat liver. Biochem J 1977; 162: 135-142.

van Zwieten PA. The central action of antihypertensive drugs, mediated via central  $\alpha_1$ -receptors. J Pharm Pharmacol 1973; 25: 89-95.

Vaz R, Senior B, Morris M, Binkiewicz A. Adrenal effects of beclomethasone inhalation therapy in asthmatic children. J Pediatr 1982; 100: 660-663.

Venter JC. Muscarinic cholinergic receptor structure. Receptor size, membrane orientation, and absence of major phylogenetic structural diversity. J Biol Chem 1983; 258: 4842-4848.

Vermeire PA, Vanhoutte PM. Inhibitory effects of catecholamines in isolated canine bronchial smooth muscle. J Appl Physiol 1979; 46: 787-791.

Vestal RE, Wood AJ, Shand DG. Reduced  $\beta$ -adrenoceptor sensitivity in the elderly. Clin Pharmacol Ther 1979; 26: 181-186.

Vidruk EH, Hahn HL, Nadel JA, Sampson SR. Mechanisms by which histamine stimulates rapidly adapting receptors in dog lungs. J Appl Physiol 1977; 43: 397-402.

Vlasses PH, Ferguson RK, Koplin JR, Clementi RA, Green PJ. Adrenocortical function after chronic inhalation of fluocortinbutyl and beclomethasone dipropionate. Clin Pharmacol Ther 1981; 29: 643-649.

Wagner HR, Crutcher KA, Davies JN. Chronic estrogen treatment decreases  $\beta$ -adrenergic responses in rat cerebral cortex. Brain Res 1979; 171: 147-151.

Wagner HR, Davies JN. Decreased  $\beta$ -adrenergic responses in the female rat brain are eliminated by ovariectomy: correlation of [ $^3$ H]dihydroalprenolol binding and catecholamine stimulated cyclic AMP levels. Brain Res 1980; 201: 235-239.

Walden SM, Bleecker ER, Chahal K, Britt EJ, Mason P, Permutt S. Effect of alpha-adrenergic blockade on exercise-induced asthma and conditioned cold air. Am Rev Respir Dis 1984; 130: 357-362.

Ward MJ, Fentem PH, Roderick Smith WH, Davies D. Ipratropium bromide in acute asthma. Br Med J 1981; 282: 598-600.

Warren JB, Dalton N. A comparison of the bronchodilator and vasopressor effects of exercise levels of adrenaline in man. Clin Sci 1983; 64: 475-479.

Warren JB, Dalton N, Turner C, Clark TJH. Protective effect of circulating epinephrine within the physiologic range on the airway response to inhaled histamine in nonasthmatic subjects. J Allergy Clin Immunol 1984; 74: 683-686.

Warren JB, Keynes RJ, Brown MJ, Jenner DA, McNicol MW. Blunted sympathoadrenal response to exercise in asthmatic subjects. Br J Dis Chest 1982; 76: 147-150.

Wasserman MA. Bronchopulmonary responses to prostaglandin  $F_{2\alpha}$  histamine and acetylcholine in the dog. Eur J Pharmacol 1975; 32: 146-155.

Weiner N. Atropine, scopolamine and related antimuscarinic drugs. In: Goodman and Gilman's The Pharmacological Basis of Therapeutics. Gilman AG, Goodman LS, Gilman A, eds. MacMillan Publishing Co Inc., New York: 1980; 120-137.

Weiss SJ, McKinney JS, Putney JW Jr. Receptor-mediated net breakdown of phosphatidylinositol 4,5-bisphosphate in parotid acinar cells. Biochem J 1982; 206: 555-560.

Welborn TA, Fraser TR. The double antibody immunoassay of insulin. Diabetologia 1965; 1: 211-218.

Weller HH, van der Straeten M, Vermeulen A, Obie NGM. Hormonal pattern in bronchial asthma. Scand J Respir Dis 1968; 49:163-184.

Wheeler T, Watkins PJ. Cardiac denervation in diabetes. Br Med J 1973; 4: 584-586.

Whitsett JA, Darovec-Beckerman C, Adams K, Pollinger J, Needelman H. Thyroid dependent maturation of  $\beta$ -adrenergic receptors in the rat lung. Biochem Biophys Res Commun 1980; 97: 913-917.

Widdicombe J The neural reflexes in the airways. Eur J Respir Dis 1985; 68 (Suppl 144): 1-33.

Widdicombe JG. Action potentials in parasympathetic and sympathetic efferent fibres to the trachea and lungs of dogs and cats. J Physiol (Lond) 1966; 186: 56-88.

Widdicombe JG. Regulation of tracheobronchial smooth muscle. Physiol Rev 1963; 43: 1-37.

Widdicombe JG, Sterling GM. The autonomic nervous system and breathing. Arch Intern Med 1970; 126: 311-329.

Wieling W, van Brederode JFM, de Rijk LG, Borst C, Dunning AJ. Reflex control of heart rate in normal subjects in relation to age: A data base for cardiac vagal neuropathy. Diabetologia 1982; 22: 163-166.

Wilken-Jensen K. The effect of inhalations of beclomethasone dipropionate on plasma cortisol levels and growth hormone production. Postgrad Med J 1975; (Suppl 4): 31-32.

Wilkinson PC. Chemotaxis of phagocytic cells towards proteins: The effect of protein denaturation. In: The Reticuloendothelial System and Immune Phenomena. Di Luzio NR, Flemming K, eds. Plenum Press, New York, London: 1971; 59-70.

Williams H, Read GF, Verrier-Jones ER, Hughes IA. Effect of inhaled beclomethasone dipropionate on saliva cortisol concentrations. Arch Dis Child 1984; 59: 553-556.

Wilson NM, Barnes PJ, Vickers H, Silverman M. Hyperventilation-induced asthma: evidence for two mechanisms. Thorax 1982; 37: 657-662.

Woenne R, Kattan M, Orange RP, Levison H. Bronchial hyperreactivity to histamine and methacholine in asthmatic children after inhalation of SCH 1000 and chlorpheniramine maleate. J Allergy Clin Immunol 197; 62: 119-124.

Wood CL, Arnett CD, Clarke WR, Tsai BS, Lefkowitz RJ. Subclassification of alpha-adrenergic receptors by direct binding studies. Biochem Pharmacol 1979; 28: 1277-1282.

Wortsman J, Frank S, Cryer PE. Adrenomedullary response to maximal stress in humans. Am J Med 1984; 77: 779-784.

Wyatt R, Waschek J, Weinberger M, Sherman B. Effects of inhaled beclomethasone dipropionate and alternate-day prednisone on pituitary-adrenal function in children with chronic asthma. N Engl J Med 1978; 299: 1387-1392.

Yamada S, Yamamura HI, Roeske WR. Alterations in cardiac autonomic receptors following 6-hydroxydopamine treatment in rats. *Mol Pharmacol* 1980; 18: 185-192.

Vernault J-C, Leclercq R, Schandevyl W, Virasoro E, De Coster A, Copinschi G. The endocrinometabolic effects of beclomethasone dipropionate in asthmatic patients. *Chest* 1977; 71: 698-702.

Yu DYC, Galant SP, Gold WM. Inhibition of antigen-induced bronchoconstriction by atropine in asthmatic patients. *J Appl Physiol* 1972; 32: 823-828.

Zaid G, Beall GN. Bronchial response to beta-adrenergic blockade. *N Engl J Med* 1966; 275: 580-584.

Zechman F, Hall FG, Hull WE. Effects of graded resistance to tracheal air flow in man. *J Appl Physiol* 1957; 10: 356-362.

Zuskin E, Mitchell CA, Bouhuys A. Interaction between effects of beta blockade and cigarette smoke on airways. *J Appl Physiol* 1974; 36: 449-452.

## APPENDIX "A"

### METHODS OF BIOCHEMICAL ANALYSIS AND HORMONE ASSAY

#### 1. PLASMA GLUCOSE

Plasma glucose levels were measured either with the Beckman Glucose Analyser I (Beckman Instruments, Fullerton, California, USA) or with an automated state routine analyser (Astra 8, Beckman Instruments, Brea, California, USA).

#### 2. PLASMA POTASSIUM

Plasma potassium levels were measured either with a flame photometer (IL543, Instrumentation Laboratories, Milan, Italy) or with an automated state routine analyser (Astra 8, Beckman Instruments, Brea, California, USA).

#### 3. PLASMA GROWTH HORMONE ASSAY

##### Principle

The principle of the assay is based on the competition for a fixed and limited number of antibody sites between growth hormone labelled with iodine<sup>125</sup> and growth hormone contained in standards or in the specimens to be assayed.

Assay precision and sensitivity are increased by pre-incubating the unlabelled growth hormone in the specimens and standards with the antiserum before adding <sup>125</sup>I-labelled growth hormone to the reaction mixture. After incubation the amount of

labelled growth hormone bound to the antibody is inversely related to the amount of unlabelled growth hormone in the sample. The antibody-growth hormone complex is allowed to react with a second antibody, and after a further incubation period, distilled water is added to the mixture.

The tubes are centrifuged and the supernatant fluids aspirated. The radioactivity in each tube is measured with a gamma scintillation counter. A graph is drawn of the percentage of relative binding obtained for each standard (y-axis) against its hormone concentration (x-axis) using semilog paper. The amount of growth hormone (ng/ml) in each sample is read by interpolating its percentage binding on the standard curve.

Expected Values:    Adults      0-1 ng/ml.  
                         Children    0-10 ng/ml

Molecular Weight:    21500.

Conversion Factor to SI Units:     $\times 0.047 = \text{nmol/l}$ .

Sensitivity:    The assay has a lower limit of sensitivity of 0,2-0,4 ng/ml and a measurable range up to 20 ng/ml.

Specificity:    The percentage of cross-reaction measured at 50% binding shows the specificity of the growth hormone antiserum used in the radioimmunoassay kit to be 100% for growth hormone, 0,48% for prolactin

and absent up to high concentrations of various other hormones.

Precision: Each sample was analysed in duplicate and the mean of the two radioactivity counts used to calculate the final value. For each study, all samples were tested in a single assay to avoid inter-assay variability.

Laboratory control pooled plasma was included in the growth hormone assays to determine the intra-assay and inter-assay variability.

In the present studies, measurements of the laboratory control pooled plasma gave the following results:

|                               | Intra-assay<br>variability | Inter-assay<br>variability |
|-------------------------------|----------------------------|----------------------------|
| Mean (ng/ml):                 | 4.7                        | 4.4                        |
| Standard deviation:           | 0.16                       | 0.36                       |
| Coefficient of variation (%): | 3.4                        | 8.1                        |

Manufactured by: Serono Biodata, Milan, Italy.

Radioactivity: Approximately 1,5 uCi (55,5 k Bq) per kit.

Reference: Melani et al. 1968.

#### 4. PROLACTIN ASSAY

##### Principle

The principle of the assay is based on the competition for a fixed and limited number of antibody binding sites between prolactin labelled with iodine<sup>125</sup> and prolactin contained in standards or in the specimens to be assayed.

After incubation the amount of labelled prolactin bound to the antibody is inversely related to the amount of unlabelled prolactin present in the sample. The antibody-prolactin complex is allowed to react with a second antibody and a solution of polyethylene glycol.

Separation is effected by centrifugation and aspiration of the supernatant fluids. The radioactivity in each tube is measured with a gamma scintillation counter.

A graph is drawn of the percentage of relative binding obtained for each standard (y-axis) against its hormone concentration (x-axis) using semilog paper. The amount of prolactin (ng/ml) in each sample is read by interpolating its percentage binding on the standard curve.

|                         |       |            |
|-------------------------|-------|------------|
| <u>Expected Values:</u> | Women | 0-15 ng/ml |
|                         | Men   | 0- 9 ng/ml |

Molecular Weight: 23000

Conversion Factor to SI Units: X0.043 = nmol/l

Sensitivity: The assay has a lower limit of sensitivity of 2 ng/ml and a measurable range up to 150 ng/ml.

Specificity: The specificity of the antiserum used in the radioimmunoassay kit is such that high levels of various hormones result in measured levels of prolactin of less than 2ng/ml.

Precision: Each sample was analysed in duplicate and the mean of the 2 radioactivity counts used to calculate the final value. All samples were tested in a single assay to avoid inter-assay variability. Laboratory control pooled plasma was included in the prolactin assay to determine the intra-assay variability. In the present study, measurements of the laboratory control pooled plasma gave the following results:

Intra-assay variability

|                               |      |
|-------------------------------|------|
| Mean (ng/ml):                 | 10.3 |
| Standard deviation:           | 0.63 |
| Coefficient of variation (%): | 6.1  |

Manufactured by: Serono Biodata, Milan, Italy

Radioactivity: Approximately 1,7  $\mu\text{Ci}$  (63 k Bq) per kit.

Reference: Friesen et al. (1973).

## 5. INSULIN ASSAY

### Principle

The principle of the assay is based on the competition for a fixed and limited number of antibody sites between porcine insulin labelled with iodine<sup>125</sup> and insulin contained in standards or in the specimens to be assayed.

After incubation the amount of labelled insulin bound to the antibody is inversely related to the amount of unlabelled insulin present in the sample.

The bound and free fractions of plasma insulin are separated by the use of an immuno-precipitating reagent in which an excess of a second antibody is preprecipitated.

After a further incubation period, the tubes are centrifuged and the supernatant fluids aspirated. The radioactivity in each tube is measured with a gamma scintillation counter.

A graph is drawn of the percentage of relative binding obtained for each standard (y-axis) against its hormone concentration (x-axis) using semilog paper. The amount of insulin (uU/ml) in each sample is read by interpolating its percentage binding on the standard curve.

Expected Values: 5-25 uU/ml.

Molecular Weight: 5807.

Conversion Factor to SI Units: Report as mU/l (no multiplication required).

Sensitivity: The assay has a lower limit of sensitivity of 0.78 uU/ml and a measurable range up to 200 uU/ml.

Specificity: Because of the highly specific nature of the immunological reactions, most anti-insulin sera fail to combine equally strongly with insulin prepared from different species, with the result that the standard lines obtained with these different hormones are not superimposable. The anti-insulin serum used in the present study yields similar results with porcine and human insulin.

Precision: Each sample was analysed in duplicate and the mean of the 2 radioactivity counts used to calculate the final values. All samples were tested in a single assay to avoid inter-assay variability. Laboratory control pooled plasma was included in the insulin assay to determine the intra-assay variability and the inter-assay variability (between the present study and other studies).

In the present study, measurements of the laboratory control pooled plasma gave the following results:

|                               | Intra-assay<br>variability | Inter-assay<br>variability |
|-------------------------------|----------------------------|----------------------------|
| Mean (/uU/ml):                | 19.4                       | 19.8                       |
| Standard deviation:           | 1.3                        | 0.6                        |
| Coefficient of variation (%): | 6.7                        | 3.0                        |

Manufacturers: Wellcome Research Laboratories, Beckenham, England, and Behringwerke AG, Marburg, Germany.

Radioactivity: Approximately 377 kBq  $^{125}\text{I}$  per kit.

Reference: Welborn and Fraser (1965).

## 6. ASSAY OF NON-ESTERIFIED FATTY ACIDS

### Principle

The principle of the assay is based on a titrimetric technique in which the non-esterified fatty acids are selectively extracted and titrated with dilute alkali using an ultramicroburette while nitrogen gas is bubbled through the solution.

Reference standards of known concentrations are similarly extracted and titrated, and a graph is drawn of the concentrations of standard (y-axis) and the volume of titrant used (x-axis) using linear graph paper.

The amount of non-esterified fatty acids (uEq/l) in each sample is read by interpolating the amount of titrant used on the standard graph.

Expected Values:     Fasting 315-1210 uEq/l.

Sensitivity:     The assay has a lower limit of sensitivity of 150 uEq/l and a measurable range up to 2400 uEq/l.

Precision:     Each sample was analysed in duplicate and the arithmetic mean of the results used as the final value.

Laboratory control pooled plasma was included in every assay to determine the intra-assay and inter-assay variability.

In the present study, measurements of the laboratory control pooled plasma gave the following results:

|                               | Intra-assay<br>variability | Inter-assay<br>variability |
|-------------------------------|----------------------------|----------------------------|
| Mean ( $\mu$ Eq/l):           | 1015                       | 1020                       |
| Standard deviation:           | 52                         | 62                         |
| Coefficient of variation (%): | 5.1                        | 6.1                        |

Reference: Kelley (1965).

## 7. CORTISOL ASSAY

### Principle

The principle of the assay is based on the competition for a fixed and limited number of antibody binding sites between cortisol labelled with iodine<sup>125</sup> and cortisol contained in standards or in the specimens to be assayed.

After incubation, the amount of labelled cortisol bound to the antibody is inversely related to the amount of unlabelled cortisol present in the sample.

The bound and free fractions of plasma cortisol are separated by the use of antibody-coated tubes. The supernatant fluid is aspirated and the radioactivity in each tube is measured with a gamma scintillation counter.

A graph is drawn of the percentage of relative binding obtained for each standard (y-axis) against its hormone concentration (x-axis) using linear graph paper. The amount of cortisol (ng/ml) in each sample is read by interpolating its percentage binding on the standard curve.

Expected Values:      8 am = 60-230ng/ml  
                             4 pm = 10- 85 ng/ml

Molecular Weight:    363.5

Conversion Factor to SI Units:   x 2.75 = nmol/l

Sensitivity: The assay has a lower limit of sensitivity of 4ng/ml and a measurable range of up to 800ng/ml.

Specificity: The percentages of cross-reaction of the cortisol antiserum used in the radioimmunoassay kit is 100% for cortisol, 51% for prednisolone and less than 3% for various other glucocorticoid hormones (calculated according to Abraham et al. 1977).

Precision: Each sample was analysed in duplicate and the mean of the 2 radioactivity counts used to calculate the final value. All the samples were tested in a single assay to avoid inter-assay variability. Laboratory control pooled plasma was included in the cortisol assay to determine the intra-assay variability and the inter-assay variability (between the present study and other studies).

In the present study, measurements of the laboratory control pooled plasma gave the following results:

|                               | Intra-assay<br>variability | Inter-assay<br>variability |
|-------------------------------|----------------------------|----------------------------|
| Mean (ng/ml):                 | 82                         | 84                         |
| Standard deviation:           | 2.7                        | 2.9                        |
| Coefficient of variation (%): | 3.2                        | 3.5                        |

Manufactured by: Sorin Biomedica, Italy.

Radioactivity: Approximately 3,4 uCi (126 k Bq) per kit.

References: Abraham et al. (1977), Rolleri et al. (1976).

## 8. ADRENOCORTICOTROPIC HORMONE (ACTH) ASSAY

### Principle

The principle of the assay is based on the competition for a fixed and limited number of antibody sites between ACTH labelled with iodine<sup>125</sup> and ACTH contained in standards or in the specimens to be assayed.

After incubation, the amount of labelled ACTH bound to the antibody is inversely related to the amount of unlabelled ACTH present in the sample.

The bound and free fractions of plasma ACTH are separated by the use of an immuno-precipitating reagent in which an excess of a second antibody has been preprecipitated.

After a second incubation period, the tubes are centrifuged and the supernatant fluid aspirated. The radioactivity in each tube is measured with a gamma scintillation counter.

A graph is drawn of the percentage of relative binding obtained for each standard (y-axis) against its hormone concentration (x-axis) using semilog paper. The amount of ACTH (pg/ml) in each sample is read by interpolating its percentage binding on the standard curve.

Expected Values: The normal range for the hormone assay laboratory of the Department of Medicine of the University of the Witwatersrand is 50-150pg/ml.

Molecular Weight: 4500.

Conversion Factor to SI Units:  $\times 0.22 = \text{pmol/l}$ .

Sensitivity: The assay has a lower limit of sensitivity of 10 pg/ml and a measurable range up to 1000 pg/ml.

Specificity: The percentage of cross-reaction of the ACTH antiserum used in the radioimmunoassay kit is 100% for ACTH with negligible cross-reactivity for other pituitary hormones (calculated according to Abraham et al. 1977).

Precision: Each sample was analysed in duplicate and the mean of the 2 radioactivity counts used to calculate the final value. All samples were tested in a single assay to avoid inter-assay variability. Laboratory control pooled plasma was included in the ACTH assay to determine the intra-assay variability and the inter-assay variability (between the present study and other studies).

In the present study, measurement of the laboratory control pooled plasma gave the following results:

|                               | Intra-assay<br>variability | Inter-assay<br>variability |
|-------------------------------|----------------------------|----------------------------|
| Mean (pg/ml):                 | 99                         | 65                         |
| Standard deviation:           | 7.4                        | 5.4                        |
| Coefficient of variation (%): | 3.3                        | 8.4                        |

Manufactured by: Compagnie ORIS Industrie SA, France.

Radioactivity: Approximately 2 uCi (74 k Bq) per kit.

References: Lee et al. (1961); Berson and Yalow, (1968);  
Abraham et al. (1977).

## 9. PLASMA CATECHOLAMINE ASSAYS

### Principle

The principle of the assay is based on the catalytic conversion of norepinephrine and epinephrine to their corresponding meta  $^3\text{H}$ -methoxy-derivatives by the enzyme catechol-O-methyl-transferase in the presence of S-adenosyl-L-methionine- $[^3\text{H-methyl}]$ .

The resulting products are extracted and separated by thin layer chromatography. The isolated catecholamine derivatives,  $^3\text{H}$ -normetanephrine and  $^3\text{H}$ -metanephrine are converted by periodate oxidation, to  $^3\text{H}$ -vanillin and extracted.

The radioactivity attributable to each catecholamine hormone is measured with a beta scintillation counter.

A catecholamine standard of known concentration is added to a second aliquot of each sample, and is extracted and separated in the same way to provide an internal standard value. This is used to calculate the concentrations of the individual catecholamines.

|                         |                |               |
|-------------------------|----------------|---------------|
| <u>Expected Values:</u> | Norepinephrine | 200-300 pg/ml |
|                         | Epinephrine    | 20- 50 pg/ml  |

|                           |                |     |
|---------------------------|----------------|-----|
| <u>Molecular Weights:</u> | Norepinephrine | 169 |
|                           | Epinephrine    | 183 |

Conversion Factors to SI Units:

|                |                   |
|----------------|-------------------|
| Norepinephrine | x 0.0059 = nmol/l |
| Epinephrine    | x 0.0055 = nmol/l |

Sensitivity: The assay has a lower limit of sensitivity of 2-5 pg/ml for norepinephrine and epinephrine, with a measurable range up to 3000 pg/ml/.

Specificity: The extent of cross-over of one catecholamine hormone into the assay of another (including dopamine) is less than 0.25% with the exception of epinephrine cross-over to norepinephrine which is 1.3%.

Precision: Each sample was analysed in duplicate and the mean of the 2 radioactivity counts used to calculate the final value. A control plasma specimen, provided by the manufacturers, with artificially high levels of catecholamines was included in each catecholamine assay to assess the efficiency of the kit by obtaining results within the ranges specified by the manufacturers, and to determine the inter-assay variability.

In the present study, the measurements of the control plasma specimen gave the following results:

|                               | Inter-assay variability |                |
|-------------------------------|-------------------------|----------------|
|                               | Epinephrine             | Norepinephrine |
| Mean (pg/ml):                 | 671                     | 729            |
| Standard deviation:           | 71                      | 52             |
| Coefficient of variation (%): | 10.6                    | 7.1            |

Manufacturer: Upjohn Diagnostics, Kalamazoo, Michigan, USA.

Radioactivity: Approximately 500 uCi  $^3\text{H}$  per kit.

References: Cryer et al. (1974), Passon and Peuler, (1973).

## APPENDIX "B"

### STUDIES OF SERUM NEUTROPHIL CHEMOTACTIC ACTIVITY

#### 1. Preparation of Polymorphonuclear Leukocytes (PMNL's)

PMNL's were prepared from blood obtained from normal adult donors which was treated with preservative-free heparin (5 units/ml blood). The PMNL's were separated from the mononuclear leukocytes by centrifugation at 400g for 15 minutes of the heparinised blood on cushions of Ficoll (Pharmacia, Uppsala, Sweden) metrizoate. The resultant erythrocyte/PMNL fraction was sedimented with 3% gelatin for 30 minutes at 37°C to remove most of the erythrocytes. The PMNL-rich supernatant was centrifuged at 250g for 10 minutes and the residual erythrocytes in the cell pellet lysed by exposure to 0.85% ammonium chloride. The PMNL's were centrifuged, washed once and resuspended at a concentration of  $10^7$ /ml in Hank's balanced salt solution (HBSS, Grand Island Biological Co., New York, USA) - pH 7.4, supplemented with 0.1% bovine serum albumin (BSA).

#### 2(a) Polymorphonuclear Leukocyte Migration Assay

The leukotactic potential of each serum specimen was measured using modified Boyden Chambers (Wilkinson, 1971). Two hundred microlitres of PMNL suspension ( $6 \times 10^5$  PMNL) were placed in the upper compartment of the Boyden chamber and separated from the lower compartment containing 20% serum diluted in BSA-supplemented HBSS by a 5,um pore-size micropore filter (Sartorius

membrane filters, Göttingen, Federal Republic of Germany). After incubation at 37°C the filters were removed and processed and the number of cells that had migrated through the entire thickness of the filter were counted (4 fields/filter). The results were expressed as PMNL/microscope high-powered field. Corresponding controls for random migration in the absence of serum were included and subtracted from each result.

2(b) Activation of PMNL membrane-associated oxidative metabolism

The effects of each serum specimen on PMNL membrane-associated oxidative metabolism were measured by a myeloperoxidase-mediated iodination assay (Root and Stossel, 1974) as an alternative method of detection of serum-associated leukoattractants (Becker et al. 1974).  $10^6$  PMNL were incubated with 20% serum, 0.06  $\mu$ Ci of Na  $I^{125}$  (17 Ci/mmol, New England Nuclear Corp., Boston, Mass., USA) and 5mg BSA in a final volume of 1ml HBSS in 5ml polypropylene test tubes which were incubated on a rotator for 30 minutes at 37°C. The reactions were terminated and the protein precipitated by the addition of ice-cold 10% trichloroacetic acid. After washing, the  $I^{125}$  in the protein precipitates was measured in a solid state gamma counter. The results were expressed in fmoles  $I^{125}$ .

Each serum specimen was coded prior to storage and the code broken only once the results were available.

**Author** Kallenbach J m

**Name of thesis** The Autonomic Nervous Sytem and Bronchial Asthma 1987

***PUBLISHER:***

University of the Witwatersrand, Johannesburg

©2013

***LEGAL NOTICES:***

**Copyright Notice:** All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed, or otherwise published in any format, without the prior written permission of the copyright owner.

**Disclaimer and Terms of Use:** Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.

**Author** Kallenbach J m

**Name of thesis** The Autonomic Nervous Sytem and Bronchial Asthma 1987

***PUBLISHER:***

University of the Witwatersrand, Johannesburg

©2013

***LEGAL NOTICES:***

**Copyright Notice:** All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed, or otherwise published in any format, without the prior written permission of the copyright owner.

**Disclaimer and Terms of Use:** Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.