

LIST OF FIGURES

Figure Page

CHAPTER 1

1	Diagrammatic representation of the cyclical changes in the female sex hormones that characterise the various menstrual phases....	4
2	Summary of the effects of progesterone (A) and oestrogen (B) on carbohydrate metabolism.....	26
3	Summary of the proposed influence of oestrogen and progesterone on fat metabolism during submaximal exercise as supported by most published observations.....	38
4	Summary of gender variation in fat metabolism during exercise.....	39
5	Serum creatine kinase activity at rest and following 20 min of downhill running in eumenorrhoeic women who participated during one of three menstrual phases and in male subjects.....	50
6	Necessity of the venous-arterial sampling model for constant infusion tracer estimates of FFA Ra.....	59
7	Background carbon-13 enrichment in expired carbon dioxide samples at rest and during prolonged moderate intensity cycling exercise in a non fasted female subject, with corresponding RER values at each time interval.....	64
8	Diagrammatic representation of Sidossis et al. (1995b) Figure 5, presenting the rate of decay of the carbon label in plasma palmitate	

and in expired carbon dioxide after stopping the infusion of a carbon-labelled palmitate, carbon-labelled sodium bicarbonate and carbon-labelled acetate tracer during exercise.....	65
9 The main possible sites of carbon label fixation in metabolic intermediate reactions or points of recovery as carbon dioxide following entry of the carbon label into the tricarboxylic acid cycle.....	67
10 Relationship between the acetate correction factor and energy expenditure during exercise as described by Sidossis et al. (1995a).....	71

CHAPTER 2

1 The respiratory exchange ratio at rest and during 90 min of submaximal exercise during the early follicular, late follicular and mid-luteal phases of the menstrual cycle.....	85
--	----

CHAPTER 3

1 (a) The relationship between the change in glycerol concentration over early follicular (EF) phase values and log oestrogen in the second menstrual phase. (b) Plasma glycerol concentration over time during exercise in the EF and mid-luteal (ML) phase.....	104
2 (a) Relationship of the change in average FFA concentration from EF phase and log oestrogen (E). (b) Plasma free fatty acid (FFA) concentration over time during exercise in the EF and mid-luteal (ML) phase.....	105
3 The relationship between the change in (a) Ra, (b) Rd and (c) MCR from EF phase values and log of the ratio of oestrogen to progesterone (E/P).....	106

4	Observational comparison of the change in palmitate and total free fatty acid (FFA) (a) rate of appearance (Ra), (b) rate of disappearance (Rd) and (c) metabolic clearance rate (MCR) over EF phase values.....	107
5	(a) Epinephrine and (b) norepinephrine concentration over time during exercise in the early follicular (EF) and mid-luteal (ML) phase and the relationship between norepinephrine concentration and (c) log oestrogen or (d) log progesterone.....	109
6	(a) Fractional contribution of carbohydrate (CHO), fat and protein to total energy expenditure (TEE). The relationship between (b) the change in the percentage of CHO contribution to TEE and the increase in progesterone (P-fold) from the EF phase and (c) the change in the percentage of protein contribution to TEE and the increase in oestrogen (E-fold) from the EF phase.....	110
7	The relationship between total carbohydrate (CHO) oxidation and log progesterone (P) across the menstrual cycle.....	111
8	The relationship between the change in total protein utilised from the EF phase and (a) log oestrogen (E), (b) the increase in oestrogen over EF phase levels (E-fold), and (c) the log oestrogen to progesterone ratio....	112

CHAPTER 4

1	Carbon-13 enrichment in expired carbon dioxide during continuous infusion of either $K^+[1-^{13}C]$ palmitate bound to HP- β -CD or $K^+[1-^{13}C]$ palmitate bound to human serum albumin during submaximal exercise.....	135
---	--	-----

CHAPTER 5

- 1 Comparison of the following ventilatory parameters: minute ventilation; respiratory rate; tidal volume; and oxygen consumption, at rest and during prolonged submaximal exercise between either the early follicular (EF) and late follicular (LF) or EF and mid-luteal (ML) phases..... 146
- 2 Significant correlations identified between the change in minute ventilation from the early follicular (EF) to mid-luteal (ML) phase during submaximal exercise and the magnitude of increase in oestrogen concentration from the EF to ML phase (E-fold) (a); logarithmic function of oestrogen concentration (b); and logarithmic function of progesterone concentration (c) in the ML phase..... 147
- 3 Significant correlations identified between the change in respiratory rate from the early follicular (EF) to mid-luteal (ML) phase during submaximal exercise and the logarithmic function of progesterone concentration (a) and the logarithmic function of the oestrogen to progesterone (E/P) ratio (b) in the ML phase..... 147

CHAPTER 6

- 1 Time to complete the a) 15 km and b) 30 km cycling time trial during the early follicular, late follicular and mid-luteal phase, in the untrained and trained group, respectively..... 155
- 2 Combined trained and untrained cycling time trial finishing times during each menstrual phase..... 155