

Combined explanation of LHC multilepton, diphoton, and top-quark excesses

Guglielmo Coloretti^{1,2,*}, Andreas Crivellin^{1,2,†} and Bruce Mellado^{3,4,‡}

¹*Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland*

²*Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland*

³*School of Physics and Institute for Collider Particle Physics, University of the Witwatersrand, Johannesburg, Wits 2050, South Africa*

⁴*iThemba LABS, National Research Foundation, P.O. Box 722, Somerset West 7129, South Africa*



(Received 11 January 2024; accepted 7 September 2024; published 1 October 2024)

In this article, we propose the Δ 2HDMS as a combined explanation of several excesses and anomalies observed at the LHC. Especially, $t\bar{t}$ differential distributions point towards the associated production of new electroweak scale Higgs bosons decaying into bottom quarks and W bosons ($> 5\sigma$) with masses consistent with the diphoton excesses at ≈ 95 GeV and ≈ 151.5 GeV (3.8σ and 4.9σ , respectively). Furthermore, CMS found indications for resonant $t\bar{t}$ production at ≈ 400 GeV (3.5σ) and both ATLAS and CMS reported elevated four-top and $t\bar{t}W$ cross sections. The Δ 2HDMS is obtained by supplementing the SM Higgs (H_2) with a second scalar doublet (H_1), real scalar singlet (S), and a Higgs triplet with zero hypercharge (Δ). We fix the masses of the neutral tripletlike and the singletlike scalars by the diphoton excesses, i.e., $m_{\Delta^0} = 151.5$ GeV and $m_S = 95$ GeV, respectively. H , the CP -even component of H_1 , is produced via gluon fusion from a top-loop and decays dominantly to $S + \Delta^0$ whose subsequent decays to W^+W^- and $b\bar{b}$ explain the differential $t\bar{t}$ distributions for $\sigma(pp \rightarrow H \rightarrow S\Delta^0) \approx 5$ pb. Choosing the top-Yukawa of H_1 accordingly, the CP -odd Higgs boson A turns out to have the right production cross section to account for the resonant $t\bar{t}$ excess at 400 GeV, while the top-associated production of H and A results in new physics pollution of Standard Model $t\bar{t}W$ and $t\bar{t}t\bar{t}$ cross sections, as preferred by the data.

DOI: [10.1103/PhysRevD.110.073001](https://doi.org/10.1103/PhysRevD.110.073001)

I. INTRODUCTION

The Standard Model (SM) of particle physics describes the known fundamental constituents of matter and their interactions (excluding gravity) at subatomic scales. It has been extensively tested and verified by a plethora of measurements [1]. However, the minimality of the SM Higgs sector [2–5] remains unexplained: While we know that the observed 125 GeV boson [6,7] has properties and decay rates compatible with the SM expectations [8,9], no fundamental principle excludes the existence of additional scalar bosons.

In fact, resonant diphoton searches show intriguing indications for new Higgs bosons at the electroweak scale, both in the sidebands of SM Higgs analyses and in

dedicated searches at 95 [10–14] and 151.5 GeV, with global significances of 3.8σ and 4.9σ [15,16], respectively. While the production mechanism for the 95 GeV scalar candidate is currently unknown, data imply that the 151.5 GeV boson is mainly produced in association, in particular with missing energy [17], but also leptons and (b) jets [18].

The mechanism of associated production provides the connection to the “LHC multilepton anomalies” [19–24]. These are processes involving multiple leptons and missing energy, with and without (b) jets, with SM signatures such as W^+W^- , WWW , Wh , tW , $t\bar{t}$, $t\bar{t}W$, and $t\bar{t}t\bar{t}$ (see Refs. [24,25] for a review) where deviations from the SM expectations have been observed. In particular, SM Higgs to W^+W^- analyses [26,27] prefer a new physics (NP) contribution, which is compatible with a ≈ 150 GeV scalar that is produced in association with missing energy [20,28]. Furthermore, the latest ATLAS analysis of the $t\bar{t}$ differential cross sections [29] prefers a NP contribution from the decay of a heavy new scalar into two lighter ones, which subsequently decay into W^+W^- and $b\bar{b}$ (see Fig. 1), over the SM hypothesis by at least 5.8σ [30]. Again, the masses are compatible with the hints for narrow resonances at 95 and 151.5 GeV which decay dominantly to $b\bar{b}$ and

*Contact author: guglielmo.coloretti@physik.uzh.ch

†Contact author: andreas.crivellin@cern.ch

‡Contact author: bmellado@mail.cern.ch

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.




FIG. 1. Feynman diagrams showing the leading SM contribution (left) to top pair production and decay as well as the NP effect mimicking this process (right).

W^+W^- , respectively. Since there is no indication for a 151.5 GeV boson decaying to ZZ , this suggests that it is the neutral component of a real $SU(2)_L$ triplet.

In this article, we propose a novel UV complete model that gives rise to these signatures and explains the $t\bar{t}$ differential distributions: the Δ 2HDMS. It contains two scalar doublets, a real singlet, and a real triplet Higgs. We will show that it predicts the experimentally preferred $\gamma\gamma$ signal strengths for the 95 and the 151.5 GeV Higgs bosons and is compatible with the suggested signal in W^+W^- (with a full jet veto) if one includes decays of the 95 GeV state to invisible. Furthermore, it can account for the resonant $t\bar{t}$ excess at ≈ 400 GeV (3.5σ locally) [31] as well as the elevated four top, $t\bar{t}W$ and $t\bar{t}Z$ cross sections and predicts a positive shift in the W mass as suggested by the global fit.

II. MODEL

As outlined in the introduction, we propose the Δ 2HDMS model. It contains two Higgs doublets with hypercharge $Y = +1/2$ (H_1 and H_2) and a real singlet (S). This means it corresponds to a 2HDMS [19,32–49], plus a real $SU(2)_L$ triplet (Δ) [50–57].¹ As in the 2HDMS, one can employ two Z_2 symmetries, one under which S and H_2 are odd, and one under which H_1 and Δ are odd, while all other fields are even. This removes trilinear coupling and thus most potential sources of CP violation.² Furthermore, this allows only one term which contains an odd number of any field³:

$$\mathcal{L}_{SH_1\Delta H_2} = -\lambda SH_1^\dagger \Delta H_2 + \text{H.c.} \quad (1)$$

¹A model with the same field content has been studied in Ref. [58] where it was shown that it can lead to electroweak baryogenesis. Supersymmetric versions of the model were considered in Refs. [59,60]. Reference [58] finds that electroweak baryogenesis can be achieved in a wide range of parameters, including the one studied here. Note that our LHC analysis is fully complementary to the one of Ref. [58]: Ref. [58] proposed the two-step mechanism for electroweak baryogenesis while we consider in detail the LHC phenomenology in light of the anomalies.

²Alternatively, one can employ a $U(1)'$ symmetry [61,62].

³The terms $\mu H_1 H_2^\dagger$ softly breaks the Z_2 symmetries but is usually included for phenomenological reasons.

TABLE I. Fields and their respective components in the limit of vanishing mixing angles. h corresponds to the SM Higgs and G^0 and $G^\pm$ to the Goldstone bosons. Note that, following the common 2HDM conventions [65], the SM Higgs boson h is contained within H_2 while the heavy doublet Higgs bosons H and A originate from H_1 .

	S	H_1	H_2	Δ
$SU(2)_L$	1	2	2	3
$U(1)_Y$	0	1/2	1/2	0
Z_2^1	+	−	+	−
Z_2^2	−	+	−	+
Components	S	$H, A, H^\pm$	$h, G^0, G^\pm$	$\Delta^0, \Delta^\pm$

After electroweak symmetry breaking, in the approximation $v_\Delta, v_S, v_1 \ll v_2$ (where v_X is the vacuum expectation value of the field X), H_1 , H_2 , S , and Δ contain the mass eigenstates as given in Table I. Note that the term in Eq. (1) does not lead to mixing among the scalars in this limit, but it gives rise to the trilinear interactions $\mathcal{L}_{\phi^3} = -\frac{v}{4}\lambda\Delta^0 HS + \frac{v}{2}\lambda\Delta^+ H^- S + \text{H.c.}$ (assuming λ to be real and $v = \sqrt{v_1^2 + v_2^2} \approx v_2 \approx \sqrt{2}174$ GeV) where Δ decomposed according to the conventions of Refs. [63,64]. The former term leads to the desired decay of H into $S\Delta^0$ needed to obtain the associated production mechanism.

For the Yukawa Lagrangian, we take the right-handed up-type and down quarks to be odd under Z_2^2 . Therefore, H_2 couples to all fermions, i.e., the Yukawa sector is the one of a type-I 2HDM. However, we also assume that the Z_2 symmetries of the Yukawa Lagrangian are broken in the top sector by the term

$$-\mathcal{L}_Y^t = \mu_t \sqrt{2} \frac{m_t}{v} \bar{Q}_3 \tilde{H}_1 u_3. \quad (2)$$

We defined this term such that μ_t is the relative coupling strength compared to the SM Higgs.⁴ Therefore, H and A can be produced from this coupling via gluon fusion induced by a top-quark loop. Furthermore, this coupling will lead to the top-associated production of H and A (see Fig. 2) as well as the tb associated production of $H^\pm$. Finally, we add neutral fermions χ (singlets under the SM gauge group) with the Yukawa coupling $Y_\chi S \bar{\chi} \chi$ to get a nonzero decay rate of S to invisible.

III. PHENOMENOLOGY

Here we study the phenomenology of our model in the limit of negligibly small mixing angles, i.e.,

⁴Note that if one defines this coupling in the down basis (i.e., Cabibbo Kobayashi Maskawa rotations appear associated with left-handed up quarks), the charged Higgs does not generate flavor violation in the down sector.




FIG. 2. Feynman diagrams showing the production of A via gluon fusion (left) and $t\bar{t}$ associated production (right) and its decays to top quarks.

$v_S, v_\Delta, v_1 \ll v_2$.⁵ Note that v_Δ is naturally small due to the additional symmetry of the potential in case it goes to zero. Furthermore, v_Δ leads to a positive-definite shift in the W mass [64,66–73]. In fact, $v_\Delta \approx 2$ GeV is preferred by the current global fit [74–76] (driven by the CDF-II result [77]).⁶

In this limit, the only possible unsuppressed decays (i.e., tree-level decays via order one couplings) of H (A) are $t\bar{t}$ and $S\Delta^0$ ($t\bar{t}$, ZH and $W^\pm H^\mp$) while $H^\pm$ decays dominantly to $t\bar{b}$ and possibly to $\Delta^\pm S$.⁷ In the following, we will assume $m_S < m_{\Delta^0} \ll m_H < m_A \approx m_{H^\pm}$ with m_H being below the top threshold. Note that a small value of v_Δ is preferred by the average [74,75,79] of the W mass measurements [77,77,80–82] (see Refs. [64,68–73,83–87]). Therefore, Δ^0 and $\Delta^\pm$ are quasidegenerate and being tripletlike, decay dominantly to W^+W^- and WZ , respectively. Since S is singletlike, it decays mainly ($\approx 86\%$) to $b\bar{b}$ and has $\text{Br}(S \rightarrow \gamma\gamma) = 1.4 \times 10^{-3}$ [88–103].

The main production mechanisms of H and A are gluon fusion and $t\bar{t}$ associated production, $H^\pm$ is produced in $t\bar{b}$ associated production while S and Δ^0 ($\Delta^\pm$) are produced from the decay of H ($H^\pm$). We fix m_S and m_{Δ^0} (and therefore $m_{\Delta^\pm}$) to the values suggested by the indications for narrow resonances, i.e., 95 and 151.5 GeV, respectively.

A. $t\bar{t}$ differential distributions

Next, we consider the production of H via top-quark-induced gluon fusion [88,104–113] and its subsequent

decay to $\Delta^0 S$. Note that this is the only unsuppressed decay of H if its mass is below the $t\bar{t}$ threshold. With $S \rightarrow b\bar{b}$ and $\Delta^0 \rightarrow W^+W^-$, the resulting final state has the same signature as top-quark pair production and decay in the SM (see Fig. 1). In fact, as shown in Ref. [30], the tensions between the measured $t\bar{t}$ (normalized) differential cross sections observed by ATLAS [29] and the corresponding SM predictions (which are compatible with the earlier CMS results [114]),⁸ can be removed if this NP process is added to the SM.⁹ Using the most conservative estimate for the NP effect, i.e., the SM simulation among the set performed by ATLAS that is in the least tension with data,¹⁰ a 5.8σ preference for NP with a best fit value of

$$\sigma(pp \rightarrow H \rightarrow \Delta^0 S \rightarrow W^+W^-b\bar{b}) \approx 5 \text{ pb}, \quad (3)$$

is found. Note that this result is approximately independent of m_H for $250 \text{ GeV} < m_H < 300 \text{ GeV}$. This translates to the preferred region (blue bands) for μ_t in Fig. 3.

B. Resonant $t\bar{t}$, $t\bar{t}t\bar{t}$, and $t\bar{b}H^\pm \rightarrow t\bar{b}t\bar{t}$

Because the CP -odd scalar A is (mainly) contained in H_1 together with H , they both have the same coupling strength to top quarks. However, since A couples axially, its production cross section from gluon fusion is ≈ 1.5 higher than the one of H for equal masses (see, e.g., Refs. [116,117]). If CP is conserved, as we assume, then A can neither decay to a pair of gauge bosons nor into two lighter Higgs bosons such that

$$\begin{aligned} \Gamma(A \rightarrow t\bar{t}) &= \frac{3G_F m_A m_t^2 \mu_t^2}{4\sqrt{2}\pi} \sqrt{1 - 4\frac{m_t^2}{m_A^2}}, \\ \Gamma(A \rightarrow HZ) &= \frac{G_F m_A^3}{8\sqrt{2}\pi} \lambda^{3/2}(m_H^2, m_Z^2; m_A^2), \\ \Gamma(A \rightarrow H^\pm W^\mp) &= \frac{G_F m_A^3}{4\sqrt{2}\pi} \lambda^{3/2}(m_{H^\pm}^2, m_{W^\mp}^2; m_A^2), \end{aligned} \quad (4)$$

⁵Note that while Δ^0 has a very small mixing with the SM Higgs boson due to the Z_2 symmetries, S can in principle have a bigger mixing with h and thus relevant gluon fusion and Z Strahlung production.

⁶To be more specific, taking into account only the real Higgs triplet (i.e., disregarding the usually small effect of the mass splitting of the doublet components), the global fit [74–76] including the CDF-II measurement [77] prefers $v_\Delta = 3.4 \pm 1.0$ GeV, while excluding the CDF-II measurement one finds $v_\Delta = 2.3 \pm 1.7$ GeV.

⁷For a dominant decay $H \rightarrow S\Delta^0$, a coupling λ of order one is necessary. However, note that also the Higgs self-coupling in the SM is of order one and that this does not cause any theoretical problem related to perturbativity, unitarity, etc. Furthermore, the mass splitting among the components of the doublet is small enough to be consistent with perturbativity and vacuum stability as well [78].

⁸These findings are also compatible with the latest differential $t\bar{t}$ results of CMS [115]. However, since we are performing a Monte Carlo–based analysis, including the new CMS result would not have a significant impact because the SM prediction is the limiting factor and CMS performs a less detailed analysis than ATLAS in this respect. Furthermore, note that the $t\bar{t}$ differential distributions only provide the normalization of the total NP effect and a smaller effect would even be welcomed by data.

⁹Note that while in Ref. [30] we focus on the $m^{e\mu}$ and $|\Delta\phi^{e\mu}|$ distributions, we checked that the other (normalized) distributions that are in reasonable agreement with the SM predictions are only marginally affected by the NP contribution (see Fig. 4 of Ref. [30]).

¹⁰This is the Powheg+Pythia8 (rew) simulation, including the reweighting of the top p_T by using the NLO predictions (see Ref. [29] for details).




FIG. 3. Constraints and preferred regions in the m_H - μ_t plane from $t\bar{t}$ differential distributions, resonant $t\bar{t}$ production and $t\bar{t}A \rightarrow t\bar{t}t\bar{t}$ (see main text for details).

with

$$\lambda(x, y; z) = \left(1 - \frac{x}{z} - \frac{y}{z}\right)^2 - 4 \frac{xy}{z^2} \quad (5)$$

are the dominant decay modes, with $\tan(\beta) = v_2/v_1$ and G_F being the Fermi constant.

To explain the CMS excess in resonant $t\bar{t}$ production at ≈ 400 GeV (3.5σ locally) [31] we set $m_A = 400$ GeV (see Fig. 2) and find¹¹

$$\mu_t^2 = (0.82 \pm 0.23)/\text{Br}(A \rightarrow t\bar{t}). \quad (6)$$

This results in the violet band in Fig. 3.¹²

The $t\bar{t}$ associated production of A is constrained by the ATLAS limit on $\sigma(pp \rightarrow A t\bar{t} \rightarrow t\bar{t}t\bar{t})$ [119] (see Fig. 2). Here, a 2HDM of type II resulted in a 95% CL lower limit of $1 \lesssim \tan\beta$ for $m_A \approx 400$ GeV and thus $\mu_t \lesssim 1/\text{Br}(A \rightarrow t\bar{t})$ (gray excluded region in Fig. 3). However, also the total four-top cross section $\sigma(pp \rightarrow t\bar{t}t\bar{t})$ at 13 TeV is measured by ATLAS [120] and CMS [121] to be $22.5^{+6.6}_{-5.5}$ fb and $17.7^{+6.0}_{-5.4}$ fb, respectively. In our

¹¹Note that CMS only reported the p value as a function of μ_t while we assume a gaussian error for the cross section (and thus μ_t^2). The corresponding CMS analysis [118] is done with less integrated luminosity and reaches only down to 500 GeV. However, in the low mass region, a weaker-than-expected limit is observed.

¹²Note that the differential lepton distributions of resonant $t\bar{t}$ production via A are similar to the shapes within the SM. Their effect drops out to a good approximation in the normalized differential distributions studied above.

model, the NP contribution to the total $t\bar{t}t\bar{t}$ cross section induced by $pp \rightarrow A t\bar{t} \rightarrow t\bar{t}t\bar{t}$ can be obtained from the ATLAS analysis of $\sigma(pp \rightarrow A t\bar{t} \rightarrow t\bar{t}t\bar{t})$: for $m_A \approx 400$ GeV, the 95% CL upper limit on the total cross section of 14.4 fb, corresponds to $\tan(\beta) > 1$ (see Fig. 8 in Ref. [119]). Therefore, $\sigma(pp \rightarrow A t\bar{t} \rightarrow t\bar{t}t\bar{t}) \approx 14\mu_t^2$ fb and thus nicely compatible with the experimentally preferred NP contribution to the cross section.

ATLAS searches for the process $pp \rightarrow A \rightarrow ZH$ with $H \rightarrow t\bar{t}$ [122]. At the point in parameter space closest to our mass region, i.e., $m_A \approx 450$ GeV and $m_H \approx 350$ GeV, a limit of ≈ 600 fb is observed, compared to an expected limit of ≈ 400 fb. Even though we have $H \rightarrow W^+W^-b\bar{b}$ instead of $H \rightarrow t\bar{t}$, the signatures are similar and one can use these results to estimate a limit on $pp \rightarrow A \rightarrow ZH$ in our model (see yellow and pink region in Fig. 3).¹³ Note that the top associated production of H leads to an effect in $t\bar{t}W$. While this effect goes in the direction preferred when comparing the SM prediction to the experimental value, it only gives a moderate effect.

CMS [124] and ATLAS [125] searched for $t\bar{b}H^\pm \rightarrow t\bar{b}b\bar{t}$. They found a lower bound on $\tan(\beta)$ [corresponding to an upper bound on $Y_t^2 = 1/\tan^2(\beta)$] as a function of $m_{H^\pm}$. For $m_{H^\pm}$ the upper limit on μ_t is around 0.6. However, depending on the coupling λ , a sizable decay rate for $H^\pm \rightarrow \Delta^\pm S$ is possible. With $\Delta^\pm \rightarrow W^\pm Z$ and $S \rightarrow b\bar{b}$, this results in $t\bar{t}Z$ and $t\bar{t}W$ -like signature. In fact, given that there is a preference for a ≈ 200 fb contribution in the latest $t\bar{t}W$ analysis of ATLAS [126], a NP contribution of around ≈ 400 fb is allowed at the 2σ level. Since the production cross section for $t\bar{b}H^\pm \rightarrow t\bar{b}b\bar{t}$ is 600 fb for $m_{H^\pm} = 400$ GeV and $\mu_t = 1$, it is possible to evade the $t\bar{b}H^\pm \rightarrow t\bar{b}b\bar{t}$ bound, which is therefore only shown as a dashed line in Fig. 3.

C. 151.5 and 95 GeV signal strength

The main indications for a new resonance at ≈ 151.5 GeV are in inclusive $\gamma\gamma$ searches [17,127], $\gamma\gamma + (E_T^{\text{miss}} > 90 \text{ GeV})$ [128,129] and W^+W^- (without jets) [26,27], where E_T^{miss} stands for missing transverse energy. Reference [15] found $\sigma(pp \rightarrow \Delta^0 \rightarrow \gamma\gamma) \approx (6 \pm 3)$ fb and $\sigma(pp \rightarrow \Delta^0 \rightarrow \gamma\gamma + (E_T^{\text{miss}} > 90 \text{ GeV})) \approx (0.6 \pm 0.2)$ fb. The latter is obtained within the simplified model $pp \rightarrow (270 \text{ GeV}) \rightarrow (151.5 \text{ GeV} \rightarrow \gamma\gamma)(151.5^* \text{ GeV} \rightarrow \tilde{\chi}\tilde{\chi})$ where $m_{\tilde{\chi}} = 50$ GeV. The process W^+W^- with a full jet veto was studied for the direct production of a new scalar in

¹³Depending on m_H , we can have the decay $A \rightarrow ZH$ with $H \rightarrow \Delta^0 S \rightarrow W^+W^-b\bar{b}$ which could potentially contaminate the $t\bar{t}Z$ cross section measurement. The most precise measurement of the inclusive $t\bar{t}Z$ cross section to date stems from the ATLAS experiment [123], which is consistent with the SM. That said, all channels, except for one (four leptons with different flavor) rely on the use of machine learning. This will suppress the sensitivity to non- $t\bar{t}Z$ final states, such as $A \rightarrow ZH, H \rightarrow W^+W^-b\bar{b}$, in particular, for $m_H < m_{\tilde{t}}$.




FIG. 4. Preferred regions from signal strength measurements of $\gamma\gamma$ (inclusive), $\gamma\gamma + (E_T^{\text{miss}} > 90 \text{ GeV})$ and W^+W^- (jet veto) of the 151.5 GeV boson Δ^0 as a function of its branching ratio to $\gamma\gamma$ and the branching ratio of the 95 GeV boson S to invisible, for $m_H = 290 \text{ GeV}$.

Ref. [28] finding $\approx 2\sigma$ significance for a nonzero cross section of $\approx 1 \text{ pb}$ at $\approx 151.5 \text{ GeV}$. In our model, Δ^0 is produced in association with S and therefore mainly with b jets, taus, and leptons. Thus, a nonzero branching ratio of $S \rightarrow \text{invisible}$, $\text{Br}(S \rightarrow \bar{\chi}\chi)$, is needed to pass the full jet veto and to give rise to $\gamma\gamma + E_T^{\text{miss}}$. We simulated both $\gamma\gamma + E_T^{\text{miss}}$ and $W^+W^- + E_T^{\text{miss}}$ in our model to obtain the efficiency and the significances for $S \rightarrow E_T^{\text{miss}}$, i.e., $m_\chi < 95/2 \text{ GeV}$.¹⁴ The results for the three regions $\gamma\gamma$ (inclusive), $\gamma\gamma + (E_T^{\text{miss}} > 90 \text{ GeV})$, and W^+W^- + 0 jets are shown in Fig. 4 for $m_H = 290 \text{ GeV}$. One can see that all three signal strengths are nicely compatible for $\text{Br}(S \rightarrow \bar{\chi}\chi) \approx 5\%$ and $\text{Br}(\Delta^0 \rightarrow \gamma\gamma) \approx 0.1\%$. Note that the latter value is within the natural expectations for the scalar triplet with hypercharge $Y = 0$ [133].

Finally, the diphoton signal strength for S ($\approx 95 \text{ GeV}$) induced via $pp \rightarrow H \rightarrow \Delta^0 S$ [30] of $\mu_{\gamma\gamma} \approx 0.1$ (with respect

¹⁴For the simulations, we employed MadGraph5aMC@NLO [130] with the parton shower performed by Pythia8.3 [131] and the detector simulation carried out with Delphes [132]. We validated our analysis of $\gamma\gamma + E_T^{\text{miss}}$ by comparing it against the ATLAS simulation in Ref. [17]. To correct for our fast simulation vs the full detector simulation of ATLAS, we added a 15 GeV smearing on the missing transverse energy (excluding jets) and on the jet energies. Note that also Drell-Yan production, i.e., $pp \rightarrow W^* \rightarrow \Delta^\pm \Delta^0$ with $\Delta^0 \rightarrow ZW$ and $Z \rightarrow \nu\nu$ and/or $W \rightarrow \ell\nu$, contributes to $\gamma\gamma + (E_T^{\text{miss}} > 90)$ which is responsible for a nonvanishing cross section at $\text{Br}(S \rightarrow \bar{\chi}\chi) = 0$ in Fig. 4.




FIG. 5. Predicted deviations from the SM predictions in the invariant mass distribution of the $b\bar{b}e\mu$ system for our benchmark point. Note that these results are in agreement with the preliminary ATLAS results of Ref. [143].

to a hypothetical SM Higgs-like boson with the same mass) is at the low side of the preferred range $\mu_{\gamma\gamma} = 0.24^{+0.09}_{-0.08}$ [134]. However, one could allow for a moderate mixing angle with the SM Higgs boson of around 0.2, which is possible if v_S is non-negligible. In this setup, one would be close to the central value while being consistent with SM Higgs boson signal strength measurements and at the same time explaining (partially) the LEP excess in Z Strahlung as well as the indications for $\tau\tau$ [135] and W^+W^- [28].

IV. CONCLUSIONS AND OUTLOOK

We propose the $\Delta 2\text{HDMS}$ as a combined explanation of the LHC multilepton anomalies, the resonant diphoton ($\approx 95 \text{ GeV}$ and $\approx 151.5 \text{ GeV}$) and top quark excesses ($\approx 400 \text{ GeV}$) with significances of $> 5\sigma$, $3\sigma - 5\sigma$, and 3.4σ , respectively. The model is obtained by extending the SM with a second Higgs boson doublet as well as a real $SU(2)_L$ singlet and a real $SU(2)_L$ triplet and naturally predicts (at tree level) a positive shift in the W mass in agreement with the current global fit for $v_\Delta = O(1 \text{ GeV})$. In particular, a benchmark points with

$$\begin{aligned} m_{\Delta^\pm} \approx m_{\Delta^0} &= 151.5 \text{ GeV}, & m_S &= 95 \text{ GeV}, \\ m_{H^\pm} \approx m_A &\approx 400 \text{ GeV}, & v_\Delta &\approx 2 \text{ GeV}, & v_S &= O(\text{GeV}), \\ m_H &= 290 \text{ GeV}, & \mu_t &= 0.7, \lambda = O(1), & \tan\beta &\gtrsim 5 \end{aligned} \quad (7)$$

is consistent with all signals and bounds.

While we provided proof of principle that all these anomalies can be explained without violating other bounds, in the future, a detailed examination of the NP effects in the differential $t\bar{t}W$ and $t\bar{t}Z$ distributions would be very interesting. Additionally, note that, e.g., the disappearance of the resonant $t\bar{t}$ would not falsify the model: if A were lighter, i.e., below the top threshold, then it could contribute to the differential $t\bar{t}$ distributions via $pp \rightarrow A \rightarrow W^\pm H^\mp$ with $H^\pm \rightarrow tb$, which would supplement the effect of $pp \rightarrow H \rightarrow \Delta^0 S \rightarrow W^+W^- b\bar{b}$. This

would allow for an explanation of the anomaly with a small top Yukawa coupling and H could be lighter since $\text{Br}(A \rightarrow ZH)$ would be smaller, and thus the $t\bar{t}Z$ -like signal is reduced.

Our model can be tested in multiple ways with forthcoming measurements. For example, it predicts deviations from the SM predictions in the invariant mass of the $b\bar{b}e\mu$ system in the differential $t\bar{t}$ analyses (see Fig. 5) and in asymmetric Higgs signals, e.g., in $\gamma\gamma b\bar{b}$. Furthermore, it opens a window of opportunity for the exploration and observation of new bosons at the electroweak scale at future e^+e^- accelerators, in particular the charged component of the triplet ($\Delta^\pm$), such as the Circular Electron-Positron Collider [136,137], the Compact Linear Collider [138], the

Future Circular Collider [139,140], and the International Linear Collider [141,142].

ACKNOWLEDGMENTS

We thank Saiyad Ashanujjaman, Sumit Banik, and Gino Isidori for useful discussions. The work of A. C. and G. C. is supported by a professorship grant from the Swiss National Science Foundation (Grant No. PP00P2_211002). B. M. gratefully acknowledges the South African Department of Science and Innovation through the SACERN program, the National Research Foundation, and the Research Office of the University of the Witwatersrand for various forms of support.

-
- [1] P. A. Zyla *et al.* (Particle Data Group Collaboration), Review of particle physics, *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020).
 - [2] P. W. Higgs, Broken symmetries, massless particles and gauge fields, *Phys. Lett.* **12**, 132 (1964).
 - [3] F. Englert and R. Brout, Broken symmetry and the mass of gauge vector mesons, *Phys. Rev. Lett.* **13**, 321 (1964).
 - [4] P. W. Higgs, Broken symmetries and the masses of gauge bosons, *Phys. Rev. Lett.* **13**, 508 (1964).
 - [5] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, Global conservation laws and massless particles, *Phys. Rev. Lett.* **13**, 585 (1964).
 - [6] G. Aad *et al.* (ATLAS Collaboration), Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012).
 - [7] S. Chatrchyan *et al.* (CMS Collaboration), Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
 - [8] J. M. Langford (ATLAS, CMS Collaborations), Combination of Higgs measurements from ATLAS and CMS: Couplings and k - framework, *Proc. Sci. LHCP2020* (2021) 136.
 - [9] ATLAS Collaboration, Combined measurements of Higgs boson production and decay using up to 139 fb⁻¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS experiment, Technical Report No. ATLAS-CONF-2021-053, 2021.
 - [10] R. Barate *et al.* (LEP Working Group for Higgs boson searches, ALEPH, DELPHI, L3, OPAL Collaborations), Search for the standard model Higgs boson at LEP, *Phys. Lett. B* **565**, 61 (2003).
 - [11] A. M. Sirunyan *et al.* (CMS Collaboration), Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV, *Phys. Lett. B* **793**, 320 (2019).
 - [12] CMS Collaboration, Searches for additional Higgs bosons and vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, Technical Report No. CMS-PAS-HIG-21-001, 2022.
 - [13] CMS Collaboration, Search for a new resonance decaying to two scalars in the final state with two bottom quarks and two photons in proton-proton collisions at $\sqrt{s} = 13$ TeV, Technical Report No. CMS-PAS-HIG-21-011, 2022.
 - [14] ATLAS Collaboration, Search for diphoton resonances in the 66 to 110 GeV mass range using 140 fb⁻¹ of 13 TeV pp collisions collected with the ATLAS detector, Technical Report No. ATLAS-CONF-2023-035, 2023.
 - [15] A. Crivellin, Y. Fang, O. Fischer, S. Bhattacharya, M. Kumar, E. Malwa, B. Mellado, N. Rapheeha, X. Ruan, and Q. Sha, Accumulating evidence for the associated production of a new Higgs boson at the LHC, *Phys. Rev. D* **108**, 115031 (2023).
 - [16] S. Bhattacharya, G. Coloretti, A. Crivellin, S.-E. Dahbi, Y. Fang, M. Kumar, and B. Mellado, Growing excesses of new scalars at the electroweak scale, [arXiv:2306.17209](https://arxiv.org/abs/2306.17209).
 - [17] G. Aad *et al.* (ATLAS Collaboration), Search for dark matter in events with missing transverse momentum and a Higgs boson decaying into two photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **10** (2021) 013.
 - [18] G. Aad *et al.* (ATLAS Collaboration), Model-independent search for the presence of new physics in events including $H \rightarrow \gamma\gamma$ with $\sqrt{s} = 13$ TeV pp data recorded by the ATLAS detector at the LHC, *J. High Energy Phys.* **07** (2023) 176.
 - [19] S. von Buddenbrock, N. Chakrabarty, A. S. Cornell, D. Kar, M. Kumar, T. Mandal, B. Mellado, B. Mukhopadhyaya, R. G. Reed, and X. Ruan, Phenomenological signatures of additional scalar bosons at the LHC, *Eur. Phys. J. C* **76**, 580 (2016).
 - [20] S. von Buddenbrock, A. S. Cornell, A. Fadol, M. Kumar, B. Mellado, and X. Ruan, Multi-lepton signatures of

- additional scalar bosons beyond the standard model at the LHC, *J. Phys. G* **45**, 115003 (2018).
- [21] S. Buddenbrock, A. S. Cornell, Y. Fang, A. Fadol Mohammed, M. Kumar, B. Mellado, and K. G. Tomiwa, The emergence of multi-lepton anomalies at the LHC and their compatibility with new physics at the EW scale, *J. High Energy Phys.* **10** (2019) 157.
- [22] S. von Buddenbrock, R. Ruiz, and B. Mellado, Anatomy of inclusive $t\bar{t}W$ production at hadron colliders, *Phys. Lett. B* **811**, 135964 (2020).
- [23] Y. Hernandez, M. Kumar, A. S. Cornell, S.-E. Dahbi, Y. Fang, B. Lieberman, B. Mellado, K. Monnagotla, X. Ruan, and S. Xin, The anomalous production of multi-lepton and its impact on the measurement of Wh production at the LHC, *Eur. Phys. J. C* **81**, 365 (2021).
- [24] O. Fischer *et al.*, Unveiling hidden physics at the LHC, *Eur. Phys. J. C* **82**, 665 (2022).
- [25] A. Crivellin and B. Mellado, Anomalies in particle physics, *Nat. Rev. Phys.* **6**, 294 (2024).
- [26] A. Tumasyan *et al.* (CMS Collaboration), Measurements of the Higgs boson production cross section and couplings in the W boson pair decay channel in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **83**, 667 (2023).
- [27] G. Aad *et al.* (ATLAS Collaboration), Measurements of Higgs boson production by gluon-gluon fusion and vector-boson fusion using $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decays in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **108**, 032005 (2023).
- [28] G. Coloretti, A. Crivellin, S. Bhattacharya, and B. Mellado, Searching for low-mass resonances decaying into W bosons, *Phys. Rev. D* **108**, 035026 (2023).
- [29] G. Aad *et al.* (ATLAS Collaboration), Inclusive and differential cross-sections for dilepton $t\bar{t}$ production measured in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector, *J. High Energy Phys.* **07** (2023) 141.
- [30] S. Banik, G. Coloretti, A. Crivellin, and B. Mellado, Uncovering new Higgses in the LHC analyses of differential $t\bar{t}$ cross sections, [arXiv:2308.07953](https://arxiv.org/abs/2308.07953).
- [31] A. M. Sirunyan *et al.* (CMS Collaboration), Search for heavy Higgs bosons decaying to a top quark pair in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **04** (2020) 171; **03** (2022) 187(E).
- [32] X.-G. He, T. Li, X.-Q. Li, J. Tandean, and H.-C. Tsai, Constraints on scalar dark matter from direct experimental searches, *Phys. Rev. D* **79**, 023521 (2009).
- [33] B. Grzadkowski and P. Osland, Tempered two-Higgs-doublet model, *Phys. Rev. D* **82**, 125026 (2010).
- [34] H. E. Logan, Dark matter annihilation through a lepton-specific Higgs boson, *Phys. Rev. D* **83**, 035022 (2011).
- [35] M. S. Boucenna and S. Profumo, Direct and indirect singlet scalar dark matter detection in the lepton-specific two-Higgs-doublet model, *Phys. Rev. D* **84**, 055011 (2011).
- [36] X.-G. He, B. Ren, and J. Tandean, Hints of standard model Higgs boson at the LHC and light dark matter searches, *Phys. Rev. D* **85**, 093019 (2012).
- [37] Y. Bai, V. Barger, L. L. Everett, and G. Shaughnessy, Two-Higgs-doublet-portal dark-matter model: LHC data and Fermi-LAT 135 GeV line, *Phys. Rev. D* **88**, 015008 (2013).
- [38] X.-G. He and J. Tandean, Low-mass dark-matter hint from CDMS II, Higgs boson at the LHC, and darkon models, *Phys. Rev. D* **88**, 013020 (2013).
- [39] Y. Cai and T. Li, Singlet dark matter in a type II two Higgs doublet model, *Phys. Rev. D* **88**, 115004 (2013).
- [40] C.-Y. Chen, M. Freid, and M. Sher, Next-to-minimal two Higgs doublet model, *Phys. Rev. D* **89**, 075009 (2014).
- [41] J. Guo and Z. Kang, Higgs naturalness and dark matter stability by scale invariance, *Nucl. Phys. B* **898**, 415 (2015).
- [42] L. Wang and X.-F. Han, A simplified 2HDM with a scalar dark matter and the galactic center gamma-ray excess, *Phys. Lett. B* **739**, 416 (2014).
- [43] A. Drozd, B. Grzadkowski, J. F. Gunion, and Y. Jiang, Extending two-Higgs-doublet models by a singlet scalar field—the Case for Dark Matter, *J. High Energy Phys.* **11** (2014) 105.
- [44] R. Campbell, S. Godfrey, H. E. Logan, A. D. Peterson, and A. Poulin, Implications of the observation of dark matter self-interactions for singlet scalar dark matter, *Phys. Rev. D* **92**, 055031 (2015); **101**, 039905(E) (2020).
- [45] A. Drozd, B. Grzadkowski, J. F. Gunion, and Y. Jiang, Isospin-violating dark-matter-nucleon scattering via two-Higgs-doublet-model portals, *J. Cosmol. Astropart. Phys.* **10** (2016) 040.
- [46] A. Arhrib, R. Benbrik, M. El Kacimi, L. Rahili, and S. Semlali, Extended Higgs sector of 2HDM with real singlet facing LHC data, *Eur. Phys. J. C* **80**, 13 (2020).
- [47] I. Engeln, P. Ferreira, M. M. Muhlleitner, R. Santos, and J. Wittbrodt, The dark phases of the N2HDM, *J. High Energy Phys.* **08** (2020) 085.
- [48] D. Azevedo, P. Gabriel, M. Muhlleitner, K. Sakurai, and R. Santos, One-loop corrections to the Higgs boson invisible decay in the dark doublet phase of the N2HDM, *J. High Energy Phys.* **10** (2021) 044.
- [49] T. Biekötter, S. Heinemeyer, and G. Weiglein, Excesses in the low-mass Higgs-boson search and the W -boson mass measurement, *Eur. Phys. J. C* **83**, 450 (2023).
- [50] D. A. Ross and M. J. G. Veltman, Neutral currents in neutrino experiments, *Nucl. Phys. B* **95**, 135 (1975).
- [51] J. F. Gunion, R. Vega, and J. Wudka, Higgs triplets in the standard model, *Phys. Rev. D* **42**, 1673 (1990).
- [52] P. H. Chankowski, S. Pokorski, and J. Wagner, (Non) decoupling of the Higgs triplet effects, *Eur. Phys. J. C* **50**, 919 (2007).
- [53] T. Blank and W. Hollik, Precision observables in $SU(2) \times U(1)$ models with an additional Higgs triplet, *Nucl. Phys. B* **514**, 113 (1998).
- [54] J. R. Forshaw, A. Sabio Vera, and B. E. White, Mass bounds in a model with a triplet Higgs, *J. High Energy Phys.* **06** (2003) 059.
- [55] M.-C. Chen, S. Dawson, and T. Krupovnickas, Higgs triplets and limits from precision measurements, *Phys. Rev. D* **74**, 035001 (2006).
- [56] R. S. Chivukula, N. D. Christensen, and E. H. Simmons, Low-energy effective theory, unitarity, and non-decoupling behavior in a model with heavy Higgs-triplet fields, *Phys. Rev. D* **77**, 035001 (2008).

- [57] P. Bandyopadhyay and A. Costantini, Obscure Higgs boson at colliders, *Phys. Rev. D* **103**, 015025 (2021).
- [58] S. Inoue, G. Ovanessian, and M. J. Ramsey-Musolf, Two-step electroweak baryogenesis, *Phys. Rev. D* **93**, 015013 (2016).
- [59] P. Bandyopadhyay, C. Coriano, and A. Costantini, Probing the hidden Higgs bosons of the $Y = 0$ triplet- and singlet-extended Supersymmetric standard model at the LHC, *J. High Energy Phys.* **12** (2015) 127.
- [60] P. Bandyopadhyay, C. Coriano, and A. Costantini, Perspectives on a supersymmetric extension of the standard model with a $Y = 0$ Higgs triplet and a singlet at the LHC, *J. High Energy Phys.* **09** (2015) 045.
- [61] P. Ko, Y. Omura, and C. Yu, A resolution of the flavor problem of two Higgs doublet models with an extra $U(1)_H$ symmetry for Higgs flavor, *Phys. Lett. B* **717**, 202 (2012).
- [62] S. Banik, A. Crivellin, S. Iguro, and T. Kitahara, Asymmetric di-Higgs signals of the next-to-minimal 2HDM with a $U(1)$ symmetry, *Phys. Rev. D* **108**, 075011 (2023).
- [63] S. Ashanujjaman, S. Banik, G. Coloretti, A. Crivellin, B. Mellado, and A.-T. Mulaudzi, $SU(2)_L$ triplet scalar as the origin of the 95 GeV excess?, *Phys. Rev. D* **108**, L091704 (2023).
- [64] A. Crivellin, M. Kirk, and A. Thapa, Minimal model for the W -boson mass, $(g-2)_\mu$, $h \rightarrow \mu^+\mu^-$ and quark-mixing-matrix unitarity, *Phys. Rev. D* **108**, L031702 (2023).
- [65] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, Theory and phenomenology of two-Higgs-doublet models, *Phys. Rep.* **516**, 1 (2012).
- [66] A. Strumia, Interpreting electroweak precision data including the W -mass CDF anomaly, *J. High Energy Phys.* **08** (2022) 248.
- [67] I. Dorsner and I. Mocioiu, Predictions from type II see-saw mechanism in $SU(5)$, *Nucl. Phys.* **B796**, 123 (2008).
- [68] P. Fileviez Perez, H. H. Patel, and A. D. Plascencia, On the W mass and new Higgs bosons, *Phys. Lett. B* **833**, 137371 (2022).
- [69] Y. Cheng, X.-G. He, F. Huang, J. Sun, and Z.-P. Xing, Electroweak precision tests for triplet scalars, *Nucl. Phys.* **B989**, 116118 (2023).
- [70] T. G. Rizzo, Kinetic mixing and dark Higgs triplets, and m_W , *Phys. Rev. D* **106**, 035024 (2022).
- [71] J.-W. Wang, X.-J. Bi, P.-F. Yin, and Z.-H. Yu, Electroweak dark matter model accounting for the CDF W -mass anomaly, *Phys. Rev. D* **106**, 055001 (2022).
- [72] M. Chabab, M. C. Peyranère, and L. Rahili, Probing the Higgs sector of $Y = 0$ Higgs triplet model at LHC, *Eur. Phys. J. C* **78**, 873 (2018).
- [73] Y. Shimizu and S. Takeshita, W boson mass and grand unification via the type-II seesaw-like mechanism, *Nucl. Phys.* **B994**, 116290 (2023).
- [74] E. Bagnaschi, J. Ellis, M. Madigan, K. Mimasu, V. Sanz, and T. You, SMEFT analysis of m_W , *J. High Energy Phys.* **08** (2022) 308.
- [75] J. de Blas, M. Pierini, L. Reina, and L. Silvestrini, Impact of the recent measurements of the top-quark and W -boson masses on electroweak precision fits, *Phys. Rev. Lett.* **129**, 271801 (2022).
- [76] R. L. Workman *et al.* (Particle Data Group Collaboration), Review of particle physics, *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [77] T. Aaltonen *et al.* (CDF Collaboration), High-precision measurement of the W boson mass with the CDF II detector, *Science* **376**, 170 (2022).
- [78] A. Broggio, E. J. Chun, M. Passera, K. M. Patel, and S. K. Vempati, Limiting two-Higgs-doublet models, *J. High Energy Phys.* **11** (2014) 058.
- [79] P. Athron, M. Bach, D. H. J. Jacob, W. Kotlarski, D. Stöckinger, and A. Voigt, Precise calculation of the W boson pole mass beyond the standard model with FlexibleSUSY, *Phys. Rev. D* **106**, 095023 (2022).
- [80] ATLAS Collaboration, Improved W boson mass measurement using 7 TeV proton-proton collisions with the ATLAS detector, Technical Report No. ATLAS-CONF-2023-004, 2023.
- [81] R. Aaij *et al.* (LHCb Collaboration), Measurement of the W boson mass, *J. High Energy Phys.* **01** (2022) 036.
- [82] S. Schael *et al.* (ALEPH, DELPHI, L3, OPAL, LEP Electroweak Collaborations), Electroweak measurements in electron-positron collisions at W -boson-pair energies at LEP, *Phys. Rep.* **532**, 119 (2013).
- [83] T.-K. Chen, C.-W. Chiang, and K. Yagyu, Explanation of the W mass shift at CDF II in the extended Georgi-Machacek model, *Phys. Rev. D* **106**, 055035 (2022).
- [84] W. Chao, M. Jin, H.-J. Li, and Y.-Q. Peng, Axion-like dark matter from the type-II seesaw mechanism, *Phys. Rev. D* **109**, 115027 (2024).
- [85] G. Lazarides, R. Maji, R. Roshan, and Q. Shafi, Heavier W boson, dark matter, and gravitational waves from strings in an $SO(10)$ axion model, *Phys. Rev. D* **106**, 055009 (2022).
- [86] G. Senjanović and M. Zantedeschi, $SU(5)$ grand unification and W -boson mass, *Phys. Lett. B* **837**, 137653 (2023).
- [87] T.-K. Chen, C.-W. Chiang, and K. Yagyu, CP violation in a model with Higgs triplets, *J. High Energy Phys.* **06** (2023) 069; **07** (2023) 169(E).
- [88] D. de Florian *et al.* (LHC Higgs Cross Section Working Group Collaboration), Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector (CERN, Geneva, 2016), Vol. 2.
- [89] E. Braaten and J. P. Leveille, Higgs boson decay and the running mass, *Phys. Rev. D* **22**, 715 (1980).
- [90] N. Sakai, Perturbative QCD corrections to the hadronic decay width of the Higgs boson, *Phys. Rev. D* **22**, 2220 (1980).
- [91] T. Inami and T. Kubota, Renormalization group estimate of the hadronic decay width of the Higgs boson, *Nucl. Phys.* **B179**, 171 (1981).
- [92] S. G. Gorishnii, A. L. Kataev, and S. A. Larin, The width of Higgs boson decay into hadrons: Three loop corrections of strong interactions, *Sov. J. Nucl. Phys.* **40**, 329 (1984), <https://inspirehep.net/literature/193688>.
- [93] S. G. Gorishnii, A. L. Kataev, S. A. Larin, and L. R. Surguladze, The analytical four loop corrections to the QED Beta function in the $\overline{MS}$ scheme and to the QED ψ function: Total reevaluation, *Phys. Lett. B* **256**, 81 (1991).
- [94] S. G. Gorishnii, A. L. Kataev, S. A. Larin, and L. R. Surguladze, Corrected three loop QCD correction

- to the correlator of the quark scalar currents and $\Gamma^{\text{tot}}(H^0 \rightarrow \text{hadrons})$, *Mod. Phys. Lett. A* **05**, 2703 (1990).
- [95] S. G. Gorishnii, A. L. Kataev, S. A. Larin, and L. R. Surguladze, Scheme dependence of the next to next-to-leading QCD corrections to $\Gamma^{\text{tot}}(H^0 \rightarrow \text{hadrons})$ and the spurious QCD infrared fixed point, *Phys. Rev. D* **43**, 1633 (1991).
- [96] A. Djouadi, P. Gambino, and B. A. Kniehl, Two loop electroweak heavy fermion corrections to Higgs boson production and decay, *Nucl. Phys.* **B523**, 17 (1998).
- [97] G. Degrandi and F. Maltoni, Two-loop electroweak corrections to the Higgs-boson decay $H \rightarrow \gamma\gamma$, *Nucl. Phys.* **B724**, 183 (2005).
- [98] G. Passarino, C. Sturm, and S. Uccirati, Complete two-loop corrections to $H \rightarrow \gamma\gamma$, *Phys. Lett. B* **655**, 298 (2007).
- [99] S. Actis, G. Passarino, C. Sturm, and S. Uccirati, NNLO computational techniques: The cases $H \rightarrow \gamma\gamma$ and $H \rightarrow gg$, *Nucl. Phys.* **B811**, 182 (2009).
- [100] A. Djouadi, M. Spira, J. J. van der Bij, and P. M. Zerwas, QCD corrections to $\gamma\gamma$ decays of Higgs particles in the intermediate mass range, *Phys. Lett. B* **257**, 187 (1991).
- [101] K. G. Chetyrkin, Correlator of the quark scalar currents and $\Gamma^{\text{tot}}(H \rightarrow \text{hadrons})$ at $O(\alpha_s^3)$ in pQCD, *Phys. Lett. B* **390**, 309 (1997).
- [102] P. A. Baikov, K. G. Chetyrkin, and J. H. Kuhn, Scalar correlator at $O(\alpha_s^4)$, Higgs decay into bottom quarks and bounds on the light-quark masses, *Phys. Rev. Lett.* **96**, 012003 (2006).
- [103] M. Spira, A. Djouadi, and P. M. Zerwas, QCD corrections to the $HZ\gamma$ coupling, *Phys. Lett. B* **276**, 350 (1992).
- [104] D. Graudenz, M. Spira, and P. M. Zerwas, QCD corrections to Higgs boson production at proton proton colliders, *Phys. Rev. Lett.* **70**, 1372 (1993).
- [105] M. Spira, A. Djouadi, D. Graudenz, and P. M. Zerwas, Higgs boson production at the LHC, *Nucl. Phys.* **B453**, 17 (1995).
- [106] C. Anastasiou and K. Melnikov, Higgs boson production at hadron colliders in NNLO QCD, *Nucl. Phys.* **B646**, 220 (2002).
- [107] R. V. Harlander and W. B. Kilgore, Next-to-next-to-leading order Higgs production at hadron colliders, *Phys. Rev. Lett.* **88**, 201801 (2002).
- [108] R. V. Harlander and W. B. Kilgore, Soft and virtual corrections to proton proton $\rightarrow H + x$ at NNLO, *Phys. Rev. D* **64**, 013015 (2001).
- [109] U. Aglietti, R. Bonciani, G. Degrandi, and A. Vicini, Analytic results for virtual QCD corrections to Higgs production and decay, *J. High Energy Phys.* **01** (2007) 021.
- [110] H.-L. Li, P.-C. Lu, Z.-G. Si, and Y. Wang, Associated production of Higgs boson and $t\bar{t}$ at LHC, *Chin. Phys. C* **40**, 063102 (2016).
- [111] C. Anastasiou, S. Beerli, S. Bucherer, A. Daleo, and Z. Kunszt, Two-loop amplitudes and master integrals for the production of a Higgs boson via a massive quark and a scalar-quark loop, *J. High Energy Phys.* **01** (2007) 082.
- [112] R. Harlander and P. Kant, Higgs production and decay: Analytic results at next-to-leading order QCD, *J. High Energy Phys.* **12** (2005) 015.
- [113] V. Ravindran, J. Smith, and W. L. van Neerven, NNLO corrections to the total cross-section for Higgs boson production in hadron hadron collisions, *Nucl. Phys.* **B665**, 325 (2003).
- [114] A. M. Sirunyan *et al.* (CMS Collaboration), Measurements of $t\bar{t}$ differential cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV using events containing two leptons, *J. High Energy Phys.* **02** (2019) 149.
- [115] CMS Collaboration, Differential cross section measurements for the production of top quark pairs and of additional jets using dilepton events from pp collisions at $\sqrt{s} = 13$ TeV, [arXiv:2402.08486](https://arxiv.org/abs/2402.08486).
- [116] S. Dawson, S. Dittmaier, and M. Spira, Neutral Higgs boson pair production at hadron colliders: QCD corrections, *Phys. Rev. D* **58**, 115012 (1998).
- [117] M. Spira, Higgs boson production and decay at hadron colliders, *Prog. Part. Nucl. Phys.* **95**, 98 (2017).
- [118] M. Aaboud *et al.* (ATLAS Collaboration), Search for heavy particles decaying into a top-quark pair in the fully hadronic final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **99**, 092004 (2019).
- [119] G. Aad *et al.* (ATLAS Collaboration), Search for $t\bar{t}H/A \rightarrow t\bar{t}t\bar{t}$ production in the multilepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **07** (2023) 203.
- [120] G. Aad *et al.* (ATLAS Collaboration), Observation of four-top-quark production in the multilepton final state with the ATLAS detector, *Eur. Phys. J. C* **83**, 496 (2023).
- [121] A. Hayrapetyan *et al.* (CMS Collaboration), Observation of four top quark production in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* **847**, 138290 (2023).
- [122] G. Aad *et al.* (ATLAS Collaboration), Search for a CP -odd Higgs boson decaying into a heavy CP -even Higgs boson and a Z boson in the $\ell^+\ell^-\bar{t}t$ and $\nu\bar{\nu}b\bar{b}$ final states using 140 fb $^{-1}$ of data collected with the ATLAS detector, *J. High Energy Phys.* **02** (2024) 197.
- [123] G. Aad *et al.* (ATLAS Collaboration), Inclusive and differential cross-section measurements of $t\bar{t}Z$ production in pp collisions at $\sqrt{s} = 13$, TeV with the ATLAS detector, including EFT and spin-correlation interpretations, *J. High Energy Phys.* **07** (2024) 163.
- [124] A. M. Sirunyan *et al.* (CMS Collaboration), Search for a charged Higgs boson decaying into top and bottom quarks in events with electrons or muons in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **01** (2020) 096.
- [125] G. Aad *et al.* (ATLAS Collaboration), Search for charged Higgs bosons decaying into a top quark and a bottom quark at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **06** (2021) 145.
- [126] ATLAS Collaboration, Measurement of the total and differential cross-sections of $t\bar{t}W$ production in pp collisions at 13 TeV with the ATLAS detector, Technical Report No. ATLAS-CONF-2023-019, 2023.
- [127] A. M. Sirunyan *et al.* (CMS Collaboration), Measurements of Higgs boson production cross sections and couplings in the diphoton decay channel at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **07** (2021) 027.
- [128] ATLAS Collaboration, Measurement of the properties of Higgs boson production at $\sqrt{s} = 13$ TeV in the $H \rightarrow \gamma\gamma$ channel using 139 fb $^{-1}$ of pp collision data with the

- ATLAS experiment, Technical Report No. ATLAS-CONF-2020-026, 2020.
- [129] A. M. Sirunyan *et al.* (CMS Collaboration), Search for dark matter produced in association with a Higgs boson decaying to $\gamma\gamma$ or $\tau^+\tau^-$ at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **09** (2018) 046.
 - [130] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
 - [131] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, An introduction to Pythia 8.2, *Comput. Phys. Commun.* **191**, 159 (2015).
 - [132] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi (Delphes 3 Collaboration), Delphes 3, A modular framework for fast simulation of a generic collider experiment, *J. High Energy Phys.* **02** (2014) 057.
 - [133] S. Ashanujjaman, S. Banik, G. Coloretti, and A. Crivellin, Anatomy of the Higgs triplet model with zero hypercharge (to be published).
 - [134] T. Biekötter, S. Heinemeyer, and G. Weiglein, The 95.4 GeV di-photon excess at ATLAS and CMS, *Phys. Rev. D* **109**, 035005 (2024).
 - [135] A. Tumasyan *et al.* (CMS Collaboration), Searches for additional Higgs bosons and for vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **07** (2023) 073.
 - [136] M. Dong *et al.* (CEPC Study Group Collaboration), CEPC conceptual design report: Volume 2—physics & detector, [arXiv:1811.10545](#).
 - [137] F. An *et al.*, Precision Higgs physics at the CEPC, *Chin. Phys. C* **43**, 043002 (2019).
 - [138] T. K. Charles *et al.* (CLICdp, CLIC Collaboration), The Compact Linear Collider (CLIC)—2018 Summary Report, [arXiv:1812.06018](#).
 - [139] A. Abada *et al.* (FCC Collaboration), FCC-ee: The lepton collider: Future circular collider conceptual design report volume 2, *Eur. Phys. J. Special Topics* **228**, 261 (2019).
 - [140] A. Abada *et al.* (FCC Collaboration), FCC physics opportunities: Future circular collider conceptual design report volume 1, *Eur. Phys. J. C* **79**, 474 (2019).
 - [141] ILC Collaboration, The International Linear Collider Technical Design Report—Volume 2: Physics, [arXiv:1306.6352](#).
 - [142] The International Linear Collider Technical Design Report—Volume 3.I: Accelerator R&D in the Technical Design Phase, [arXiv:1306.6353](#).
 - [143] B. Rottler, Search for Higgs-boson pair-production in the $b\bar{b}\ell\ell + ET_{\text{miss}}$ final state with the ATLAS detector at $\sqrt{s} = 13$ TeV, Ph.D. thesis Freiburg University, 2023.