

https://dspace.org/

Metamorphic studies in the Vredefort Dome, South Africa

Ogilvie, Paula

2010-09-08
http://hdl.handle.net/10539/8628

Downloaded from a DSpace repository. https://dspace.org/

METAMORPHIC STUDIES IN THE VREDEFORT DOME, SOUTH AFRICA

Paula Ogilvie

A dissertation submitted to the Faculty of Science, University of the Witwatersrand, in fulfilment of the requirements for the degree of Doctor of Philosophy

Johannesburg, 2010

Statement of original contribution

I declare that this dissertation is my own, unaided work. It is being submitted for the Degree of Doctor of Philosophy in the University of the Witwatersrand, Johannesburg, South Africa. It has not been submitted before for any degree or examination in any other University.

Signed this 8th day of June 2010



_____

Paula Ogilvie

Abstract

Metasedimentary granulites in the core of the Vredefort Dome present textural and chemical evidence for three discrete metamorphic events. These include a peak Archaean anatectic event (M_1), shock metamorphism (M_2) with impact at 2.02 Ga and post-shock metamorphism (M_3) of the target rocks related to Dome formation and non-adiabatic loading of the crust.

Regional granulite facies metamorphism (M_1) occurred between 3.10 Ga and 3.08 Ga with tectonomagmatic thickening of the crust attributable to easterly- to northeasterly-directed subduction of an oceanic slab beneath the Kaapvaal craton. Phase equilibria modelling in THERMOCALC of highly restitic pelitic granulites constrains peak conditions of M_1 metamorphism at 870 - 885 ºC and 7.1 - 7.7 kbar. Slightly lower peak conditions of 858 °C and 7.1 kbar were obtained for a more melt-rich granulite, reflecting back-reaction with a melt on the suprasolidus retrograde path. The prograde up-pressure trajectory is dominated by heating from 6.5 kbar, 700 °C to 7.5 kbar, 850 °C. Phase equilibria constraints on the prograde and suprasolidus retrograde evolution are consistent with a clockwise Archaean *P-T* path for the M_1 event.

Overprinting the M_1 peak assemblage are shock-induced, extreme disequilibrium deformation features (irregular shock fractures, planar fractures, planar deformation features, isotropization and shock melting) that formed instantaneously during meteorite impact at 2.02 Ga (M_2). Reconstruction of the shock pressure and post-shock temperature distribution across the central core of the Vredefort Dome from observed shock effects in component phases from the pelitic granulites required an experimental study to constrain shock effects in an analogous, complex, polymineralic, pelitic granulite from the Etivé aureole, with a significant proportion of hydrous and ferromagnesian minerals. Shock experiments were performed at 12.5, 25, 34, 40 and 56 GPa at 25 °C, and at 18 and 25 GPa at 400 °C to investigate the roles of both increasing shock pressure and pre-shock temperature on shock deformation features in major minerals. Both the shock experiments and Vredefort granulites are characterised by heterogeneous distribution of shock effects in minerals on intragranular and intergranular scales. Shock heterogeneity compromises estimates of absolute shock pressures based exclusively on observed

shock effects in minerals. Independent constraints on shock pressures are obtained from post-shock metamorphic conditions and range from > 35 GPa to > 40 GPa at 8 and 5 km from the centre of the Dome, respectively.

The Vredefort granulites underwent unusually rapid and highly variable M_3 heating, exhumation and cooling associated with the 2.02 Ga meteorite impact event. The short-lived nature of the thermal event and restitic bulk rock compositions owing to melt loss during the Archaean M_1 event, led to diffusion-controlled reaction and the growth of coronas around garnet. Coronas display a strongly sectoral development indicative of highly localized compositional domains. Grain size, sectoral complexity and thickness of coronas all increase toward the centre of the Dome, indicating strong temperature control on the extent of reaction. This sectoral complexity is unique to Vredefort coronas compared to coronas reported from regional and contact metamorphic terranes and affords the opportunity to evaluate controls on extent of corona development and degree of equilibration. Minimum peak M_3 temperatures were 980 °C at 2.5 – 3.0 kbar, between 8 and 5 km from the centre of the Dome.

Open-system diffusion and phase equilibria modelling of the Vredefort coronas has established a relationship between equilibration in granulites at the micrometre-scale as a function of temperature and melt fertility of the corona bulk composition. Higher melt modes and solidus depression in fertile corona bulk compositions enhance component diffusion and equalization of chemical potential gradients throughout the equilibration volume. Coronas are characterized by non-linear open-system metasomatic exchange of components with adjacent domains. Selective and variable open-system metasomatic exchange of components with the matrix or with contiguous domains is required to reproduce observed mineral modes and compositions. Reaction may be induced in chemically inert corona domains through open-system diffusive communication with a hydrous matrix, thereby fluxing the solidus and elevating melt modes. A better understanding of the textural and compositional evolution of coronas requires a shift from closed-system or linear phase equilibria modelling to non-linear, open-system modelling.

Dedication

This thesis is dedicated to my parents, Colin and Edna, and my brother and sister, Colin and Adrienne, for their endless support and encouragement.

Acknowledgements

I would like to express my deepest gratitude to my supervisor, Prof. Roger Gibson for his infinite reserves of patience and all his encouragement over the years. Without his support, guidance and insight, none of this would have been possible. Prof. Gary Stevens and Prof. Uwe Reimold are also thanked for their invaluable scientific contributions throughout the study. Prof. Carl Anhaeusser is also thanked for many inspiring chats and discussions. The National Research Foundation is acknowledged for financial support for the duration of my full-time studies.

I would also like to thank Dr. Jochen Schweitzer, Phil Lambert, Dr. Richard Stewart, Dave Stewart, Gokhan Güler, Vanessa Vermaak, Fatima Vogt and all the staff at Shango Solutions for their humbling support and guidance throughout the duration of my studies. Adele Grobler is also thanked for assistance with logistics and printing.

Dr Johann Diener and Prof. Richard White are thanked for their THERMOCALC technical support. Special acknowledgement must go to Johann Diener for the assistance and time invested in teaching me melt-integration in THERMOCALC.

Leonie Reyneke and the staff at Kumba Resources Mineralogy department are thanked for allowing me access to their microscopes during the study. Prof. Alex Deustch is thanked for performing the shock experiments in this study and preparation of the thin sections of the shocked specimens. At Stellenbosch University, Dr. Esme Spicer and Madelaine Frazenburg are also thanked for technical support during acquisition of SEM and XRF data. Dr. Rudolph Erasmus of the NRF Centre of Excellence in Strong Materials at Wits University is thanked for assistance during acquisition of Raman Spectra. Thanks also go to Peter Gräser from the University of Pretoria for his meticulous work and assistance with microprobe analysis. I would like to express my gratitude to Alex Mathebula for help with numerous thin sections and sample preparation. Safia Cannell, Dalena Blithental, Matt Kitching, Mitch Miles and Melody van Wyngaard are gratefully acknowledged for logistical support throughout this study.

I have been fortunate to have made some wonderful friends during my time at Wits, who have also contributed greatly to this study through their fantastic support,

encouragement and technical input. Thanks go to my good friend Louise Coney, Luke Longridge, Lindi Richer, Anika Solanki, Guy Freemantle, James Roberts, Charlie Seabrook and Christine Reinaud. For the last two years, I have been able to complete my studies in the Garden Route and I would like to thank my awesome drumming friends and teachers, Nidhi, Richelle, Anthea, Marissa, Jude, Karen, Sheila, Pat and Kaya, for teaching me to speak ‘djembe’.

Finally, my deepest and most heart-felt gratitude goes to my parents, brother and sister for all their love and support over the years.

TABLE OF CONTENTS

Declaration ii
Abstract iii
Dedication v
Acknowledgements vi
List of figures xiii
List of tables xxi

VOLUME 1

Chapter 1: Introduction 1

1.1 General introduction 1
1.2 Geology of the Vredefort Dome 7
1.3 Geology of the Archaean Basement Complex 12
1.4 Metamorphic evolution of the Vredefort Dome 14
1.4.1 M_1: peak anatectic event (3.1 Ga) 16
1.4.2 M_2: shock metamorphism (2.02 Ga) 17
1.4.3 M_3, post-shock metamorphism (2.02 Ga) 19
1.5 Thesis outline 27

Chapter 2: Archaean metamorphic evolution of metapelites and metagreywackes of the Archaean Basement Complex in the core of the Vredefort Dome 31

2.1 Introduction 31
2.2 Mesoscopic sample description 35
2.2.1 Metapelites (SK6C, SK9-CLM) 35
2.2.2 Metagreywackes (SK4A) 36
2.3 Petrography 40
2.3.1 Metapelites (SK6C, SK9-CLM) 40
2.3.2 Metagreywackes (SK4A) 50
2.4 Bulk-rock compositions 52
2.5 Mineral chemistry 63
2.6 Thermobarometry and *P-T* path constraints 84
2.6.1 Estimation of peak metamorphic conditions 89

2.6.2 Prograde metamorphic evolution 107
2.6.3 Retrograde metamorphic evolution 113
2.7 Discussion 117
2.7.1 Towards a new model 118

Chapter 3: Corona textures in granulites - a review of formation mechanisms and implications for the metamorphic evolution of terranes 122

3.1 Introduction 122
3.2 Single-stage, steady-state, diffusion-controlled corona growth (SSDC) 125
3.3 Sequential, diffusion-controlled corona growth (SEQ) 130
3.4 Determining a sequential or single-stage model of corona growth? 137
3.5 Controls on corona development in pelitic and mafic granulites 139
3.5.1 Pressure, temperature and $a\mathrm{H_2O}$ 139
3.5.2 Reaction mechanism: single-stage vs. sequential development 143
3.5.3 Reactant compositions 143
3.5.4 Diffusion kinetics 148
3.5.5 Deformation and strain 150
3.6 Conditions of corona formation 163
3.7 Corona microstructure and compositional zonation 165
3.8 Quantitative modelling of coronas 171
3.9 Discussion 178

Chapter 4: Petrography of complex pelitic granulite coronas in the Vredefort Dome 180

4.1 Introduction 180
4.2 Petrography of coronas from the NW group 186
4.2.1 Garnet-quartz corona domain 191
4.2.2 Garnet-biotite corona domain 197
4.2.3 Garnet-K-Feldspar corona domain 204
4.2.4 Garnet-plagioclase corona domain 210
4.2.5 Garnet-cordierite corona domain 216
4.2.6 Cordierite-biotite-sillimanite pseudomorph 221
4.2.7 Fracture symplectite 224
4.2.8 Summary 227

4.3 Petrography of coronas from the SE group 232
4.3.1 Garnet-quartz corona domain 236
4.3.2 Garnet-biotite corona domain 242
4.3.3 Garnet-K-Feldspar corona domain 248
4.3.4 Garnet-plagioclase corona domain 254
4.3.5 Garnet-cordierite corona domain 260
4.3.6 Cordierite-biotite-sillimanite pseudomorph 264
4.3.7 Fracture symplectite 267
4.3.8 Summary 270
4.4 Comparison between the NW and SE groups 275
4.5 Discussion 281

Chapter 5: Open-system, metasomatic, diffusion modelling of complex coronas in pelitic granulites from the Vredefort Dome 285

5.1 Introduction 285
5.2 Mineral chemistry 293
5.2.1 Compositional variation in corona product phases 312
5.2.2 Evidence for open-system behaviour in corona phase compositions – implications for calculation of the overall corona reaction 318
5.3 Open-system diffusion modelling 320
5.3.1 Corona overall reactions 337
5.3.2 L-ratios 342
5.3.3 Corona product proportions 344
5.3.4 Component fluxes 346
5.3.5 Chemical potential gradients 375
5.3.6 Chemical potential relationships and corona structure 383
5.3.7 Gibbs free energy gradients of corona phases 390
5.3.8 Stability criterion 392
5.3.9 Non-equilibrium thermobarometry 396
5.3.10 Reaction affinity 415
5.4 Discussion 417

VOLUME II

Chapter 6: Phase equilibria modelling of complex coronas 422

6.1 Introduction 422
6.2 Bulk composition of corona domains 429
6.3 Phase equilibria modelling of the matrix contribution to the garnet-core compositional domain 437
6.4 Phase equilibria modelling of the garnet-reactant domains 443
6.4.1 Garnet-quartz domain 444
6.4.2 Garnet-biotite domain 452
6.4.3 Garnet-feldspar domain 461
6.4.4 Garnet-cordierite domain 470
6.5 Phase equilibria modelling of communication between contiguous corona domains 474
6.5.1 Garnet-quartz vs. garnet-biotite domains 480
6.5.2 Garnet-quartz vs. garnet-K-Feldspar/plagioclase domains 483
6.5.3 Garnet-K-Feldspar/plagioclase vs. garnet-biotite domains 488
6.5.4 Garnet-K-Feldspar vs. garnet-plagioclase domain 492
6.6 Phase equilibria modelling of cordierite-biotite-sillimanite compositional domains 495
6.7 Non-linear, metasomatic modification and evolution of corona bulk composition with prolonged reaction 498
6.7.1 Modal mismatch in garnet-feldspar and garnet-biotite domains 501
6.7.2 Modal mismatch in garnet-quartz domains 503
6.7.3 Melt loss and corona textural and compositional evolution 511
6.8 Extreme melt loss and the origin of the aluminous granofelses from the Inlandsee Pan 517
6.9 Discussion 526

Chapter 7: Experimental investigation of shock metamorphic effects in a metapelitic granulite 540

7.1 Introduction 540
7.2 Methodology 546
7.3 Optical and scanning electron microscopy 550
7.3.1 Quartz 550
7.3.2 Plagioclase 559
7.3.3 K-feldspar 567

7.3.4 Cordierite 575
7.3.5 Biotite 581
7.3.6 Garnet 588
7.3.7 Orthopyroxene 588
7.3.8 Spinel 593
7.3.9 Ilmenite and pyrite 593
7.4 PDF development in quartz 601
7.5 Raman spectroscopy 605
7.5.1 Quartz 605
7.5.2 Plagioclase 608
7.5.3 K-feldspar 611
7.5.4 Cordierite 614
7.5.5 Biotite 617
7.5.6 Garnet 620
7.5.6 Orthopyroxene 620
7.6 Mineral chemistry 626
7.7 Discussion 646
7.7.1 Summary of diagnostic shock effects in minerals 646
7.7.2 Role of pre-shock temperature on shock-induced deformation features 654
7.7.3 Shock heterogeneity 655
7.7.4 Shock pressure barometry in the core of the Vredefort Dome 658
7.7.5 Shock metamorphism and post-shock corona development 673

Chapter 8: Conclusions 676

8.1 M_1 peak anatectic metamorphism: approaching full equilibration 678
8.2 M_2 shock metamorphism: disequilibrium 681
8.3 M_3 Post-shock metamorphism: partial equilibration 685
8.3.1 Controls on corona formation 685
8.3.2 Thermobarometric estimates of corona formation 689
8.3.3 Limitations to phase equilibria modelling 690
8.4 A new frontier in metamorphism 692

References 694

Appendix A: Derivation of open-system diffusion models 721
Appendix B: Individual phase compositions for diffusion modelling of coronas 738

LIST OF FIGURES

Figure 1.1 The equilibration continuum 6
Figure 1.2 Schematic cross-section through the Vredefort complex crater 7
Figure 1.3 The Theophilus lunar impact structure 8
Figure 1.4 Simplified geological map showing the location of the Witwatersrand basin and Vredefort Dome 8
Figure 1.5 Geological map and cross-section of the Vredefort Dome 9
Figure 1.6 Shock pressure distribution throughout the Vredefort Dome 10
Figure 1.7 Numerical models for shock pressure and post-shock temperature distribution throughout the Vredefort Dome 11
Figure 1.8 Post-impact thermal evolution of target rocks 20
Figure 1.9 Pre- and post-impact *P-T* trajectories defined by thermobarometry on the cores of peak metamorphic phases and symplectite mineral pairs respectively Perchuk et al. (2002) 26
Figure 1.10 P-T diagram demonstrating *P-T* conditions determined by previous workers for each metamorphic event in the Vredefort Dome. 28
Figure 2.1 Geological map of the Jagers Vrede and Steynskraal farms 33
Figure 2.2 Metasedimentary garnet granulites from Steynskraal and Jagers Vrede 34
Figure 2.3 Mesoscopic images of SK6C 37
Figure 2.4 Mesoscopic images of SK9-CLM 38
Figure 2.5 Mesoscopic images of SK4A 39
Figure 2.6 Scanned thin section from SK6C mesosome 41
Figure 2.7 Microscopic images of SK6C 42
Figure 2.8 SK9-CLM scanned thin section 46
Figure 2.9 Microscopic images of SK9-CLM 47
Figure 2.10 SK4A scanned thin section 50
Figure 2.11 Microscopic images of SK4A 51
Figure 2.12 AFM projection from quartz, plagioclase and K-feldspar of bulk rock samples from the Steynskraal traverse 55
Figure 2.13 FeO+MgO vs. $K_2O/(K_2O+Na_2O)$ plot of Steynskraal metasedimentary granulites 56
Figure 2.14 Isocon plots for the bulk rock compositions of Steynskraal metasedimentary granulites 59
Figure 2.15 Isocon plots for the melt melt-reintegrated bulk rock compositions of Steynskraal metasedimentary granulites 60
Figure 2.16 Geochemical plots of bulk rock compositions of Steynskraal ultramafic and mafic granulites 61
Figure 2.17 AFM projection of all measured mineral compositions and respective bulk rock compositions of Steynskraal metasedimentary granulites 66
Figure 2.18 Garnet EMPA profiles for sample SK9-CLM 67
Figure 2.19 Garnet EMPA profiles for sample SK6C 68
Figure 2.20 Garnet EMPA profiles for sample SK4A 69
Figure 2.21 Cordierite EMPA profiles for sample SK9-CLM 70
Figure 2.22 Cordierite EMPA profiles for sample SK6C 71
Figure 2.23 Orthopyroxene EMPA profiles for sample SK4A 72
Figure 2.24 *P-T* pseudosection constructed for SK6C peak metamorphic conditions 92

Figure 2.25 *P-T* pseudosection constructed for SK6C peak metamorphic conditions, contoured for mineral composition 93
Figure 2.26 *P-T* pseudosection constructed for SK6C peak metamorphic conditions, contoured for mineral modes 94
Figure 2.27 *P-T* pseudosection constructed for SK9-CLM peak metamorphic conditions 96
Figure 2.28 *P-T* pseudosection constructed for SK9-CLM peak metamorphic conditions, contoured for mineral composition 97
Figure 2.29 *P-T* pseudosection constructed for SK9-CLM peak metamorphic conditions, contoured for mineral modes 98
Figure 2.30 *P-T* pseudosection constructed for SK4A peak metamorphic conditions 101
Figure 2.31 *P-T* pseudosection constructed for SK4A peak metamorphic conditions, contoured for mineral composition 102
Figure 2.32 *P-T* pseudosection constructed for SK4A peak metamorphic conditions, contoured for mineral modes 103
Figure 2.33 Combined reaction equilibria from peak pseudosections for SK4A, SK6C and SK9-CLM 106
Figure 2.34 *P-T* pseudosection constructed for SK6C prograde metamorphic evolution based on an estimated protolith composition 108
Figure 2.35 *P-T* pseudosection constructed for SK9-CLM prograde metamorphic evolution based on an estimated protolith composition 109
Figure 2.36 *T-X* pseudosection for SK9-CLM demonstrating the effect of melt loss on the equilibrium assemblage 111
Figure 2.37 *P-T* pseudosection constructed for SK4A prograde metamorphic evolution based on an estimated protolith composition 112
Figure 2.38 Supra-solidus retrograde *P-T* evolution derived from peak pseudosections for SK6C and SK9-CLM 115
Figure 2.39 *P-T* pseudosection constructed for SK4A peak metamorphic conditions with superimposed peak conditions and proposed *P-T* path. 116
Figure 2.40 Comparison of *P-T* estimates from previous studies and those attained in this study 120
Figure 3.1 Chemographic relationships and chemical potential saturation surfaces for local transient equilibria at corona boundaries during incipient stages of SSDC corona growth 126
Figure 3.2 Open-system, single-stage, steady-state, diffusion-controlled growth of prograde corona layers between olivine and plagioclase 129
Figure 3.3 Sequential layer development in a corona between olivine and plagioclase 131
Figure 3.4 Sequential model of corona layer growth at constant *P* and *T* 134
Figure 3.5 Sequential model of corona layer growth between plagioclase and olivine through varying component fluxes across the corona bands and later due to decompression 136
Figure 3.6 Common corona textures developed in mafic granulites. 140
Figure 3.7 Common corona textures developed in pelitic granulites. 146
Figure 3.8 Summary of *P-T* conditions of formation for coronas reviewed in this study 164
Figure 3.9 Variation in corona microstructure in mafic and pelitic bulk rock compositions for coronas reviewed in this study 166
Figure 3.10 Magnitude of compositional zonation in product corona bands for coronas reviewed in this study 170
Figure 3.11 Sketch of a typical corona developed between plagioclase and olivine in metagabbros 174
Figure 3.12 Isocon plot of Al/Si ratios in symplectites and the adjacent reactant plagioclase 175

Figure 4.1 Geological map of the Jagers Vrede and Steynskraal farms 183
Figure 4.2 Samples from the NW group, in the northwest of the Steynskraal traverse 184
Figure 4.3 Samples from the SE group, in the southeast of the Steynskraal traverse 185
Figure 4.4 Corona development after garnet in metasedimentary granulites from the NW group of the Steynskraal traverse 188
Figure 4.5 Photomicrographs demonstrating variation in corona development after garnet with respect to grain size of the garnet and enclosing matrix in the NW group 190
Figure 4.6 Schematic representation of coronas developed between garnet and quartz from the NW group rocks 193
Figure 4.7 Core coronas developed between garnet and quartz inclusions in the NW group of the Steynskraal traverse 195
Figure 4.8 Coronas developed between garnet and quartz in the NW group 196
Figure 4.9 Schematic representation of coronas developed between garnet and biotite from the NW group rocks 199
Figure 4.10 BSE images of SK6C garnet-biotite core coronas 201
Figure 4.11 Rim garnet-biotite coronas developed in the NW group rocks 202
Figure 4.12 Schematic representation of coronas developed between garnet and K-feldspar from the NW group rocks 206
Figure 4.13 Coronas developed between garnet and K-feldspar in the NW group 208
Figure 4.14 BSE images of a SK6C garnet-K-feldspar rim corona adjacent to matrix plagioclase and biotite 209
Figure 4.15 Schematic representation of coronas developed between garnet and plagioclase from the NW group rocks 212
Figure 4.16 Core coronas developed between garnet and plagioclase in the NW group 214
Figure 4.17 Rim coronas developed between garnet and plagioclase in the NW group 215
Figure 4.18 Schematic representation of coronas developed between garnet and cordierite in SK6C 217
Figure 4.19 BSE images of SK6C garnet-cordierite core coronas 219
Figure 4.20 BSE images of SK6C garnet-cordierite rim coronas 220
Figure 4.21 Schematic representation of reaction textures developed after cordierite, biotite and sillimanite 221
Figure 4.22 Symplectite development after cordierite, biotite and sillimanite in the NW group 223
Figure 4.23 BSE images of cordierite-orthopyroxene-spinel symplectite in fractures within the garnet cores 225
Figure 4.24 BSE images demonstrating the transition from cordierite-orthopyroxene-spinel symplectite in fractures within the garnet core to cordierite-dominated symplectites at the garnet rim 226
Figure 4.25 Variation of corona product modes between core and rim coronas for the NW group 228
Figure 4.26 Variation of corona layer thickness between core and rim coronas for the NW group 229
Figure 4.27 Variation of orthopyroxene vermicule size between core and rim coronas for the NW group 230
Figure 4.28 Variation of orthopyroxene vermicule spacing between core and rim coronas for the NW group 231
Figure 4.29 Corona development in the SE group rocks of the Steynskraal traverse 233
Figure 4.30 Photomicrographs demonstrating variation in corona development after garnet with respect to grain size of the garnet and enclosing matrix in the SE group 235
Figure 4.31 Schematic representation of coronas developed between garnet and quartz from the SE group rocks 238

Figure 4.32 Core coronas developed between garnet and quartz in the SE group of the Steynskraal traverse 240
Figure 4.33 Rim coronas developed between garnet and quartz in the SE group 241
Figure 4.34 Schematic representation of coronas developed between garnet and biotite from the SE group 244
Figure 4.35 Core corona development between garnet and biotite in the SE group 246
Figure 4.36 Rim corona development between garnet and biotite in the SE group 247
Figure 4.37 Schematic representation of coronas developed between garnet and K-feldspar from the SE group 250
Figure 4.38 Core coronas developed between garnet and K-feldspar in the SE group 252
Figure 4.39 Rim coronas developed between garnet and K-feldspar in the SE group 253
Figure 4.40 Schematic representation of coronas developed between garnet and plagioclase from the SE group 256
Figure 4.41 Core corona development between garnet and plagioclase in the SE group 258
Figure 4.42 Rim coronas developed between garnet and plagioclase in the SE group 259
Figure 4.43 Schematic representation of coronas developed between garnet and cordierite from the SE group 261
Figure 4.44 Corona development between garnet and cordierite in the SE group 263
Figure 4.45 Schematic representation of reaction textures developed after cordierite, biotite and sillimanite in the SE group 264
Figure 4.46 Symplectite development after cordierite, biotite and sillimanite in the SE group 266
Figure 4.47 Fracture symplectite in the core of garnet from SK9-CLM 268
Figure 4.48 EMPA mapping of fracture symplectite in the core of garnet from SK9-CLM. 269
Figure 4.49 Variation of corona product modes between core and rim coronas for the SE group 271
Figure 4.50 Variation of corona layer thickness between core and rim coronas for the SE group 272
Figure 4.51 Variation of orthopyroxene vermicule size between core and rim coronas the SE group 273
Figure 4.52 Variation of orthopyroxene vermicule spacing between core and rim coronas for the SE group 274
Figure 4.53 Variation in microstructure and modes for the garnet-quartz domain from NW group to SE group coronas 276
Figure 4.54 Variation in microstructure and modes for the garnet-biotite domain from NW group to SE group coronas 277
Figure 4.55 Variation in microstructure and modes for the garnet- K-feldspar domain from NW group to SE group coronas 278
Figure 4.56 Variation in microstructure and modes for the garnet-plagioclase domain from NW group to SE group coronas 279
Figure 4.57 Variation in microstructure and modes for the garnet-cordierite domain from NW group to SE group coronas 280
Figure 4.58 Comparison of average Vredefort corona width with maximum corona thickness observed in coronas described in the literature 282
Figure 4.59 Comparison of maximum corona vermicule size of Vredefort coronas width with maximum vermicule size observed in coronas described in the literature 283
Figure 5.1 Distribution of corona sample localities SK8C, SK12A and SK9A, B, C in the Steynskraal area 289
Figure 5.2 BSE images of the coronas investigated in this study 290
Figure 5.3 AFM projection from quartz, plagioclase and K-feldspar for corona mineral assemblages. 295

Figure 5.4 SK8C gridded and contoured mineral compositions for orthopyroxene 298
Figure 5.5 SK8C gridded and contoured mineral compositions for layer 1 cordierite 299
Figure 5.6 SK12A gridded and contoured mineral compositions for orthopyroxene 300
Figure 5.7 SK12A gridded and contoured mineral compositions for layer 1 cordierite 301
Figure 5.8 SK9A gridded and contoured mineral compositions for orthopyroxene 302
Figure 5.9 SK9A gridded and contoured mineral compositions for layer 1 cordierite 303
Figure 5.10 SK9A gridded and contoured mineral compositions for layer 2 plagioclase 304
Figure 5.11 SK9B gridded and contoured mineral compositions for orthopyroxene 305
Figure 5.12 SK9B gridded and contoured mineral compositions for layer 1 cordierite 306
Figure 5.13 SK9B gridded and contoured mineral compositions for layer 2 plagioclase 307
Figure 5.14 SK9B gridded and contoured mineral compositions for layer 2 K-feldspar 308
Figure 5.15 SK9C gridded and contoured mineral compositions for orthopyroxene 309
Figure 5.16 SK9C gridded and contoured mineral compositions for layer 1 cordierite 310
Figure 5.17 SK9C gridded and contoured mineral compositions for layer 2 plagioclase 311
Figure 5.18 Standard deviations for layer 1 cordierite-orthopyroxene symplectite phase compositions 314
Figure 5.19 Standard deviations for layer 2 plagioclase compositions 315
Figure 5.20 Standard deviations for layer 3 orthopyroxene compositions 315
Figure 5.21 Standard deviations for all orthopyroxene compositions 316
Figure 5.22 Comparison of zonation in Vredefort corona phases with coronas reviewed in the literature 317
Figure 5.23 Isocon plot of Al/Si ratios in modelled symplectites compared with symplectites in the literature 319
Figure 5.24 AFM diagram demonstrating the evolution of corona bulk composition with prolonged reaction 338
Figure 5.25 True and apparent metasomatic fluxes for open-system diffusion models reported in this study 341
Figure 5.26 L-ratios for stable diffusion models for coronas SK8C, SK12A, SK9A, SK9B and SK9C. 343
Figure 5.27 Observed proportions of corona product phases determined by image analysis 345
Figure 5.28 Development of positive fluxes over a corona band in a model system 348
Figure 5.29 Development of negative fluxes over a corona band in a model system 349
Figure 5.30 Model component fluxes for each component in corona layers 351
Figure 5.31 Development of negative fluxes for SiO_2 over corona layers in SK9C 355
Figure 5.32 Development of negative fluxes for SiO_2 over corona layers in SK9B 356
Figure 5.33 Development of negative fluxes for SiO_2 over corona layers in SK8C 357
Figure 5.34 Development of positive fluxes for $AlO_{3/2}$ over corona layers in SK9A 359
Figure 5.35 Development of positive fluxes for $AlO_{3/2}$ over corona layers in SK9C 360
Figure 5.36 Development of positive fluxes for FeO over corona layers in SK12A 361
Figure 5.37 Development of positive fluxes for FeO over corona layers in SK9B 362
Figure 5.38 Development of negative fluxes for MgO over corona layers in SK12A 365
Figure 5.39 Development of negative fluxes for MgO over corona layers in SK9B 366
Figure 5.40 Development of fluxes for CaO over corona layers in SK12A. 369
Figure 5.41 Development of fluxes for CaO over corona layers in SK9B 370
Figure 5.42 Development of fluxes for $NaO_{1/2}$ over corona layers in SK12A 372
Figure 5.43 Development of fluxes for $NaO_{1/2}$ over corona layers in SK9B 373
Figure 5.44 Development of fluxes for $KO_{1/2}$ over corona layers in SK9B 374

Figure 5.45 Schematic plot demonstrating the nature of chemical potential gradients during corona growth....376
Figure 5.46 Model chemical potential gradients for coronas SK8C, SK12A, SK9A, SK9B and SK9C. 378
Figure 5.47 Model chemical potential gradients in coronas SK8C, SK12A, SK9A, SK9B and SK9C....379
Figure 5.48 Relationship between positive component flux, phase composition and negative chemical potential gradients....381
Figure 5.49 Relationship between negative component flux, phase composition and positive chemical potential gradients....382
Figure 5.50 Qualitative $\mu_{FeO} - \mu_{SiO_2}$ section in NCFAS....385
Figure 5.51 Evolving chemical potential gradients and corona structure with prolonged reaction and variable diffusion rates....388
Figure 5.52 Model Gibbs free energy gradients for all corona phases across layers in coronas SK8C, SK12A, SK9A, SK9B and SK9C....391
Figure 5.53 Model free energy differences of phases at boundary q from those at the boundary where phase is present....394
Figure 5.54 Evaluation of reaction affinity at reference boundary A (q_j>q) and B (q_j<q)....398
Figure 5.55 Non-equilibrium thermobarometry – SK8C....410
Figure 5.56 Non-equilibrium thermobarometry – SK12A....411
Figure 5.57 Non-equilibrium thermobarometry – SK9A....412
Figure 5.58 Non-equilibrium thermobarometry – SK9B....413
Figure 5.59 Non-equilibrium thermobarometry – SK9C....414
Figure 5.60 Model reaction affinity for coronas investigated....416
Figure 6.1 Phase equilibria modelling of coronas....426
Figure 6.2 Harker plots demonstrating compositional differences between corona end-member EBCs....433
Figure 6.3 Si – Fe – Mg cation plot comparing effective bulk compositions for corona end-member domains....434
Figure 6.4 *T-X* pseudosections for fracture symplectite in the NW group....441
Figure 6.5 *T-X* pseudosections for fracture symplectite in the SE group....442
Figure 6.6 Combined garnet-matrix, garnet-quartz and garnet *T-X* pseudosections for the NW group....448
Figure 6.7 Combined garnet-matrix, garnet-quartz and garnet *T-X* pseudosections for the SE group....450
Figure 6.8 Combined garnet-matrix, garnet-biotite and garnet *T-X* pseudosections for the NW group....455
Figure 6.9 Combined garnet-matrix, garnet-biotite and garnet *T-X* pseudosections for the SE group....457
Figure 6.10 SK6C – Suprasolidus fields of garnet-matrix vs. garnet-biotite *T-X* pseudosection....458
Figure 6.11 SK8C – Suprasolidus fields of garnet-matrix vs. garnet-biotite *T-X* pseudosection....459
Figure 6.12 SK9-CLM – Suprasolidus fields of garnet-matrix vs. garnet-biotite *T-X* pseudosection....460
Figure 6.13 Combined garnet-matrix, garnet-plagioclase and garnet *T-X* sections for the NW group....464
Figure 6.14 Combined garnet-matrix, garnet-K-feldspar and garnet *T-X* sections for the NW group....466
Figure 6.15 Combined garnet-matrix, garnet-plagioclase/K-feldspar and garnet *T-X* sections for the SE group....468
Figure 6.16 Combined garnet-matrix, garnet-cordierite and garnet *T-X* sections for the NW and SE groups....472
Figure 6.17 Modal and textural relationships between contiguous corona domains in the NW group....475
Figure 6.18 Modal and textural relationships between contiguous corona domains in the SE group....478
Figure 6.19 *T-X* sections for the compositional range between end-member garnet-quartz and adjacent end-member garnet-biotite EBCs....481

Figure 6.20 *T-X* sections for the compositional range between end-member garnet-quartz and adjacent end-member garnet-K-feldspar EBCs 485
Figure 6.21 *T-X* sections for the compositional range between end-member garnet-quartz and adjacent garnet-plagioclase EBCs 487
Figure 6.22 *T-X* sections for the compositional range between end-member garnet-K-feldspar and adjacent end-member garnet-biotite EBCs 489
Figure 6.23 *T-X* pseudosection for the compositional range between end-member garnet-plagioclase and adjacent garnet-biotite EBCs 491
Figure 6.24 *T-X* sections for the compositional range between end-member garnet-K-feldspar and adjacent garnet-plagioclase EBCs 493
Figure 6.25 *T-X* sections for the compositional range between sillimanite-dominated, biotite+cordierite and a sillimanite-absent, biotite+cordierite EBCs 497
Figure 6.26 Comparison of observed and predicted Opx/Crd ratios for corona products from aluminous corona compositional domains 499
Figure 6.27 Comparison of observed and predicted Opx/Crd and Pl/Crd ratios for corona products for the garnet-quartz corona compositional domain 500
Figure 6.28 *T-X* pseudosection simulating the effect of open-system metasomatic loss of FeO and MgO from a garnet-dominated EBC 502
Figure 6.29 Sectoral development of the garnet-quartz corona 504
Figure 6.30 Electron microprobe element mapping for garnet-quartz and garnet-plagioclase domains in SK9B'3 505
Figure 6.31 *T-X* pseudosection for the compositional range between garnet-dominated, garnet-quartz EBCs and quartz-dominated, garnet-quartz EBCs 506
Figure 6.32 *T-X* pseudosection for the compositional range between garnet-dominated, garnet-quartz EBCs and quartz-plagioclase EBCs 507
Figure 6.33 *T-X* pseudosection accommodating open-system metasomatic exchange of FeO, MgO, and CaO with adjacent domains 508
Figure 6.34 Metasomatic exchange between compositional domains 510
Figure 6.35 *T-X* pseudosection for SK9-CLM garnet-quartz domain with a matrix component (28 mol%) showing the effect of 80% melt loss at 908°C on corona mineral modes and solidus elevation 512
Figure 6.36 *T-X* pseudosection for SK9-CLM garnet-plagioclase domain (20 mol% matrix) showing the effect of 100% melt loss at 1050 °C on corona mineral modes and solidus elevation. 514
Figure 6.37 Photomicrographs showing textures in VT484 518
Figure 6.38 *P-T* pseudosection for VT484 520
Figure 6.39 *P-T* pseudosection for VT484, a pelitic cordierite-K-feldspar-bearing granofels from the Inlandsee terrane, contoured for cordierite, K-feldspar and spinel mode 521
Figure 6.40 *T-X* pseudosection simulating residual cumulate and associated liquid compositions with progressive melt loss 523
Figure 6.41 Model compositional evolution with progressive melt extraction from a melt-cumulate mixture in a (Fe*+Ti+Mg)-(Na+Ca)-K cation percentage plot 524
Figure 6.42 Comparison of model liquid and residuum bulk compositions with progressive melt extraction from a melt-cumulate mixture and actual granofels and granite compositions 525
Figure 6.43 Variation in EBC range in which equilibration is approached in all corona domains from the NW and SE group rocks determined from *T-X* sections 527

Figure 6.44 Variation in melt modes predicted for a post-shock temperature of 1000 °C for the EBC range in which equilibration is approached in all corona domains from the NW and SE group rocks 529
Figure 6.45 Range of solidus temperatures for the EBC ranges in which equilibration is approached in all corona domains from the NW and SE group rocks 530
Figure 6.46 T-X_{Mg} pseudosection for SK9-CLM, demonstrating stable phase equilibria in the garnet-quartz domain with variable X_{Mg} 532
Figure 6.47 Variation in EBC ranges and solidus temperatures for adjacent corona domains in the NW group 535
Figure 6.48 Variation in EBC ranges and solidus temperatures for adjacent corona domains in the SE group 536
Figure 6.49 Variation in melt modes for adjacent corona domains in the NW and SE groups 537
Figure 7.1 Schematic shock pressure (P) - volume (V) Hugoniot and release adiabatic curves for a target mineral 542
Figure 7.2 Pressure-temperature fields of endogenic metamorphism and shock metamorphism 543
Figure 7.3 Unshocked pelitic granulite from the Etivé aureole, Scotland 548
Figure 7.4 Experimental apparatus used to generate shock waves in the specimen under investigation 549
Figure 7.5 Shock effects in quartz 552
Figure 7.6 Shock effects in plagioclase 561
Figure 7.7 Shock effects in K-feldspar 569
Figure 7.8 Shock effects in cordierite 576
Figure 7.9 Shock effects in biotite 582
Figure 7.10 Shock effects in garnet 589
Figure 7.11 Shock effects in orthopyroxene 592
Figure 7.12 Shock effects in spinel 594
Figure 7.13 Shock effects in ilmenite and pyrite 596
Figure 7.14 PDF development in quartz at 25 GPa (25 °C) 602
Figure 7.15 PDF development in quartz at 18 GPa (400 °C) 603
Figure 7.16 PDF development in quartz at 25 GPa (400 °C) 604
Figure 7.17 Unprocessed Raman spectra of quartz 606
Figure 7.18 Processed Raman spectra of quartz 607
Figure 7.19 Unprocessed Raman spectra of plagioclase 609
Figure 7.20 Processed Raman spectra of plagioclase 610
Figure 7.21 Unprocessed Raman spectra of K-feldspar 612
Figure 7.22 Processed Raman spectra of K-feldspar 613
Figure 7.23 Unprocessed Raman spectra of cordierite 615
Figure 7.24 Processed Raman spectra of cordierite 616
Figure 7.25 Unprocessed Raman spectra of biotite 618
Figure 7.26 Processed Raman spectra of biotite 619
Figure 7.27 Unprocessed Raman spectra of garnet 621
Figure 7.28 Processed Raman spectra of garnet 622
Figure 7.29 Processed Raman spectra of garnet of isotropic and birefringent garnet cores at 56 GPa 623
Figure 7.30 Unprocessed Raman spectra of orthopyroxene 624
Figure 7.31 Processed Raman spectra of orthopyroxene 625
Figure 7.32 Variation in garnet X_{Mg} between core and rim with increasing shock pressure and preheating. 639

Figure 7.33 Variation in biotite composition between core and rim with increasing shock pressure and preheating. 640
Figure 7.34 Variation in cordierite composition between core and rim with increasing shock pressure and preheating. 641
Figure 7.35 Variation in X_{Or} and X_{Ab} of K-feldspar between core and rim with increasing shock pressure and preheating....642
Figure 7.36 Variation in K/Si, and Al/Si of K-feldspar between core and rim with increasing shock pressure and preheating. 643
Figure 7.37 Variation in X_{Ab}, and X_{An} of plagioclase between core and rim with increasing shock pressure and preheating....644
Figure 7.38 Variation in Na/Si, and Al/Si of plagioclase between core and rim with increasing shock pressure and preheating. 645
Figure 7.39 Summary of shock effects in quartz in this study and results from previous studies 647
Figure 7.40 Summary of shock effects in plagioclase and K-feldspar in this study and results from previous 649
Figure 7.41 Summary of shock effects in biotite and cordierite in this study compared with results from previous studies 651
Figure 7.42 Summary of shock effects in orthopyroxene and garnet in this study compared with results from previous studies....653
Figure 7.43 Strain localization owing to shock impedance contrast between garnet and cordierite....658
Figure 7.44 Results of numerical models by Ivanov (2005) for shock pressure and post-shock temperature distribution throughout the Vredefort Dome....660
Figure 7.45 Photomicrograph montage (crossed polarisers) showing shock-induced microdeformation features in component minerals from SK11 (SE group) 662
Figure 7.46 Shock-induced microdeformation features in SK11 663
Figure 7.47 Photomicrograph montage (crossed polarisers) showing shock induced microdeformation features in component minerals from VT466B (NW group) 666
Figure 7.48 Shock-induced microdeformation features in VT466B 667
Figure 7.49 Shock-induced deformation in Etivé pelitic granulite at 25 GPa and 400 °C 670
Figure 7.50 Schematic representation of shock metamorphism of the Steynskraal pelitic granulites. 671
Figure 8.1 Schematic cartoon demonstrating the capacity for preservation of the peak assemblage as a function of melt loss in granulites.....680

LIST OF TABLES

Table 2.1 XRF Bulk rock compositions: Metagreywackes....53
Table 2.2 XRF Bulk rock compositions: Metapelites....54
Table 2.3 XRF Bulk rock compositions: Mafic granulites 62
Table 2.4 Representative mineral analyses SK9-CLM 73
Table 2.5 Representative mineral analyses SK6C 77
Table 2.6 Representative mineral analyses SK4A 81
Table 3.1 Summary of prograde corona occurrences in the literature....152
Table 3.2 Summary of retrograde corona occurrences in the literature 157

Table 4.1 Mineral modal compositions of the NW group granulites 186
Table 4.2 Summary of garnet-quartz corona petrography from the NW group 194
Table 4.3 Summary of garnet-biotite corona petrography from the NW group 200
Table 4.4 Summary of garnet-K-feldspar corona petrography from the NW group 207
Table 4.5 Summary of garnet-plagioclase corona petrography from the NW group 213
Table 4.6 Summary of garnet-cordierite corona petrography from the NW group 218
Table 4.7 Summary of sillimanite-cordierite-biotite pseudomorph petrography from the NW group 222
Table 4.8 Mineral modal compositions of the SE group granulites 232
Table 4.9 Summary of garnet-quartz corona petrography from the SE group 239
Table 4.10 Summary of garnet-biotite corona petrography from the SE group 245
Table 4.11 Summary of garnet-K-feldspar corona petrography from the SE group 251
Table 4.12 Summary of garnet-plagioclase corona petrography from the SE group 257
Table 4.13 Summary of garnet-cordierite corona petrography from the SE group 262
Table 4.14 Summary of sillimanite-cordierite-biotite pseudomorph petrography from the SE group 265
Table 5.1 Inferred THERMOCALC bulk compositions for SK8C, SK12A , SK9A, SK9B and SK9C coronas 288
Table 5.2 Average mineral chemistry for coronas SK8C, SK12A and SK9A 296
Table 5.3 Summary statistics of mineral chemistry for coronas SK8C, SK12A, SK9A, SK9B and SK9C 313
Table 5.4 Number of analytical points for each phase in all coronas 314
Table 5.5 Modelling results for corona SK8C 322
Table 5.6 Modelling results for corona SK12A 325
Table 5.7 Modelling results for corona SK9A 328
Table 5.8 Modelling results for corona SK9B 331
Table 5.9 Modelling results for corona SK9C 334
Table 5.10 Molar bulk compositions for SK8C, SK12A, SK9A, SK9B and SK9C coronas 339
Table 5.11 Component fluxes through layers and molar amounts of components liberated and consumed at layer boundaries 352
Table 5.12 Chemical potential relationships for components and end-members across corona layers – SK8C 400
Table 5.13 Chemical potential relationships for components and end-members across corona layers – SK12A 401
Table 5.14 Chemical potential relationships for components and end-members across corona layers – SK9A 402
Table 5.15 Chemical potential relationships for components and end-members across corona layers – SK9B 403
Table 5.16 Chemical potential relationships for components and end-members across corona layers – SK9C 404
Table 5.17 Model independent end-member reaction affinities 407
Table 6.1 Summary of coronal assemblage and textural trends 424
Table 6.2 End-member bulk compositions for phase equilibria modelling of corona domains (NCKFMASHTO) 430
Table 6.3 Observed and predicted corona product modes with attendant solidus temperatures for the fracture symplectite 440
Table 6.4 Observed and predicted corona product modes with attendant solidus temperatures for the garnet-quartz domain 447

Table 6.5 Observed and predicted corona product modes with attendant solidus temperatures for the garnet-biotite domain 454
Table 6.6 Observed and predicted corona product modes with attendant solidus temperatures for the garnet-feldspar domains 463
Table 6.7 Observed and predicted corona product modes with attendant solidus temperatures for the garnet-cordierite domain 471
Table 6.8 Best-fit model EBC ranges for contiguous corona domains 494
Table 6.9 Observed and predicted corona product modes with attendant solidus temperatures and melt modes for the cordierite-biotite-sillimanite domain 496
Table 6.10 Changes in the garnet-quartz corona assemblage with melt loss 513
Table 6.11 Changes in the garnet-plagioclase corona assemblage with melt loss 515
Table 7.1 Shock pressures and diagnostic shock effects in quartz, feldspar and biotite 545
Table 7.2 Average EMP analyses of garnet 629
Table 7.3 Average EMP analyses of biotite 631
Table 7.4 Average EMP analyses of cordierite 633
Table 7.5 Average EMP analyses of K-feldspar 635
Table 7.6 Average EMP analyses of plagioclase 637
Table 7.7 Intragranular and intergranular heterogeneity of shock effects developed in component phases 656