

https://dspace.org/

Hearing function in children with chronic renal dysfunction.

Lau, Jennifer

2013-04-02
http://hdl.handle.net/10539/12614

Downloaded from a DSpace repository. https://dspace.org/

Hearing Function in Children with Chronic Renal Dysfunction

Student: Jennifer Lau

Submitted to the

Discipline of Speech Pathology and Audiology

School of Human and Community Development

Faculty of Humanities

University of the Witwatersrand

in fulfilment of the requirements for the

Degree of Master of Audiology

July 2012

Declaration

I, Jennifer Lau, hereby declare that this submission is my own original work and that the assistance which I have received is detailed in the Acknowledgements of this report. To the best of my knowledge and belief, it contains no material which has been accepted for the award of any other degree or diploma at any other university or other institute of higher learning, except where due acknowledgement has been made in the text. I am responsible for the study and conclusions reached.

_____ _____

JENNIFER LAU Date

Acknowledgements

I would like to express my sincerest gratitude to the following people:

- Professor Katijah Khoza-Shangase, my supervisor, for her encouragement, advice and assistance throughout the supervision of my research.
- The participants with chronic renal disorder and their caregivers, who so willingly gave of their time to expand my knowledge.
- The Chief Executive Officer, Medical Superintendent and Heads of Department for both the Renal Department and Speech and Hearing Department at Charlotte Maxeke Johannesburg Academic Hospital, for granting me the opportunity to conduct this research.
- Dr. Levy, for his time and knowledge that he shared with me regarding renal disorders as well as the advice that he gave me regarding the research. I would also like to thank all the other doctors, nurses and administrative staff from the Renal Department, who so willingly helped me during the data collection phase of the study.
- The Head of Department as well as my fellow friends and colleagues from the Speech and Hearing Department, for their continual support, friendship, encouragement and assistance during my academic endeavours.
- The Ear, Nose and Throat Specialists at Charlotte Maxeke Johannesburg Academic Hospital, for their support and encouragement.
- Professor Peter Fridjhon, Dr. Michael Greyling, and Dr. Charles Chimedza, statisticians from the University of the Witwatersrand, for all their valuable time, knowledge and guidance regarding data analysis of the study. Their input and guidance is much appreciated.
- Mr. Herman Tesner, for his most timeous input, guidance and support with regards to the final check and editing of the research report.
- All hospital speech and hearing therapists and university staff who provided audiological input and emotional encouragement during this study.
- My family, for their unconditional love and support in the realization of my dreams, and standing by me through thick and thin.
- My friends, for their encouragement, interest and patience in the carrying out of my research study.

May God Bless You Always!

Abstract

The primary aim of the research was to describe hearing function in a group of children with chronic renal dysfunction receiving treatment in an academic hospital in Johannesburg, South Africa. Specific objectives in the study were to determine the prevalence of hearing loss in paediatric patients with chronic renal dysfunction; to describe the type, degree and configuration of the hearing loss; and to establish if there was a relationship between the presenting hearing loss and the severity of renal dysfunction, the different treatment regimens, duration of renal dysfunction, and the duration of treatment.

One hundred children between the ages five -18 years participated in the study and comprised 65 males and 35 females. The mean age of the participants was 11.68 years.

A cross-sectional, descriptive, quantitative research design was employed. All participants underwent a case history interview and a full audiological examination which included an otoscopic examination, immittance testing (tympanometry and ipsilateral acoustic reflex testing), pure tone audiometry including extended high frequency testing up to 16 kilohertz as well as diagnostic distortion product otoacoustic emission testing. A record review was also done.

Both descriptive and inferential statistics were used to analyse the collected data. Inferential statistics included parametric measures using multiple regression measures as well as non parametric measures using the Kruskal-Wallis statistical analysis.

Results revealed that there was a high prevalence of hearing loss in children with chronic renal dysfunction. Results from the extended high frequency pure tone testing as well as the diagnostic distortion product testing revealed that the most common hearing loss was a low and high to ultrahigh frequency mild sensorineural hearing loss. The study showed that there was no relationship between the severity of hearing loss and the severity of renal dysfunction, or the duration of renal dysfunction and the duration of treatment. However, the study showed that there was a relationship between the severity of hearing loss and certain treatments, that is,

haemodialysis and the use of ototoxic medication such as loop diuretics, tuberculosis medication, and antimalarial medication.

As the potential to miss hearing loss in this population is high, the research highlighted the importance of extended high frequency audiometry as well as diagnostic distortion product otoacoustic emission testing for the use of ototoxic monitoring in patients with chronic renal dysfunction. The research also highlighted the need for further research in this area as well as the need for educating medical personnel and caregivers working with children with chronic renal disease.

Key words: chronic renal dysfunction; hearing loss; distortion product; otoacoustic emissions; extended high frequency audiometry.

Table of Contents

	Pages
Acknowledgements	**iii**
Abstract	**iv-v**
List of Tables	**viii-ix**
List of Figures	**x**
List of Appendices	**xi**
Introduction	**1-22**
Methodology	**23-40**
Aims	**23**
Research Design	**24**
Participants	**24-25**
Research instrumentation	**26-27**
Ethical considerations	**27-28**
Research procedure	**28-30**
Audiological testing	**30-37**
Validity and reliability	**37**
Analysis of data	**37-40**
Results	**41-74**
Description of participants	**41-42**
Data analysis in terms of case history	**43-46**
Description of hearing function	**47-56**
Prevalence of hearing loss	**56-57**
Type, degree and configuration of hearing loss	**57-61**
Relationship between hearing loss and severity of chronic renal disease	**61-63**
Relationship between hearing loss and the different treatment regimens	**64-66**
Relationship between hearing loss and the duration of renal dysfunction	**66-67**
Relationship between hearing loss and the duration of treatment for renal dysfunction	**68-70**
Additional results obtained of the study	**70-74**
Discussion of results	**75-82**
Conclusion, limitations and recommendations	**83-88**

References **89-97**

Appendices **98-113**

List of tables

		Pages
Table 1:	Stages of Chronic Kidney Disease	**2**
Table 2:	Prevalence Figures of Hearing Loss for Sub-Saharan Africa	**12**
Table 3:	Published Studies on the Effect of Haemodialysis on Hearing Function	**16-17**
Table 4:	Results of Loop Diuretic Effects on Hearing Function in Animal Studies	**20-21**
Table 5:	Interpretation of Tympanometry Results	**32**
Table 6:	Classification of Hearing Loss	**34-35**
Table 7:	Mean Pure Tone Audiometry Results, Standard Deviations and Median Results for the Right and Left Ear (in dB HL)	**51**
Table 8:	Pure Tone Audiometry Results for the Right and Left Ear	**52-53**
Table 9:	Mean DPOAE Results and Standard Deviations for the Right and Left Ear	**54-55**
Table 10:	Type and degree of hearing loss between ears	**60-70**
Table 11:	Parametric analysis showing the relationship between severity of hearing loss and stage of chronic renal dysfunction	**62**
Table 12:	Non-parametric statistical analysis showing the relationship between degree of hearing loss and severity of chronic renal dysfunction	**63**
Table 13:	Parametric Statistical Analysis of the independency of variable severity of hearing loss and treatment methods	**64**
Table 14:	Non-Parametric Statistical Analysis of the relationship between degree of hearing loss and different ototoxic medications	**65**
Table 15:	Parametric logistic regression statistical analysis of the relationship between degree of hearing loss and the duration of chronic renal dysfunction	**66**
Table 16:	Non-parametric Kruskal-Wallis statistical analysis of the relationship between degree of hearing loss and the duration of chronic renal dysfunction	**67**

Table 17: Parametric logistic regression statistical analysis of the relationship between degree of hearing loss and the duration of treatment for chronic renal dysfunction **68**

Table 18: Non-parametric Kruskal-Wallis statistical analysis of the relationship between degree of hearing loss and duration of treatment for chronic renal dysfunction **69**

Table 19: Parametric multiple regression statistical analysis of the relationship between degree of hearing loss, duration of renal dysfunction and duration of treatment **70**

List of figures

		Page
Figure 1:	Gender distribution of participants with chronic renal dysfunction	**41**
Figure 1:	Distribution of stages of chronic kidney disease	**42**
Figure 2:	Number of participants with a family history of kidney disease	**43**
Figure 3:	Treatment regimens for participants with chronic renal disorder	**44**
Figure 4:	Medications used by the participants	**45**
Figure 5:	Otoscopic examination for each participant	**47**
Figure 6:	Tympanometry results for each participant	**48**
Figure 7:	Ipsilateral acoustic reflexes for the right and left ear	**49**
Figure 8:	Mean pure tone audiometry results for the right and left ear for each frequency (in dBHL)	**50**
Figure 9:	Mean DPOAE results for the right and left ear for each frequency	**54**
Figure 10:	DPOAE responses for the right and left ear	**56**
Figure 11:	Prevalence of hearing loss in a sample of paediatric participants with chronic kidney disease	**57**
Figure 12:	Types of hearing loss	**58**
Figure 13:	Degree of hearing loss	**59**
Figure 14:	Perceived hearing loss by the participants and their caregivers	**71**
Figure 15:	Perceived onset of hearing loss	**72**
Figure 16:	Perceived symmetry of hearing loss	**73**
Figure 17:	Other related ear symptoms	**74**

List of appendices

		Page
Appendix A:	Participant information sheet for parents/legal guardians	**98-99**
Appendix B:	Participant information sheet for paediatric patients	**100-101**
Appendix C:	Consent form for parent/legal guardian of the participants in the study	**102**
Appendix D:	Assent form for the participants in the study	**103**
Appendix E:	Hospital information sheet	**104-105**
Appendix F:	Case history questionnaire for parents/legal guardians of paediatric patients with renal dysfunction (Biographical Information)	**106**
Appendix G:	Case history questionnaire for parents/legal guardians of paediatric patients with renal dysfunction (Renal and Hearing Information)	**107**
Appendix H:	Letter of permission for the study from the chief executive officer of Charlotte Maxeke Johannesburg Academic Hospital	**108**
Appendix I:	Letter of permission for the study from the head of the paediatric renal department of Charlotte Maxeke Johannesburg Academic Hospital	**109**
Appendix J:	Letter of permission for the study from the head of the speech and hearing department of Charlotte Maxeke Johannesburg Academic Hospital	**110**
Appendix K:	Letter of permission for the study from Human Research Ethical Committee	**111**
Appendix L:	Certificate of calibration for the GSI Tympstar tympanometer	**112**
Appendix M:	Certificate of calibration for the Audera distortion product otoacoustic emission meter	**113**