

https://dspace.org/

Cardiovascular risk profile of kidney transplant recipients at the Charlotte Maxeke Johannesburg Academic Hospital.

Muhammad, Aminu Sakajiki

2014-04-25
http://hdl.handle.net10539/14587

Downloaded from a DSpace repository. https://dspace.org/

CARDIOVASCULAR RISK PROFILE OF KIDNEY TRANSPLANT RECIPIENTS AT THE CHARLOTTE MAXEKE JOHANNESBURG ACADEMIC HOSPITAL

MUHAMMAD AMINU SAKAJIKI


A dissertation submitted to the Faculty of Health Sciences University of Witwatersrand, Johannesburg, in fulfillment of the requirements for the degree of

Master of Science in Medicine

Johannesburg, 2013

DECLARATION

I, Muhammad Aminu Sakajiki, declare that this dissertation is my own work. It is being submitted for the degree of Master of Science in Medicine in the Division of Nephrology in the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at this or any other University.



MUHAMMAD AMINU SAKAJIKI

July, 2013

DEDICATION

I dedicate this work to:

The end-stage renal disease patients in sub-Saharan Africa and their relatives who struggle to pay for renal replacement therapy that is not readily available and where available, is not affordable.

My wife and children, who endured through my absence from the house in the course of this work.

PUBLICATIONS AND PRESENTATIONS ARISING FROM THIS STUDY

1. Aminu MS, Naicker S, Naidoo S, Maranga BM, Britz R, Manga P, Nazir MS. Carotid intima-media thickness and cardiovascular risk factors in kidney transplant recipients. World Congress of Nephrology, Hong Kong, June 2013 (accepted for presentation).

2. Aminu MS, Naicker S, Naidoo S, Maranga BM, Britz R, Manga P, Nazir MS. Obesity and graft outcome among kidney transplant recipients in a South African public hospital. World Congress of Nephrology, Hong Kong, June 2013 (accepted for presentation).

3. Aminu MS, Naicker S, Naidoo S, Maranga BM, Britz R, Manga P, Nazir MS. Left ventricular hypertrophy in kidney transplant recipients at Charlotte Maxeke Johannesburg Academic Hospital, South Africa. World Congress of Nephrology, Hong Kong, June 2013 (accepted for presentation).

4. Aminu MS, Naicker S, Naidoo S, Maranga BM, Britz R, Manga P, Nazir MS. Proteinuria, graft outcome and cardiovascular risk among kidney transplant recipients in a South African public hospital. Nigerian Association of Nephrology Conference (NANCOF), Abuja, Feb 2013 (accepted for presentation).

ABSTRACT

INTRODUCTION

Cardiovascular diseases (CVD) are more common in kidney transplant recipients (KTRs) than in the general population. The high incidence of CVD in the KTRs can be attributed to traditional risk factors, additional risk factors associated with graft dysfunction and those specifically related to transplantation.

Carotid intima-media thickness (cIMT) is a proven surrogate of atherosclerosis; it correlates with vessel pathology and is precisely imaged using ultrasound technology.

This study was aimed at determining the prevalence and predictors of cardiovascular risk among KTRs at the Charlotte Maxeke Johannesburg Academic Hospital (CMJAH) and to examine the relationship between cardiovascular risk factors and carotid intima media thickness.

METHODS

Patients aged 18 years and above who received a kidney transplant at the CMJAH between January 2005 and December 2009 were recruited. A questionnaire that captured cardiovascular risk factors was administered. Patients records were assessed for information on their post transplant follow up. All patients had echocardiography and carotid doppler done for measurement of intima-media thickness. The Framingham Risk Score was used to categorize patients into low, moderate, high risk and very high risk groups. Results were analyzed using statistical package for social sciences (SPSS) version 17, p value of 0.05 was considered significant.

RESULTS

One hundred (KTRs) 63 male (63%) and 37 female (37%) were recruited ranging in age from 19 to 70 years, with a mean age of 42.2 ± 12.42. Thirty six patients (36%) were found to have high cardiovascular

risk. Multiple regression showed proteinuria ($p = 0.022$), higher cumulative steroid dosage ($p = 0.028$), elevated serum triglycerides ($p = 0.04$) and the presence of plaques in the carotid artery ($p = 0.012$) as predictors of higher cardiovascular risk.

Carotid intima-media thickness correlates with higher CVD risk. Fourteen patients (14%) had a carotid artery plaque. Twenty five patients (25%) had cIMT of >0.7 mm.

CONCLUSION

Kidney transplant recipients in CMJAH were found to have high cardiovascular risk (36%) and carotid intima-media thickness correlates with this high CVD risk. Routine follow up of KTRs should include measurement of cIMT as it provides a simple noninvasive assessment of subclinical atherosclerosis.

ACKNOWLEDGEMENTS

I wish to extend my sincere appreciation to:

- Professor Saraladevi Naicker my supervisor, for her initiative, support and encouragement during this study.
- Dr Sagren Naidoo my second supervisor for his assistance and support.
- Professor P Manga, Dr Nazir MS and other members of the Division of Cardiology of the Charlotte Maxeke Johannesburg Academic Hospital for assisting with the echocardiography and allowing me to use their machine for carotid artery sonar.
- Consultants, fellows and other staff of the Division of Nephrology CMJAH for their support and assistance during the course of my stay.
- The kidney transplant recipients of the CMJAH for enthusiastically participating in this study.

TABLE OF CONTENTS **Page**

TITLE		i
DECLARATION		ii
DEDICATION		iii
PUBLICATIONS AND PRESENTATIONS		iv
ABSTRACT		v
ACKNOWLEDGEMENTS		vii
TABLE OF CONTENTS		viii
LIST OF FIGURES		xiv
LIST OF TABLES		Xv
NOMENCLATURE		xvii
1.0	**INTRODUCTION AND LITERATURE REVIEW**	**1**
1.1	Epidemiology of cardiovascular disease in KTRs	2
1.2	Framingham risk score	5
1.3	Spectrum of CVD in KTRs	7
1.3.1	Alterations in cardiac geometry	8
1.3.2	Atherosclerosis	9
1.3.3	Arteriosclerosis	10
1.4	Risk factors for CVD in KTRs	10
1.4.1	Traditional cardiovascular risk factors	12
1.4.1.1	Hypertension	12

1.4.1.2 Dyslipidaemia 14

1.4.1.3 Diabetes Mellitus 16

1.4.1.4 Obesity 18

1.4.1.5 Smoking 19

1.4.1.6 Age 20

1.4.1.7 Physical inactivity 21

1.4.1.8 Gender 21

1.4.2 Non-traditional cardiovascular risk factors 21

1.4.2.1 Graft function and cardiovascular disease 21

1.4.2.2 Proteinuria 24

1.4.2.3 Anaemia 24

1.4.2.4 Carotid intima-media thickness 25

1.4.2.5 Measurement of carotid intima-media thickness 26

1.4.2.6 Left ventricular hypertrophy 27

1.4.2.7 Rejection 27

1.4.2.8 Donor type 28

1.4.3 Immunosuppression and cardiovascular risk 28

1.4.3.1 Effects on renal function 29

1.4.3.2 Effects on diabetes 29

1.4.3.3 Effects on hypertension 30

1.4.3.4 Effects on dyslipidaemia 31

1.4.3.5 Effects on anaemia 32

1.4.3.6 Duration and type of dialysis 33

1.4.4 C-reactive protein 33

1.4.5 Homocysteine 34

1.5 Management of CVD in KTRs 35

1.5.1 Hypertension 35

1.5.1.1 Life style modification for blood pressure control 35

1.5.1.2 Drug therapy 36

1.5.2 Diabetes mellitus 38

1.5.2.1 Life style modification for the prevention and treatment of NODAT 39

1.5.3 Smoking cessation 41

1.5.4 Dyslipidaemia treatment 42

1.5.4.1 Lifestyle modification 42

1.5.4.2 Drug therapy 42

1.5.5 Treatment of anaemia 45

1.5.6 Aspirin therapy in the management of CVD 45

1.6 Study justification 46

1.7 Study objectives 46

2.0 MATERIALS AND METHODS 47

2.1 Study population 47

2.1.1 Inclusion criteria 47

2.1.2 Exclusion criteria 47

2.2 Methods 47

2.2.1 Blood pressure 48

2.2.2 Echocardiography 49

2.2.3 Carotid intima-media thickness 50

2.2.4 Laboratory test 51

2.2.4.1 Serum creatinine 52

2.2.4.2 Serum lipids 52

2.2.4.3 Complete blood count 53

2.2.4.4 Urine protein 53

2.2.5 Cardiovascular disease risk score 54

2.2.6 Statistics 54

3.0 RESULTS 55

3.1 Demographic data of the study population 55

3.2 Aetiology of End-stage renal disease 58

3.3 Traditional cardiovascular disease risk factors 60

3.3.1 History of cardiovascular disease 60

3.3.2 Hypertension 61

3.3.3 Diabetes mellitus 62

3.3.4 Obesity 63

3.3.5 Dyslipidaemia 64

3.3.6 Smoking 65

3.3.7 Exercise 66

3.3.8 Gender 66

3.3.9 Age 67

3.4	Transplant-associated cardiovascular risk factors	68
3.4.1	Graft dysfunction	68
3.4.2	Proteinuria	69
3.4.3	Anaemia	69
3.4.4	Rejection	69
3.4.5	Immunosuppressive medication	70
3.4.5.1	Calcineurin inhibitors	70
3.4.5.2	Steroids	70
3.4.5.3	Rapamycin	70
3.5	Left ventricular hypertrophy	71
3.6	Carotid intima-media thickness	74
4.0	**DISCUSSION**	78
4.1	Demographic data	78
4.2	Cardiovascular risk profile of the study population	79
4.3	Traditional cardiovascular risk factors	79
4.3.1	Hypertension	79
4.3.2	Serum lipids	79
4.3.3	Diabetes mellitus	80
4.3.5	Obesity	81
4.3.6	Smoking	81
4.3.7	Age	81
4.3.8	Physical exercise	82

4.3.9 Gender 82

4.4 Non-traditional cardiovascular risk factors 83

4.4.1 Graft function 83

4.4.2 Proteinuria 83

4.4.3 Anaemia 84

4.4.4 Rejection 84

4.4.5 Immunosuppressive medication 85

4.4.5.1 Calcineurin inhibitors 85

4.4.6 Carotid intima-media thickness 86

4.4.7 Left ventricular hypertrophy 86

4.5 Limitations of the study 88

5.0 CONCLUSIONS AND RECOMMENDATIONS 89

5.1 Conclusions 89

5.2 Recommendation for clinical practice 89

5.3 Recommendations for future work 89

APPENDICES 91

APPENDIX A Participant information sheet 91

APPENDIX B Participant consent form 93

APPENDIX C Data collection form 94

APPENDIX D Ethics clearance certificate 99

APPENDIX E Framingham 10-year risk assessment chart 100

REFERENCES **102**

LIST OF FIGURES

Figure **Page**

3.1 HLA Matching of study population 56

3.2 Source of Donor Kidneys 57

3.3 Causes of ESRD in the study population 58

3.4 Cardiovascular Risk category of the study population 59

LIST OF TABLES

Table		Page
1.1	Risk factors for the development of CVD after kidney transplant	5
1.2	Decreased kidney function after transplant is significantly associated with increased cardiac risk	23
1.3	Classes of antihypertensive agents commonly used in KTRs	37
1.4	Management of dyslipidaemia in KTRs: NKF working group guidelines	44
3.1	Age distribution by gender and race	55
3.2	Mean ± SD blood pressures of the study population	61
3. 3	BMI and waist circumference of the study population.	63
3.4	Serum cholesterol and HDL cholesterol of the study population	64
3.5	Smoking habit of the study population	65
3.6	Levels of physical exercise of the study population	66
3.7	Gender and CVD risk group	66
3.8	Age and CVD risk group	67
3.9	CVD risk group and eGFR	68
3.10	Echocardiographic values of the study population	72

3.11 LVMI and its correlation with some CV risk variables 73

3.12 Risk factors for increased LVMI 74

3.13 Mean ± SD CIMT of the study population 75

3.14 Correlation of CIMT with some CV risk variables 76

NOMENCLATURE

AVF	Arteriovenous Fistula
ALERT	Assessment of Lescol in Renal Transplantation
AMVL	Anterior mitral Valve Leaflet
AT	Acceleration Time
BSA	Body Surface Area
CVD	Cardiovascular Disease
CNI	Calcineurin Inhibitors
CIMT	Carotid Intima-Media Thickness
CAD	Coronary Artery Disease
CRP	C - reactive protein
CMJAH	Charlotte Maxeke Johannesburg Academic Hospital
DBP	Diastolic Blood Pressure
DT	Deceleration Time
ESRD	End Stage Renal Disease
FHS	Framingham Heart Study
GFR	Glomerular Filtration Rate
HF	Heart Failure
HDL	High Density lipoprotein
HbA1C	Glycosylated Haemoglobin
IHD	Ischaemic Heart Disease
IVSTD	Inter Ventricular Septal Thickness in Diastole
JNC	Joint National Committee

KTR	Kidney Transplant Recipients
KDOQI	Kidney Disease Outcome Quality Initiative
LDL	Low Density Lipoprotein Cholesterol
LVH	Left Ventricular Hypertrophy
LVEDD	Left Ventricular Internal Dimension in End – Diastole
LVESD	Left Ventricular Internal Dimension in End Systole
LVPWTD	Left Ventricular Posterior wall thickness in diastole
LVM	Left Ventricular Mass
LVMI	Left Ventricular Mass Index
MI	Myocardial Infarction
MAP	Mean Arterial Pressure
NRT	Nicotine Replacement Therapy
NCEP III	National Cholesterol Education Project Plan III
NHLS	National Health Laboratory Service
PVD	Peripheral Vascular Disease
RWT	Relative Wall Thickness
SPSS	Statistical Package for Social Science
SD	Standard Deviation
SBP	Systolic Blood Pressure
T2DM	Type 2 Diabetes Mellitus
T1DM	Type 1 Diabetes Mellitus
TOD	Target Organ Damage
UPCR	Urine Protein Creatinine Ratio

VLDL Very Low Density Lipoprotein