

APPENDICES

Appendix A1: Protein standard curve (Bradford, 1976)




Figure A1: Protein standard curve. Bradford's reagent was used to determine the concentration of P450 Protein at A595nm. The curve had an R^2 value of 0.9994 and the equation used was $y = 0.2956x - 4E-05$

Appendix A2 - Reproducibility of qRT-PCR tested for CYP6P9 on the Light Cycler® 480




Figure A2-1: CYP6P9 amplification profile, melting peak and standard curve - Although there is a difference in the melting peaks – there is a peak observed at 58°C when performed on the BioRad CFX96™ Real-Time PCR Detection System (bottom image) but not on the LC480 (top image) – there was no statistical difference ($P < 0.05$) between the data performed on the two machines.

Appendix A3: pA259 [pGEM-T easy::17 α -CYP6P9 (FANG), with no Histidine tag]

GGAACCCCTAAAGGGAGCCCCGATTTAGAGCTTGACGGGGAAAGCCGGCGAACGTGGCGAGAAAG
GAAGGGAAGAAAGCGAAAGGAGCGGGCGCTAGGGCGCTGGCAAGTGTAGCGGTACGCTGCGCGT
AACCACCACACCCGCCGCGCTTAATGCGCCGCTACAGGGCGCGTCCATTTCGCCATTCAGGCTGCGC
AACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCTATTACGCCAGCTGGCGAAAGGGGGATGT
GCTGCAAGGCGATTAAGTTGGGTAACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGACGGCC
AGTGAATTGTAATACGACTCACTATAGGGCGAATTGGGCCCCGACGTCGCATGCTCCCGGCCGCCAT
GGCCGCGGGATTTTTCATATCGCTCTGTTATTAGCAGTTTTTGGCCGCTTCATCTTCGTAGTGTGCGC
AGTGTACCTTTTCATTTCGGAACAAACATAATTACTGGAAAGACGATGGATTCCCGTATGCGCCGAA
CCCACATTTTCTGTTTCGGACACGCGAAAGGACAGGCCAGACAAGGCATGCGGCCGACATCCATCT
GGAAGTGTACAAAAAATTCAAGCAGCGCCGTGAGCGGTACGTTGGTGTGAGCCAGTTCATGATACC
TTCATTGCTCGTGATCGATCCAGAGCTGGTGAAAACGATCCTAGTAAAGGACTTTAATGTATTCCAC
GATCATGGTGTATTCAATAATGCAAGAGACGACCCGCTGTCCGCACATCTTTTTGCGCTTGAAGGTA
ACCCATGGCGCTTGTTCGTCAGAAGCTCACGCCAACGTTACCTCAGGTCGCATGAAGCAAATGT
TTGGTACACTATGGGATGTAGCACTTGAGCTGGACAAGTATATGGAAGAAAACATCGTCAGCCGG
ATATTGAGATGAAGGATGTGCTAGGTTCGGTTTACGACAGATGTGATTGGCACCTGTGCATTTCGGGA
TTGAGTGTAAACGCTTAAGACACCCGACTCGGAATTCGCAAAATACGGCAACAAAGCGTTTGAGT
TTAATCTGATGATTATTCTAAAAACTTTCTTAGCATCGGCTTATCCGTCACCTGTGCGGAAACCGCG
AATGAAGATAACATTCGACGATGTGGAACAGTTTTTCTTAAAAATTGTTAAGGAAACGGTAGAATA
TCGAGAAAGTAACAACATTAAACGAAACGACTTCATGAACCTGCTGTTGCAGATTAAGAATAAGGG
TAAGCTGGACGACAGCGATGATGGGAGTGTGGCAAGGGTGAAGTAGGAATGACACAACGAGAAC
TAGCGGCACAGGCATTCATTTTCTTCTTGGCCGGTTTCGAGACATCATCCACGACGCAAAGCTTCTG
TCTGTACGAGTTGGCAAAGAACCCTGACATCCAGGAGCGCCTTAGACAAGGGATCAACCAAGCAAT
CGAGGAGAATGACGGCCAGGTGACGTACGATGTCGCCATAAACATACAGTATCTGGACAATGTGAT
AAACGAAACACTTCGCAAGTACCCACCGGTAGAATCGTTGAGTCGTGTGCCGTCTGTTGACTATGTC
ATCCCAGGTACGAAACATGTCATTCCCAAACGAACGTTAGTGCAAATTCCGGTTCACGCCATTCAA
CATGATCCTGAGCACTGTCCCGATCCAGAACGTTTCGACCCGGATCGCTTCTCACCGAGGAAGTG
AAGAAGCGACATCCCTTCACGTTCTCCCATTCGGTGAGGGGCCACGCGTTTGCAATTGGGCTTCGGT
TTGGTGTGATGCAGACGAAGGTAGGATTGATAACGCTGTTGAGAAAGTTCCGCTTCTCACCGTCAG
CGCGTACACCAGATTGTGTAAAGTTCGATCCGAAAATGATCATTCTGTCACCGATCGCGGGTAATTA
CTTGAAGGTGAAAAAGTTGTAGTCTAGAAAAATCACTAGTGCGGCCGCCTGCAGGTCGACCATAG
GGAGAGCTCCCAACGCGTTGGATGCATAGCTTGAGTATTCTATAGTGTACCTAAATAGCTTGGCGT
AATCATGGTCATAGCTGTTTCTGTGTGAAATTGTTATCCGCTCACAAATCCACACAACATACGAGC
CGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCG
CTCACTGCCCCGCTTTCCAGTCGGGAAACCTGTCTGTGCCAGCTGCATTAATGAATCGGCCAACGCGC
GGGAGAGGCGGTTTTCGCTATTGGGCGCTCTTCGCTTCTCGCTCACTGACTCGCTGCGCTCGGTTCG
TTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGA
ATAACGCAGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAGGCC


PURPLE = 17 α hydroxylase sequence (MALLLAVF) with

GREEN = NdeI restriction site


BLUE = XbaI restriction site

APPENDIX A4 – Kinetic raw data and calculations, graphs and plots


| concentration (uM) | -ve peak one area mAU*min | +ve peak one area mAU*min |
|--------------------|---------------------------|---------------------------|
| 16 | 6.9228 | 5.8531 |
| 8 | 3.3554 | 2.5063 |
| 4 | 1.5817 | 1.2167 |
| 2 | 0.8157 | 0.5937 |
| 1 | 0.4026 | 0.2881 |
| 0.5 | 0.2105 | 0.1436 |



| concentration (uM) | -ve peak two area mAU*min | +ve peak two area mAU*min |
|--------------------|---------------------------|---------------------------|
| 16 | 6.2474 | 5.5257 |
| 8 | 3.022 | 2.4074 |
| 4 | 1.4269 | 1.2107 |
| 2 | 0.7558 | 0.5917 |
| 1 | 0.361 | 0.2818 |
| 0.5 | 0.2222 | 0.1441 |




| concentration (uM) | Combined -ve peaks | combined +ve peaks |
|--------------------|--------------------|--------------------|
| 16 | 6.5851 | 5.6894 |
| 8 | 3.1887 | 2.45685 |
| 4 | 1.5043 | 1.2137 |
| 2 | 0.78575 | 0.5927 |
| 1 | 0.3818 | 0.28495 |
| 0.5 | 0.21635 | 0.14385 |




| concentration (uM) | difference | turnover per min |
|--------------------|------------|------------------|
| 16 | 13.93472 | 2.065275 |
| 8 | 6.090633 | 1.909367 |
| 4 | 3.074011 | 0.925989 |
| 2 | 1.567095 | 0.432905 |
| 1 | 0.820311 | 0.179689 |
| 0.5 | 0.477918 | |

| concentration (uM) | Combined -ve peaks |
|--------------------|--------------------|
| 16 | 6.5851 |
| 8 | 3.1887 |
| 4 | 1.5043 |
| 2 | 0.78575 |
| 1 | 0.3818 |
| 0.5 | 0.21635 |




| concentration (uM) | turnover per min |
|--------------------|------------------|
| 16 | 1.032638 |




| | |
|-----|----------|
| 8 | 1.909367 |
| 4 | 0.925989 |
| 2 | 0.432905 |
| 1 | 0.179689 |
| 0.5 | 0.022082 |

**Excluding
16uM**



Reciprocal plot (i.e. 1/concentration versus 1/turnover per min)



2.806624 Vmax = (1/0.3563)


0.061267 (=0.3563/5.8155)
16.32192 Km (uM) (= 1/0.061267)

Calculated
using
sigmaPlot
11:


| Parameter | Value | StdErr | Dependencies |
|-----------------|-------------|--------|--------------|
| a (Vmax) | 2.28 | 0.25 | 0.950839 |
| b (Km) | 6.90 | 1.30 | 0.950839 |

Second run


| concentration (uM) | -ve peak one area mAU*min | +ve peak one area mAU*min |
|--------------------|---------------------------|---------------------------|
| 16 | 7.9937 | 7.6858 |
| 8 | 4.054 | 3.288 |
| 4 | 2.056 | 1.5012 |
| 2 | 0.9712 | 0.7007 |
| 1 | 0.453 | 0.3539 |
| 0.5 | 0.2402 | 0.1833 |




| concentration (uM) | -ve peak two area mAU*min | +ve peak two area mAU*min |
|--------------------|---------------------------|---------------------------|
| 16 | 7.1672 | 7.255 |
| 8 | 3.647 | 3.1634 |
| 4 | 1.8616 | 1.4482 |
| 2 | 0.8598 | 0.695 |
| 1 | 0.3917 | 0.3273 |
| 0.5 | 0.1941 | 0.1586 |




| concentration (uM) | combined-ve peaks | combined +ve peaks |
|--------------------|-------------------|--------------------|
| 16 | 7.58045 | 7.4704 |
| 8 | 3.8505 | 3.2257 |
| 4 | 1.9588 | 1.4747 |
| 2 | 0.9155 | 0.69785 |
| 1 | 0.42235 | 0.3406 |
| 0.5 | 0.21715 | 0.17095 |




| concentration (uM) | Combined -ve peaks |
|--------------------|--------------------|
| 16 | 7.58045 |
| 8 | 3.8505 |
| 4 | 1.9588 |
| 2 | 0.9155 |
| 1 | 0.42235 |
| 0.5 | 0.21715 |



| concentration (uM) | turnover per min |
|--------------------|------------------|
| 16 | 0.150755 |
| 8 | 1.207721 |
| 4 | 0.881662 |
| 2 | 0.511645 |
| 1 | 0.261225 |
| 0.5 | 0.117184 |






2.392917 Vmax

0.137309

7.282843 Km (uM)

Calculated with sigmaPlot 11:

| Parameter | Value | StdErr | CV(%) | Dependencies |
|-----------------|-------------|--------|-------|--------------|
| a (Vmax) | 2.28 | 0.25 | 10.76 | 0.950839 |
| b (Km) | 6.90 | 1.30 | 18.77 | 0.950839 |



SIGMA PLOT 11 DATA:

| 1-concentration | 2-rate |
|-----------------|--------------------|
| 8 | 1.2077 0.5 |
| 4 | 0.8817 0.5293 |
| 2 | 0.5116 0.5586 |
| 1 | 0.2612 0.5879 |
| 0.5 | 0.1172 0.6172 |

Nonlinear Regression

Data Source: Data 1 in Notebook1

Equation: Hyperbola, Single Rectangular, 2 Parameter

$f = a * x / (b + x)$

| R | Rsq | Adj Rsqr | Standard Error of Estimate |
|----------|------------|-----------------|-----------------------------------|
| 0.9973 | 0.9945 | 0.9927 | 0.0383 |

| | Coefficient | Std. Error | t | P |
|---|--------------------|-------------------|----------|----------|
| a | 2.2823 | 0.2456 | 9.2941 | 0.0026 |
| b | 6.8953 | 1.2945 | 5.3265 | 0.0129 |

Analysis of Variance:

| | DF | SS | MS |
|------------|-----------|-----------|-----------|
| Regression | 2 | 2.5753 | 1.2876 |
| Residual | 3 | 0.0044 | 0.0015 |
| Total | 5 | 2.5797 | 0.5159 |

Corrected for the mean of the observations:

| | DF | SS | MS | F | P |
|------------|-----------|-----------|-----------|----------|----------|
| Regression | 1 | 0.7999 | 0.7999 | 545.7456 | 0.0002 |
| Residual | 3 | 0.0044 | 0.0015 | | |
| Total | 4 | 0.8043 | 0.2011 | | |

Statistical Tests:

Normality Test (Shapiro-Wilk) Passed (P = 0.3531)

W Statistic= 0.8892 Significance Level = 0.0500

Constant Variance Test Passed (P = 0.0500)

Fit Equation Description:

[Variables] x = col(1); y = col(2)

reciprocal_y = 1/abs(y)

reciprocal_ysquare = 1/y^2

[Parameters] a = max(y) "Auto {{previous: 2.28228}}"; b = x50(x,y,0.1) "Auto {{previous: 6.89532}}"

[Equation] f=a*x/(b+x); fit f to y; "fit f to y with weight reciprocal_y; "fit f to y with weight reciprocal_ysquare

[Constraints] [Options] tolerance=1e-10 ; stepsize=1; iterations=200

Number of Iterations Performed = 9

Nonlinear Regression

Data Source: Data 1 in Notebook1

Equation: Hyperbola, Single Rectangular, 2 Parameter

f=a*x/(b+x)

| R | Rsq | Adj Rsqr | Standard Error of Estimate |
|----------|------------|-----------------|-----------------------------------|
| 0.9973 | 0.9945 | 0.9927 | 0.0383 |

| | Coefficient | Std. Error | t | P |
|---|--------------------|-------------------|----------|----------|
| a | 2.2823 | 0.2456 | 9.2941 | 0.0026 |
| b | 6.8953 | 1.2945 | 5.3265 | 0.0129 |

Analysis of Variance:

| | DF | SS | MS |
|------------|-----------|-----------|-----------|
| Regression | 2 | 2.5753 | 1.2876 |
| Residual | 3 | 0.0044 | 0.0015 |
| Total | 5 | 2.5797 | 0.5159 |

Corrected for the mean of the observations:

| | DF | SS | MS | F | P |
|------------|-----------|-----------|-----------|----------|----------|
| Regression | 1 | 0.7999 | 0.7999 | 545.7456 | 0.0002 |
| Residual | 3 | 0.0044 | 0.0015 | | |
| Total | 4 | 0.8043 | 0.2011 | | |

Statistical Tests:**Normality Test (Shapiro-Wilk)**

Passed (P = 0.3531)

W Statistic= 0.8892

Significance Level = 0.0500

Constant Variance Test

Passed (P = 0.0500)

Fit Equation Description:


[Variables] x = col(1); y = col(2); reciprocal_y = 1/abs(y); reciprocal_ysquare = 1/y^2

[Parameters] ; a = max(y) "Auto" { {previous: 2.28228} } ; b = x50(x,y,0.1) "Auto" { {previous: 6.89532} }

[Equation] f=a*x/(b+x) ; fit f to y; "fit f to y with weight reciprocal_y" ; "fit f to y with weight reciprocal_ysquare"

[Constraints] [Options] ; tolerance=1e-10; stepsize=1 ; iterations = 200

Number of Iterations Performed = 9

**Data 2:**

| | | | |
|----|--------|--------|--------|
| 16 | 1.0326 | 0.5 | 0.2025 |
| 8 | 1.9094 | 0.5605 | 0.224 |
| 4 | 0.926 | 0.6211 | 0.245 |
| 2 | 0.4329 | 0.6816 | 0.2654 |
| 1 | 0.1797 | 0.7422 | 0.2853 |

Nonlinear Regression

Data Source: Data 2 in 10Aug_results.JNB

Equation: Hyperbola, Single Rectangular, 2 Parameter

f=a*x/(b+x)

| R | Rsq | Adj Rsqr | Standard Error of Estimate |
|--------|--------|----------|----------------------------|
| 0.8274 | 0.6846 | 0.6058 | 0.4360 |

| | Coefficient | Std. Error | t | P |
|---|-------------|------------|--------|--------|
| a | 1.8390 | 0.8165 | 2.2523 | 0.0874 |
| b | 4.0417 | 4.6439 | 0.8703 | 0.4332 |

Analysis of Variance:

| | DF | SS | MS |
|------------|----|--------|--------|
| Regression | 2 | 5.0293 | 2.5147 |
| Residual | 4 | 0.7603 | 0.1901 |
| Total | 6 | 5.7897 | 0.9649 |

Corrected for the mean of the observations:

| | DF | SS | MS | F | P |
|------------|----|--------|--------|--------|--------|
| Regression | 1 | 1.6503 | 1.6503 | 8.6824 | 0.0421 |
| Residual | 4 | 0.7603 | 0.1901 | | |
| Total | 5 | 2.4107 | 0.4821 | | |

Statistical Tests:

Normality Test (Shapiro-Wilk) Passed (P = 0.0708)

W Statistic= 0.8091 Significance Level = 0.0500

Constant Variance Test Passed (P = 0.0600)

Fit Equation Description:

[Variables]; x = col(1); y = col(2)

reciprocal_y = 1/abs(y)


reciprocal_ysquare = 1/y^2

[Parameters]; a = max(y) "Auto {{previous: 1.83897}}"; b = x50(x,y,0.1) "Auto {{previous: 4.04169}}"

[Equation] ; f=a*x/(b+x) ; fit f to y ; "fit f to y with weight reciprocal_y ;"fit f to y with weight reciprocal_ysquare

[Constraints] [Options] ; tolerance=1e-10 ; stepsize=1; iterations=200

Number of Iterations Performed = 9



Appendix A5: Peptide synthesis and production of monoclonal antibody CYP6P9

A) 2D-Electrophoresis

I. *First dimension: Isoelectric focusing (IEF)*

- Samples (proteins expressed in tested cell lines or purified proteins) were prepared in Urea lysis buffer (*Appendix A5 – 2D ELFO solutions*) and run in IEF on an Immobiline DryStrip, pH 3-10 NL (Amersham-Pharmacia-Biotech).
- Sample (30µl; 100µg of total protein concentration per DryStrip) plus 95µl of reswelling buffer (*Appendix A5 – 2D ELFO solutions*) was loaded and the DryStrip was covered with mineral oil and incubated overnight at room temperature (RT).
- IEF was run under the following conditions: 6min/200V, 100min/1750V, 60min/3500V.

II. *Second dimension: SDS-Polyacrylamide Gel Electrophoresis (SDS-PAGE)*

- The Immobiline DryStrip was equilibrated in SDS equilibration buffer (*Appendix A5 – 2D ELFO solutions*) with addition of 2% DTT:
- For 15 minutes and with addition of 2.5% iodoacetamide and 0.01% bromophenol blue
- For another 7 minutes (*Appendix A5 – 2D ELFO solutions*)
- The proteins from IEF were separated by SDS-PAGE using 7.5%-15% gel. Strips were loaded using 0.5% agarose dissolved in SDS-PAGE running buffer. Separated proteins were transferred onto nitrocellulose membrane in either Bio-Rad Mini Trans-Blot Electrophoretic Transfer Cell for 2hrs (4°C, 150mA) in transfer buffer or in Trans-Blot SD Semi-Dry Transfer-Cell unit (BioRad) for 1hr (RT, 200mA) using transfer buffer.
- The pre-stained molecular weight markers (Bio-Rad) were run in parallel.

2D ELFO solutions:

| | |
|----------------------|-------------|
| Lysis buffer: | 25ml |
| 9 M Urea | 13.5135g |
| 2% NP-40 | 500µl |
| 3% CHAPS | 750µl |
| 70 mM DTT | 270mg |

| | |
|--------------------------------------|--------------------------------|
| <u>Before use:</u> Aprotinin 1 µg/ml | |
| Leupeptin 1 µg/ml | Protease inhibitor kit (Roche) |
| 2 mM Na ₃ VO ₄ | |
| 1mM PMSF | 200µl |
| 25mM NaF | 26.24mg |
| 2% Ampholyte pH 9-11 | 20 µl/1ml buffer |

| | |
|--|------------------|
| Reswelling buffer for IEF strips: | 25 ml |
| 7M Urea | 10.5105g |
| 0.5% Triton X-100 | 125µl |
| 10mM DTT | 38.55mg |
| 2M Thiourea | 3.806g |
| 4% CHAPS | 1g |
| 0.003% Bromphenol Blue | 150µl (0.5%) |
| <u>Add before use:</u> 1 % Ampholyte pH 3-10 | 10 µl/1ml buffer |

| | |
|---|--------------|
| SDS Equilibration buffer after IEF (5 ml for 1 strip): | 100ml |
| 50mM Tris, pH 6.8 | 5ml (1M) |
| 6M Urea | 36g |
| 30% Glycerol | 30ml |
| 2% SDS | 2g |

Incubations:

- A. Add 2% DTT (0.2g/10ml) and incubate 15min/RT.
- B) Add 2.5% iodoacetamide (0.25g/10ml) and 0.01% bromophenol blue (200µl, 0.5%/10ml) and incubate 7min/RT.

| | |
|----------------------------|--------------|
| SDS stock solution: | 500ml |
| 20% SDS | 100g |

| | |
|------------------------------|--------------|
| Acrylamide mix (30%): | 100ml |
| 30% Acrylamide | 30g |
| 0.8% Bis-acrylamide | 0.8g |

| | |
|----------------------|--------------|
| Tris Buffers: | 100ml |
| 1M Tris-HCl, pH 8.8 | 12.1g |
| 1M Tris-HCl, pH 6.8 | 12.1g |

| | | | | |
|----------------------------|-------------|------------|--------------|------------|
| Running gel (30ml): | 7.5% | 10% | 12.5% | 15% |
| Acrylamide mix (30%) | 7.5ml | 10ml | 12.5ml | 15ml |
| 1M Tris-HCl pH 8.8 | 11.2ml | 11.2ml | 11.2ml | 11.2ml |
| 20% SDS | 0.15ml | 0.15ml | 0.15ml | 0.15ml |
| Deionized H ₂ O | 13.7ml | 11.2ml | 8.7ml | 3.7ml |

Add before use: 80µl of 10% ammonium persulfate (APS) and 20µl of TEMED

| | | |
|-----------------------------|-----------|-----------|
| Stacking gel (30ml): | 5% | 3% |
| Acrylamide mix (30%) | 1.67ml | 1.0ml |
| 1 M Tris-HCl pH 6.8 | 1.25ml | 1.25ml |
| 20% SDS | 0.05ml | 0.05ml |
| Deionized H ₂ O | 7.03ml | 7.7ml |

Add before use: 50µl of 10% ammonium persulfate (APS) and 10µl of TEMED

B) ELISA assay (Two captured assay)

- A 96-well microtitre plate (Becton Dickinson - FALCON 3912) was incubated overnight at RT with 50µl per well of 10-20µg/ml Ig of purified antibody (monoclonal or polyclonal) diluted in buffer E1 (*Appendix A5 – ELISA buffers*).
- The plate was rinsed in PBS (1X) and blocked for 2hrs in 3% bovine serum albumin (BSA, SIGMA, A-7030) in PBS (200µl per well, RT).
- A washing step: PBS (1X), PBS + 0.1% NP-40 (2X), PBS (1X) followed. The extract (antigen) was added – 50µl of serial two-fold dilutions either in lysis buffer or PBS per well (minimal total protein concentration 1 mg/ml) and allowed to incubate for 2hrs at 4°C.
- The washing step: PBS (1X), PBS + 0.1% NP-40 (2X), PBS (1X) followed.
- Antibody was added (50µl of 1µg/ml of antibody per well) (if coated by monoclonal add polyclonal; if coated by polyclonal add monoclonal) diluted in 1% BSA in PBS and incubated for 2hrs at 4°C.
- The washing step as before then followed.
- Peroxidase-conjugated swine antiserum was added to rabbit immunoglobulin or rabbit antiserum was added to mouse immunoglobulin (DAKO), diluted 1:1000 in 1% BSA in PBS; 50µl per well. This was left to incubate for 1hr at 4°C, followed by the wash step as before.
- Visualization was carried out with tetramethylbenzidine: 6ml of buffer E2 (*Appendix A5 – ELISA buffers*) + 60µl of solution E3 (*Appendix A5 – ELISA buffers*) + 15µl of H₂O₂ - aliquot 50µl per well. This was left for 20-30 min and the reaction stopped with 1M H₂SO₄ (50µl per well).
- The results were monitored – O.D. 450nm.

ELISA Buffers

Buffer E1: Buffer for coating plates (340ml):

0.1M CO_3/HCO_3 pH 9-9.8

0.42g Na_2CO_3 (SIGMA)

2.52g NaHCO_3 (SIGMA)

Buffer E2: Substrate buffer; 0.1M citrate/phosphate, pH 6.0 (650ml):

7.05g Na_2HPO_4 (SIGMA)

3.15g citric acid (SIGMA)

Buffer E3: Substrate: 3,3', 5,5'– tetramethylbenzidine (TMB: SIGMA – T-2885)

10mg/ml TMB in DMSO (SIGMA)

C) Immunization of the mice

Mice BALB/C was used for the immunization protocol.

For each protein 5 mice were used.

Immunization protocol:

- 1) 1st injection – **day 1**: intraperitoneal injection of 10–50µg of purified protein or peptide coupled to KLH (in complete Freund's adjuvants).
- 2) 2nd injection – **day 21**: intraperitoneal injection of 10–50µg of purified protein or peptide coupled to KLH (in incomplete Freund's adjuvants).
- 3) 3rd injection – **day 42**: same as day 21
- 4) 4th injection – **day 45**: intraperitoneal injection of 10µg of purified protein or peptide coupled to KLH (in PBS).
- 5) 5th injection – **day 46**: same as day 45
- 6) Fusion – **day 47** (*production of monoclonal antibodies*)

D) Affinity purification of mouse sera on peptide used as immunogen

Day 1: Pierce 6x reactigel beads (Cat. No: 20259) (0.5ml) were washed with 0.1M Sodium Borate buffer (pH 9.0) in an eppendorf tube (to exchange the acetone). Peptide (5mg or 1mg - minimum) was added to the beads and borate buffer was added to make up a final volume of 1.5ml. The peptide and beads were allowed to mix overnight in a wheel at RT.

Day 2: The beads were spun down, supernatant removed and 1ml of 1M Tris HCl pH 8.8 was added and allowed to mix throughout the day. Residual peptide was washed away very well [about 10-15 washes in PBS-Tween-20 (0.1%) depending upon the volume of each wash] and the beads (containing 1-5mg of coupled peptide) were added to a 50ml tube containing up to 50ml of serum. This was allowed to mix overnight.

Day 3: The beads were spun and serum saved. Serum proteins were washed away with PBS-Tween-20 (0.1%) and the beads were transferred to an eppendorf tube and 1ml of 0.1M Glycine (pH 2.5) was then added to it to elute the IgG. Beads were mixed in 0.1M Glycine (pH 2.5) for 30min. The supernatant (1ml) was removed and 0.15ml of 1.5M Tris-HCl pH 8.8 was added to it to neutralise the acid. Another 1ml of 0.1M Glycine (pH 2.5) was added to the beads and mixed for 30min, and this step was repeated another three times. A 10% or 12% SDS gel with 10µl of each fraction was run to see which fractions the IgG came off in (generally fractions 1, 2, and 3). The Ig concentration was measured using Bio-Rad Protein Assay (to do standard curve use bovine gamma globulin).

Note: The beads can be reused - Reapply the 1x used serum to the beads and repeat the purification steps to see if you adsorbed all the IgG from the serum.

E) Mass spec (MS) and HPLC analysis for sample CVK 70301 (CVKFDPKMIL)




Figure E-1: MALDI-TOF MS analysis report showing purity of sample CVK 70301 (Sequence CVKFDPKMIL) >86%. Y-axis = atomic mass units (amu); X-axis – mass/charge ratio (m/z). MH⁺ (theoretical) = 1306.738. In MS analysis, ions generally have a single charge, therefore the m/z value is equivalent to mass itself. Each bar represents an ion having a specific m/z ratio and the length of the bar indicates the relative abundance of the ion. The highest mass = the molecular ion; the lower mass ions = fragments from the molecular ion assuming the sample is a single compound.




Figure E-2: HPLC Analysis of sample CVK 70301. Injection volume was 100 μ l, with UV detection at Ch1/220nm. Sample was run over a time period of 40min. Peak 6 showed a retention time of 23.585min, Area = 5093407, height = 662588, Area % = 86.001 and height % = 83.843.

Appendix A6: Immunizations and dot blots




Figure A6-1: Mouse sera to CVK after 2nd immunization dot blot showing mouse number versus dilution to CVK.



Figure A6-2: Mouse sera to CVK after 3rd immunization dot blot showing mouse number versus dilution to CVK.




Figure A6-3: Supernatants of fusion CVK dot blot. 8B2 shows the strongest fusion reaction. 5D1, 5G10, 8B2, 8B5, 8B8 and 8C11 were selected for use in the final hybridoma analysis.

| Specific IgG detection | | | | Positive hybridomas | | |
|------------------------|-----|-----|-----|---------------------|-------------------------------|--------------------|
| 8B2 | CVK | CVK | CVK | Positive hybridomas | Positive clones after cloning | Name of the clones |
| 5G10 | | | | 5D1 | 5D1 / E6 | CVK 1.1 |
| 8B5 | | | | | 5D1 / F7 | CVK 1.2 |
| 8B8 | | | | 8B8 | 8B8 / D5 | CVK 2.1 |
| 5D1 | | | | | 8B8 / D7 | CVK 2.2 |
| 8C11 | | | | | | |
| | | | | 8B5 | 8B5 / D4 | CVK 5.1 |
| | | | | | | CVK 5.2 |
| | | | | 8C11 | 8C11 / B10 | CVK 6.1 |
| | | | | | | CVK 6.2 |
| | | | | 8B2 | 8B2 / E6 | CVK 3.1 |
| | | | | | 8B2 F8 | CVK 3.2 |
| | | | | 5G10 | 5G10 / E12 | CVK 4.1 |
| | | | | | | CVK 4.2 |

Figure A6-4: CVK 3.1 from hybridoma 8B2/E6 was the only positive hybridoma for IgG1, Kappa. The others (CVK 1.1, CVK 2.1, CVK 4.1, CVK 5.1 and CVK 6.1 are all IgM, Kappa).

Appendix A7: Other buffers and reagents used in this study

10X MOPS running buffer (RNA gel)

0.4M MOPS

0.1M sodium acetate

0.01M EDTA (disodium)

Adjust pH to 7.0 using NaOH or Glacial Acetic acid CH_3COOH

10X loading RNA buffer Recipe

0.35% (w/v) Orange G

30% (w/v) Ficoll 400

1mM EDTA (disodium)

For electrophoresis buffer: dilute 10X running buffer to make 1X running buffer

RNA agarose gel (1%, 20ml), prepared with 1X running buffer and 7% formaldehyde:

0.2g agarose

2ml 10X running buffer

14.2ml ultrapure water

Microwave for 45 seconds; then allow to cool. Add 3.8ml of 37% formaldehyde and then pour into casting tray with combs.

To prepare RNA for loading onto wells combine the following:

3.5 μl of each RNA sample with:

10 μl formamide

3.5 μl 37 % formaldehyde

1 μl 10X running buffer

2 μl 10X loading buffer

Denature RNA in buffers for 10 minutes at 65-75 °C. Then load into gel wells.

LB broth for ligations and transformations:

Liquid broth

2.5 g Bacto-Tryptone

1.25 g Bacto-yeast extract

1.25 g NaCl

Make up to 250 ml with dH₂O

For Solid Broth, mix the same as for Liquid Broth, but include 3.75g Agar (1.5%) before making up with dH₂O.

Liquid and solid broths are then autoclaved, allowed to cool and then:

To the Solid broth, 100 µg/ml Ampicillin is added, and

To the Liquid broth 50 µg/ml Ampicillin is added.

SDS-PAGE (Laemmli) Buffer system

Stock solutions and buffers

1. **Bis/Acrylamide (30% T, 2.67% C)**

2. **10% (w/v) SDS**

Dissolve 10g SDS in 90ml water with gentle stirring and bring to 100ml with deionized water. Alternatively, 10% SDS solution (250ml) can be used (Bio-Rad cat no: 161-0416).

3. **1.5M Tris-HCl, pH 8.8 (resolving gel buffer)**

27.23g Tris base (18.15g/100ml)

80ml deionised water

Adjust to pH 8.8 with 6N HCl. Bring total volume to 150ml with deionized water and store at 4°C.

4. **0.5M Tris-HCl, pH 6.8 (Stacking gel buffer)**

6g Tris base

50ml deionised water

Adjust to pH 6.8 with 6N HCl. Bring total volume to 150ml with deionized water and store at 4°C.

5. **Laemmli 2x sample buffer (mixed 1:1)**

3.55ml deionized water

1.25ml 0.5M Tris-HCl, pH 6.8

2.5ml glycerol

2.0ml 10% (w/v) SDS

0.2ml 0.5% (w/v) bromophenol blue

9.5ml total volume

Store at room temperature

Before use: add 50µl β-mercaptoethanol to 950µl sample buffer. Dilute the sample at least 1:2 with sample buffer and heat at 95°C for 4 minutes.

6. **10X Electrode (Running) buffer, pH 8.3 (makes 1L)**

30.3g Tris-base

144.0g glycine

10.0g SDS

Dissolve and bring total volume up to 1L with deionized water. Do not adjust pH with acid or base. Store at 4°C.

Before use: dilute 50ml of 10X stock with 450ml deionized water for each electrophoresis run. Mix thoroughly.

7. **10% (w/v) APS (fresh daily)**

100mg ammonium persulphate

Dissolve in 1ml of deionized water.

8. **Gel formulations (10ml)**

- a. Prepare the monomer solution by mixing all reagents except the TEMED and 10% APS. Degas the mixture for 15 minutes.

| Percent gel | DDI H ₂ O (ml) | 30% Degassed Acrylamide/Bis (ml) | Gel buffer * (ml) | 10% w/v SDS (ml) |
|-------------|---------------------------|----------------------------------|-------------------|------------------|
| 5% | 5.7 | 1.7 | 2.5 | 0.1 |
| 12% | 3.4 | 4.0 | 2.5 | 0.1 |
| 15% | 2.4 | 5.0 | 2.5 | 0.1 |

* Resolving gel buffer – 1.5M Tris-HCl, pH 6.8

* Stacking gel buffer – 0.5M Tris-HCl, pH 8.8

- b. Immediately prior to pouring gel, add: (*For 10ml monomer solution*)

Resolving gel: 50µl 10% APS and 5µl TEMED

Stacking gel: 50µl 10% APS and 10µl TEMED

9. **Transfer buffer (1L)**

25mM Tris-base

190mM glycine

20% methanol

Store at 4°C

10. **Coomassie Stain (1L)**

1g Coomassie R250

100ml glacial acetic acid

400ml methanol

500ml ddH₂O

Store at room temperature in a sealable container

11. **De-Stain for Coomassie (1L)**

200ml methanol

100ml glacial acetic acid

700ml ddH₂O