

TO ESTABLISH NORMATIVE DATA USING
THE JEBSEN TAYLOR HAND FUNCTION
TEST (JHFT) FOR NORMAL, ETHNICALLY
DIVERSE SOUTH AFRICANS AGED
BETWEEN 20 AND 59 YEARS.

Paran Govender

A research report submitted to the Faculty of
Health Sciences, University of the
Witwatersrand, Johannesburg, in partial
fulfilment of the requirements for the degree
of
Master of Science in Occupational Therapy

Johannesburg, 2008

I, Paran Govender, declare that this research report is my own work. It is being submitted for the degree of Master of Science in Occupational Therapy in the University of Witwatersrand, Johannesburg. It has not been submitted before for any degree of examination at this or any other university.



10th June 2008

For my parents

Abstract

The use of outcome measures by occupational therapists to establish the effectiveness of treatment is important. The Jebsen-Taylor Hand Function Test (JHFT) was identified as a standardised assessment suitable for the evaluation of outcomes related to the treatment of hand function. This study established norms for this test on 120 normal, ethnically diverse South Africans between the ages of 20 and 59 years.

Statistically significant differences were found between the original norms published by Jebsen et al ⁽¹⁾ and the South African sample for males and females, dominant and non-dominant hands and ethnic groupings. It is postulated that gender and cultural factors may influence the speed with which different groups complete functional tasks and these must be factored in when using the JHFT as an outcome measure in South Africa. The test appears to have limited application as a norm referenced test in South Africa, but may be valuable in determining progress in therapy or research.

Acknowledgements

Thank you to all who have helped make this work possible and especially to Denise, Mags, Rahila and James.

Table of Contents

	Page
Title	i
Declaration	ii
Dedication	iii
Abstract	iv
Acknowledgements	v
Table of Contents	vi
List of Tables	ix
List of Figures	x
Operational Definitions and Abbreviations	xi
 <u>Chapter 1 Introduction</u>	
1.1 Introduction	1
1.2 Statement of the Problem	3
1.3 Purpose of this Study	4
1.4 Aim of this study	4
1.4.1 Objectives of this Study	4
1.5 Justification of the Study	4
 <u>Chapter 2 Review of Literature</u>	
2.1 Importance of Standardised Tests of Hand Function	6
2.2 Evaluation of the Jebsen-Taylor Hand Function Test	9
2.2.1 Clinical Utility	9
2.2.1.1 Clinical Applicability	9
2.2.1.2 Specificity	10
2.2.1.3 Availability	10
2.2.1.4 Time Demands	11
2.2.1.5 Acceptability to clients	11
2.2.2 Standardisation of the Jebsen-Taylor Hand Function Test	11
2.2.3 Purpose of the Test	12
2.2.4 Psychometric Properties	13
2.2.4.1 Item Construction	13
2.2.4.2 Reliability	13
2.2.4.3 Validity	13
2.2.4.4 Norms	14
2.2.5 The Patient's Perspective	15
2.3 Limitations of the Jebsen-Taylor Hand Function Test	16
2.4 The Value of the Test as an Outcome Measure in Research	16
2.5 Factors Affecting Hand Function	19
2.5.1 Genetic Factors	19
2.5.2 Dominance	19
2.5.3 Culture and hand Preference	20
2.5.4 Gender and Age	20

Chapter 3 Methodology

3.1 Research Design	22
3.2 Subject Selection	22
3.2.1 Inclusion criteria	23
3.2.2 Exclusion criteria	23
3.2.3 Sample size	24
3.3 Measurement Instrument	24
3.3.1 Pilot Study of inter tester and intra tester reliability	24
3.4 Research Procedure	25
3.5 Data Collection	27
3.6 Ethical Considerations	28
3.7 Data Analysis	28

Chapter 4 Results

4.1 Demographics	30
4.1.1 Age	30
4.1.2 Handedness	31
4.2 Differences in results between genders	32
4.3 Comparison of Test results by dominant hand and non dominant hand	33
4.3.1 Comparison to Jebesen et al's means for the dominant hand	34
4.3.2 Comparison to Jebesen et al's means for the non-dominant hand	35
4.3.3 Comparison to Jebesen et al's means using Confidence Intervals	36
4.4 Comparison of test results by population group	37
4.5 Summary of results	40

Chapter 5 Discussion

5.1 Comparison to the Jebesen et al Study	41
5.2 Gender	42
5.3 Genetic Factors	44
5.4 Dominant and Non Dominant hand	44
5.5 Ethnicity	44
5.6 The Use of the Jebesen-Taylor hand Function Test in South Africa	47
5.6.1 Clinical utility	47
5.6.2 Standardisation	48
5.6.3 The patient's perspective	49

Chapter 6 Conclusion

References	54
-------------------	----

Appendices

A. Method of Administration of the JHFT	62
B. Interrater reliability - Mean times and Standard deviations for normal subjects	67

C. Data Collection Form	68
D. Ethics Approval	69
E. Subject Information and Informed Consent Form	70
F. Bonferroni Statistically Significant Results	71
G. Results of ANOVA	72
H. Results of Bonferroni Correction	74

List of Tables

		Page
Table 1	Age demographics of Sample according to Population Group and Gender	30
Table 2.	Handedness in the Sample and Population Groups	31
Table 3.	Composite findings for all genders across the items on the Jebsen-Taylor Hand Function Test	32
Table 4.	Norms for the South African Sample for the Jebsen-Taylor Hand Function Test	33
Table 5.	Confidence Intervals for the means found by Jebsen et al and the means in this study	37
Table 6.	Composite results for all population groups across the items on the Jebsen-Taylor Hand Function Test	38
Table 7.	Composite results for Sample of all Population Groups for all subtests	39
Table 8	Statistically significant results post Bonferroni Correction	71
Table 9	ANOVA for Sample Population	72
Table 10	Post Hoc Tests for Sample Population	74

List of Figures

		Page
Figure 1	Comparison of means between this study and Jebsen et al's	
34	means for females dominant hand	
Figure 2	Comparison of means between this study and Jebsen et al's	
35	means for males dominant hand	
Figure 3	Comparison of means between this study and Jebsen et al's	
35	means for females non dominant hand	
Figure 4	Comparison of means between this study and Jebsen et al's	
36	means for males non dominant hand	
Figure 5	Jebsen-Taylor Hand Function Test	
62		

OPERATIONAL DEFINITIONS

Hand Function

- the ability of a patient to use the hands effectively in everyday activity, dependent on anatomical integrity, mobility, muscle strength , sensation and coordination ⁽¹⁾.

Standardised Tests

- are tests given to a group in a similar setting under similar diagnoses in order to determine and evaluate against a "norm"⁽³³⁾

Norm referenced

- norms against which a patients performance can be measured ⁽¹⁾.

Functional limitation

- restriction or lack of ability to perform an action in the manner or within the range consistent with the purpose of an organ and organ system ⁽²⁾ .

ABBREVIATIONS

ADL	Activities of daily living
DASH	The Disability of Arm, Shoulder, and Hand questionnaire
FDT	Functional Dexterity Test
GP	general practitioner
ICF	International Classification of Functioning, Disability and Health.
JHFT	Jebsen-Taylor Hand Function Test. – In the original article by Jebsen et al ⁽¹⁾ the test was not named. The name of the test varies in the literature published since then and abbreviations for the test vary and are reflected as JTHFT ^(12, 39) , JT-HF ⁽⁸⁷⁾ with Webster ⁽⁸⁸⁾ abbreviating the test to JTT. The abbreviation JHFT was adopted for use in this study ^(10, 16) .
NYPDG	New York Presbyterian glove
OT	Occupational Therapy/ist
PEDI	The Paediatric Evaluation of Disability Inventory
SWM	Semmes Weinstein monofilaments
Wee FIM	Wee Functional Independence Measure for Childre