

https://dspace.org/

Identification and characterisation of bacteria isolated from a platinum mine

Sodi, Koketso

2023

https://hdl.handle.net/10539/35667

Downloaded from a DSpace repository. https://dspace.org/



Identification and characterisation of bacteria isolated from a platinum mine

by

Koketso Sodi

(1354286)

Dissertation

Submitted in fulfilment of the requirements for the degree

Master of Science

in

Molecular and Cell Biology

in the Faculty of Science, University of the Witwatersrand, Johannesburg, South Africa

Supervisor: Karl Rumbold

Co-supervisor: Sanchia Moodley

November 2022

DECLARATION

I, Koketso Elaine Sodi am a student registered for the degree of Master of Science (Dissertation) in the academic year 2022.

I hereby declare the following:

- I am aware that plagiarism (the use of someone else's work without their permission and/or without acknowledging the original source) is wrong.
- I confirm that the research proposal submitted for assessment for the above degree is my own unaided work except where explicitly indicated otherwise and acknowledged.
- I have followed the required conventions in referencing the thoughts and ideas of others.
- I understand that the University of the Witwatersrand may take disciplinary action against me if there is a belief that this is not my own unaided work or that I have failed to acknowledge the source of the ideas or words in my writing.

Signed at: Randburg Date: 04/11/2022 Signature: 

ABSTRACT

Bioleaching involves the solubilisation of metals from solid insoluble substrates such as minerals using microorganisms. Bacteria isolated from mines could have properties that are desirable in the metal bioleaching field. These include iron and sulphur-oxidation and organic acid production. The aim of this study was to isolate, identify and characterise bacteria from a platinum mine. Bacteria were isolated from the concentrator plant of a platinum mine and enumerated using traditional microbiological techniques. The bacteria were then identified using Sanger sequencing and NCBI-BLAST. The colony morphology of the bacteria was noted, and the cellular morphology of the bacteria was analysed using gram staining. To find the optimum growth media for the bacteria, growth curves were conducted in Luria-Bertani, nutrient and 9K broth. Organic acid production by the bacteria was analysed using high-performance liquid chromatography. The results showed that the isolated indigenous bacteria grew well in soils with a pH between 6-9. The colony-forming units of bacteria from each sampling site showed that there was a substantial abundance of bacteria in and around the concentrator plant. A total of 50 bacterial isolates were identified and Proteobacteria was the dominant bacterial phylum. Six bacterial isolates namely *Bacillus* sp., *Pseudoxanthomonas mexicana*, *Pseudomonas stutzeri*, *Arthrobacter globiformis*, *Pseudomonas aeruginosa* and *Pseudomonas brassicacearum* had desirable bioleaching characteristics and were selected for further characterisation. The bacteria were able to produce gluconic, citric, lactic, propionic, butyric, formic, and indole-3-acetic acid. Therefore, they can potentially be used for the bioleaching of platinum group metals.

DEDICATION

Dedicated to my family and friends for being my rock.

ACKNOWLEDGEMENTS

I would like to thank my supervisor, Prof Karl Rumbold, for taking me under his wing and guiding me through my masters.

I would also like to thank my co-supervisor, Dr Sanchia Moodley, for her advice, guidance, and support. Thank you for encouraging me and helping me believe in myself. I am eternally grateful.

To Michael Tobin and Naledi Nkabinde, our lab technicians, thank you for your advice and assistance throughout this project. I would like to express my gratitude to Matshidiso Tsotetsi, Letta Ntuli and Motlatsi Phali for their patience and assistance.

A special thank you to my colleagues in the BIORECOVER project, Prof Ndlovu, Dr Yah, Dr Chipise and Alan Shemi for their assistance throughout this project.

To my Industrial Microbiology and Biotechnology Lab colleagues, thank you so much for your help and motivation throughout my project. Kiara, thank you for your friendship, for your advice and the countless phone calls, you have made my MSc journey a whole lot brighter.

To my family and friends thank you for your love, support, and encouragement throughout this journey. None of this would have been possible without you. To Melony Groening, you have supported me in more ways than I can count, thank you for your love, advice, and encouragement, I will forever be grateful. A special thank you to Siwelile Mkhwanazi for allowing me to stay at your place while I completed my dissertation.

To my mom, uncle, and aunt, who have always been guiding and cheering me on from above, thank you. I love and miss you.

Above all, to God, thank you for your grace, guidance, strength, and peace that helped me through this journey.

My sincere gratitude to the Nation Research Foundation (NRF) for providing me with funding.

CONTENTS

DECLARATION ii

ABSTRACT iii

DEDICATION iv

ACKNOWLEDGEMENTS v

CONTENTS vi

LIST OF FIGURES ix

LIST OF TABLES x

LIST OF SYMBOLS xi

LIST OF ABBREVIATIONS xii

CHAPTER ONE: INTRODUCTION 1

1.1 Introduction 2

1.1.1 General introduction and problem statement 2

1.1.2 Research problem 4

1.1.3 Research aims and objectives 4

CHAPTER 2: LITERATURE REVIEW 5

2.1 The platinum group metals mining industry 6

2.1.1 Platinum group metals 6

2.1.2 Current platinum group mining methods: An overview 7

2.1.3 An overview of the platinum group metal concentrate process 9

2.2 Bacterial diversity in mines 9

2.2.1 Introduction 9

2.2.2 Bacteria found at mining sites 10

2.2.3 Bacteria isolated from platinum mines 13

2.3 Bacterial Bioleaching 14

2.3.1 Introduction 14

2.3.2 Bacteria used in bioleaching 14

2.3.2.1 Chemolithotrophic bacteria..... 14

2.3.2.2 Heterotrophic bacteria..... 14
2.3.2.3 Mesophilic bacteria 15
2.3.3 Mechanisms of bioleaching 15
2.3.3.1 Redoxylysis 16
2.3.3.2 Organic acids 17
2.4 Bacterial bioleaching of platinum group metal ores..... 18
CHAPTER THREE: IDENTIFICATION OF BACTERIA 20
3.1 Introduction 21
3.2 Methods and Materials 22
3.2.1 Sample collection 22
3.2.2 Determination of sample pH 22
3.2.3 Isolation of bacteria 22
3.2.4 Genomic DNA extraction 23
3.2.5 DNA sequencing 23
3.3 Results and Discussion 24
3.3.1 Bacterial enumeration 24
3.3.2 Identification of bacterial isolates 26
3.4 Summary and conclusions 31
CHAPTER FOUR: CHARACTERISATION OF BACTERIA 33
4.1 Introduction 34
4.2 Materials and Methods 35
4.2.1 Characterisation of cellular and colony morphology 35
4.2.2 Growth curves 35
4.2.3 High-performance liquid chromatography 37
4.2.4 Statistical analysis. 38
4.3 Results and Discussion 39
4.3.1 Phylogeny and Colony and cellular morphology 39
4.3.2 Growth curves 42

4.3.3 Organic acid production..... 49
4.4 Summary and conclusions 53
CHAPTER 5: CONCLUSIONS AND RECOMMENDATIONS 55
5.1 Conclusions 56
5.2 Recommendations 57
APPENDIX A 58
A1 Preparation of media used in the isolation and enumeration of bacterial species. 58
A1.1 Luria-Bertani Agar Medium 58
A1.2 9K Agar Medium 58
A2 Preparation of culture medium used for the characterisation of the six bacterial isolates. 59
A2.1 Luria-Bertani Agar Medium 59
A2.2 Nutrient Broth (Merck) 59
A2.3 9K Broth Medium 59
APPENDIX B 61
B1 DNA Extraction Protocol 61
REFERENCES 63

LIST OF FIGURES

Figure 2.1: A general overview of PGM ore processing…….....8

Figure 2.2: Bacterial and Archaeal diversity in terms of phyla found at mine sites…….10

Figure 2.3: Bacterial mechanisms of bioleaching…….16

Figure 2.4: Overview of metal solubilisation by organic acids through acidolysis and complexolysis……..18

Figure 4.1: Phylogenetic analysis based on a partial sequence of the 16S rRNA gene inferred using the maximum likelihood method, showing the relationship of the six bacterial species and bacterial isolates of the same species.

Figure 4.2: Morphological characteristics shown by the six bacterial isolates.….42

Figure 4.3: Growth curves of A) *Arthrobacter globiformis* and B) *Pseudoxanthomonas mexicana* grown in 9K, LB and NB broth……..43

Figure 4.4: Growth curves of A) *Pseudomonas brassicacearum* and B) *Bacillus sp.* C) *Pseudomonas aeruginosa* grown in 9K, LB and NB broth…….44

Figure 4.5: Growth curves of A) *Pseudomonas stutzeri* and B) mixed culture grown in 9K, LB and NB broth……..…….....45

Figure 4.6: Optical density (600 nm) of *Bacillus* sp., *Pseudoxanthomonas mexicana*, *Pseudomonas stutzeri*, *Arthrobacter globiformis*, *Pseudomonas aeruginosa*, *Pseudomonas brassicacearum* and the mixed culture grown in nutrient broth…….…….48

Figure 4.7: The colony forming units (CFUs) of *Bacillus* sp., *Pseudoxanthomonas mexicana*, *Pseudomonas stutzeri*, *Arthrobacter globiformis*, *Pseudomonas aeruginosa*, *Pseudomonas brassicacearum* grown in nutrient agar…….49

LIST OF TABLES

Table 2.1: Summary of the bacteria isolated from copper, gold, and coal mines..12

Table 2.2: Bacteria isolated from platinum mining sites……...13

Table 3.1: Universal primer sets used for PCR amplification……...….23

Table 3.2: Sampling sites within the Anglo Platinum Mogalakwena mine in Limpopo South Africa and the pH and colony forming units per gram/ milliliter of each sample. ……..24

Table 3.3: Identity of bacteria and GenBank Accession numbers of closely related species or species with highest sequence similarity percentage…….....26

Table 4.1: Six bacterial isolates isolated from a platinum mine, the GenBank Accession numbers of closely related species or species with the highest sequence similarity percentage and studies showing their organic acid production and iron and sulphur oxidation capabilities. ……36

Table 4.2: Morphological characteristics of the bacterial isolates when grown on LB agar……...38

Table 4.3: The growth rates of *Arthrobacter globiformis, Pseudoxanthomonas mexicana, Pseudomonas stutzeri, Pseudomonas brassicacearum, Bacillus sp., Pseudomonas aeruginosa* and mixed culture grown in LB and NB media……....46

Table 4.4: Organic acid production of the six bacterial isolates…….50

LIST OF SYMBOLS

°C	Degree Celsius
<	Less than
(-)	Negative
%	Percent
(+)	Positive

LIST OF ABBREVIATIONS

2,4-DAPG	2,4-Diacetylphloroglucinol
$(NH_4)_2SO_4$	Ammonium sulphate
ANOVA	Analysis of variance
AAS	Atomic Absorption Spectroscopy
BLAST	Basic Local Alignment Search Tool
$Ca(NO_3)_2$	Calcium nitrate
CFU	Colony-forming units
CFU/g	Colony-forming units per gram
CFU/mL	Colony-forming units per millilitre
DNA	Deoxyribonucleic acid
K_2HPO_4	Dipotassium hydrogen phosphate
EPS	Extracellular polysaccharide
Fe^{3+}	Ferric iron
Fe^{2+}	Ferrous iron
i.e.,	For example
g	gram
g/L	Grams per litre
Cr(VI)	Hexavalent chromium
HPLC	High-Performance Liquid Chromatography
H	Hydrogen
ID	Identity
IAA	Indole-3-acetic acid
IMBL	Industrial Microbiology and Biotechnology
Ir	Iridium
FeS_2	Iron disulphide
LB	Luria-Bertani
$MgSO_4$	Magnesium sulphate
µL	microlitres
µm	micromoles
μhr^{-1}	Micros per hour
mg/mL	Milligrams per millilitre
mL	millilitre
mM	millimolar
M	Molar
ng/g	Nanograms per gram
NCBI	National Center for Biotechnology
NGS	Next Generation Sequencing
NB	Nutrient Broth
100X	One hundred times
OD_{600}	Optical density at 600 nanometres wavelength
Os	Osmium
O_2	Oxygen
Pd	Palladium
Pt	Platinum

PGM	Platinum Group Metal
PCR	Polymerase Chain Reaction
KCl	Potassium chloride
p	Probability
rpm	Revolutions per minute
Rh	Rhodium
rDNA	Ribosomal Deoxyribonucleic acid
rRNA	Ribosomal ribonucleic acid
Ru	Ruthenium
NaCl	Sodium chloride
SO_4	Sulphate
H_2SO_4	Sulphuric acid
UG2	Upper Ground 2
H_2O	Water

CHAPTER ONE: INTRODUCTION

1.1 Introduction

1.1.1 General introduction and problem statement

Platinum group metals (PGM's) are a group of scarce metals that includes platinum, iridium, palladium, osmium, rhodium, and ruthenium. The Circular Economy of the European Union has declared that platinum is a scarce raw material (Keersemaker, 2020). There is a high demand for PGMs due to their desirable properties and applications in several industries, including the transportation and energy industries, such as for vehicle catalytic converters and fuel cells. An increase in the demand for more sustainable and environmentally friendly technologies in both the transportation and energy industries has also resulted in high demand for PGMs. These properties include thermal stability, resistance to corrosion and chemical inactivity (Granados-Fernández *et al.*, 2021). PGMs, more especially platinum, have a distinctive catalytical property that plays an important role in chemical and electrochemical processes that are important in catalytic converters found in vehicles (Yousif, 2019).

In the earth, platinum occurs at an average concentration of 0.4 ng/g and very few large economically prospective deposits have been identified, making it extremely rare (Reith *et al.*, 2014). Due to the high demand and rarity of platinum, it has become a commodity for several countries. The main natural deposits of PGMs are found in South Africa, Zimbabwe, and Russia (Oberthür, 2018), with South Africa having one of the largest PGM's industries in the world. The most commonly used metallurgical process for the recovery of PGMs from pristine ores is floatation and it has achieved platinum recoveries greater than 85 % (Sefako, Sekgarametso & Sibanda, 2017). In South Africa, PGMS are mined from the Bushveld Complex which consists of the Merensky reef, Upper Group 2 (UG2) reef and the Platreef (Cawthorn, 2010). The fast exhaustion of the Merensky reef has forced PGM producers to mine the upper group 2 (UG2) reef or the Platreef. During the milling of UG2 or Platreef ores, a low-quality concentrate with high chromite content is produced (Mwase, Petersen & Eksteen, 2012). This low-grade concentrate increases the smelter cost and decreases the smelter integrity because of the increase in chromite (Mwase, Petersen & Eksteen, 2014). Increased chromite in the

concentrate causes an accumulation of heat resistant chromite spinel layers in the furnaces which affect the functioning and availability of the furnaces (Valenta & Mapheto, 2010). This slows production down resulting in a financial loss. Consequently, the low-grade concentrate leads to an increase in processing cost and a decrease in PGM production.

Given that the recovery of PGMs is expensive, not sustainable, or environmentally friendly, alternative approaches such as biohydrometallurgical methods must be explored. PGMs are primarily contained in either PGM accessory minerals or in base metal sulphide minerals such as chalcopyrite ($CuFeS_2$), pyrrohotite (FeS) and pentandite [$(NiFe_9)S_8$] (Kraemer, Junge & Bau, 2017). These methods include bioleaching with sulphur and iron oxidising microorganisms as well as organic/inorganic acid-producing microorganisms (Schippers *et al.,* 2014; Kraemer *et al.,* 2015). Despite South Africa having several PGM mines very little work has been done on studying the microbial diversity at these mining sites. Microorganisms found at these PGMs mines could have characteristics that make them suitable for use in the bioleaching of PGMs. Bioleaching has thus far been applied on an industrial level in the recovery of nickel, copper, zinc, and cobalt from sulphide bearing ores (Hedrich *et al*., 2020). Mesophilic microorganisms such as *Acidithiobacillus ferrooxidans, Acidithiobacillus thiooxidans* and *Leptospirillum ferrooxidans* have been mainly used for the bioleaching of these metals (Simate, 2009; Nguyen *et al*., 2018). Two studies have been successfully carried out where a thermophilic microorganism, *Metallosphaera hakonensis*, has been used for the base metal bioleaching of a PGM low-grade concentrate and a coarse ore (Mwase, Petersen & Eksteen, 2012, 2014).

1.1.2 Research problem

The bacterial community and diversity found at several mine sites such as gold and copper mines have been extensively studied. However, very few studies have been done on the bacterial community and diversity found at PGM mines. It is theorised that organic acid-producing, and iron and sulphur-oxidising bacteria may be isolated from a PGM mine, in this study.

1.1.3 Research aims and objectives

This study aims to isolate, identify, and characterise bacteria from a platinum concentrator plant.

Objectives

- Isolation and enumeration of bacteria.
- Molecular identification of bacteria.
- Characterisation of bacteria.

CHAPTER 2: LITERATURE REVIEW

2.1 The platinum group metals mining industry

2.1.1 Platinum group metals

Platinum group metals (PGMs) are rare precious metals and consist of six metals namely: platinum (Pt), palladium (Pd), osmium (Os) rhodium, iridium (Ir) and ruthenium (Ru) (Seymour & O'Farrelly, 2012). In contrast to gold, which is primarily used in jewellery and dentistry, PGMs are used in a variety of industrial applications such as dental alloys, jewellery, catalytic converters, and thermocouples. The most commercially important metals within this group are platinum and palladium, with the automotive industry being their most substantial industrial application, where they are used in vehicle catalytic converter units (Gunn, 2013). These metals are extremely rare due to their low natural abundance and the complexity of their extraction and refining processes (Seymour & O'Farrelly, 2012).

The PGMs are transition metals and are chemically similar. They all have siderophile and chalcophile propensities where they can associate or bond with iron and sulphur and have also been shown to associate with nickel and copper (Seymour & O'Farrelly, 2012). The physical properties of the PGMs differ considerably, but they all show properties that are consistent with metals such as forming alloys, conducting heat and electricity and a certain degree of malleability and ductility (Seymour & O'Farrelly, 2012). Platinum is exceptionally resistant to corrosion due to its high oxidation potential, making it stable in the air even at high temperatures (Reith *et al.*, 2014). In addition to this, it is also soft and ductile. These properties make it a highly sort out metal with applications in dentistry, thermometers, jewellery, anticancer drugs, prosthetics, and catalytic converter units (Reith *et al*., 2014).

Platinum group metals are very rare and occur at mean concentration of 0.4 ng/g. As a result, there are very few minable deposits that have been identified (Reith *et al*., 2014). Currently, the major minable PGM deposits include the Bushveld Indigenous complex in South Africa, the Great Dyke mine in Zimbabwe, the Stillwater complex in the United States of America and placer deposits in Russia

(Birke *et al*., 2018). Placer deposits are mineral deposits, formed by surface weathering of primary rocks, that accumulate due to wind, water or ice action (Haldar, 2013). Approximately 90 % of the world's PGM deposits are found in South Africa. It is the largest producer of platinum in the world, with an approximate production of 80 % of the world's annual platinum production (Birke *et al*., 2018). Within the South African Bushveld Indigenous complex, platinum is mined from three sections, namely the Merensky Reef, UG2 Reef and Platreef.

2.1.2 Current platinum group mining methods: An overview

The South African platinum industry has many underground mines and a couple of opencast mines. Figure 2.1, shows the conventional steps involved in the beneficiation of PGMs which include; comminution, electrical smelting, converting, base metal refining and PGM refining and separation (Safarzadeh, Horton & Van Rythoven, 2018). It is important to note that the mineralogy of the ore will determine the beneficiation process and there are often deviations from and additions to this general process depending on the type of ore. Comminution involves the treatment of the PGM ore in primary and secondary crushers resulting in fine fragments or particles. It is then milled in rod and ball milling circuits (Thethwayo, 2018). There is a wide array of comminution circuits used in the PGM industry, with the two-staged circuit as the most common. Thereafter, the milled PGM ore undergoes flotation where it is treated using flotation cells and gravity separators. Flotation involves the separation of mineral particles according to their hydrophobicity (Sahu *et al*., 2022). The flotation circuit typically has a rougher stage and two cleaner stages (Cramer, 2001). The common reagents used during flotation are dithiophosphate and xanthate collectors with pH between 7.5-9 (Cramer, 2001). The resulting product from comminution is a sulphide-rich PGM concentrate. Iron and sulphur-oxidising bacteria can potentially be used to for the bioleaching of the sulphide-rich PGM concentrate. The dried concentrate is then sent to an electrical smelter where the oxide bearing gauge (silicates) is removed from the PGM rich sulphides using high temperatures between 600-900°C. The resulting matte can either be processed directly into converters or it can be granulated before it is converted, depending on the producer.

The resulting matte goes through a process called converting, where iron sulphide and sulphur are converted to ferrous oxide and sulphur oxide, respectively. The oxidation is performed through the treatment of the matte in Pierce-Smith converters or the Ausmelt process where the iron dioxide is separated as a fayalitic slag, and the sulphur dioxide is separated as an off-gas. After cooling the matte is then milled and treated in a base metal refinery that includes the removal of impurities such as zinc, lead and selenium, cobalt removal and recovery, copper and nickel electrowinning and the disposal of sulphur (Cramer, 2001). The final stage of platinum mining is purification which involves the use of several hydrometallurgical processes such as solvent extraction, leaching, ion exchange and molecular recognition technology (Cramer, 2008).



Figure 2.1: A general overview of PGM ore processing (Sinisalo & Lundström, 2018; Thethwayo, 2018).

2.1.3 An overview of the platinum group metal concentrate process

In the PGM mining industry, several flotation circuits are currently employed to concentrate the PGMs that have been mined, such as the "South African MF2/MF3" Flowsheet, Northam Flowsheet, Noril'sk Flowsheet, Stillwater Flowsheet (Cole & Ferron, 2002). The South African PGM mining industry mainly uses the "South African MF2/MF3" Flowsheet and Northam Flowsheet (O'Connor & Alexandrova, 2021). The type of flotation circuit used depends on the ore mineralogy of the mined PGM. The process of PGM concentrating begins with the PGM ore being fed into a semi-autogenous mill which grinds the ore, thus providing primary grinding (Cramer, 2001). Thereafter, the ground ore is processed into a ball mill, which provides secondary grinding. The ore is then put through a process called flash flotation where approximately 60% of the PGM is recovered (Cole & Ferron, 2002). Primary flotation takes place in a rougher or scavenger flotation bank. The rougher bank contains a cell/tank that can produce a final grade concentrate that does not require further processing in the concentrator (Cole & Ferron, 2002). The scavenger bank produces a concentrate that may need to undergo primary flotation again. The remaining rougher bank concentrate is then processed into the cleaner circuit which is a series of three column cells/tanks that each recover approximately 30% of PGMs (Cole & Ferron, 2002). The scavenger concentrate is either processed to the rougher bank or the cleaner circuit. The final concentrate is then made into a slurry through a concentrate thickener and is then subsequently processed to a smelter (Cramer, 2001). At this stage, the produced concentrate can potentially be bioleached using iron and sulphur oxidising or organic acid producing bacteria.

2.2 Bacterial diversity in mines

2.2.1 Introduction

The diversity of microorganisms found at various mines is dependent on several factors which include physiochemical factors such as pH, redox potential, and electrical conductivity. It is also dependent on soil structure, nutrients, temperature, heavy metal concentration, and the soil ecosystem (Marais, 2012). The dominant phyla of microorganisms commonly found at mining sites are Proteobacteria which includes *Pseudomonas* and *Acidovorax* species (Kamika & Momba, 2014; Pereira,

Vicentini & Ottoboni, 2014; Pradhan *et al*., 2022). Some of the bacterial species found at mining sites are extremophilic and can grow at a pH as low as 2 and at high heavy metal concentrations. The bacterial diversity at mining sites differs across regions and is primarily dependent on the type of extractive process used at a particular site (Fernandes *et al*., 2018).

2.2.2 Bacteria found at mining sites

The study of microorganisms at mining sites, which are considered complex environments, has become possible due to the application of PCR amplification of the 16S rRNA genes. Proteobacteria, Bacteroidetes, Acidobacteria and Firmicutes are dominantly found in mining sites (Thavamani *et al*., 2017; Fashola, Ngole-Jeme & Babalola, 2020). Figure 2.2 provides a brief overview of the bacterial diversity found at mining sites.



Figure 2.2: Bacterial and Archaeal diversity in terms of phyla found at mine sites (Thavamani *et al*., 2017).

Bacteria found at mine sites have adapted to the high heavy metal concentration characteristic of mine sites and are typically iron and sulphur-oxidising, as shown in Figure 2.2. This involves the oxidation of sulphur and ferrous iron to sulphate and ferric iron and the ions generated from these reactions are subsequently used by the bacteria as electron donors to obtain energy (Ranalli, Zanardini & Sorlini, 2009; Korehi *et al.*, 2013; Sultan & Faisal, 2014; Hassanshahian & Ghoebani, 2018). Examples of iron and sulphur-oxidising bacteria found at mine sites include *Acidithiobacillus thiooxidans*, *Acidithiobacillus ferrooxidans*, *Leptospirillum* sp. and *Sulfobacillus* sp.

Bacteria have been isolated and studied from several mines around the world including copper, iron, silver, coal, platinum, and gold mines. Table 2.1 provides a summary of the different bacteria isolated from various mines. Studies by He *et al*. (2007); Kuang *et al.*(2013) and Santofimia *et al*. (2013) that focused on analysing microbial communities at copper mines in China and Spain showed that the microbial diversity is directly affected by soil elements, pH, and temperature. Xie *et al*. (2007), also found that there was a diverse distribution of *A. ferrooxidans* and *Leptospirillum* sp. which are iron-oxidising bacteria. The distribution of these bacterial species can be attributed to the high concentration of elements such as copper, sulphur, and iron at the mine. *Acidithiobacillus ferrooxidans* is a widely studied bacterial species in studies of acid mine drainage and it has been considered as a model organism in bioleaching research. A study by Domingues *et al.*(2020), found that Firmicutes was the dominant phylum found at a copper mine in Brazil, followed by Proteobacteria and Actinobacteria, with bacteria from the *Bacillus*, *Lysinibacillus* and *Alcaligenes* genera as the most dominant. Pepper *et al.* (2012), found that Proteobacteria and Actinobacteria were the most dominant phyla found at an Arizona copper mine. These studies show that there is a variety of bacteria at copper mines.

Bacteria isolated from gold mines have also been shown to dominantly be from the Proteobacteria, Actinobacteria and Firmicutes phyla. These bacteria also exhibit heavy metal tolerance. *Bacillus cereus* and *Enterobacter asburiae* have been isolated from gold mine tailings and these bacteria exhibit high heavy metal

tolerance suggesting that they have developed specific mechanisms to endure heavy metal stress (Fashola *et al*., 2020). A study by Fashola *et al*. (2020), identified and characterised seventeen bacterial isolates with *Bacillus sp.* and *Enterobacter sp*. as the most dominant. Bacteria isolated from coal mines are also dominantly from the Proteobacteria, Firmicutes and Actinobacteria phyla, however, Adeleke, Cloete & Khasa (2012), found that Bacteroidetes was also a dominant phylum. These bacteria were able to produce organic acids that could assist in iron ore solubilisation. A study by Ekpa *et al.* (2017), isolated *Acidithiobacillus* , *Pseudomonas* and *Leptospirillum* bacterial species. These bacteria can oxidise ferrous iron thereby generating energy and can locally concentrate ferric iron by absorbing it to microbial surfaces. They are also mesophilic acidophiles and exhibit good bioleaching properties.

Table 2.1: Summary of the bacteria isolated from copper, gold, and coal mines.

Mines	Dominant microorganisms	Reference
Copper mine	Proteobacteria, Actinobacteria, Firmicutes, *Acidithiobacillus ferrooxidans*, *Leptospirillum sp*., *Bacillus sp*., *Lysininbacillus sp*. and *Alcaligenes*.	(Xie *et al*., 2007; Pepper *et al*., 2012; Domingues *et al*., 2020)
Gold mine	Proteobacteria, Actinobacteria, Firmicutes *Bacillus sp*., *Enterobacter sp*., *Bacillus subtilis*, *Bacillus cereus*, *Arthrobacter sp*.,	(Fashola, Ngole-Jeme & Babalola, 2020)
Coal mine	Proteobacteria, Actinobacteria, Firmicutes, Bacteroidetes, *Acidithiobacillus sp.*, *Pseudomonas sp.*, *Leptospirillum sp.*	(Adeleke, Cloete & Khasa, 2012; Ekpa *et al*., 2017)

2.2.3 Bacteria isolated from platinum mines

Several studies have focused on identifying and characterising bacteria isolated from platinum mines. These studies focused on heavy metal of the isolated bacteria. Marais (2012), investigated bacteria found at a platinum mine tailing dam. Nine species were identified, as shown in Table 2.2, with *Paenibacillus lautus* as the most widely isolated. Some of the *Paenibacillus* isolates identified in the study were able to produce endospores, which enables the bacteria to endure environmental stress. These bacterial strains have a high heavy metal tolerance with *Bacillus subtilis* showing the highest tolerance. A study by Rauwane (2008), also found *Paenibacillus lautus* to be the most abundant isolate found at a platinum mining site.

Table 2.2: Bacteria isolated from platinum mining sites.

Bacterial isolate	**Reference**
Paenibacillus lautus, Paenibacillus lautus strain DS19, *Paenibacillus* sp. C15, uncultured *Paenibacillaceae, Stenotrophomonas maltophilia, Bacillus* sp. KDNB5, *Bacillus cereus, Alcaligenes* sp. DJWH 146-2,	(Marais, 2012)
Bacillus barbaricus	(Rauwane, 2008)

2.3 Bacterial Bioleaching

2.3.1 Introduction

There is an increasing demand for PGMs due to an increase in their industrial applications. However, there is also a decrease in the easily exploited and highgrade PGM deposits. Therefore, the use of new biotechnological methods to recover PGMs from low-grade ores is essential to meeting this demand. One such biotechnological method is bioleaching. Bioleaching or microbial leaching can be defined as the process where metals are solubilized from an insoluble substrate (Schippers *et al*., 2014). This process normally occurs because of the metabolism of the microorganisms. Bioleaching provides a sustainable, and cost-effective alternative to conventional metallurgical extraction methods.

2.3.2 Bacteria used in bioleaching

The bacteria that are important in the bioleaching process involved with metal extraction can be divided according to their nutritional requirements i.e., chemolithotrophic and heterotrophic bacteria and their growth temperature i.e., mesophilic, and thermophilic bacteria.

2.3.2.1 Chemolithotrophic bacteria

Chemolithotrophic bacteria can produce energy through the oxidation of inorganic molecules through aerobic or anaerobic respiration for biosynthesis (Enrich-Prast *et al*., 2014). A major group of this category are iron and sulphur-oxidising bacteria. Chemolithotrophic bacteria commonly used for bioleaching include *A. ferrooxidans* which can oxidise iron and sulphur, *A. thiooxidans* and *A. caldus* which can oxidise sulphur and *L. ferrooxidans* and *Leptospirillum ferriphilum* which are iron oxidisers. Bioleaching using chemolithotrophic bacteria has been commercially applied for the extraction of copper (Brierley & Brierley, 2013; Yin *et al*., 2018), gold (Van Hille, Wyk & Harrison, 2011; Gericke, 2012) and nickel (Puhakka, Kaksonen & Riekkola-Vanhanen, 2007; Riekkola-Vanhanen, 2007).

2.3.2.2 Heterotrophic bacteria

Heterotrophic bacteria obtain energy and carbon from organic compounds (Van Aken, 2011). These bacteria produce metabolic by-products that can interact with the mineral surface resulting in metal dissolution. These by-products include

complexing organic acids such as citric, acetic, and oxalic acid, exopolysaccharides as well as proteins that can aid in metal solubilisation (Schippers *et al*., 2013). Heterotrophic bacteria that have been used for bioleaching include *Pseudomonas aeruginosa* and *Bacillus sp*. (Jain & Sharma, 2004; Irannajad *et al*., 2014; Barnett, Palumbo-Roe & Gregory, 2018). Heterotrophic bacteria have also been used for the bioleaching of copper and zinc (Shabani *et al*., 2013; Irannajad *et al.*, 2014).

2.3.2.3 Mesophilic bacteria

Mesophilic bacteria grow at temperatures between 25°C and 35°C. Examples of chemolithotrophic mesophilic bacteria used in bioleaching include *Acidithiobacillus* spp., which are aerobic, iron and sulphur-oxidising, these include *Acidithiobacillus caldus*, *albertis*, *acidophilus*, *concretivorus* and *prosperous* (Kumar e*t al*., 2019). These bacteria can grow at a low pH, making them suitable for bioleaching. *Leptospirillum* spp. are also mesophilic and can tolerate higher concentrations of molybdenum, silver and uranium when compared to *A. ferrooxidans* (Kumar *et al*., 2019). *Leptospirillum* spp. can grow between temperatures of 20°C and 45°C (Rawlings, 2013). However, bioleaching at higher temperatures is more desirable. Examples of mesophilic heterotrophic bacteria used in bioleaching include *Bacillus* spp. and *Pseudomonas* spp. These bacteria can grow between 20°C to 42°C (Chakravarty & Anderson, 2015; Kumar *et al*., 2019).

2.3.3 Mechanisms of bioleaching

Leaching reactions in metal leaching use ferric iron and protons from organic or inorganic acids to solubilise metals (Rawlings, 2005). Metal leaching relies on three processes, acidolysis, redoxylysis and complexolysis (Wu & Ting, 2006; Sethurajan, 2015). When microorganisms attach to a surface, they start forming an exopolysaccharide layer and within this layer the bio-oxidation reactions involved in bioleaching occur (Rawlings, 2005). The microorganisms provide both the space and leaching chemicals for bioleaching. In previous literature, bioleaching was described as either direct or indirect bioleaching as shown in Figure 2.3 (Bosecker, 1997; Sand *et al*., 1999; Tributsch, 1999). The direct mechanism involves the attachment of bacteria to the mineral and the indirect mechanism the bacteria oxidise iron resulting in mineral dissolution.



Figure 2.3: Bacterial mechanisms of bioleaching. In direct bioleaching, bacteria attach to the mineral surface and the oxidation of sulphur results in metal dissolution. In indirect bioleaching, bacteria consume oxygen and oxidise iron (Mishra & Rhee, 2014).

2.3.3.1 Redoxylysis

Bioleaching by chemolithotrophic bacteria is achieved through iron and sulphur oxidation. Mineral dissolution reactions and the oxidation intermediates differ depending on the mineralogy of the sulphide bearing mineral. Schippers & Sand (1999), proposed that there are two mechanisms involved in mineral dissolution, a thiosulphate mechanism, and a polysulphide mechanism. Metals like pyrite and molybdenite which are acid insoluble use the thiosulphate mechanism for oxidation and sphalerite, galena and chalcopyrite use the polysulphide mechanism (Tao & Dongwei, 2014). For the thiosulphate mechanism, thiosulphate is the primary intermediate and sulphate is the primary end-product, and the metal is solubilised when ferric iron acts on the metal sulphides (Schippers *et al.*, 2014). In the polysulphide mechanism, sulphur is the primary intermediate and the insoluble metal is acted on by both ferric iron and protons. The bacteria involved in the

solubilisation of the metals provide the protons for a proton attack in the form of sulphuric acid and re-oxidise the iron keeping it in its oxidised ferric form for oxidation of the mineral.

2.3.3.2 Organic acids

Organic acids produced by heterotrophic bacteria can solubilise metals such as copper and nickel. These acids are produced through the metabolic activity of bacteria where sugar substrates such as glucose and sucrose are converted into organic acids (Barnett, Palumbo-Roe & Gregory, 2018). Organic acids solubilise metals, as shown in Figure 2.4, through mineral acidolysis by providing protons (hydrogen) or by acting as complexing agents and complexing with the metals in the minerals (Giese, 2021). The metal dissolution reaction is catalysed by organic acids where the cation-oxygen bonds of the metal are weakened by the formation of metal-organic complexes. Organic acids also aid in metal dissolution by lowering the pH of the solution which is ideal for bioleaching (Barnett, Palumbo-Roe & Gregory, 2018). Bacteria that are used in organic acid bioleaching include *Bacillus* sp., *Pseudomonas* sp., and *Erwinia* sp. The organic acids that can potentially be used in bioleaching include but are not limited to citric, lactic, oxalic and gluconic acid. Several studies have shown that PGMs can be dissolved by organic acids (Mwase, 2009; Trinh *et al*., 2017). The use of organic acids in bioleaching is advantageous because they can be biodegraded easily, and they are cost-effective.



Figure 2.4: Overview of metal solubilisation by organic acids through acidolysis and complexolysis (Mwase, 2009).

2.4 Bacterial bioleaching of platinum group metal ores

Bioleaching has been accepted as a cost-effective, green, and sustainable alternative to traditional hydrometallurgical and pyrometallurgical extraction methods. To date, only gold sulphide ores have been bioleached on an industrial scale (Brierley, 2016), and on a lab scale. The bioleaching of sulphide platinum ore concentrates has been demonstrated (Mwase, Petersen & Eksteen, 2012, 2014). Mwase, Petersen & Eksteen (2012), conducted heap leaching using thermophilic and mesophilic bacteria to bioleach base metals and PGMs from a low-grade PGM concentrate. The experiment resulted in the extraction of 85% cobalt, 52% copper and 95% nickel after 30 days. The remaining concentrate from the heap leaching experiment was leached with cyanide resulting in the extraction of 46% rhodium, 20.3% platinum and 87% palladium. In 2014, Mwase, Petersen & Eksteen, studied a heap leaching method that was tested on crushed coarse platinum ore using iron and sulphur oxidising bacteria that are both thermophilic and acidophilic. The bioleaching

experiment resulted in the extraction of 75% nickel, 93% copper and 53% cobalt after 304 days. A subsequent cyanide leach resulted in the extraction of 90.3% gold, 57.8% platinum and 99.7% palladium after 60 days. Both studies showed that using bioleaching to extract PGMs from PGM ores or concentrates is a viable and effective alternative method to the costly and problematic smelter process.

CHAPTER THREE: IDENTIFICATION OF BACTERIA

3.1 Introduction

The number of studies that have identified and characterised bacteria from platinum mines are very few and have focused solely on bacteria from platinum mine tailings. The study of bacterial diversity at mine sites has become more common. Literature has shown that the microbial community found in heavy metal contaminated soils is complex and well-adapted (Halter *et al*., 2011; Gillan *et al*., 2015). The bacteria found at these sites play environmentally important roles such as biogeochemical cycling and heavy metal bioaccumulation (Fernandes *et al*., 2018). Thus, it is important to acquire comprehensive knowledge about these bacteria and their communities.

The increase in demand for bacteria capable of bioleaching and bioremediation has motivated studies on bacteria that inhabit mine sites. Most of the studies conducted on these bacteria have focused on their heavy metal tolerance/resistance (Wei *et al*., 2009; Marais, 2012; Ndeddy Aka & Babalola, 2017). This study aimed to isolate and identify bacteria from a platinum mine concentrator plant and its surroundings.

3.2 Methods and Materials

3.2.1 Sample collection

Thirty-nine samples were collected from the Anglo Platinum Mogalakwena Mine (23.9709° S, 28.9166° E), in Limpopo, South Africa. The samples were collected from the concentrator plant, the pond outside the concentrator plant and in front of the crusher and concentrator and consisted of a mixture of liquid, slurry, and soil samples. They were placed in labelled 50 mL Greiner Bio-One polypropylene conical centrifuge tubes except for sample 39 which was placed in a polypropylene plastic bag. The samples were transported to the laboratory (Industrial Microbiology and Biotechnology Lab, University of the Witwatersrand) and stored at 4°C.

3.2.2 Determination of sample pH

For the measurement of the soil samples, 1 g of each soil sample was suspended in 10 mL of autoclaved distilled water. The pH of the samples was measured using a calibrated glass electrode Crison pH-meter.

3.2.3 Isolation of bacteria

For the isolation of bacteria from the Anglo samples, a 10^{-1} - $10^{-4}$and 10^{-1} - 10^{-5} dilution series was performed for plating onto Luria-Bertani (LB) and 9K agar, respectively. Luria-Bertani agar was composed of 5 g/L yeast extract, 5 g/L NaCl, 10 g/L tryptone and 15 g/L agar. 9K agar was composed of 5 g/L $MgSO_4 \cdot 7H_2O$, 0.05 g/L $Ca(NO_3)_2$, 3 g/L $(NH_4)_2SO_4$, 0.5 g/L K_2HPO_4, 0.1 g/L KCl and 15 g/L of agar. For the dilution series, 1 g or 1 mL of each sample was suspended in 9 mL of 0.9% saline (NaCl) solution. Thereafter, aliquots of 100 µL from each dilution were spread plated onto their respectively labelled LB or 9K agar plates. The plates were incubated at 37°C for 48 hours. After the incubation period, the plates were observed, colonies were counted, and distinct colonies were subcultured, using the streak plate method until pure cultures were obtained.

3.2.4 Genomic DNA extraction

The extraction of DNA was performed on bacteria pure cultures isolated from the samples. Prior to DNA extraction, a colony of each bacterial isolate was resuspended in 300 µL of autoclaved distilled water, vortexed and 250 µL of the mixture was used for DNA extraction. DNA extraction was performed using the ZymoBiomics™ DNA Miniprep Kit (Catalog No D4301) as per manufacturer's protocol (Appendix B). The quantity and quality of the DNA was analysed using the NanoDrop® ND-1000 spectrophotometer and stored at 4°C until further analysis.

3.2.5 DNA sequencing

The DNA was sent to Inqaba Biotec™ (Pretoria, South Africa) where PCR, agarose gel electrophoresis and Sanger sequencing were carried out. The 16S target region was amplified using OneTaq® Quick-Load® 2X Master Mix (NEB) with the universal primers 27F and 1492R shown in Table 3.1 (Lane *et al*., 1991; Turner *et al*., 1999). The PCR products were electrophoresed on an agarose gel and extracted using a Zymoclean™ Gel DNA Recovery Kit. The extracted fragments were sequenced in the forward and reverse direction (Nimagen, BrilliantDye™ Terminator Cycle Sequencing Kit V3.1, BRD3-100/1000) and purified (Zymo Research, ZR-96 DNA Sequencing Clean-up Kit) to remove any impurities, unincorporated dNTPs, DNA polymerase enzymes and unbound primers. The ABI 3500xl Genetic Analyzer (Applied Biosystems, ThermoFisher Scientific) was used to analyse the purified fragments. The .ab files produced by the ABI 3500XL Genetic Analyzer were analysed by the CLC Bio Main Workbench v7.6 and the results were analysed through a Basic Local Alignment Search Tool (BLAST) search [National Centre for Biotechnology Information (NCBI)] **Table 3.1**: Universal primer set used for PCR amplification.

Name of Primer	Target	Sequence (5' to 3')
16S-27F	16S rDNA sequence	AGAGTTTGATCMTGGCTCAG
16S-1492R	16S rDNA sequence	CGGTTACCTTGTTACGACTT

3.3 Results and Discussion

3.3.1 Bacterial enumeration

Prior to bacterial isolation and enumeration, the pH of each sample was measured, and the results are shown in Table 3.2 below. The pH for the samples taken from the concentrator plant, which were all liquid samples, ranges from 7.35 to 8.69. For the samples from the pond outside the concentrator plant, the pH ranged from 6.39 to 9.11. The pH was between 7.90 and 9.27 for the samples taken from in front of the crusher. The soil sample collected from in front of the concentrator had a pH of 8.49. Therefore, the bacteria were isolated from liquid, slurry and soil samples with a pH that ranged from 6.39-9.11.

Soil chemistry is correlated to changes in the soil microorganisms' communities across different spaces. Studies have shown that the bacterial composition of microbial communities corresponds to soil pH (Shen *et al*., 2013; Qi *et al*., 2018; Cho *et al*., 2019).Therefore, the pH of the samples and isolation plates would influence the bacterial species isolated and their CFUs. The pH of the isolation plates was 7 therefore, acidophilic bacteria were not selected.

Table 3.2: Sampling sites within the Anglo Platinum Mogalakwena mine in Limpopo South Africa and the pH and colony forming units per gram/ milliliter of each sample.

Sample number	Sample	Site Collection point	Sample pH	CFU/mL or CFU/g of sample
1	Rougher Feed	Concentrator plant	8.69	8×10^3
2	Rougher Feed	Concentrator plant	8.61	5.2×10^4
3	Rougher Tail A	Concentrator plant	8.15	8.9×10^5
4	Rougher Tail A	Concentrator plant	8.17	2.6×10^5
5	Rougher Tail B	Concentrator plant	7.69	5.7×10^4
6	Rougher Tail B	Concentrator plant	7.68	2.04×10^5
7	Scavenger Feed	Concentrator plant	8.51	1.5×10^5
8	Scavenger Feed	Concentrator plant	8.51	1.8×10^5
9	Scavenger Tail A	Concentrator plant	7.35	4.4×10^5
10	Scavenger Tail A	Concentrator plant	7.40	1.9×10^4
11	Scavenger Tail B	Concentrator plant	8.55	1.4×10^5
12	Scavenger Tail B	Concentrator plant	8.53	2.3×10^5
13	Cleaner Feed	Concentrator plant	8.27	6×10^4
14	Cleaner Feed	Concentrator plant	8.40	6.8×10^4

Sample number	Sample	Site Collection point	Sample pH	CFU/mL or CFU/g of sample
15	Cleaner Scavenger Tail A	Concentrator plant	8.23	5.4×10^4
16	Cleaner Scavenger Tail A	Concentrator plant	8.45	5.6×10^4
17	Cleaner Scavenger Tail B	Concentrator plant	8.36	1.42×10^5
18	Cleaner Scavenger Tail B	Concentrator plant	8.42	8.9×10^4
19	Dry soil	Pond outside of the plant	8.63	1.33×10^5
20	Dry soil	Pond outside of the plant	9.11	8.2×10^5
21	Dry soil	Pond outside of the plant	8.30	1.3×10^5
22	Dry soil	Pond outside of the plant	8.25	7.8×10^4
23	Soil and liquid	Pond outside of the plant	7.27	6.6×10^5
24	Soil and liquid	Pond outside of the plant	6.96	9×10^3
25	Soil and liquid	Pond outside of the plant	6.90	1.43×10^4
26	Soil and liquid	Pond outside of the plant	7.53	4×10^4
27	Soil and liquid	Pond outside of the plant	6.87	7.2×10^3
28	Soil and liquid	Pond outside of the plant	7.39	1×10^5
29	Liquid	Pond outside of the plant	6.86	9.3×10^4
30	Liquid	Pond outside of the plant	6.90	7×10^4
31	Liquid	Pond outside of the plant	6.78	6×10^4
32	Soil	In front of the crusher	9.27	5×10^5
33	Soil	In front of the crusher	9.21	1.09×10^3
34	Soil	In front of the crusher	8.96	1.94×10^5
35	Soil	In front of the crusher	8.78	2.01×10^5
36	Soil and liquid	In front of the crusher	7.90	6.1×10^4
37	Soil and liquid	In front of the crusher	8.27	1.6×10^4
38	Soil and liquid	In front of the crusher	8.32	2.9×10^5
39	Soil	In front of the concentrator	8.49	4.5×10^6

Sample 39 (Table 3.2), taken in front of the concentrator plant, has the highest CFU followed by sample 3 from the concentrator plant and samples 20 and 23 from the pond outside the concentrator plant. Sample 27, from the pond outside the concentrator plant, had the lowest CFU followed by sample 1, from the concentrator plant and 24 from the pond outside the concentrator plant. In general, soil samples had the highest CFU's compared to the liquid and slurry samples.

Bacterial community composition is affected by the soil where the bacterial community is found. The high CFU's seen in sample 39 could be due to the nature of the sample since it is a soil sample. A study by Kemp and Aller (2004),

investigated the bacterial diversity in various environments including aquatic and soil environments using rDNA libraries. They found that the phylogenetic richness of soils seems to be higher than the aquatic environments, indicating that it had higher bacterial diversity. Several studies have shown that soil has one of the highest bacterial diversity compared to other environments (Thompson *et al*., 2017; Bahram *et al*., 2018; Louca *et al*., 2019) The higher CFUs of the soil samples compared to the liquid samples could also be attributed to the number of organic nutrients available in the soil compared to the water found at the concentrator plant.

The difference in CFU's between the soil and water samples could also be due to the storage of the samples. Ahammed (2003), showed that the storage at 4°C of bacterial water samples results in a decrease in the CFUs of the bacteria after 12 hours. A study by Lee *et al*. (2021), showed that the CFUs of bacteria in soil samples was not significantly affected when stored at 4°C over a period of time.

3.3.2 Identification of bacterial isolates

The DNA sequences obtained from sequencing the bacterial isolates were matched with sequences of a similar nature on the NCBI website using the BLAST tool. The BLAST query shown in Table 3.3 demonstrated that a total of 50 bacterial species were isolated from the concentrator plant and the areas surrounding the plant. Bacterial species from the genus Bacillus and Pseudomonas were dominant, however, there was a wide variety of different genera.

Table 3.3: Identity of bacteria and GenBank Accession numbers of closely related species or species with highest sequence similarity percentage.

Isolate no.	Mine sample no.	BLAST prediction	GenBank Accession	Read length	E-value	% Identity	Phylum
15	26	*Acidovorax* sp.	AB646302.1	1124	0	100	
26	32	*Pseudomonas aeruginosa*	HQ018745.1	870	0	78,04	
27	26	*Pseudomonas brassicacearum*	KT695844.1	1381	0	99,71	
29	3	*Pseudomonas stutzeri*	CP071899.1	1439	0	99,93	
30	1	*Pseudomonas alcaliphila*	CP016162.1	1435	0	99,16	

33	5	*Pseudomonas stutzeri*	MT027239.1	1435	0	99,93	
35	4	*Pseudoxanthomonas mexicana*	JX860406.1	1442	0	99,65	
36	13	*Pseudoxanthomonas mexicana*	CP060731.1	1445	0	99,93	
37	31	*Comamonas aquatica*	KJ395470.1	1426	0	99,93	
38	21	*Salmonella enterica*	CP053585.1	1441	0	97,43	
41	36	*Pseudomonas otitidis*	AP022642.1	855	0	99,88	
43	9	*Pseudoxanthomonas mexicana*	JX860406.1	1394	0	99,93	
45	14	*Alishewanella fetalis*	MN809397.1	1419	0	99,15	
46	6	*Pseudomonas alcaligenes*	KT998857.1	1440	0	99,93	
49	10	*Methylobacterium adhaesivum (98% match)*	AB698743.1	826	0	98,42	
7	25	*Bacillus* sp.	MN961108.1/ CP050183.1	1459/1458	0/0	99,86/99,93	Firmicutes
10	18	*Bacillus* sp.	CP053938.1/ CP053997.1	1457/2908	0/0	99,86/99.86	
16	3	*Staphylococcus capitis*	CP053957.1	1399	0	99,71	
25	24	*Bacillus* sp.	MT510153.1/ MT490862.1	1401/1402	0/0	100/100	
28	24	*Bacillus* sp.	MT534569.1/ MT527538.1	1229/1229	0/0	99,92/99,92	
31	15	*Bacillus* sp.	MT538261.1/ MT184866.1/ CP040334.1	1448/1448/1448	0/00	99,93/99,93/99,93	
32	37	*Bacillus* sp.	MN581187.1/ MN006152.1	1440/1440	0/0	100\100	
34	16	*Bacillus* sp.	CP053938.1/ CP053997.1	1447/1447	0/0	99,86/99,93	
39	38	*Bacillus halotolerans*	CP054584.1	1445	0	99,93	
40	30	*Paenibacillus alvei*	MT498465.1	1437	0	99,51	
42	34	*Bacillus* sp.	CP054568.1/ CP053997.1	1133/1133	0/0	99,74/99,74	
5	35	*Arthrobacter globiformis*	EU221407.1	803	0	100	Actinobacteria
9	39	*Georgenia satyanarayanai*	MN326714.1	1409	0	99,01	
17	22	*Rhodococcus* sp.	MK443065.1/ MG833299.1	1090/1086	0/0	99,91/100	

Isolate no.	Mine sample no.	BLAST prediction	GenBank Accession	Read length	E-value	% Identity	Phylum
18	5	*Rhodococcus pyridinivorans*	MN179918.1	1423	0	99,86	
20	35	*Arthrobacter globiformis*	EU221407.1	842	0	99,76	
48	7	*Pseudarthrobacter phenanthrenivorans*	MN658730.1	835	0	99,52	
50	23	*Pseudarthrobacter defluvii*	KY882049.1	835	0	99,28	
14	23	*Microbacterium* sp.	CP027929.1, CP018134.1, KX404917.1	1361;1361;1361	0;0;0	99,85; 99,85; 99,85	Actinomycetota
21	20	*Microbacterium* lacusdiani	MG947593.1	1426	0	99,79	
22	30	*Microbacterium foliorum*	KP208636.1	830	0	99,75	
23	22	*Leucobacter chromiiresistens*	NR_117509.1	1369	0	100	
24	19	*Nocardia cyriacigeorgica*	LR215973.1	1406	0	100	
44	20	*Nocardia farcinica*	MT317187.1	1417	0	100	
19	23	*Chryseobacterium takakiae*	KC560016.1	575	0	98,43	Bacteroidetes
47	39	*Mucilaginibacter polysacchareus*	KM019772.1	1359	0	99,93	

The Proteobacteria phylum consisted of bacterial species such as *Pseudomonas*, *Acidovorax* and *Pseudoxanthomonas* (Table 3.3). *Pseudomonas* species have been shown to grow in heavy metal contaminated soils, they require minimum nutrients and some get energy from iron and sulphur oxidation (facultative chemolithotrophic) (Williams, Surridge & Cloete, 2008; Ranalli, Zanardini & Sorlini, 2009; Choudhary & Sar, 2011; Dixit, 2018). *Acidovorax* species have also been shown to be heavy metal tolerant, making them suitable for use in bioremediation (Ghane, 2017). In this study, the Firmicutes phylum consisted of mostly *Bacillus* species. *Bacillus* species have been isolated from various mines worldwide including a South African platinum mine (Marais, 2012). Some *Bacillus* species can detoxify heavy metals (Syed & Chinthala, 2015) and oxidise sulphur

and iron (Zhou *et al*., 2018), enabling this species to inhabit environments with high metal concentrations such as a platinum concentrator plant.

The isolation method in this study selected for mesophilic bacteria (Table 3.3). Some of the bacterial species have not been previously reported as isolates from mines, this includes, *Chelatococcus daeguensis, Pseudoxanthomonas mexicana, Georgenia satyanarayanai, Staphylococcus capitis, Chryseobacterium takakiae, Microbacterium* sp., *Leucobacter chromiiresistens, Nocardia cyriacigeorgica, Comamonas aquatica, Salmonella enterica, Bacillus halotolerans, Paenibacillus alvei, Nocardia farcinica, Alishewanella fetalis, Mucilaginibacter polysacchareus, Pseudarthrobacter phenanthrenivorans, Methylobacterium adhaesivum and Pseudarthrobacter defluvii*. This could be because little research has been conducted on identifying and characterising bacteria from platinum mines.

Some of the bacteria such as *G. satyanarayanai* and *Microbacterium* sp. have been isolated from other environments and have bioleaching capabilities.
Georgenia satyanarayanai has been isolated from a soda lake in India and a study by Shivlata & Tulasi (2015), showed that it has the potential to be used for bioleaching of metals at alkaline sites. Three *Microbacterium* spp. were isolated in this study, from the pond outside the concentrator plant. *Microbacterium* spp. have been previously isolated from soils contaminated with heavy metals (Brown *et al*., 2012), and some species have been shown to reduce metals such as chromium and alter heavy metal motility in contaminated soils (Soni *et al.*, 2014; Kumar & Saini, 2019). It has been postulated that *M. adhaesivum* has the potential to solubilise zinc and produce iron-chelating compounds but more research is required (Yadav & Nath Yadav, 2018). Therefore, these isolated *G. satyanarayanai* and *Microbacterium* bacterial species can potentially be used for PGM bioleaching.

Several of the isolated bacteria have been previously isolated from mines such as copper, coal, iron, and platinum mines. This includes *Arthrobacter globiformis*, *Bacillus* sp., *Pseudomonas stutzeri, Acidovorax* sp., *Rhodococcus* sp., *Rhodococcus pyridinivorans, Pseudomonas aeruginosa, Pseudomonas brassicacearum,*

Pseudomonas alcaliphila, Pseudomonas otitidis and Pseudomonas alcaligenes. Four *P. stutzeri* bacterial species were isolated from the concentrator plant. Isolate 8 comes from the Rougher Feed, 13 from the Cleaner Scavenger Tail A, 29 and 33 from Rougher Tail A and B respectively. Palanivel *et al.* (2020), found that *P. stutzeri* has copper bioremoval and bioaccumulation properties. These properties are important during the base metal bioleaching of PGMs. Three *Acidovorax* sp. (11,12 and 15) were isolated from the pond outside of the concentrator plant. A study by Xie *et al*. (2007), on the microbial population at several acid mineral bioleaching systems at a copper mine found that *Acidovorax* sp. were present in these populations, thus suggesting that they have high metal tolerance and could potentially be used for bioleaching. *Acidovorax* sp. have also been shown to be present in the bacterial community involved in chalcopyrite bioleaching in a shake flask system (He *et al*., 2010).

Two *Rhodococcus* bacterial species have been identified in this study, one from the concentrator plant and another from the pond outside the concentrator plant. *Rhodococcus* species including *R. pyridinivorans* have been isolated from copper and gold mines and some of these species are also capable of copper biosorption (Neeratanaphan, Rungsunsombut & Phonimdaeng, 2015; Sulistinah, Riffiani & Sunarko, 2015; Baltazar *et al.*, 2019). *Pseudomonas aeruginosa* has low nutritional requirements and has been isolated from low nutrient environments including uranium mine waste and iron mine (Williams, Surridge & Cloete, 2008; Choudhary & Sar, 2011). In this study *P. aeruginosa* was isolated from a soil sample collected in front of the crusher. It has previously been shown to be tolerant to copper, nickel and chromium and has the potential for use in bioleaching and bioremediation (Abdelbary, Elgamal & Farrag, 2019).

Eight *Bacillus* sp. isolates were identified, isolates 31 and 34 come from the Cleaner Scavenger Tail A and isolate 10 from Cleaner Scavenger Tail B samples taken from the concentrator plant. *Bacillus* sp. isolates 7, 25 and 28 come from a soil and liquid sample taken from the pond outside the concentrator plant and isolates 32 and 42 come from a soil sample taken from in front of the crusher. *Bacillus* sp. was the

most identified bacterial species in this study and inhabited 3 out of the 4 collection sites. *Bacillus* sp. have previously been isolated from several areas including platinum, coal and chromite mines (Dhal *et al*., 2010; Marais, 2012; Rekha, Dhanalakshmi & Sudha, 2013; Banerjee e*t al*., 2019). One *P. brassicacearum* bacterial isolate was identified from the pond outside of the concentrator plant it has previously been isolated from copper mines (Hassanshahian *et al*., 2019). Two *A. globiformis* bacterial isolates were identified, both come from a soil sample taken from in front of the crusher. *A. globiformis* is commonly found in the rhizosphere of graminaceous crops and has been isolated from an abandoned mine (Sharma *et al*., 2016).

In addition to organic acid-producing bacteria, iron and sulphur oxidising bacteria can be used for metal bioleaching. *Pseudomonas aeruginosa*, *Bacillus* sp., *P. stutzeri* and *P. brassicacearum* can oxidise iron and sulphur (Mason & Kelly, 1988; Cornelis & Dingemans, 2013; Xia *et al*., 2017; Zhou *et al*., 2018; Hassanshahian *et al*., 2019). Six *P. mexicana* bacterial isolates were identified from the concentrator plant. Isolates 4 and 35 come from the Rougher Feed, isolates 2 and 43 from Scavenger Tail A, isolate 36 from the cleaner feed and isolate 3 from the Cleaner Scavenger Tail A. *Pseudomonas mexicana* has previously been isolated from an anaerobic digester and the rhizosphere of a leguminous plant can oxidise sulphur (Thierry *et al*., 2004; Ghosh & Roy, 2007; Krishnani *et al*., 2010). These bacterial isolates have the potential to be used for the bioleaching of PGMs through the oxidation of sulphur and iron.

3.4 Summary and conclusions

The bacteria prefer growing at a pH between 6-9. Therefore, bioleaching should be done between pH 6-9.-The soil samples had a higher number of bacteria compared to the liquid and slurry samples.

A total of 50 mesophilic bacterial isolates were identified from the platinum mine concentrator plant and its surrounding. The bacteria belong to five phyla namely, Proteobacteria, Actinobacteria, Firmicutes, Actinomycetota and Bacteroidetes, with

Proteobacteria being the most dominant. *Pseudomonas* and *Bacillus* species were the most dominant. Several of the isolated bacteria are sulphur oxidising bacteria such as *P. mexicana* and some are iron oxidisers such as *P. stutzeri* and *P. aeruginosa*. Some *Bacillus* sp. can oxidise both iron and sulphur. *Bacillus* sp., *P. mexicana, P. stutzeri, P. aeruginosa* and *P. brassicacearum* can produce organic acids. Several of the identified bacteria have previously been isolated from mines such as copper and platinum mines and these include *P. stutzeri, P. brassicacearum, P. aeruginosa* and *Bacillus* sp. Bioleaching experiments have been conducted using *Acidovorax* sp., *P. aeruginosa* and *P. brassicacearum*.

CHAPTER FOUR: CHARACTERISATION OF BACTERIA

4.1 Introduction

The characterisation of bacteria isolated from mines has focused on their heavy metal tolerance/resistance. However, the bacteria from these mines could have properties desirable for the metal bioleaching process. Bacteria have been used for biotechnological processes, due to their ability to grow fast and produce desirable metabolites. Before using bacteria for any biotechnological process, it is important to characterise the bacteria. This includes their preferred growth conditions, growth rate, and metabolite production.

When growing bacteria to produce metabolites it is important to note what medium results in optimum growth. There are many media available, and the composition of the media varies. Different media provide different carbon sources for the bacteria which can result in different growth rates and metabolite production (Carlson *et al.*, 2020).

Organic acids such as citric, lactic, butyric, propionic, gluconic and formic acids are gaining more attention for use in bioleaching. These acids are biodegradable and effective in metal dissolution (Pathak, Vinoba & Kothari, 2021). In some cases, the organic acid leaching efficiency has been shown to be greater than the typically used inorganic acids (Al-Sheeha *et al.*, 2013). The organic acids dissolve metals through complexation and acidolysis. Bacteria have been shown to produce these organic acids (Anastassiadis *et al.*, 2008; Delvasto *et al*., 2008; Juturu & Wu, 2016; Gonzalez-Garcia *et al.*, 2017). This study focused on characterising six bacterial isolates based on morphology, growth, and organic acid production.

4.2 Materials and Methods

4.2.1 Characterisation of cellular and colony morphology

A total of 50 bacterial isolates were identified, however multiples of the same bacterial species were identified (Table 3.3). In these cases, one of the repeated bacterial species was selected for characterisation, in total 31 bacterial isolates were characterised for cell and colony morphology. The cell and colony morphology were investigated as a general characterisation of the identified bacteria and to see whether the identified bacteria where different, morphologically, to already classified bacteria of the same species. The colony morphology of the bacteria isolated from the Anglo Platinum Mogalakwena Mine was determined. The bacteria grown on LB (pH 7) agar plates were analysed and the shape and colour of the colonies were recorded. The cellular morphology of each isolate was determined using the Gram staining technique according to Coico (2006).

4.2.2 Growth curves

Six bacterial isolates (Table 4.1), *Bacillus* sp., *P. mexicana*, *P. stutzeri*, *A. globiformis*, *P. aeruginosa* and *P. brassicacearum* were isolated from the Anglo Platinum Mogalakwena Mine. An extensive literature review (Table 4.1) was done, and these six bacterial species were selected for characterisation based on their potential to produce organic acids and oxidise iron and sulphur.

Table 4.1: Six bacterial isolates isolated from a platinum mine, the GenBank Accession numbers of closely related species or species with the highest sequence similarity percentage and studies showing their organic acid production and iron and sulphur oxidation capabilities.

Bacterial Isolate	GenBank Accession	Studies
Bacillus sp. (isolate 10)	CP053938.1/ CP053997.1	(Xia *et al*., 2017; Saeid, Prochownik & Dobrowolska-Iwanek, 2018; Zhou *et al*., 2018)
Pseudomonas stutzeri	LR134482.1	(Lalucat et al., 2006; Hao *et al*., 2007; Mahmood *et al*., 2009; Prasanna *et al*., 2010; Li, Li & Li, 2017)
Pseudoxanthomonas mexicana	MK942423.1	(Thierry *et al*., 2004; Krishnani *et al*., 2010)
Arthrobacter globiformis	EU221407.1	(Vijayalakshmi, 2011)
Pseudomonas aeruginosa	HQ018745.1	(Mason & Kelly, 1988; Sankaralingam *et al*., 2014)
Pseudomonas brassicacearum	KT695844.1	(Achouak *et al*., 2000; Kong & Glick, 2017; Hassanshahian *et al*., 2019)

To evaluate the growth and find the optimum growth media of the 6 selected bacterial isolates, growth curves were carried out in triplicates for 45 hours. The 6 isolates were each inoculated into 250 mL of Luria Bertani broth (5 g/L yeast extract, 5 g/L NaCl, 10 g/L tryptone), 9K broth [5 g/L $MgSO_4 \cdot 7H_2O$, 0.05 g/L $Ca(NO_3)_2$, 3 g/L $(NH_4)_2SO_4$, 0.5 g/L K_2HPO_4, 0.1 g/L KCl] and nutrient broth (16 g/L) (NB) and incubated at their respective growth temperatures. The growth media used was all at a pH of 7. The growth media and temperatures were selected based on the growth media used in the studies (Table 4.1) showing organic acid production and iron and sulphur oxidation. *Pseudomonas mexicana*, *P. aeruginosa*, *Bacillus* sp., and the mixed culture, which consisted of all the 6 selected bacterial isolates,

were incubated at 37°C (Thierry *et al*., 2004; Sakil Munna *et al*., 2015; Nur Indah Koni *et al*., 2017). A mixed culture was included to see how the bacteria would grow together. Studies have shown that bioleaching using a mixed culture of bacteria results is more efficient in bioleaching compared to using single cultures (Akcil, Ciftci & Deveci, 2007; Fu *et al.*, 2008; Peng *et al*., 2021) *Pseudomonas stutzeri* was incubated at 35°C (Lalucat *et al*., 2006) and *A. globiformis* and *P. brassicacearum* at 30°C (Achouak *et al*., 2000; Busse & Wieser, 2014). After 3 and 6 hours, 1 mL was sampled from each culture and the optical density (OD_{600}) was measured. One millilitre of LB, 9K and NB were used as blanks. After 20 hours and every 2 hours thereafter, the optical density was measured for 45 hours.

To test for organic acid production during the exponential phase, the 6 bacterial isolates were inoculated into 250 mL of nutrient broth (pH 7) and incubated at their respective temperatures for 16 hours. Thereafter, 1 mL was sampled for optical density to monitor bacterial growth and high-performance liquid chromatography (HPLC) analysis. The organic acids were not extracted prior to analysis, the broth containing the metabolites was analysed. The samples for HPLC analysis were centrifuged at 10000 rpm for 10 minutes. The supernatant was then filtered using 0.45 µm GVS cellulose acetate syringe filters and stored at 4°C until further analysis.

4.2.3 High-performance liquid chromatography

For the analysis of the organic acids, standards of gluconic, lactic, citric, acetic, butyric, propionic, formic, and indole-3-acetic acid were prepared with concentrations ranging from 1mM to 5 mM. The standards were filtered using 0.45 µm GVS cellulose acetate syringe filters. The organic acids were analysed using an Agilent-1200 series HPLC fitted with a Bio-rad fermentation column and refractive index detector. Separation was performed at a column temperature of 65°C and 20% H_2SO_4 was used as the eluent and was delivered at a rate of 1mL/min for 13 minutes.

4.2.4 Statistical analysis.

A one-way ANOVA was performed to analyse statistically significant differences between the mean growth of the 6 bacterial isolates in the three different media, namely 9K, LB and NB.

4.3 Results and Discussion

4.3.1 Phylogeny and Colony and cellular morphology

The Gram stain, cell shape and colony morphology of the isolated bacterial species are shown in Table 4.2 below.

Table 4.2: Morphological characteristics of the bacterial isolates grown on LB agar.

Bacterial Isolates	Gram stain	Cell shape	Colony morphology
Chelatococcus daeguensis	-	Rod	Circular, white opaque
Pseudoxanthomonas mexicana	-	Rod	Irregular, light brown
Arthrobacter globiformis	+	Rod/cocci	Circular, transparent white
Bacillus sp.	+	Rod	Circular, white
Pseudomonas stutzeri	-	Rod	Irregular, cream white
Georgenia satyanarayanai	+	Rod	Circular, yellow
Acidovorax sp.	-	Rod	Irregular, cream white
Microbacterium sp.	+	Rod	Irregular, cream white
Staphylococcus capitis	+	Cocci	Circular, white
Rhodococcus sp.	+	Rod/cocci	Circular, pale pink
Rhodococcus pyridinivorans	+	Rod/cocci	Circular, pale pink
Chryseobacterium takakiae	-	Rod	Circular, translucent yellow
Microbacterium lacusdiani	+	Rod	Circular, yellow
Microbacterium foliorum	+	Rod	Circular, pale yellow
Leucobacter chromiiresistens	+	Rod	Circular, yellow
Nocardia cyriacigeorgica	+	Rod	Circular, white
Pseudomonas aeruginosa	-	Rod	Circular, opaque white
Pseudomonas brassicacearum	-	Rod	Circular, white
Pseudomonas alcaliphila	-	Rod	Irregular, translucent white
Comamonas aquatica	-	Curved rod	Circular, white
Salmonella enterica	-	Rod	Irregular, translucent white
Bacillus halotolerans	+	Rod	Circular, cream white
Paenibacillus alvei	+	Rod	Circular, white
Pseudomonas otitidis	-	Rod	Irregular, cream white
Nocardia farcinica	+	Rod	Circular, cream white
Alishewanella fetalis	-	Rod	Circular, cream white
Pseudomonas alcaligenes	-	Rod	Circular, translucent white
Mucilaginibacter polysacchareus	-	Rod	Circular, light pink
Pseudarthrobacter phenanthrenivorans	+	Rod/cocci	Circular, cream white
Methylobacterium adhaesivum	-	Rod	Circular, light pink
Pseudarthrobacter defluvii	+	Rod/cocci	Circular, cream white

In this study, the number of Gram-positive and Gram-negative bacterial species was similar as shown in Table 4.2. Despite this, more Gram-negative bacterial species were isolated from the concentrator plant compared to the other sampling sites. The concentrator plant could have a higher concentration of heavy metals when compared to the other sampling sites as shown by a study performed by Ekosse (2006), which showed that the heavy metal concentration of the samples increased closer to the concentrator plant. This finding supports the theory that Gram-negative bacteria are more tolerant to heavy metals compared to their Gram-positive counterparts. Piotrowska-Seget, Cycoń & Kozdrój, (2005), found that metal tolerance has been more associated with Gram-positive bacteria such as those in the genera *Corynebacterium*, *Bacillus* and *Arthrobacter*. Several of the bacteria isolated in our study were identified as *Bacillus*, which corresponds with the findings of Ellis *et al*. (2003), that there was a higher abundance of *Bacillus* sp. in most soils contaminated with heavy metals.

Several studies have shown that Gram-negative bacteria are better adapted and more tolerant to heavy metal environments compared to Gram-positive bacteria (Hassen *et al.*, 1998; Wenderoth & Reber, 1999; Bennisse *et al.*, 2004). Benyehuda *et al*. (2003), stated that Gram-negative bacteria are more tolerant to heavy metals due to the complex structure of their cell wall, which forms a more effective barrier of permeability against the heavy metals thereby limiting the access of metals into the cells. They also argued that the surface structures that form the cell wall can interact with metal ions and detoxify the ions (Benyehuda *et al*., 2003). Another study by Mounaouer, Nesrine & Hassen (2014), also showed that Gram-negative bacteria are more heavy metal tolerant when compared to Gram-positive bacteria. It was stated that in addition to their cell wall, Gram-negative bacteria can produce low molecular weight proteins that can bind metals resulting in a conformational modification of the molecules into higher molecular weight molecules that are then unable to diffuse through the plasma membrane and enter the cell (Mounaouer, Nesrine & Hassen, 2014). Some Gram-negative bacteria do not have metal protein transporters which thus prevents or stops the entry of metals into the cell thus increasing its resistance or tolerance to heavy metals (Mounaouer, Nesrine & Hassen, 2014).



Figure 4.1: Phylogenetic analysis based on a partial sequence of the 16S rRNA gene inferred using the maximum likelihood method, showing the relationship of the six bacterial species and bacterial isolates of the same species.

Figure 4.1 shows the phylogenetic relationship of the six selected bacterial species and bacterial isolates of the same species identified in chapter 3. The six bacterial isolates from the present study showed a variety of morphological characteristics after incubation for 48 hours on nutrient agar as shown in Figure 4.2. The *Bacillus* sp. had circular, white colonies which is consistent with what was found in our study (Figure 4.2D). *Pseudoxanthomonas mexicana* had irregularly shaped, cream white colonies. *Pseudomonas aeruginosa* shown in Figure 4.2E, had irregularly shaped, opaque colonies. *Pseudomonas brassicacearum* had small white colonies as shown in Figure 4.2C. *Arthrobacter globiformis* has small white circular colonies (Figure 4.2A) and has previously been isolated from an abandoned mine (Sharma *et al.*, 2016). In Figure 4.1F, *P. stutzeri* has cream white wrinkle-like, irregularly shaped colonies.



Figure 4.2: Morphological characteristics shown by the six bacterial isolates. The bacteria were identified as A) Arthrobacter *globiformis*, B) *Pseudoxanthomonas mexicana*, C) *Pseudomonas brassicacearum*, D) *Bacillus sp.*, E) *Pseudomonas aeruginosa*, and F) *Pseudomonas stutzeri.*

4.3.2 Growth curves

To determine which media is optimal for the growth of the 6 bacterial isolates, growth curves were conducted in 9K, LB and NB media. All 6 bacterial isolates were shown to grow best in NB media compared to 9K and LB media as shown in Figures 4.3, 4.4 and 4.5. There was little to no growth in 9K media and an

unsubstantial amount of growth in LB. For *Bacillus* sp., *P. stutzeri* and the mixed culture, the exponential phase was between hours 3 and 20. The exponential phase for *A. globiformis*, *P. mexicana* and *P. aeruginosa* was between 6 and 20 hours. *Pseudomonas brassicacearum* had slower growth in NB compared to the other isolates, with an exponential phase starting after 20 hours.



Figure 4.3: Growth curves of A) *Arthrobacter globiformis* and B) *Pseudoxanthomonas mexicana* grown in 9K, LB and NB broth. The error bars represent the standard deviations of the three bacterial replicates. The linear breaks on the x-axis indicate the missing sampling periods. The statistical significance was analysed using one-way ANOVA with $p < 0.05$. The bacteria showed the greatest growth in NB media and the least growth in 9K media.



Figure 4.4: Growth curves of A) *Pseudomonas brassicacearum* and B) *Bacillus sp.* C) *Pseudomonas aeruginosa* grown in 9K, LB and NB broth. The error bars represent the standard deviations of the three bacterial replicates. The linear breaks on the x-axis indicate the missing sampling periods. The statistical significance was analysed using one-way ANOVA with $p < 0.05$. The bacteria showed the greatest growth in NB media and the least growth in 9K media.

A)



B)



Figure 4.5: Growth curves of A) *Pseudomonas stutzeri* and B) mixed culture grown in 9K, LB and NB broth**.** The error bars represent the standard deviations of the three bacterial replicates. The linear breaks on the x-axis indicate the missing sampling periods. The statistical significance was analysed using one-way ANOVA with $p < 0.05$. The bacteria showed the greatest growth in NB media and the least growth in 9K media.

The growth rates of each of the 6 bacterial isolates were calculated during the exponential phase using the slope of the log of the growth curve. The following equation was used to calculate the respective bacterial growth rates: $\mu = \frac{2.303(\log OD_2 - \log OD_1)}{t_2 - t_1}$

$\mu = 2.303(\log OD_2 - \log \mu = \frac{2.303(\log OD_2 - \log OD_1)}{t_2 - t_1.}) / t_2 - t_1.$

Where μ is the growth rate, OD_1 is the initial optical density, OD_2 is the final optical density, t_1 is the initial time and t_2 is the final time.

The growth of each bacterial isolate is shown in Table 4.3.

Table 4.3: The growth rates of *Arthrobacter globiformis*, *Pseudoxanthomonas mexicana*, *Pseudomonas stutzeri*, *Pseudomonas brassicacearum*, *Bacillus sp.*, *Pseudomonas aeruginosa* and mixed culture grown in LB and NB media. The bacteria grown in the NB medium had the highest growth rate.

Bacteria	*Bacillus sp.*		*P. mexicana*		*P. stutzeri*		*A. globiformis*		*P. aeruginosa*		*P. brassicacearum*		Mixed culture	
Media	LB	NB	LB	NB	LB	NB	LB	NB	LB	NB	LB	NB	LB	NB
Growth rate (μ/hr)	0,015	0,039	0,002	0,256	0,069	0,350	0,069	0,304	0,027	0,197	0,071	0,449	0,021	0,028

The results show that NB is the optimum growth medium for the bacterial isolates when compared to LB and 9K. The composition of NB and LB differs significantly, with the main difference being that NB contains peptone and meat extract and LB contains tryptone. Tryptone is a mixture of peptides that is produced through the enzymatic digestion of casein by trypsin. Tryptone provides nitrogen in the form of amino acids which play a role in the synthesis of protein for the bacteria (Godbey, 2014). Meat peptone found in nutrient broth is produced through the enzymatic digestion of animal bovine tissue resulting in peptides and amino acids which provides a nitrogen source for microorganisms (Sanchez-Rosario & Johnson, 2021). The amino acid composition of tryptone and meat peptone differ, which can result in a difference in growth of the bacteria in the two different media. Although theoretically, amino acids can assist the growth of a variety of bacteria, the amount and distribution of amino acids can potentially alter the growth of bacteria.

Meat extract found in nutrient broth is produced from low-fat meat such as bovine heart and tendon content. It contains a combination of vitamins, peptides, amino

acids, and minerals (Pearson, 2003). Compared to peptone, meat extract does not undergo severe protein hydrolysis treatment, as a result, it can also provide phosphates and additional energy sources that might not be found in peptone. Little to no growth was observed for the bacterial isolates grown in 9K. 9K contains no organic compounds that the bacteria can use for energy, this shows that the bacterial isolates are heterotrophic and require an organic carbon source to grow.

It has been suggested that mixed bacterial cultures grow better than single bacterial cultures. In nature, bacteria flourish in compounded communities and the survival and fitness of a single bacterium is dependent on its interactions with other bacteria in the population (West *et al.*, 2006). Studies by (Paczia *et al.*, 2012; Fiore *et al.*, 2015) have shown that bacteria in communities cooperate metabolically, where the metabolites from one bacterial species are used as energy sources for another bacterial species in the population. In this study, the mixed culture did not grow better than the other bacterial isolates as shown by the growth rates in Table 4.3. This could be caused by some of the isolates producing antimicrobial compounds that inhibit the growth of the other bacterial isolates in the culture. *Pseudomonas stutzeri* has been shown to produce an antimicrobial compound, zafrin [4betamethyl-5, 6, 7, 8 tetrahydro-1 (4beta-H)-phenanthrenone], which has been shown to act against Gram-positive and Gram-negative bacteria (Uzair *et al.*, 2008). *Pseudomonas brassicacearum* produces hydrogen cyanide and the antimicrobial compound, 2,4-DAPG. This could result in the inhibition of bacterial growth (Zhou *et al.*, 2012).

The one-way ANOVA test was used to determine if there is a statistically significant difference between the mean growth of the bacterial isolates grown in the three different media. The p-value for all the bacterial isolates was less than 0.05. Therefore, there is a significant difference between the mean growth of the bacterial isolates grown in the different media. The media used have different compositions as explained previously, thus resulting in the difference in growth.

From the growth curves it was determined that the exponential phase of the bacterial isolates except for *P. brassicacearum* was approximately between 3 and 20 hours. The organic acid production and substrate consumption during the 16–22-hour

growth period of the bacterial isolates was tested. Figure 4.6 shows the optical density of the 6 bacterial isolates and the mixed culture. During this period *Bacillus* sp. has the highest optical density followed by the mixed culture. The optical densities of *A. globiformis*, *P. mexicana*, *P. brassicacearum* and *P. aeruginosa* are low compared to the other bacterial isolates.



Figure 4.6: Optical density (600 nm) of *Bacillus* sp., *Pseudoxanthomonas mexicana*, *Pseudomonas stutzeri*, *Arthrobacter globiformis*, *Pseudomonas aeruginosa*, *Pseudomonas brassicacearum* and the mixed culture grown in nutrient broth. The error bars represent the standard deviations of the bacterial replicates.

Colony forming units of the 6 bacterial (Figure 4.7) were used to estimate the number of metabolically viable cells between 16 – 22 hours. The colonies of the mixed culture were too many to count (>300 colonies). *Bacillus* sp. has the highest CFUs followed by *P. stutzeri* this corresponds with the optical density values shown in Figure 4.6. However, after 22 hours all the bacterial isolates except *P. mexicana* had similar CFUs.



Figure 4.7: The colony forming units (CFUs) of *Bacillus* sp., *Pseudoxanthomonas mexicana*, *Pseudomonas stutzeri*, *Arthrobacter globiformis*, *Pseudomonas aeruginosa*, *Pseudomonas brassicacearum* grown in nutrient agar. The error bars represent the standard deviations of the bacterial replicates.

4.3.3 Organic acid production

The bacterial isolates were tested for organic acid production and the results are displayed in Table 4.4. All the bacterial isolates produced citric and lactic acid. Gluconic acid was only produced by *Bacillus* sp., *P. mexicana*, *P. stutzeri*, *A globiformis* and *P. aeruginosa*. Indole-3-acetic acid was produced by *Bacillus* sp., *P. mexicana* and *P. aeruginosa*. Butyric acid was produced by *P. mexicana* and *P. stutzeri*. Propionic acid was produced by *Bacillus* sp., *P. mexicana*, *A. globiformis*, and the mixed culture. Formic acid was produced by *P. mexicana*, *A. globiformis*, *P. aeruginosa* and the mixed culture. None of the bacterial isolates produced acetic acid.

Table 4.4: Organic acid production by the six bacterial isolates.

		Organic acid concentration (mM)						
Bacteria	**Time (Hours)**	**Gluconic acid**	**Citric acid**	**Indole-3acetic acid**	**Lactic acid**	**Butyric acid**	**Propionic acid**	**Formic acid**
Bacillus sp.	16	28,269±1,801		3,180±1,042	1,675±0,440			
	18	0,082±0.013						
	20		3,455±0,402		4,388±1,465			
	22		2,072±0,408		4,024±0,541		6,974±0,112	
P. mexicana	16	29,833±1,369		2,904+1,993	4,904±0,912	0,116±0.031		
	18		27,329±0,299		1,095±0,974	0,039±0,059		
	20		2,610±0,435		5,160±0,276		7,971±0,145	4,027±0,556
	22		2,506±0,211				8,174±0,871	
P. stutzeri	16	27,846±1,295						
	18					0,394±0,513		
	20							
	22		2,592±0,580		5,870±0,880			
A. globiformis	16	34,894±0,580					2,839±0,163	
	18				7.558±1,452			
	20		2,123±1,004		2,991±0,301		5,566±1,007	
	22		2,071±0,613		5,450±0,494		5,838±0,890	3,570±0,817
P. aeruginosa	16	30,173±1,611						
	18			4,162±1,722			0,914±0,515	8,644±0,702
	20		1,924±0,220		0,144±0,150		5,370±0,509	
	22		2,199±1,006		7,241±0,766		2,040±0,931	6,429±0,549
P. brassicacearum	16				2,700±0,460			
	18		0,003±0,004					
	20		3,078±1,016		16,178±1,363			
	22		2,471±0,556		9,982±0,182			
Mixed culture	16						2,301±0,631	
	18		2,821±0,258		9,307±0,850			
	20		1,985±0,300		18,973±0,085		4,877±0,283	
	22		3,705±0,859					0,103±0,126

1-Blank space indicates a concentration of 0 mM.

2- Acetic acid is not included in the table because none of the bacterial species produced it.

The use of organic acids for bioleaching has gained traction due to them being costeffective and economically friendly compared to chemical leaching. Organic acids are also biodegradable and result in little environmental pollution. The selected organic acids, except indole-3-acetic acid, have been used in bioleaching experiments. Gluconic acid has been used in the bioleaching of vanadium (Bellenberg *et al.*, 2021) and rare earth elements (Reed *et al.*, 2016). *Bacillus* sp., *P. mexicana*, *P. stutzeri*, *A. globiformis* and *P. aeruginosa* produced a gluconic acid at 16 hours. Little to no gluconic acid was detected in the 18–22-hour samples for these bacterial species therefore the presence of gluconic acid in the 16-hour sample seems to be an artifact. *Pseudomonas, Acetobacter* and *Gluconobacter* species have been shown to produce gluconic acid (Ramachandran *et al*., 2006). A study by Delvasto *et al*. (2008), found that a *Bacillus* species could produce 34 mM of gluconic acid after 7 days of growth. In our study *Bacillus* sp. produced 28,269 mM. An increase in gluconic acid production can be achieved by altering the growth temperature and medium. Nonetheless, these results show that the bacterial isolates have the potential to produce gluconic acid which can then be used in the bioleaching of PGMs.

Citric acid has been used in several bioleaching experiments. A study by Li *et al*. (2014), showed that the use of citric acid increased the metal leaching of copper tailings. Williams & Cloete (2010), investigated the use of microbial citric acid for the removal of phosphorus and potassium from an iron ore concentrate and found that 23.53% of phosphorus and potassium were removed. Another study by Barnett, Palumbo-Roe & Gregory (2018), used a *Bacillus* sp. that produced citric, gluconic and oxalic acid to bioleach rare earth elements and found that it was effective when compared to other leaching techniques. In our study, *P. mexicana* produced the most citric acid, however, to date no other known studies have shown that this bacterium can produce citric acid. *Bacillus, Pseudomonas* and *Arthrobacter* species have been shown previously to produce citric acid (Anastassiadis *et al*., 2008; Vyas & Gulati, 2009).

Indole-3-acetic (IAA) acid is a plant growth-promoting compound that is commonly produced by soil microorganisms found in the rhizosphere of plants

(Zhang *et al*., 2021).*Pseudomonas mexicana, Bacillus sp*. and *P. aeruginosa* from the current study produced IAA. The bacterial isolates have previously been isolated as endophytes which grow in close association with plants and produce plant growth-promoting compounds such as IAA (Liaqat & Eltem, 2016; Devi *et al*., 2017; Lopes *et al*., 2018).

Lactic acid has been used for the leaching of cobalt, nickel, lithium, and manganese from spent lithium-ion batteries with a recovery rate of up to 98.9% (Li *et al*., 2017). It has also been used, in conjunction with other organic acids for the dissolution of cobalt and nickel from limonite ores (Tang & Valix, 2004). *Pseudomonas mexicana* produced the most lactic acid followed by *A. globiformis*. The variance in lactic acid production for these bacterial species could be a result of the acidification of the medium during organic acid production. Lactic acid is commonly produced by lactic acid bacteria (LAB) which include *Lactobacillus*, *Leuconostoc* and *Lactococcus* species (Juturu & Wu, 2016). However, these bacteria are fastidious, have complex nutritional requirements and grow at low temperatures. *Bacillus* species have been shown to produce high amounts of lactic acid when grown at a high pH (Abedi & Hashemi, 2020). The increased production of lactic acid usually requires high temperatures because amylases, which hydrolyse starch to lactic acid, have increased activity at higher temperatures (Juturu & Wu, 2016). Thus, increasing the temperature and pH for the bacterial isolates could potentially result in higher lactic acid production.

Butyric acid has been used in the bioleaching of zinc (Wang *et al*., 2019), while acetic acid has been used for the leaching of scandium (Kiskira *et al*., 2021), zinc (Elomaa *et al*., 2019) and lead (Osasona, Oke & Adebayo, 2021). Butyric acid was only produced by two of the bacterial isolates in very low amounts. Butyric acid production is commonly an anaerobic process and is produced by strictly anaerobic bacteria such as *Clostridium* and *Butyrivibrio* species. The culture conditions favourable for butyric acid production include temperatures between 35-37°C and glucose, sucrose, and lactose as the preferred carbon sources. In this study, NB was used as the medium of choice and did not contain the necessary carbon sources for butyric acid production. The bacterial isolates were also grown at different

temperatures with *A. globiformis* and *P. brassicacearum* being grown at 30°C, which can account for these isolates not producing any butyric acid. The bacterial isolates were also grown in aerobic conditions, which is not favourable for butyric acid production and could have resulted in little to no butyric acid being produced.

Acetic acid was not produced by any of the bacterial isolates. Of the 6 bacterial isolates, only *Bacillus* sp. has been shown to produce acetic acid (Yan *et al*., 2013; Mumtaz *et al.*, 2019). Acetic acid production is typically an aerobic fermentative process and bacteria from the *Acetobacter* and *Gluconobacter* genera are the most commonly used (Xu, Shi & Jiang, 2011). The medium used to grow up the bacteria as well as the growth temperature could have resulted in no acetic acid production.

Formic acid has been used to enhance the bioleaching of copper from chalcopyrite concentrate (Arslanoğlu & Yaraş, 2020). The study also showed that formic acid could be used as a leaching agent for the leaching of precious metals containing ores. Formic acid was also used in the bioleaching of cobalt from copper/cobaltrich sulphatic mine tailings, with a recovery of 66% (Zhang *et al*., 2020). *Pseudomonas aeruginosa* produced the most formic acid in our study.

Propionic acid has been used in the chemical leaching of iron ore. It has also been used in combination with other organic acids for the bioleaching of cobalt and copper from lithium batteries (Urias *et al.*, 2021). *Pseudoxanthomonas mexicana* and *A. globiformis* produced the most propionic acid. There are no studies that have shown these bacteria to produce propionic acid. Propionic acid is produced by several bacteria including *Propionibacterium acidipropionici*, *Clostridium homopropionicum* and *Propionibacterium jensenii* (Ranaei *et al*., 2020). *Propionibacterium* is suggested as the best producer of propionic acid (GonzalezGarcia *et al.*, 2017). Wang, Jin & Yang (2015), showed that *P. acidipropionici* was able to produce up to 751.890 mM of propionic acid. *Pseudomonas mexicana* and *A. globiformis* produced a total of 16.145 and 14.243 mM of propionic acid respectively. When compared to *P. acidipropionici*, these concentrations are very low. This could be because these bacteria are not common propionic acid producers, and the culture conditions were not favourable for propionic acid production.

4.4 Summary and conclusions

For the morphology of the bacterial isolates, there was not a considerable difference in the number of Gram-positive and Gram-negative species. Most of the bacterial species were rod-shaped and only one was cocci shaped. The six selected bacterial species did not grow in 9K media and there was very little growth in LB broth. The bacteria grew best in NB. The mixed culture which was expected to have the fastest growth did not grow better than the individual bacterial isolates. The bacteria produced organic acids, with *P. aeruginosa* producing the most formic and indole3-acetic acid, *P. mexicana* producing the most lactic and propionic acid, *P. stutzeri* producing the most butyric acid, *A. globiformins* producing the most gluconic acid and the mixed culture producing the most citric acid.

From the growth curves, NB is the best medium to grow the bacteria in. The bacteria produce organic acids which can be used for bioleaching. A mixed culture of the bacteria that produced the most organic acids could be used during bioleaching. However, low amounts of organic acids are produced, and it would be necessary to explore ways to increase organic acid concentrations. Overall, the bacteria have great potential to be used in the bioleaching of PGMs.

CHAPTER 5: CONCLUSIONS AND RECOMMENDATIONS

5.1 Conclusions

This study aimed to isolate, identify, and characterise indigenous bacteria from a platinum mine concentrator plant and its surroundings. The bacteria were identified using molecular techniques and characterised morphologically, by Gram-staining and metabolically through growth curves, analysing and organic acid production. This is the first known study to identify and characterise bacteria from a platinum mine concentrator plant.

An estimation of the abundance of bacteria found at the concentrator plant was made using colony-forming units (CFUs). There was a substantial abundance of bacteria found at the concentrator plant. The pH of the soil where the bacteria inhabit has an influence on the number of bacteria found in that soil. This study showed that the isolated indigenous bacteria preferred growing in soils with a pH between 6-9. Therefore, when growing the bacteria up in the laboratory the media should be at a pH between 6-9.

The molecular identification of the bacteria showed that there were 50 different bacterial isolates. This suggests that the bacteria found at the concentrator plant and its surrounding are quite diverse and it also shows that these bacteria can adapt to unfavourable environments such as heavy metal contaminated environments. Bacteria from the Proteobacteria phylum such as *Pseudomonas, Pseudoxanthomonas* and *Acidovorax* species were the most dominant. Several of the indigenous bacteria such as *P. aeruginosa*, *Bacillus sp.*, *P. brassicacearum*, *P. stutzeri* and *P. mexicana* had some properties that are desirable for bioleaching such as iron and sulphur oxidation and organic acid production.

Of the 50 identified bacterial isolates, six were selected for further characterisation namely, *P. aeruginosa*, *P. mexicana*, *A. globiformis*, *P. stutzeri*, *P. brassicacearum* and *Bacillus sp*. These bacteria grew well in NB and reached the exponential phase within six hours, except *P. brassicacearum* which reached the exponential phase after 20 hours. These bacteria produced gluconic, citric, lactic, propionic, and formic acid which have been shown to be effective in bioleaching. In conclusion, these six indigenous bacteria have great potential to be used in bioleaching.

5.2 Recommendations

This study has shown that bacteria isolated from a platinum mine can be used to produce organic acids. It is therefore recommended that these bacteria be used for organic acid production for bioleaching purposes. It is also suggested that more organic acid production tests using these bacteria are conducted using different culture conditions.

Based on literature, few of the isolated bacteria can oxidise iron and sulphur. It is recommended that these bacteria be tested, to determine whether they are iron and sulphur-oxidising bacteria

The bacteria were isolated from a platinum mine concentrator plant and its surroundings. The presence and concentration of heavy metals at the concentrator plant and its surrounding should be measured. Thereafter performing metal tolerance tests for these bacteria is recommended.

APPENDIX A

A1 Preparation of media used in the isolation and enumeration of bacterial species.

A1.1 Luria-Bertani Agar Medium

Composition

Yeast Extract	5 g/L
Sodium Chloride	5 g/L
Tryptone	10 g/L
Agar	15 g/L

Suspend in 1 L of distilled water

Autoclave at 121°C for 15 minutes

A1.2 9K Agar Medium

Composition

$MgSO_4 \cdot 7H_2O$	5 g/L
$Ca(NO_3)_2$	0.05 g/L
$(NH_4)_2SO_4$	3 g/L
K_2HPO_4	0.5 g/L
KCl	0.1 g/L
Agar	15 g/L

Suspend in 1 L of distilled water

Autoclave at 121°C for 15 minutes

A2 Preparation of culture medium used for the characterisation of the six bacterial isolates.

A2.1 Luria-Bertani Agar Medium

Composition

Yeast Extract	5 g/L
Sodium Chloride	5 g/L
Tryptone	10 g/L

Suspend in 1 L of distilled water

Autoclave at 121°C for 15 minutes

A2.2 Nutrient Broth (Merck)

Composition

Meat Extract	1 g/L
Yeast Extract	2 g/L
5 g/L Peptone	5 g/L
Sodium Chloride	8g/L

Suspend 16 g in 1 L of distilled water

Autoclave at 121°C for 15 minutes

A2.3 9K Broth Medium

Composition

$MgSO_4 \cdot 7H_2O$	5 g/L
$Ca(NO_3)_2$	0.05 g/L
$(NH_4)_2SO_4$	3 g/L
K_2HPO_4	0.5 g/L
KCl	0.1 g/L

Suspend in 1 L of distilled water

Autoclave at 121°C for 15 minutes

APPENDIX B

B1 DNA Extraction Protocol

1. Add sample to a **ZR BashingBead™ Lysis Tubes (0.1 & 0.5 mm)**. Add 750 µL **ZymoBIOMICS™ Lysis Solution** to the tube and cap tightly. ***Note****: For samples stored and lysed in* ***DNA/RNA Shield™ Lysis Tubes****, do not add ZymoBIOMICS™ Lysis Solution and proceed to Step 2.*

Sample Type	**Maximum Input**
Feces	200 mg
Soil	250 mg
Liquid Samples1 and Swab Collections2	250 µL
Cells (isotonic buffer, *e.g*., PBS)	50-100 mg (wet weight) (109 bacterial and 108 yeast cells)
Samples in DNA/RNA Shield™,3	≤ 1 mL

2. Secure in a bead beater fitted with a 2 mL tube holder assembly and process using optimized beat beating conditions (speed and time) for your device.
3. Centrifuge the **ZR BashingBead™ Lysis Tubes (0.1 & 0.5 mm)** in a microcentrifuge at ≥ 10,000 x *g* for 1 minute.
4. Transfer up to 400 µL supernatant to the **Zymo-Spin™ III-F Filter** in a **Collection Tube** and centrifuge at 8,000 x *g* for 1 minute. Discard the Zymo-Spin™ III-F Filter.
5. Add 1,200 µL of **ZymoBIOMICS™ DNA Binding Buffer** to the filtrate in the Collection Tube from Step 4. Mix well.
6. Transfer 800 µL of the mixture from Step 5 to a **Zymo-Spin™ IICR Column** in a Collection Tube and centrifuge at 10,000 x *g* for 1 minute.

7. Discard the flow through from the Collection Tube and repeat Step 6.

8. Add 400 µL **ZymoBIOMICS™ DNA Wash Buffer 1** to the Zymo-Spin™ IICR Column in a new Collection Tube and centrifuge at 10,000 x *g* for 1 minute. Discard the flow-through.

9. Add 700 µL **ZymoBIOMICS™ DNA Wash Buffer 2** to the Zymo-Spin™ IICR Column in a Collection Tube and centrifuge at 10,000 x *g* for 1 minute. Discard the flow-through.

10. Add 200 µL **ZymoBIOMICS™ DNA Wash Buffer 2** to the Zymo-Spin™ IICR Column in a Collection Tube and centrifuge at 10,000 x *g* for 1 minute.

11. Transfer the Zymo-Spin™ IICR Column to a clean 1.5 mL microcentrifuge tube and add 100 µL (50 µL minimum) **ZymoBIOMICS™ DNase/RNase Free Water** directly to the column matrix and incubate for 1 minute. Centrifuge at 10,000 x *g* for 1 minute to elute the DNA5, 6.

12. Place a **Zymo-Spin™ III-HRC Filter** in a new Collection Tube and add 600 µL **ZymoBIOMICS™ HRC Prep Solution**. Centrifuge at 8,000 x *g* for 3 minutes.

13. Transfer the eluted DNA (Step 11) to a prepared Zymo-Spin™ III-HRC Filter in a clean 1.5 mL microcentrifuge tube and centrifuge at exactly 16,000 x *g* for 3 minutes.

REFERENCES

1. Abdelbary, S., Elgamal, M.S. & Farrag, A. 2019. Trends in Heavy Metals Tolerance and Uptake by Pseudomonas aeruginosa, in D. Sriramulu (ed.). *Pseudomonas Aeruginosa - An Armory Within*. IntechOpen, London. DOI: 10.5772/intechopen.85875.

2. Abedi, E. & Hashemi, S.M.B. 2020. Lactic acid production – producing microorganisms and substrates sources-state of art. *Heliyon*. 6(10):e04974. DOI: 10.1016/j.heliyon.2020.e04974.

3. Achouak, W., Sutra, L., Heulin, T., Meyer, J.M., Fromin, N., Degraeve, S., Christen, R. & Gardan, L. 2000. *Pseudomonas brassicacearum* sp. nov. and *Pseudomonas thivervalensis* sp. nov., two root-associated bacteria isolated from Brassica napus and Arabidopsis thaliana. *International Journal of Systematic and Evolutionary Microbiology*. 50(1):9–18. DOI: 10.1099/00207713-50-1-9.

4. Adeleke, R., Cloete, T.E. & Khasa, D.P. 2012. Culturable microorganisms associated with Sishen iron ore and their potential roles in biobeneficiation. *World Journal of Microbiology and Biotechnology*. 28(3):1057–1070. DOI: 10.1007/s11274-011-0904-2.

5. Ahammed, M.M. 2003. Effect of holding time and temperature on bacterial counts. *Indian Journal of Environmental Health*. 45(3):209–212.

6. Akcil, A., Ciftci, H. & Deveci, H. 2007. Role and contribution of pure and mixed cultures of mesophiles in bioleaching of a pyritic chalcopyrite concentrate. *Minerals Engineering*. 20(3):310–318. DOI: 10.1016/j.mineng.2006.10.016.

7. Al-Sheeha, H., Marafi, M., Raghavan, V. & Rana, M.S. 2013. Recycling and Recovery Routes for Spent Hydroprocessing Catalyst Waste. *Industrial & Engineering Chemistry Research*. 52(36):12794–12801. DOI: 10.1021/ie4019148.

8. Anastassiadis, S., Morgunov, I., Kamzolova, S. & Finogenova, T. 2008. Citric Acid Production Patent Review. *Recent Patents on Biotechnology*. 2(2):107–123. DOI: 10.2174/187220808784619757.

9. Arslanoğlu, H. & Yaraş, A. 2020. Chalcopyrite Leaching with Hydrogen Peroxide in Formic Acid Medium. *Transactions of the Indian Institute of Metals*. 73(3):785–792. DOI: 10.1007/s12666-020-01896-x.

10. Bahram, M., Hildebrand, F., Forslund, S.K., Anderson, J.L., Soudzilovskaia, N.A., Bodegom, P.M., Bengtsson-Palme, J., Anslan, S., *et al*. 2018. Structure and function of the global topsoil microbiome. *Nature*. 560(7717):233–237. DOI: 10.1038/s41586-018-0386-6.

11. Baltazar, M. dos P.G., Gracioso, L.H., Avanzi, I.R., Karolski, B., Tenório, J.A.S., do Nascimento, C.A.O. & Perpetuo, E.A. 2019. Copper biosorption by *Rhodococcus erythropolis* isolated from the Sossego Mine – PA – Brazil. *Journal of Materials Research and Technology*. 8(1):475–483. DOI: 10.1016/j.jmrt.2018.04.006.

12. Banerjee, S., Misra, A., Chaudhury, S. & Dam, B. 2019. A Bacillus strain TCL isolated from Jharia coalmine with remarkable stress responses, chromium reduction capability and bioremediation potential. *Journal of Hazardous Materials*. 367:215–223. DOI: 10.1016/j.jhazmat.2018.12.038.

13. Barnett, M., Palumbo-Roe, B. & Gregory, S. 2018. Comparison of Heterotrophic Bioleaching and Ammonium Sulfate Ion Exchange

Leaching of Rare Earth Elements from a Madagascan Ion-Adsorption Clay. *Minerals*. 8(6):236. DOI: 10.3390/min8060236.

14. Bellenberg, S., Turner, S., Seidel, L., van Wyk, N., Zhang, R., Sachpazidou, V., Embile, R.F., Walder, I., *et al*. 2021. Towards Bioleaching of a Vanadium Containing Magnetite for Metal Recovery. *Frontiers in Microbiology*. 12:693615. DOI: 10.3389/fmicb.2021.693615.

15. Bennisse, R., Labat, M., el asli, A., Brhada, F., Chandad, F., Liebgott, P.-P., Hibti, M. & Qatibi, A.-I. 2004. Rhizosphere bacterial populations of metallophyte plants in heavy metal-contaminated soils from mining areas in semiarid climate. *World Journal of Microbiology and Biotechnology*.

20:759–766. DOI: 10.1007/s11274-004-5812-2.

16. Benyehuda, G., Coombs, J., Ward, P.L., Balkwill, D. & Barkay, T. 2003. Metal resistance among aerobic chemoheterotrophic bacteria from the deep terrestrial subsurface. *Canadian Journal of Microbiology*. 49(2):151–156. DOI: 10.1139/w03-012.

17. Birke, M., Rauch, U., Stummeyer, J., Lorenz, H. & Keilert, B. 2018. A review of platinum group element (PGE) geochemistry and a study of the changes of PGE contents in the topsoil of Berlin, Germany, between 1992 and 2013. *Journal of Geochemical Exploration*. 187:72–96. DOI:

10.1016/j.gexplo.2017.09.005.

18. Bosecker, K. 1997. Bioleaching: metal solubilization by microorganisms. *FEMS Microbiology Reviews*. 20(3–4):591–604. DOI: 10.1111/j.15746976.1997.tb00340.x.

19. Brierley, C.L. 2016. Biological Processing: Biological Processing of Sulfidic Ores and Concentrates—Integrating Innovations. In *Innovative Process Development in Metallurgical Industry: Concept*

to Commission. V.I. Lakshmanan, R. Roy, & V. Ramachandran, Eds. Cham: Springer International Publishing. 109–135. DOI: 10.1007/978-3-319-21599-0_6.

20. Brierley, C.L. & Brierley, J.A. 2013. Progress in bioleaching: part B: applications of microbial processes by the minerals industries. *Applied*

Microbiology and Biotechnology. 97(17):7543–7552. DOI: 10.1007/s00253-013-5095-3.

21. Brown, S.D., Palumbo, A.V., Panikov, N., Ariyawansa, T., Klingeman, D.M., Johnson, C.M., Land, M.L., Utturkar, S.M., et al. 2012. Draft Genome Sequence for *Microbacterium laevaniformans* Strain OR221, a Bacterium Tolerant to Metals, Nitrate, and Low pH. *Journal of Bacteriology*. 194(12):3279–3280. DOI: 10.1128/JB.00474-12.

22. Busse, H.-J. & Wieser, M. 2014. The Genus *Arthrobacter*. In *The Prokaryotes: Actinobacteria*. E. Rosenberg, E.F. DeLong, S. Lory, E.

Stackebrandt, & F. Thompson, Eds. Berlin, Heidelberg: Springer. 105–132. DOI: 10.1007/978-3-642-30138-4_204.

23. Carlson, H.K., Lui, L.M., Price, M.N., Kazakov, A.E., Carr, A.V., Kuehl, J.V., Owens, T.K., Nielsen, T., *et al*. 2020. Selective carbon sources influence the end products of microbial nitrate respiration. *The ISME Journal*. 14(8):2034–2045. DOI: 10.1038/s41396-020-0666-7.

24. Cawthorn, R. 2010. The Platinum Group Element Deposits of the Bushveld Complex in South Africa. *Platinum Metals Review*. 54:205–215. DOI: 10.1595/147106710X520222.

25. Chakravarty, S. & Anderson, G. 2015. The Genus *Pseudomonas*. 321–344. DOI: 10.1201/b17871-24.

26. Cho, H., Tripathi, B.M., Moroenyane, I., Takahashi, K., Kerfahi, D., Dong, K. & Adams, J.M. 2019. Soil pH rather than elevation determines bacterial phylogenetic community assembly on Mt. Norikura. *FEMS microbiology ecology*. 95(3). DOI: 10.1093/femsec/fiy216.

27. Choudhary, S. & Sar, P. 2011. Uranium biomineralization by a metal resistant *Pseudomonas aeruginosa* strain isolated from contaminated mine waste. *Journal of Hazardous Materials*. 186(1):336–343. DOI: 10.1016/j.jhazmat.2010.11.004.

28. Coico, R. 2005. Gram Staining. *Current Protocols in Microbiology*. DOI: 10.1002/9780471729259.mca03cs00.

29. Cole, S. & Ferron, C.J. 2002. A Review of the Beneficiation and Extractive Metallurgy of the Platinum Group Elements, Highlighting Recent Process Innovations. SGS Minerals Services Technical Paper 2002-03, 1–43.

30. Cornelis, P. & Dingemans, J. 2013. *Pseudomonas aeruginosa* adapts its iron uptake strategies in function of the type of infections. *Frontiers in Cellular and Infection Microbiology*. 3. Available: https://www.frontiersin.org/article/10.3389/fcimb.2013.00075 [2022, January 27].

31. Cramer, L.A. 2001. The extractive metallurgy of south africa's platinum ores. *JOM*. 53(10):14–18. DOI: 10.1007/s11837-001-0048-1.

32. Cramer, L.A. 2008. What is your PGM concentrate worth? In *Third International Platinum Conference 'Platinum in Transformation'*; The Southern African Institute of Mining and Metallurgy: Johannesburg, South Africa, 2008; pp. 387–394.

33. Delvasto, P., Valverde, A., Ballester, A., Muñoz, J.A., González, F., Blázquez, M.L., Igual, J.M. & García-Balboa, C. 2008. Diversity and activity of phosphate bioleaching bacteria from a high-phosphorus iron ore.

Hydrometallurgy. 92(3):124–129. DOI: 10.1016/j.hydromet.2008.02.007.

34. Devi, K.A., Pandey, G., Rawat, A.K.S., Sharma, G.D. & Pandey, P. 2017. The Endophytic Symbiont—*Pseudomonas aeruginosa* Stimulates the Antioxidant Activity and Growth of *Achyranthes aspera* L. *Frontiers in Microbiology*. 8:1897. DOI: 10.3389/fmicb.2017.01897.

35. Dhal, B., Thatoi, H., Das, N. & Pandey, B.D. 2010. Reduction of hexavalent chromium by *Bacillus* sp. isolated from chromite mine soils and characterization of reduced product. *Journal of Chemical Technology & Biotechnology*. 85(11):1471–1479. DOI: 10.1002/jctb.2451.

36. Dixit, S.J. 2018. Genetic characterization of sulphur and iron oxidizing bacteria in manganese mining area of Balaghat and Chhindwara, Madhya Pradesh, India. *Indian Journal of Biotechnology*. 17(4): 595-601

37. Domingues, V.S., de Souza Monteiro, A., Júlio, A.D.L., Queiroz, A.L.L. & dos Santos, V.L. 2020. Diversity of Metal-Resistant and TensoactiveProducing Culturable Heterotrophic Bacteria Isolated from a Copper Mine in Brazilian Amazonia. *Scientific Reports*. 10(1):6171. DOI: 10.1038/s41598-020-62780-8.

38. Ekosse, G.-I. 2006. Heavy metals concentrations in the biophysical environment around the Ni-Cu mine and the smelter/concentrator plant,

Selebi Phikwe, Botswana. *Global Journal of Environmental Sciences*. 4:97110. DOI: 10.4314/gjes.v4i2.2448.

39. Ekpa, E., Adaji, A., Odiba, J., Iyere, E., Ohi, P., Dorcas, U. & Abubakar, A. 2017. Isolation, identification and characterization of some bacteria from soil samples of agbaja iron ore mining site of kogi state. *Journal of*

Bacteriology & Mycology: Open Access. 4(3): 79-84. DOI: 10.15406/jbmoa.2017.04.00092.

40. Ellis, R.J., Morgan, P., Weightman, A.J. & Fry, J.C. 2003. Cultivationdependent and -independent approaches for determining bacterial diversity in heavy-metal-contaminated soil. *Applied and Environmental*

Microbiology. 69(6):3223–3230. DOI: 10.1128/AEM.69.6.32233230.2003.

41. Elomaa, H., Seisko, S., Lehtola, J. & Lundström, M. 2019. A study on selective leaching of heavy metals vs. iron from fly ash. *Journal of Material*

Cycles and Waste Management. 21(4):1004–1013. DOI: 10.1007/s10163019-00858-w.

42. Enrich-Prast, A., Bastviken, D., Crill, P., Santoro, A.L., Signori, C.N. & Sanseverino, A.M. 2014. Chemosynthesis. In *Reference Module in Earth*

Systems and Environmental Sciences. Elsevier. 1-19. DOI: 10.1016/B9780-12-409548-9.09054-0.

43. Fashola, M.O., Ngole-Jeme, V.M. & Babalola, O.O. 2020. Physicochemical properties, heavy metals, and metal-tolerant bacteria profiles of abandoned gold mine tailings in Krugersdorp, South Africa. *Canadian Journal of Soil Science*. 100(3):217–233. DOI: 10.1139/cjss-2018-0161.

44. Fashola, O., Ngole-Jeme, V., Olubukola, · & Babalola, O. 2020. Heavy Metal Immobilization Potential of Indigenous Bacteria Isolated from Gold

Mine Tailings. *International Journal of Environmental Research*. 14:71–86.

45. Fernandes, C.C., Kishi, L.T., Lopes, E.M., Omori, W.P., Souza, J.A.M. de, Alves, L.M.C. & Lemos, E.G. de M. 2018. Bacterial communities in mining soils and surrounding areas under regeneration process in a former ore mine.

Brazilian Journal of Microbiology. 49(3):489–502. DOI: 10.1016/j.bjm.2017.12.006.

46. Fiore, C.L., Longnecker, K., Kido Soule, M.C. & Kujawinski, E.B. 2015. Release of ecologically relevant metabolites by the cyanobacterium S*ynechococcus elongatus* CCMP 1631. *Environmental Microbiology*. 17(10):3949–3963. DOI: 10.1111/1462-2920.12899.

47. Fu, B., Zhou, H., Zhang, R. & Qiu, G. 2008. Bioleaching of chalcopyrite by pure and mixed cultures of *Acidithiobacillus* spp. and *Leptospirillum ferriphilum*. *International Biodeterioration & Biodegradation*. 62:109–115. DOI: 10.1016/j.ibiod.2007.06.018.

48. Gericke, M. 2012. Review of the role of microbiology in the design and operation of heap bioleaching processes. *Journal of the Southern African Institute of Mining and Metallurgy*. 112(12):1005–1012.

49. Ghane, M. 2017. Isolation of heavy metal resistant *Acidovorax* sp. from agricultural soil and optimization of metal biosorption. Journal of Microbial World. 10(1):37-48.

50. Ghosh, W. & Roy, P. 2007. Chemolithotrophic oxidation of thiosulphate and tetrathionate by novel strains of *Azospirillum* and *Pseudoxanthomonas* isolated from the rhizosphere of an Indian tropical leguminous plant. *Current Science*. 93(11):1613–1615.

51. Giese, E.C. 2021. Influence of organic acids on pentlandite bioleaching by *Acidithiobacillus ferrooxidans* LR. *3 Biotech*. 11(4):165. DOI: 10.1007/s13205-021-02711-0.

52. Gillan, D.C., Roosa, S., Kunath, B., Billon, G. & Wattiez, R. 2015. The long-term adaptation of bacterial communities in metal-contaminated sediments: a metaproteogenomic study. *Environmental Microbiology*.
17(6):1991–2005. DOI: 10.1111/1462-2920.12627.

53. Godbey, W.T. 2014. Genetic Engineering. In *An Introduction to Biotechnology*. Elsevier. 237–274. DOI: 10.1016/B978-1-907568-28-
2.00012-5.

54. Gonzalez-Garcia, R., McCubbin, T., Navone, L., Stowers, C., Nielsen, L. & Marcellin, E. 2017. Microbial Propionic Acid Production. *Fermentation*. 3:21. DOI: 10.3390/fermentation3020021.

55. Granados-Fernández, R., Montiel, M.A., Díaz-Abad, S., Rodrigo, M.A. & Lobato, J. 2021. Platinum Recovery Techniques for a Circular Economy.
Catalysts. 11(8):937. DOI: 10.3390/catal11080937.

56. Gunn, G. 2013. Platinum-group metals. In *Critical Metals Handbook*. G.
Gunn, Ed. Oxford: John Wiley & Sons. 284–311. DOI: 10.1002/9781118755341.ch12.

57. Haldar, S.K. 2013. Economic Mineral Deposits and Host Rocks. In *Mineral Exploration*. Elsevier. 23–39. DOI: 10.1016/B978-0-12-416005-7.00002-7.

58. Halter, D., Cordi, A., Gribaldo, S., Gallien, S., Goulhen-Chollet, F., Heinrich-Salmeron, A., Carapito, C., Pagnout, C., *et al*. 2011. Taxonomic and functional prokaryote diversity in mildly arsenic-

contaminated sediments. *Research in Microbiology*. 162(9):877–887. DOI: 10.1016/j.resmic.2011.06.001.

59. Hao, J., Ma, C., Gao, C., Qiu, J., Wang, M., Zhang, Y., Cui, X. & Xu, P. 2007. *Pseudomonas stutzeri* as a novel biocatalyst for pyruvate production from DL-lactate. *Biotechnology Letters*. 29(1):105–110. DOI: 10.1007/s10529-006-9204-6.

60. Hassanien, W. & Desouky, O. 2014. Bioleaching of some Rare Earth Elements from Egyptian Monazite using Aspergillus ficuum and Pseudomonas aeruginosa. *Walailak Journal of Science and Technology*. 11:809–823. DOI: 10.2004/wjst.v11i6.481.

61. Hassanshahian, M. & Ghoebani, S. 2018. Isolation and Characterization of Iron and Sulfur-Oxidizing Bacteria from Maiduk Copper Mine at Shahrbabk Province in Iran. *Geomicrobiology Journal*. 35(4):261–265. DOI: 10.1080/01490451.2017.1348408.

62. Hassanshahian, M., Toorani, S., Roghanian, R., Emtiazi, G. & Genovese, M. 2019. Characterization of heterotrophic iron and sulfur-oxidizing bacteria from acid mine drainage in Sarcheshmeh copper mine, Iran. *Desalination and Water Treatment*. 147:264–272. DOI: 10.5004/dwt.2019.23723.

63. Hassen, A., Say, S., Cherif, M. & Boudabous, A. 1998. Resistance of environmental bacteria to heavy metal. *Bioresource Technology*. 64:7–15. DOI: 10.1016/S0960-8524(97)00161-2.

64. He, Z., Xie, X., Xiao, S., Liu, J. & Qiu, G. 2007. Microbial diversity of mine water at Zhong Tiaoshan copper mine, China. *Journal of Basic Microbiology*. 47(6):485–495. DOI: 10.1002/jobm.200700219.

65. He, Z., Gao, F., Zhao, J., Hu, Y. & Qiu, G. 2010. Insights into the dynamics of bacterial communities during chalcopyrite bioleaching. *FEMS Microbiology Ecology*. 74(1):155–164. DOI: 10.1111/j.15746941.2010.00943.x.

66. Hedrich, S., Kraemer, D., Junge, M., Marbler, H., Bau, M. & Schippers, A. 2020. Bioprocessing of oxidized platinum group element (PGE) ores as pretreatment for efficient chemical extraction of PGE. *Hydrometallurgy*. 196:105419. DOI: 10.1016/j.hydromet.2020.105419.

67. Irannajad, M., Shabani, M.A., Meshkini, M. & Azadmehr, A.R. 2014. Bioleaching of copper and zinc from oxide resources by a heterotrophic bacterium. Unpublished paper delivered at the Eighth International Chemical Engineering Congress & Exhibitions. 24-27 February, Kish, Iran.

68. Jain, N. & Sharma, D.K. 2004. Biohydrometallurgy for Nonsulfidic Minerals—A Review. *Geomicrobiology Journal*. 21(3):135–144. DOI: 10.1080/01490450490275271.

69. Juturu, V. & Wu, J.C. 2016. Microbial production of lactic acid: the latest development. *Critical Reviews in Biotechnology*. 36(6):967–977. DOI: 10.3109/07388551.2015.1066305.

70. Kamika, I. & Momba, M. 2014. Microbial Diversity of Emalahleni Mine Water in South Africa and Tolerance Ability of the Predominant Organism to Vanadium and Nickel. *PLoS ONE* 9:e86189. DOI: 10.1371/journal.pone.0086189.

71. Keersemaker, M. 2020. Critical Raw Materials. In *Suriname Revisited: Economic Potential of its Mineral Resources*. M. Keersemaker, Ed. (SpringerBriefs in Earth Sciences). Cham:

Springer International Publishing. 69–82. DOI: 10.1007/978-3-030-40268-6_9.

72. Kemp, P.F. & Aller, J.Y. 2004. Bacterial diversity in aquatic and other environments: what 16S rDNA libraries can tell us. *FEMS Microbiology Ecology*. 47(2):161–177. DOI: 10.1016/S0168-6496(03)00257-5.

73. Kiskira, K., Lymperopoulou, T., Tsakanika, L.-A., Pavlopoulos, C., Papadopoulou, K., Ochsenkühn, K.-M., Lyberatos, G. & OchsenkühnPetropoulou, M. 2021. Study of Microbial Cultures for the Bioleaching of Scandium from Alumina Industry By-Products. *Metals*. 11(6):951. DOI: 10.3390/met11060951.

74. Kong, Z. & Glick, B.R. 2017. The Role of Plant Growth-Promoting Bacteria in Metal Phytoremediation. In *Advances in Microbial Physiology*. V. 71.

Elsevier. 97–132. DOI: 10.1016/bs.ampbs.2017.04.001.

75. Korehi, H., Blöthe, M., Sitnikova, M.A., Dold, B. & Schippers, A. 2013. Metal Mobilization by Iron- and Sulfur-Oxidizing Bacteria in a Multiple Extreme Mine Tailings in the Atacama Desert, Chile. *Environmental Science & Technology*. 47(5):2189–2196. DOI: 10.1021/es304056n.

76. Kraemer, D., Junge, M., Oberthür, T. & Bau, M. 2015. Improving recoveries of platinum and palladium from oxidized Platinum-Group

Element ores of the Great Dyke, Zimbabwe, using the biogenic siderophore Desferrioxamine B. *Hydrometallurgy*. 152:169–177. DOI: 10.1016/j.hydromet.2015.01.002.

77. Kraemer, D., Junge, M. & Bau, M. 2017. Oxidized Ores as Future Resource for Platinum Group Metals: Current State of Research. *Chemie Ingenieur Technik*. 89(1–2):53–63. DOI: 10.1002/cite.201600092.

78. Krishnani, K.K., Kathiravan, V., Natarajan, M., Kailasam, M. & Pillai, S.M. 2010. Diversity of Sulfur-Oxidizing Bacteria in Greenwater System of Coastal Aquaculture. *Applied Biochemistry and Biotechnology*. 162(5):1225–1237. DOI: 10.1007/s12010-009-8886-3.

79. Kuang, J.-L., Huang, L.-N., Chen, L.-X., Hua, Z.-S., Li, S.-J., Hu, M., Li, J.-T. & Shu, W.-S. 2013. Contemporary environmental variation determines microbial diversity patterns in acid mine drainage. *The ISME Journal*. 7(5):1038–1050. DOI: 10.1038/ismej.2012.139.

80. Kumar, M. & Saini, H.S. 2019. Reduction of hexavalent chromium (VI) by indigenous alkaliphilic and halotolerant *Microbacterium* sp. M5: comparative studies under growth and nongrowth conditions. *Journal of Applied Microbiology*. 127(4):1057–1068. DOI: 10.1111/jam.14366.

81. Kumar, P., Jyoti, B., Kumar, A. & Paliwal, A. 2019. Chapter 8 - Biotechnological and microbial standpoint cahoot in bioremediation. In *Smart Bioremediation Technologies*. P. Bhatt, Ed. Academic Press. 137–158. DOI: 10.1016/B978-0-12-818307-6.00008-1.

82. Lalucat, J., Bennasar, A., Bosch, R., García-Valdés, E. & Palleroni, N.J. 2006. Biology of *Pseudomonas stutzeri*. *Microbiology and Molecular Biology Reviews*. 70(2):510–547. DOI: 10.1128/MMBR.00047-05.

83. Lane, D.J. 16S/23S rRNA sequencing. In: Stackebrandt,E.; Goodfellow, M., (Eds.). *Nucleic acid techniques in bacterial systematics*. New York: John Wiley and Sons, 1991. p. 115-175.

84. Lee, J., Kim, H.-S., Jo, H.Y. & Kwon, M.J. 2021. Revisiting soil bacterial counting methods: Optimal soil storage and pretreatment methods and comparison of culture-dependent and -independent

methods. *PLoS ONE* 16(2):e0246142. DOI: 10.1371/journal.pone.0246142.

85. Li, D., Yin, Y., Binquan, J., Song, L.-Y., Wang, Y., Qiao, J. & Xu, Z. 2014. Chemical dissolving of citric acid on bioleaching of copper mine tailings by *Acidithiobacillus ferrooxidans* ATCC 23270. *Research journal of biotechnology*. 8.

86. Li, L., Fan, E., Guan, Y., Zhang, X., Xue, Q., Wei, L., Wu, F. & Chen, R. 2017. Sustainable Recovery of Cathode Materials from Spent Lithium-Ion Batteries Using Lactic Acid Leaching System. *ACS Sustainable Chemistry & Engineering*. 5(6):5224–5233. DOI: 10.1021/acssuschemeng.7b00571.

87. Li, S., Li, X. & Li, F. 2017. Fe(II) oxidation and nitrate reduction by a denitrifying bacterium, *Pseudomonas stutzeri* LS-2, isolated from paddy soil. *Journal of Soils and Sediments*. DOI: 10.1007/s11368-017-1883-1.

88. Liaqat, F. & Eltem, R. 2016. Identification and characterization of endophytic bacteria isolated from in vitro cultures of peach and pear rootstocks. *3 Biotech*. 6(2):120. DOI: 10.1007/s13205-016-0442-6.

89. Lopes, R., Tsui, S., Gonçalves, P.J.R.O. & de Queiroz, M.V. 2018. A look into a multifunctional toolbox: endophytic *Bacillus* species provide broad and underexploited benefits for plants. *World Journal of Microbiology & Biotechnology*. 34(7):94. DOI: 10.1007/s11274-018-2479-7.

90. Louca, S., Mazel, F., Doebeli, M. & Parfrey, L.W. 2019. A census-based estimate of Earth's bacterial and archaeal diversity. *PLoS Biology*. 17(2):e3000106. DOI: 10.1371/journal.pbio.3000106.

91. Mahmood, Q., Zheng, P., Hu, B., Jilani, G., Azim, M.R., Wu, D. & Liu, D. 2009. Isolation and characterization of *Pseudomonas stutzeri*

QZ1 from an anoxic sulfide-oxidizing bioreactor. *Anaerobe*. 15(4):108–115. DOI: 10.1016/j.anaerobe.2009.03.009.

92. Majhi, K., Let, M., Halder, U. & Bandopadhyay, R. 2021. Isolation and Characterization of Copper Tolerant Bacterial Species Habituating in Copper Mines and Study of Their Potentiality as a Plant Growth Stimulator. (Preprint). In Review. DOI: 10.21203/rs.3.rs-705506/v1.

93. Marais, L. 2012. Characterization of bacteria isolated from a platinum mine tailings dam. Unpublished masters thesis. Potchefstroom: North-West University [Online] Available: http://dspace.nwu.ac.za/bitstream/handle/10394/8721/Marais_LM.pdf?sequence=1.

94. Mason, J. & Kelly, D.P. 1988. Thiosulfate Oxidation by Obligately Heterotrophic Bacteria. *Microbial Ecology*. 15(2):123–134.

95. Mishra, D. & Rhee, Y. 2014. Microbial leaching of metals from solid industrial wastes. *Journal of microbiology (Seoul, Korea)*. 52:1–7. DOI: 10.1007/s12275-014-3532-3.

96. Mounaouer, B., Nesrine, A. & Hassen, A. 2014. Identification and characterization of heavy metal-resistant bacteria selected from different polluted sources. *Desalination and Water Treatment*. 52. DOI: 10.1080/19443994.2013.823565.

97. Mumtaz, M.Z., Barry, K.M., Baker, A.L., Nichols, D.S., Ahmad, M., Zahir, Z.A. & Britz, M.L. 2019. Production of lactic and acetic acids by *Bacillus* sp. ZM20 and *Bacillus cereus* following exposure to zinc oxide: A possible mechanism for Zn solubilization. *Rhizosphere*. 12:100170. DOI: 10.1016/j.rhisph.2019.100170.

98. Mwase, J.M. 2009. Hydrometallurgical extraction of platinum group metals from a low-grade ore concentrate. Available: https://open.uct.ac.za/handle/11427/8966 [2021, October 20].

99. Mwase, J.M., Petersen, J. & Eksteen, J.J. 2012. A conceptual flowsheet for heap leaching of platinum group metals (PGMs) from a low-grade ore concentrate. *Hydrometallurgy*. 111–112:129–135. DOI: 10.1016/j.hydromet.2011.11.012.

100. Mwase, J.M., Petersen, J. & Eksteen, J.J. 2014. A novel sequential heap leach process for treating crushed Platreef ore. *Hydrometallurgy*. 141:97–104. DOI: 10.1016/j.hydromet.2013.11.005.

101. Ndeddy Aka, R.J. & Babalola, O.O. 2017. Identification and characterization of Cr-, Cd-, and Ni-tolerant bacteria isolated from mine tailings. *Bioremediation Journal*. 21(1):1–19. DOI: 10.1080/10889868.2017.1282933.

102. Neeratanaphan, L., Rungsunsombut, M. & Phonimdaeng, P. 2015. Efficient Bacterial Species for Arsenic Bioremediation of Gold Mining Soil. *Acta Biologica Malaysiana*. 4:36–45. DOI: 10.7593/abm/4.2.36.

103. Nguyen, K., Borja, D., You, J., Hong, G., Jung, H. & Kim, H. 2018. Chalcopyrite Bioleaching Using Adapted Mesophilic Microorganisms: Effects of Temperature, Pulp Density, and Initial Ferrous Concentrations. *Materials Transactions*. 59. DOI: 10.2320/matertrans.M2018247.

104. Nur Indah Koni, T., Rusman, R., Hanim, C. & Zuprizal, Z. 2017. Effect of pH and Temperature on *Bacillus subtilis* FNCC 0059 Oxalate Decarboxylase Activity. *Pakistan journal of biological sciences: PJBS*. 20(9):436–441. DOI: 10.3923/pjbs.2017.436.441.

105. Oberthür, T. 2018. The Fate of Platinum-Group Minerals in the Exogenic Environment—From Sulfide Ores via Oxidized Ores into Placers: Case Studies Bushveld Complex, South Africa, and Great Dyke, Zimbabwe. *Minerals*. 8(12):581. DOI: 10.3390/min8120581.

106. O'Connor, C. & Alexandrova, T. 2021. The Geological Occurrence, Mineralogy, and Processing by Flotation of Platinum Group Minerals (PGMs) in South Africa and Russia. *Minerals*. 11(1):54. DOI: 10.3390/min11010054.

107. Osasona, I., Oke, O.E. & Adebayo, A.O. 2021. Direct Leaching of Lead from Galena Using Acetic Acid in Iron(III) Chloride. *Journal of Hazardous, Toxic, and Radioactive Waste*. 25(3):04021021. DOI: 10.1061/(ASCE)HZ.2153-5515.0000625.

108. Paczia, N., Nilgen, A., Lehmann, T., Gätgens, J., Wiechert, W. & Noack, S. 2012. Extensive exometabolome analysis reveals extended overflow metabolism in various microorganisms. *Microbial Cell Factories*. 11(1):122. DOI: 10.1186/1475-2859-11-122.

109. Palanivel, T.M., Sivakumar, N., Al-Ansari, A. & Victor, R. 2020. Bioremediation of copper by active cells of *Pseudomonas stutzeri* LA3 isolated from an abandoned copper mine soil. *Journal of Environmental Management*. 253:109706. DOI: 10.1016/j.jenvman.2019.109706.

110. Patel, M.V. & Patel, R.K. 2014. Indole-acetic-acid (IAA) Production by Endophytic Bacteria Isolated from Saline Dessert, The Little Runn of Kutch. *CIBTech Journal of Microbiology*. 3(2):17-28.

111. Pathak, A., Vinoba, M. & Kothari, R. 2021. Emerging role of organic acids in leaching of valuable metals from refinery-spent hydroprocessing catalysts, and potential techno-economic challenges: A review. *Critical Reviews in Environmental Science and Technology*. 51(1):1–43. DOI: 10.1080/10643389.2019.1709399.

112. Pearson, A.M. 2003. MEAT | Extracts. In *Encyclopedia of Food Sciences and Nutrition (Second Edition)*. B. Caballero, Ed. Oxford:

Academic Press. 3812–3817. DOI: 10.1016/B0-12-227055-X/00758-6.

113. Peng, T., Liao, W., Wang, J., Miao, J., Peng, Y., Gu, G., Wu, X., Qiu, G., et al. 2021. Bioleaching and Electrochemical Behavior of Chalcopyrite by a Mixed Culture at Low Temperature. *Frontiers in Microbiology*. 12:663757. DOI: 10.3389/fmicb.2021.663757.

114. Pepper, I.L., Zerzghi, H.G., Bengson, S.A., Iker, B.C., Banerjee, M.J. & Brooks, J.P. 2012. Bacterial populations within copper mine tailings: long-term effects of amendment with Class A biosolids. *Journal of Applied Microbiology*. 113(3):569–577. DOI: 10.1111/j.1365-2672.2012.05374.x.

115. Pereira, L.B., Vicentini, R. & Ottoboni, L.M.M. 2014. Changes in the Bacterial Community of Soil from a Neutral Mine Drainage Channel.

PLoS ONE. 9(5):e96605. DOI: 10.1371/journal.pone.0096605.

116. Piotrowska-Seget, Z., Cycoń, M. & Kozdrój, J. 2005. Metal-tolerant bacteria occurring in heavily polluted soil and mine spoil. *Applied Soil Ecology*. 28(3):237–246. DOI: 10.1016/j.apsoil.2004.08.001.

117. Pradhan, S.K., Sahu, D.K., Singh, N.R., Kumar, U. & Thatoi, H. 2022. Unveiling of metal-tolerance bacterial consortia in chromite mine by metagenomic approaches. (Preprint). In Review. DOI: 10.21203/rs.3.rs1254880/v1.

118. Prasanna, N.D., KVijayalakshmi, Shaheen, S.K., Jyotsna, K., Revathi, G., Jayalakshmi, P. & Devika, P. 2010. Screening and isolation of phosphate solubilizing *Pseudomonas stutzeri* (EGB3) from gut of earthworm [*Eisenia foetida*: Solubilization as influenced by organic acids. *Journal of Pure and Applied Microbiology*. 4:717–723.

119. Puhakka, J., Kaksonen, A. & Riekkola-Vanhanen, M. 2007. Heap Leaching of Black Schist. *Biomining*. (January, 1):139–151. DOI: 10.1007/978-3-540-34911-2_7.

120. Qi, D., Wieneke, X., Tao, J., Zhou, X. & Desilva, U. 2018. Soil pH Is the Primary Factor Correlating with Soil Microbiome in Karst Rocky Desertification Regions in the Wushan County, Chongqing, China. *Frontiers in Microbiology*. 9:1027. DOI: 10.3389/fmicb.2018.01027.

121. Ramachandran, S., Fontanille, P., Pandey, A. & Larroche, C. 2006. Gluconic Acid: Properties, Applications and Microbial Production. *Food Technology and Biotechnology*. 44:185-195.

122. Ranaei, V., Pilevar, Z., Khaneghah, A.M. & Hosseini, H. 2020. Propionic Acid: Method of Production, Current State and Perspectives. *Food Technology and Biotechnology*. 58(2):115–127. DOI: 10.17113/ftb.58.02.20.6356.

123. Ranalli, G., Zanardini, E. & Sorlini, C. 2009. Biodeterioration – Including Cultural Heritage. In *Encyclopedia of Microbiology (Third Edition)*. M. Schaechter, Ed. Oxford: Academic Press. 191–205. DOI: 10.1016/B978-012373944-5.00132-2.

124. Rauwane, M.E. 2008. Microbial diversity and metal pollution from a platinum mine tailings dam in the North-West Province (RSA).*Unpublished masters thesis. Potchefstroom:* North-West University [Online]. Available: https://www.semanticscholar.org/paper/Microbialdiversity-and-metal-pollution-from-a-mine-Rauwane/9985cd7895c4595016ae0716a8986cefa055ce5a [2022, January 16].

125. Rawlings, D.E. 2005. Characteristics and adaptability of iron- and sulfur-oxidizing microorganisms used for the recovery of metals

from minerals and their concentrates. *Microbial Cell Factories*. 4(1):13. DOI: 10.1186/1475-2859-4-13.

126. Rawlings, D.E. 2013. *Biomining: Theory, Microbes and Industrial Processes*. Springer Science & Business Media.

127. Reed, D., Fujita, Y., Daubaras, D., Jiao, Y. & Thompson, V. 2016. Bioleaching of Rare Earth Elements from Waste Phosphors and Cracking Catalysts. *Hydrometallurgy*. 166. DOI: 10.1016/j.hydromet.2016.08.006.

128. Reith, F., Campbell, S.G., Ball, A.S., Pring, A. & Southam, G. 2014. Platinum in Earth surface environments. *Earth-Science Reviews*. 131:1–21. DOI: 10.1016/j.earscirev.2014.01.003.

129. Rekha, V., Dhanalakshmi & Sudha, K. 2013. Characterization of *Bacillus* and *Micrococcus* sp isolated from coal mine soil. *Asian Journal of Microbiology, Biotechnology and Environmental Sciences*. 15:503–506.

130. Riekkola-Vanhanen, M. 2007. Talvivaara Black Schist Bioheapleaching Demonstration Plant. *Advanced Materials Research*.
20:30–33. DOI: 10.4028/www.scientific.net/AMR.20-21.30.

131. Saeid, A., Prochownik, E. & Dobrowolska-Iwanek, J. 2018. Phosphorus Solubilization by *Bacillus* Species. *Molecules (Basel, Switzerland)*. 23(11):E2897. DOI: 10.3390/molecules23112897.

132. Safarzadeh, M.S., Horton, M. & Van Rythoven, A.D. 2018. Review of Recovery of Platinum Group Metals from Copper Leach Residues and Other Resources. *Mineral Processing and Extractive Metallurgy Review*. 39(1):1–17. DOI: 10.1080/08827508.2017.1323745.

133. Sahu, P., Dash, P., Jena, M.S. & Mandre, N.R. 2022. Flotation of Platinum Group Elements Ores: A Review. *Mineral Processing and*

Extractive Metallurgy Review. 0(0):1–23. DOI: 10.1080/08827508.2022.2027768.

134. Sakil Munna, Md., Tahera, J., Mohibul Hassan Afrad, Md., Nur, I.T. & Noor, R. 2015. Survival of *Bacillus* spp. SUBB01 at high temperatures and a preliminary assessment of its ability to protect heat-stressed *Escherichia coli* cells. *BMC Research Notes*. 8(1):637. DOI: 10.1186/s13104-015-1631-9.

135. Sanchez-Rosario, Y. & Johnson, M.D.L. 2021. Media Matters, Examining Historical and Modern *Streptococcus pneumoniae* Growth Media and the Experiments They Affect. *Frontiers in Cellular and Infection Microbiology*. 11:613623. DOI: 10.3389/fcimb.2021.613623.

136. Sand, W., Gehrke, T., Jozsa, P.-G. & Schippers, A. 1999. Direct versus indirect bioleaching. In *Process Metallurgy*. V. 9. R. Amils & A. Ballester, Eds. (Biohydrometallurgy and the Environment Toward the Mining of the 21 Century - Proceedings of the International Biohydrometallurgy Symposium). Elsevier. 27–49. DOI: 10.1016/S15724409(99)80004-2.

137. Sankaralingam, S., Eswaran, S., Balakan, B., Sundaram, V.M. & Shankar, T. 2014. Screening and growth characterization of phosphate solubilizing bacterium *Pseudomonas aeruginosa*. *Advances in Environmental Biology*. 8(13):673–680.

138. Santofimia, E., González-Toril, E., López-Pamo, E., Gomariz, M., Amils, R. & Aguilera, Á. 2013. Microbial Diversity and Its Relationship to Physicochemical Characteristics of the Water in Two Extreme Acidic Pit Lakes from the Iberian Pyrite Belt (SW Spain). *PLOS ONE*. 8(6):e66746. DOI: 10.1371/journal.pone.0066746.

139. Schippers, A. & Sand, W. 1999. Bacterial Leaching of Metal Sulfides Proceeds by Two Indirect Mechanisms via Thiosulfate or via Polysulfides and Sulfur. *Applied and Environmental Microbiology*. 65(1):319–321.

140. Schippers, A., Hedrich, S., Vasters, J., Drobe, M., Sand, W. & Willscher, S. 2013. Biomining: Metal Recovery from Ores with Microorganisms. In *Geobiotechnology I*. V. 141. A. Schippers, F. Glombitza, & W. Sand, Eds. (Advances in Biochemical Engineering/Biotechnology). Berlin, Heidelberg: Springer Berlin Heidelberg. 1–47. DOI: 10.1007/10_2013_216.

141. Schippers, A., Hedrich, S., Vasters, J., Drobe, M., Sand, W. & Willscher, S. 2014. Biomining: metal recovery from ores with microorganisms. *Advances in Biochemical Engineering/Biotechnology*. 141:1–47. DOI: 10.1007/10_2013_216.

142. Sefako, R., Sekgarametso, K. & Sibanda, V. 2017. Potential Processing Routes for Recovery of Platinum Group Metals from Southern African Oxidized PGM Ores: A Review. *Journal of Sustainable Metallurgy*. 3(4):797–807. DOI: 10.1007/s40831-017-0146-0.

143. Sethurajan, M. 2015. Metallurgical sludges, bio/leaching and heavy metals recovery (Zn, Cu).Unpublished doctoral thesis. Paris: Université Paris-Est [Online]. Avalible: https://www.semanticscholar.org/paper/Metallurgical-sludges%2Cbio%2Fleaching-and-heavy-(Zn%2C-Sethurajan/7f40830f8e9234866a3703813eca04f6b83ac37d

144. Seymour, R.J. & O'Farrelly, J. 2012. Platinum-Group Metals. In *Kirk-Othmer Encyclopedia of Chemical Technology*. American

Cancer Society. 1–37. DOI: 10.1002/0471238961.1612012019052513.a01.pub3.

145. Shabani, M.A., Irannajad, M., Azadmehr, A.R. & Meshkini, M. 2013. Bioleaching of copper oxide ore by *Pseudomonas aeruginosa*. *International Journal of Minerals, Metallurgy, and Materials*. 20:1130–1133. DOI: 10.1007/s12613-013-0845-x.

146. Sharma, M., Mishra, V., Rau, N. & Sharma, R.S. 2016. Increased iron-stress resilience of maize through inoculation of siderophoreproducing *Arthrobacter globiformis* from mine: *Arthrobacter* from mine alleviates iron stress in maize. *Journal of Basic Microbiology*. 56(7):719– 735. DOI: 10.1002/jobm.201500450.

147. Shen, C., Xiong, J., Zhang, H., Feng, Y., Lin, X., Li, X., Liang, W. & Chu, H. 2013. Soil pH drives the spatial distribution of bacterial communities along elevation on Changbai Mountain. *Soil Biology and Biochemistry*. 57:204–211. DOI: 10.1016/j.soilbio.2012.07.013.

148. Shivlata, L. & Tulasi, S. 2015. Thermophilic and alkaliphilic Actinobacteria: biology and potential applications. *Frontiers in Microbiology*. 6. Available: https://www.frontiersin.org/article/10.3389/fmicb.2015.01014 [2022, January 26].

149. Simate, G.S. 2009. Bacterial Leaching of Nickel Laterites Using Chemolithotrophic Microorganisms. Unpublished doctoral thesis. Braamfontein: University of the Witwatersrand [Online]. Available: https://core.ac.uk/download/pdf/39666104.pdf.

150. Singh, P. & Tiwary, B.N. 2016. Isolation and characterization of glycolipid biosurfactant produced by a *Pseudomonas otitidis* strain isolated from Chirimiri coal mines, India. *Bioresources and Bioprocessing*. 3(1):42. DOI: 10.1186/s40643-016-0119-3.

151. Sinisalo, P. & Lundström, M. 2018. Refining Approaches in the Platinum Group Metal Processing Value Chain—A Review. *Metals*. 8:203.

DOI: 10.3390/met8040203.

152. Soni, S.K., Singh, R., Awasthi, A. & Kalra, A. 2014. A Cr(VI)reducing *Microbacterium* sp. strain SUCR140 enhances growth and yield of Zea mays in Cr(VI) amended soil through reduced chromium toxicity and improves colonization of arbuscular mycorrhizal fungi. *Environmental Science and Pollution Research International*. 21(3):1971–1979. DOI: 10.1007/s11356-013-2098-7.

153. Sulistinah, N., Riffiani, R. & Sunarko, B. 2015. R*hodococcus pyridinivorans* strain TPIK, A Nitrile-Degrading Bacterium Isolated from Tailing Pond of Antam Gold Mining Industry, Pongkor. *Journal Teknologi Indonesia (JTI)*. 38 (2) : 61-67 DOI: 10.14203/JTI.V38I2.158.

154. Sultan, S. & Faisal, M. 2014. Isolation and Characterization of Iron and Sulfur Oxidizing Bacteria from Coal Mines. *Journal of Environmental and Erath Sciences*. 6(3):153-157.

155. Syed, S. & Chinthala, P. 2015. Heavy Metal Detoxification by Different *Bacillus* Species Isolated from Solar Salterns. *Scientifica*. 2015:319760. DOI: 10.1155/2015/319760.

156. Tang, J. & Valix, M. 2004. Leaching of Low- Grade Nickel Ores by Fungi Metabolic Acids. Unpublished paper delivered at the Engineering

Conferences International. Available: https://dc.engconfintl.org/cgi/viewcontent.cgi?article=1004&context=separations_technology_vi

157. Tao, H. & Dongwei, L. 2014. Presentation on mechanisms and applications of chalcopyrite and pyrite bioleaching in

biohydrometallurgy – a presentation. *Biotechnology Reports*. 4:107–119. DOI:
10.1016/j.btre.2014.09.003.

158. Thavamani, P., Samkumar, R.A., Satheesh, V., Subashchandrabose, S.R., Ramadass, K., Naidu, R., Venkateswarlu, K. & Megharaj, M. 2017. Microbes from mined sites: Harnessing their potential for reclamation of derelict mine sites. *Environmental Pollution*. 230:495–505. DOI:
10.1016/j.envpol.2017.06.056.

159. Thethwayo, B.M. 2018. Extraction of Platinum Group Metals. In *Noble and Precious Metals - Properties, Nanoscale Effects and Applications*. M.S. Seehra & A.D. Bristow, Eds. InTech. DOI: 10.5772/intechopen.73214.

160. Thierry, S., Macarie, H., Iizuka, T., Geißdörfer, W., Assih, E.A., Spanevello, M., Verhe, F., Thomas, P., et al. 2004. *Pseudoxanthomonas mexicana* sp. nov. and *Pseudoxanthomonas japonensis* sp. nov., isolated from diverse environments, and emended descriptions of the genus *Pseudoxanthomonas Finkmann* et al. 2000 and of its type species. *International Journal of Systematic and Evolutionary Microbiology*. 54(6):2245–2255. DOI: 10.1099/ijs.0.02810-0.

161. Thompson, L.R., Sanders, J.G., McDonald, D., Amir, A., Ladau, J., Locey, K.J., Prill, R.J., Tripathi, A., et al. 2017. A communal catalogue reveals Earth's multiscale microbial diversity. *Nature*. 551(7681):457–463. DOI: 10.1038/nature24621.

162. Tiong, B., mohd bahari, Z., Lee, N., Jaafar, J., Ibrahim, Z. & Shahir, S. 2014. Cyanide Degradation by *Pseudomonas pseudoalcaligenes* Strain W2 Isolated from Mining Effluent. *Sains Malaysiana*. (July, 25). DOI: 10.17576/jsm-2015-4402-10.

163. Tributsch, H. 1999. Direct versus indirect bioleaching. In *Process Metallurgy*. V. 9. R. Amils & A. Ballester, Eds. (Biohydrometallurgy and the Environment Toward the Mining of the 21 Century - Proceedings of the International Biohydrometallurgy Symposium). Elsevier. 51–60. DOI: 10.1016/S1572-4409(99)80005-4.

164. Trinh, H.B., Lee, J., Srivastava, R.R., Kim, S. & Ilyas, S. 2017. Ecothreat Minimization in HCl Leaching of PGMs from Spent Automobile Catalysts by Formic Acid Prereduction. *ACS Sustainable Chemistry & Engineering*. 5(8):7302–7309. DOI: 10.1021/acssuschemeng.7b01538.

165. Turner, S., Pryer, K.M., Miao, V.P.W. & Palmer, J.D. 1999. Investigating Deep Phylogenetic Relationships among Cyanobacteria and Plastids by Small Subunit rRNA Sequence Analysis. *The Journal of Eukaryotic Microbiology*. 46(4):327–338. DOI: 10.1111/j.15507408.1999.tb04612.x.

166. Urias, P.M., dos Reis Menêzes, L.H., Cardoso, V.L., de Resende, M.M. & de Souza Ferreira, J. 2021. Leaching with mixed organic acids and sulfuric acid to recover cobalt and lithium from lithium ion batteries. *Environmental Technology*. 42(25):4027–4037. DOI: 10.1080/09593330.2020.1772372.

167. Uzair, B., Ahmed, N., Ahmad, V.U., Mohammad, F.V. & Edwards, D.H. 2008. The isolation, purification and biological activity of a novel antibacterial compound produced by *Pseudomonas stutzeri*. *FEMS microbiology letters*. 279(2):243–250. DOI: 10.1111/j.15746968.2007.01036.x.

168. Valenta, M.M. & Mapheto, H. 2010. Application of fundamentals in optimizing platinum concentrator performance. *Journal of the Southern African Institute of Mining and Metallurgy*. 111(2):93-99.

169. Van Aken, B. 2011. 6.20 - Transgenic Plants and Associated Bacteria for Phytoremediation of Organic Pollutants. In *Comprehensive Biotechnology (Second Edition)*. M. Moo-Young, Ed. Burlington: Academic Press. 223–237. DOI: 10.1016/B978-0-08-088504-9.00330-5.

170. Van Hille, R., Wyk, N. & Harrison, S. 2011. Review of microbial ecology of BIOX reactors illustrates the dominance of the genus *Ferroplasma* in many commercial reactors. *Proceedings of the 19th International Biohydrometallurgy Symposium (IBS 2011)*. (January, 1).

171. Vijayalakshmi, P. 2011. Production of L-Glutamic Acid by *Arthrobacter globiformis*. MTCC 4299 Using Fruits of *Mimusops elengi* Linn. *International Journal of Applied Biology and Pharmaceutical Technology*. 2(2):167–173.

172. Vyas, P. & Gulati, A. 2009. Organic acid production in vitro and plant growth promotion in maize under controlled environment by phosphate-solubilizing fluorescent *Pseudomonas*. *BMC Microbiology*. 9:174. DOI: 10.1186/1471-2180-9-174.

173. Wang, J., Wang, Z., Zhang, Z. & Zhang, G. 2019. Comparison of Butyric Acid Leaching Behaviors of Zinc from Three Basic Oxygen Steelmaking Filter Cakes. *Metals*. 9(4):417. DOI: 10.3390/met9040417.

174. Wang, Z., Jin, Y. & Yang, S.-T. 2015. High cell density propionic acid fermentation with an acid tolerant strain of *Propionibacterium acidipropionici*. *Biotechnology and Bioengineering*. 112(3):502–511. DOI: 10.1002/bit.25466.

175. Wei, G., Fan, L., Zhu, W., Fu, Y., Yu, J. & Tang, M. 2009. Isolation and characterization of the heavy metal resistant bacteria CCNWRS33-2 isolated from root nodule of *Lespedeza cuneata* in gold mine tailings in China. *Journal of Hazardous Materials*. 162(1):50–56. DOI: 10.1016/j.jhazmat.2008.05.040.

176. Wenderoth, D. & Reber, H. 1999. Correlation between structural diversity and catabolic versatility of metal-affected prototrophic bacteria in soil. *Soil Biology and Biochemistry*. 31(3):345-352. DOI: 10.1016/S00380717(98)00132-1.

177. West, S.A., Griffin, A.S., Gardner, A. & Diggle, S.P. 2006. Social evolution theory for microorganisms. *Nature Reviews Microbiology*. 4(8):597–607. DOI: 10.1038/nrmicro1461.

178. Williams, P.J. & Cloete, T.E. 2010. The use of Aspergillus niger for the removal of phosphorous and potassium from the iron ore concentrate of the Sishen Iron Ore Mine, South Africa. *Journal of Biotechnology*. 150:372–373. DOI: 10.1016/j.jbiotec.2010.09.449.

179. Williams, P.J., Surridge, A.K.J. & Cloete, T.E. 2008. Microbial community study of the process: and groundwater of the Sishen Iron-Ore Mine, South Africa. *Water SA*. 34(5):573–578.

180. Wu, H.-Y. & Ting, Y.-P. 2006. Metal extraction from municipal solid waste (MSW) incinerator fly ash—Chemical leaching and fungal bioleaching. *Enzyme and Microbial Technology*. 38(6):839–847. DOI: 10.1016/j.enzmictec.2005.08.012.

181. Xia, Y., Lü, C., Hou, N., Xin, Y., Liu, J., Liu, H. & Xun, L. 2017. Sulfide production and oxidation by heterotrophic bacteria under aerobic conditions. *The ISME Journal*. 11(12):2754–2766. DOI: 10.1038/ismej.2017.125.

182. Xie, X., Xiao, S., He, Z., Liu, J. & Qiu, G. 2007. Microbial populations in acid mineral bioleaching systems of Tong Shankou Copper Mine, China. *Journal of Applied Microbiology*. 103(4):1227–1238. DOI: 10.1111/j.1365-2672.2007.03382.x.

183. Xu, Z., Shi, Z. & Jiang, L. 2011. Acetic and Propionic Acids. In *Comprehensive Biotechnology (Second Edition)*. M. Moo-Young, Ed.
Burlington: Academic Press. 189–199. DOI: 10.1016/B978-0-08-088504-9.00162-8.

184. Yadav, N. & Nath Yadav, A. 2018. Biodiversity and biotechnological applications of novel plant growth promoting methylotrophs. *Journal of Applied Biotechnology & Bioengineering*. 5(6).
DOI: 10.15406/jabb.2018.05.00162.

185. Yan, Z., Zheng, X.-W., Chen, J.-Y., Han, J.-S. & Han, B.-Z. 2013. Effect of different *Bacillus* strains on the profile of organic acids in a liquid culture of Daqu. *Journal of the Institute of Brewing*. 119(1–2):78–83. DOI: 10.1002/jib.58.

186. Yin, S., Wang, L., Kabwe, E., Chen, X., Yan, R., An, K., Zhang, L. & Wu, A. 2018. Copper Bioleaching in China: Review and Prospect. *Minerals*. 8(2):32. DOI: 10.3390/min8020032.

187. Yousif, A.M. 2019. Recovery and Then Individual Separation of Platinum, Palladium, and Rhodium from Spent Car Catalytic Converters
Using Hydrometallurgical Technique followed by Successive Precipitation Methods. *Journal of Chemistry*. 2019:e2318157. DOI:
10.1155/2019/2318157.

188. Zhang, B.-X., Li, P.-S., Wang, Y.-Y., Wang, J.-J., Liu, X.-L., Wang, X.-Y. & Hu, X.-M. 2021. Characterization and synthesis of indole-

3-acetic acid in plant growth promoting *Enterobacter* sp. *RSC Advances*.
11(50):31601–31607. DOI: 10.1039/D1RA05659J.

189. Zhang, R., Hedrich, S., Römer, F., Goldmann, D. & Schippers, A. 2020. Bioleaching of cobalt from Cu/Co-rich sulfidic mine tailings from the polymetallic Rammelsberg mine, Germany. *Hydrometallurgy*. 197:105443. DOI: 10.1016/j.hydromet.2020.105443.

190. Zhou, G., Yang, X.-R., Su, J.Q. & Bang-Xiao, Z. 2018. *Bacillus ferrooxidans* sp. nov., an iron(II)-oxidizing bacterium isolated from paddy soil. *Journal of Microbiology*. 56. DOI: 10.1007/s12275-018-7543-3.

191. Zhou, T., Chen, D., Li, C., Sun, Q., Li, L., Liu, F., Shen, Q. & Shen, B. 2012. Isolation and characterization of *Pseudomonas brassicacearum* J12 as an antagonist against *Ralstonia solanacearum* and identification of its antimicrobial components. *Microbiological Research*. 167(7):388–394.
DOI: 10.1016/j.micres.2012.01.003.