

A Review of the Use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre

Dr Wilhelm Herbst

671874

*Supervisors: Dr Jarrod Zamparini, Dr Halvani Moodley, Dr Shastra
Bhoora*

A research project submitted to the Faculty of Health Sciences, University of the
Witwatersrand, Johannesburg, in fulfilment of the requirements for the degree of
Master of Medicine (MMed) in Internal Medicine

i. DECLARATION

I, Wilhelm Herbst declare that this Research Report is my own, unaided work. It is being submitted for the Degree of Master of Medicine (MMed) in Internal Medicine at the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination at any other University.

A handwritten signature in dark ink, appearing to read 'Herbst', enclosed within a light grey rectangular border.

Wilhelm Herbst

On this 23th day of April 2024

ii. DECLARATION: STUDENT'S CONTRIBUTION TO ARTICLE(S) AND AGREEMENT OF CO-AUTHOR(S)

I, Wilhelm Herbst, student number 671874, declare that this Research Report is my own work and that I contributed adequately towards research findings published in the article(s) stated below which are included in my Research Report .

Signature of Student:..........Date: 23rd April 2024.

Name of Primary Supervisor: Dr Jarrod Zamparini

Signature of Primary SupervisorDate.....

Article 1: Title: A Review of the Use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre.

Journal name: Sage Obstetric Medicine

Authors	Name	Signature
1 st Author	Dr Wilhelm Herbst	
2 nd Author	Prof Shastra Bhoora	
3 rd Author	Dr Halvani Moodley	
4 th Author	Dr Ashesh Rachod	
5 th Author	Dr Tracey Westgarth-Taylor	
6 th Author	Dr Jarrod Zamparini	

iii. PUBLICATIONS AND PRESENTATIONS

This research project has been prepared in the format for a submissible article according to the author guidelines as outlined by the journal *Obstetric Medicine*. This work has never been published nor presented at any congresses.

iv. LIST OF ABBREVIATIONS

ATS – American Thoracic Society

BMI – body mass index

sBP – systolic blood pressure

CRP – C-reactive protein

CTPA – computed tomography pulmonary angiography

CT – computed tomography

ECG - electrocardiogram

HIV – Human Immunodeficiency Virus

IQR – interquartile range

MMR – maternal mortality rate

MUAC – mid upper arm circumference

RCOG – Royal College of Obstetricians and Gynaecologists

SOGC – Society of Obstetricians and Gynaecologists of Canada

VL – viral load

VTE – venous thrombo-embolism

PACS – picture archiving and communicating system

PE – pulmonary embolism

PCT - Procalcitonin

PROM – premature rupture of membranes

V. ACKNOWLEDGEMENTS

I would like to express my sincere thanks to:

- My supervisor Dr Jarrod Zamparini for his astute guidance, wholehearted support and mentorship. His drive for excellence has made this a much better study.
- My co-supervisors Professor Shastra Bhoora and Dr Halvani Moodley for their determination and meticulous assistance through every step.
- Dr Ashesh Ranchod, Dr Tracey Westgarth-Taylor and Dr Halvani Moodley for their time and effort to read the CTPA scans.

vi. DEDICATION



I dedicate this work to my wife Ilze and my son Joshua for their steadfast love and support. This would not have been possible without you.

Table of Contents

<i>i.</i>	DECLARATION.....	<i>i</i>
<i>ii.</i>	DECLARATION: STUDENT’S CONTRIBUTION TO ARTICLE(S) AND AGREEMENT OF CO-AUTHOR(S)	<i>ii</i>
<i>iii.</i>	PUBLICATIONS AND PRESENTATIONS	<i>iii</i>
<i>iv.</i>	LIST OF ABBREVIATIONS	<i>iv</i>
<i>v.</i>	ACKNOWLEDGEMENTS.....	<i>v</i>
<i>vi.</i>	DEDICATION	<i>vi</i>
	CHAPTER 1: RESEARCH PROTOCOL	2
		2
1.1	Introduction	6
	Background	6
	Risk Factors.....	6
	Clinical predictors.....	7
	D-dimer	8
	CT Pulmonary Angiography	8
	Other imaging modalities.....	9
1.2	Aim of the study.....	10
1.3	Research Question	10
1.4	Study Objectives	11
1.5	Methods	11
	Study Design.....	11
	Study Population	11
	Exclusion Criteria	11
	Data Collection	11
	Statistical Analysis	12
1.6	Limitations	12
1.7	Ethical approval	13
1.8	Funding.....	13
1.9	Timeline	14
1.10	References	15
	APPENDIX A: Data Collection Sheet	20
	APPENDIX B: Letters to HOD	24
	APPENDIX C: Letters to CMJAH Research Committee	26
	CHAPTER 2: MANUSCRIPT.....	27
2.1	TITLE PAGE	28
2.2	ABSTRACT	30
2.3	INTRODUCTION.....	31
	Risk Factors.....	31
	Clinical predictors.....	32

D-dimer	32
CT Pulmonary Angiography	32
2.4 METHODS	33
2.5 RESULTS	34
2.6 DISCUSSION	36
2.7 CONCLUSION.....	38
2.8 DECLARATIONS:.....	40
2.9 TABLES AND FIGURES:	41
2.10 REFERENCES:.....	45
APPENDIX A: ETHICAL CLEARANCE - HUMAN RESEARCH ETHICS COMMITTEE (MEDICAL).....	48
APPENDIX B: ETHICAL CLEARANCE - CHARLOTTE MAXEKE JOHANNESBURG ACADEMIC HOSPITAL	
49	
APPENDIX C: JOURNAL SUBMISSION GUIDELINES: <i>OBSTETRIC MEDICINE</i>	51
APPENDIX D: TURNITIN REPORT	68
Table 1: Alternative CT Findings	Error! Bookmark not defined.
Table 2: Demographics and pregnancy details	Error! Bookmark not defined.
Table 3: Pregnancy complications and comorbidities	44
Figure 1: Study population	41

CHAPTER 1: RESEARCH PROTOCOL

CANDIDATE'S SURNAME: HERBST [Please print]		FIRST NAME/S: WILHELM	STUDENT NUMBER: 671874
CURRENT QUALIFICATIONS: MBChB (Pret); Dip HIV MAN (SA)			
TEL:	CELL: 0798859694	E-MAIL: wherbst.herbst@gmail.com	FAX:
DEGREE FOR WHICH PROTOCOL IS BEING SUBMITTED: MMed Internal Medicine			
PART-TIME OR FULL-TIME: FULL-TIME			
FIRST REGISTERED FOR THIS DEGREE:	TERM: 1	YEAR: 2022	
DEPARTMENT: INTERNAL MEDICINE			
TITLE OF PROPOSED RESEARCH: A REVIEW OF THE USE OF CT PULMONARY ANGIOGRAPHY IN PREGNANT AND POSTPARTUM PATIENTS AT AN ACADEMIC CENTRE			
CANDIDATE'S SIGNATURE: 			DATE: 27/01/2022
SUPERVISOR 1 (NAME & SURNAME): Jarrod Zamparini			% Supervision - 40
SUPERVISOR'S QUALIFICATIONS: MBChB (Pret), FCP (SA), MMed Int Med (Wits)			
SUPERVISOR'S DEPARTMENT: INTERNAL MEDICINE			
SUPERVISOR'S ADDRESS / TEL / E-MAIL: 0722781158 jarrod.zamparini@wits.ac.za			
SUPERVISOR 2 (NAME & SURNAME): Shastra Bhoora			% Supervision - 30
SUPERVISOR'S QUALIFICATIONS: MBBCh (Wits), FCOG (SA), MMed O&G (Wits), Cert Crit Care (SA) O&G			
SUPERVISOR'S ADDRESS / TEL / E-MAIL: 0836578561 shastra.bhoora@wits.ac.za			
SUPERVISOR 3 (NAME & SURNAME): Halvani Moodley			% Supervision - 30
SUPERVISOR'S QUALIFICATIONS: MBChB (Pret), FC Rad Diag (SA), MMed Diag Rad (Wits)			
SUPERVISOR'S ADDRESS / TEL / E-MAIL: 0741020580 halvani.moodley@gmail.com			
SYNOPSIS OF RESEARCH: (Brief summary of proposed research project; between 200-300 words only; with sub-headings: an introduction and justification for study, aim/s, proposed methodology and expected outcome/s)			
Please see below.			
WITS ETHICS NOT REQUIRED:		Yes ☒ No ☐	IF Y SUPPLY ETHICS CLEARANCE CERTIFICATE AS ATTACHMENT AND INCLUDE ETHICS NUMBER HERE:
WITS ETHICS PENDING:		Yes ☐ No ☒	
WITS ETHICS APPROVED: (circle appropriate symbol)*		Yes ☐ No ☒	
*Please note the final human ethics clearance certificate or animal ethics certificate must be available prior to starting the study.			
As supervisor/s, I/we confirm that I have read the protocol, which has been submitted for assessment.			
SIGNATURE OF SUPERVISOR/S:			
SIGNATURE PG OFFICE STAFF		REGISTERED	STAMP
		YES..... NO.....	

A Review of the Use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre

Dr Wilhelm Herbst

671874

Supervisors: Dr Jarrod Zamparini, Dr Halvani Moodley, Dr Shastra Bhoora

A research protocol submitted to the Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, in fulfilment of the requirements for the degree of Master of Medicine (MMed) in Internal Medicine

1.1 Introduction

Background

The most common cause of maternal death during pregnancy and the puerperium in developed countries is venous thromboembolic events, including pulmonary embolism (PE).¹ The risk for venous thromboembolism (VTE) is significantly increased during pregnancy and the postpartum period, as these patients are in a state of hypercoagulability, are prone to venous stasis and may have superimposed endothelial damage.²

Data has shown that women have a 5-fold increased risk of developing VTE during pregnancy, as compared to their non-pregnant counterparts,³ and, according to a Scottish study, the incidence of antenatal VTE has increased over the last 26 years.⁴ Past research has observed an incidence of PE in pregnant or postpartum women of 3 in 10,000^{2,5}, with one death in every 100,000 deliveries.⁶ Some studies have found an absolute incidence of VTE in pregnancy to be as high as 1 to 2 cases per 1000 pregnancies³; a risk that is nearly five times higher than that among non-pregnant women.⁷

More than half the cases of VTE in pregnancy occur in the first trimester, before 20 weeks' gestation.⁵ Yet, 80% of VTE cases in the postpartum period have been observed to occur within the first 3 weeks following delivery.⁸ Recent studies have revealed a raised relative risk (however low absolute risk) that remains up to 12 weeks following delivery.⁹

A large meta-analysis and systematic review of seventeen studies, which included 25,339 patients, found that 2% of patients presenting to the emergency department with symptoms suggestive of PE, were pregnant.¹⁰ This translates to a 12.4% positivity rate for VTE in nonpregnant patients, compared with 4.1% in pregnant patients.¹⁰ The perceived lower yield of confirmed VTE in pregnancy can be ascribed to the low threshold physicians have to scan pregnant patients, due to the high risk of devastating sequelae of PE in pregnancy.¹⁰

Risk Factors

The clinical presentation of VTE may be masked by the characteristic symptoms of pregnancy such as peripheral pedal oedema, along with an increased respiratory rate and mild tachycardia.¹¹ Pregnancy presents certain risk factors to hypercoagulability

including immobility in late pregnancy, iliac compression by the gravid uterus, and dehydration secondary to nausea and vomiting (which affects 50-90% of pregnant women¹²), hypercoagulability (resulting from increased concentration of certain clotting factors as well as a decrease in endogenous anticoagulants such as antithrombin and protein S)¹², endothelial injury resulting from delivery trauma and venous distension^{12,13} as well as thrombophilia.¹⁴

Certain non-modifiable risk factors associated with thrombosis such as Human Immunodeficiency Virus (HIV)¹⁵, Antiphospholipid syndrome (APS)¹⁶ and Obesity¹⁷ must also be taken into account as these pre-existing conditions may be encountered in pregnancy. Gestational diabetes and multiple gestations have not been proven to be significantly associated with an increased risk for thrombosis.¹⁸

There are a number of postpartum factors which have been associated with an increased risk of VTE, such as surgical delivery, particularly in the emergency setting or during prolonged labour, post-partum haemorrhage, thrombophilia, pre-eclampsia, and puerperal sepsis.¹⁴

Normal physiological changes in pregnancy can overlap with clinical manifestations of VTE which deems the diagnosis of PE exceptionally difficult, in addition the accuracy of diagnostic imaging is notably compromised by the hyperdynamic circulatory state in pregnancy.¹⁹ Most studies have generally disregarded the pregnant population when investigating the diagnosis of PE.²⁰

Clinical predictors

Certain clinical predictors of PE, such as active malignancy or advanced age are less relevant in the pregnant population. The Wells' criteria includes tachycardia and lower limb oedema, both of which occur commonly in pregnancy, lowering the specificity in these patients.²¹ Consequently, considering an alternative diagnosis to PE will be more challenging to establish in pregnancy.²¹ The recommendations outlined by the Society of Obstetricians and Gynaecologists of Canada (SOGC) and the Royal College of Obstetricians and Gynaecologists (RCOG), advocate against clinical predictors to assess VTE in pregnant women, whilst other guidelines omit instruction in this regard.²³

A recent Irish study that included 103 pregnant and post-partum participants from a single tertiary maternity establishment, who underwent CTPA for the suspicion of PE, observed that none of these women with a retrospectively assigned Wells' score of 6 or less confirmed a diagnosis of PE, yielding a very significant negative predictive value of 100% (95% confidence interval not stipulated).²⁴ A comparable British study that was conducted with 183 pregnant women with suspected PE, all referred for ventilation perfusion (V/Q) imaging, established that PE was successfully excluded in all 107 participants with a modified Wells' score less than 5.²⁵

D-dimer

In non-pregnant patients PE can be safely excluded in the presence of a normal D-dimer in addition to a low pre-test probability. It must conversely be interpreted with caution in patients with a high pre-test probability, as a falsely negative result can miss a diagnosis of PE.²⁰ Most of the related studies that ascertained the use and shortcoming of D-dimer in diagnosing PE, have not included pregnant women as participants.²⁶

Van der Pol et al found that the pregnancy-adapted YEARS algorithm safely excluded acute PE in pregnant patients who were suspected to have PE, evidently this approach allowed CTPA to be avoided in 32 – 65% of patients, dependent on the trimester of pregnancy, without risking further harm.²⁷

According to the guidelines outlined by the Society of Thrombosis and Haemostasis Research (GTH) and the European Society of Cardiology (ESC), a negative D-dimer can justify avoiding unnecessary imaging in pregnant women suspected to have a PE.^{22,23} However the American Thoracic Society (ATS), SOGC, and RCOG recommendations state that a D-dimer should not be used in pregnancy to exclude the diagnosis of PE.²⁸ Several observational studies, evaluating the use of D-dimer to assess suspected DVT in pregnant women, revealed a high negative predictive value proposing that a negative D-dimer test using pregnancy-specific cut-offs may sufficiently exclude DVT in pregnancy.^{27,29,30}

CT Pulmonary Angiography

There is no local data to evaluate the yield of CTPA in pregnant and postpartum patients, however accepted standards in the general population indicate CTPA should detect PE in 15.4 - 37.4% of patients and alternative diagnoses in up to 56%.³¹

A study done by Shahir et al found that 5.6% of pregnant women who underwent CTPA had nondiagnostic studies, a rate similar to that in the general non-pregnant population.³² Certain physiological factors may influence the quality of CTPA such as increased circulatory volume and altered cardiac output, which effectively leads to lower pulmonary artery opacification.³² Insufficient opacification of the pulmonary vasculature during CTPA is described twice as often among pregnant women compared to those who are not pregnant.³³ This may be explained by the increase in plasma volume with consequent dilution of contrast medium, in addition the hyperdynamic circulation in pregnancy results in contrast disruption when there is accelerated mixture of opacified blood from the superior vena cava and non-opacified blood from the inferior vena cava.²¹ Evidently normal CTPA imaging in pregnancy, has been found to have a high negative predictive value.³⁴ Given the higher radiation dose associated with CTPA, exposure to the maternal breast carries carcinogenic risk on account of the radiosensitive character of the glandular breast in pregnancy, and during lactation.³⁵ Hohl et al observed that CTPA can increase the risk of breast cancer by 1.5%³⁵, however recent studies have established that this risk is negligibly low³⁶ unless there is a family history of breast cancer.³⁷ Adaptions have been made to current CT protocols to ensure lower maternal and foetal radiation doses.^{32,38}

A study by Jordan et al including 34 pregnant women, 16 postpartum women, and 50 age-matched controls, found that CTPA in pregnant patients had a significantly lower positivity rate than controls (2% compared to 16% respectively).³⁹ A 17-year retrospective study by Rotzinger et al done at a single center in Switzerland observed that CTPA rarely indicated PE in pregnancy, with an alternative diagnosis found in 30% of pregnant patients investigated.⁴⁰

Other imaging modalities

Ultrasound and lower limb doppler in pregnant women with suspected PE are recommended by four international guidelines, however factors affecting the selection criteria of patients differ.^{28,41,42} The GTH and ESC guidelines advocate for bilateral lower limb doppler for all pregnant women with a suspicion for PE, regardless of whether there are clinical evidence of DVT,^{28,41} whilst the RCOG and ATS guidelines suggest that lower limb ultrasound only be used in pregnant patients who have clinical

evidence of DVT,²³ as the diagnostic yield of lower limb ultrasound is low when DVT symptoms are absent.

Despite the role for chest radiography in patients with suspected PE, it is neither sensitive nor specific.²¹ Nevertheless, its role is vital as it can reveal other underlying pathologies that may provide an alternative explanation for the presenting symptoms.²¹ In addition, V/Q imaging can only be accurately interpreted in the presence of a normal chest X-ray.²¹

Compared to ventilation/perfusion (V/Q) imaging, CTPA has been associated with an higher radiation dose to maternal breast tissue, but lower dose to the fetus.⁴³ The dose of radiation per maternal breast is 0.11 – 0.31 mGy in perfusion scanning⁴⁴, and 10 – 70 mGy in CTPA.³⁵ V/Q imaging and CTPA in pregnancy, have a similar yield of non-diagnostic results; 12.5% (95% CI 5.3 -19.6) and 11.5% (95% CI 7.5 – 18.4) respectively in a systematic review that evaluated 22 studies with a total of 2,391 imaging investigations (1,165 V/Q scan and 1,226 CTPA).⁴⁵ However, in South Africa, CT remains the primary diagnostic modality for PE given the limited number of Nuclear Medicine units and the relative ease of access to CT facilities.⁴⁶

1.2 Aim of the study

PE is the leading cause of mortality in pregnant and postpartum patients worldwide, therefore physicians have a low threshold to investigate for it. International data established that CTPA is effective in detecting PE, as well as alternative diagnoses. In South Africa CTPA is considered the investigation of choice for PE, as VQ scanning is not available in most settings. There is however no local data to evaluate the yield of CTPA in pregnant and postpartum patients in South Africa. The aim of the study is to evaluate the yield of CTPA to diagnose PE in pregnancy and postpartum period in a South African context, and to compare clinical features in patients suspected to have PE in pregnancy and the postpartum period who underwent CTPA, as well as to ascertain the prevalence of alternative CT findings in patients without PE who underwent CTPA.

1.3 Research Question

In patients admitted to an Obstetric High Care unit what were the clinical characteristics of patients that underwent CTPA, what proportion of patients had a pulmonary embolus

and was there a difference in clinical findings in those with and without pulmonary embolus by CTPA?

1.4 Study Objectives

- To describe the clinical characteristics of patients that underwent CTPA and determine the proportion of patients undergoing CTPA that had a pulmonary embolus
- To compare the clinical features of those patients with PE to those without
- To ascertain the prevalence of alternative CT findings in patients without PE that underwent CTPA

1.5 Methods

Study Design

This study will be a retrospective cross sectional review.

Study Population

Pregnant and postpartum women admitted to the Obstetric High Care unit at Charlotte Maxeke Johannesburg Academic Hospital from 1 January to 31 December 2019, inclusive, who underwent CTPA.

Exclusion Criteria

- Patients who had a PE diagnosed by a method other than CTPA
- Patients referred from other institutions where imaging was performed elsewhere
- Patients with known PE, diagnosed prior to admission
- Patients who were not pregnant or not in the puerperium (period of 6 weeks following delivery)

Data Collection

Data will be collected from the following sources:

- The names and hospital numbers of all patients admitted to the Obstetric High Care Unit for the study period will be collected from the ward admission register.
- The names and hospital numbers above will be cross-referenced with the hospital PACS Radiology system to determine which patients underwent CTPA.
- The included patients will be anonymised and allocated a unique study number which corresponds to the included patient.

- The CTPA images which were done for the included patients will be reviewed and reported on by two radiologist to remove any interobserver bias that may exist in the interpretation of the CTPA results
- The files for the relevant patients will be collected from the archives department and clinical data as indicated on the data collection sheet (Appendix A) will be extracted.

Statistical Analysis

General Statistical considerations

This is a cross sectional retrospective analysis with the primary objective to describe the clinical characteristics of patients who underwent CTPA and to compare the clinical features of those patients with PE to those without. Data will be entered into a database which will then be used to perform statistical analysis. Descriptive statistics will be employed to analyse the data. Summary statistics with variables will include the mean, and standard deviation. Categorical variables will be presented as frequency counts and percentages. The data will be tabulated and analysed with respect to patient demographics, clinical features and laboratory investigations, and outcome of radiological investigations with appropriate statistical tests chosen depending on the distribution of data.

Analysis populations

The Obstetric High Care at CMJAH had 800 admissions in the 2019 calendar year. We anticipate, through estimations from the head of unit, that approximately 10% of these women would undergo CTPA during admission. Given this estimation and based on a 95% confidence level with a 5% margin of error we aim to include 66 women in our study population, where imaging will be reviewed and clinical characteristics will be tabulated and compared to the radiological findings.

Efficacy Analysis

Outcome of radiological imaging will be analysed. In view of the findings on imaging, the clinical characteristics of those patients with confirmed PE will be compared to those with PE excluded. The incidence of PE in patients who underwent CTPA will be compared to statistics from international data.

1.6 Limitations

- The retrospective design of the study places limitation on:

- The availability of clinical data as all required data may not have been recorded,
- The availability of radiological imaging as studies with suboptimal imaging will not be able to be repeated.
- The patient population is limited to admissions to the high care unit, however the majority of patients with suspected PE are managed in this unit.
- The imaging modality is limited to CTPA, excluding patients who received ventilation perfusion scans.
- This study will take place in a single healthcare centre, excluding patients from other centres.

1.7 Ethical approval



Ethical approval will be obtained from the Wits Human Research Ethics Committee (Medical). Permission to conduct the study at the research site will be sought from the Research Committee of Charlotte Maxeke Johannesburg Academic Hospital and the Heads of the Departments of Obstetrics & Gynaecology and Radiology, to access clinical and radiological data respectively. The study will also be registered with the National Health Research Database.

1.8 Funding

The preparation and printing costs will be self-funded, which will include the following:

- Stationery – R 1 500.
- Hard drive to contain imaging to be reviewed by Radiologist – R 1 500.

1.9 Timeline

	January 2022	February 2022	March 2022	April 2022	May 2022	June 2022	July 2022	August 2022	September 2022	October 2022
Completion of Protocol										
Submission to the Protocol Assessment Committee, Ethics Committee and Hospital Research Committee										
Protocol Assessment										
Data Collection, analysis and interpretation (depending on outcome of Protocol, Ethics and Hospital Research applications)										
Completion of study and write up										
Supervisor review										
Submission of completed project										

1.10 References

1. MBRRACE-UK Update: Key messages from the UK and Ireland Confidential Enquiries into Maternal Death and Morbidity 2017. *Obstet Gynecol.* 2018;20(1):75-79. Available from: <http://doi.org/10.1111/tog.12466>.
2. Meng K, Hu X, Peng X, Zhang Z. Incidence of venous thromboembolism during pregnancy and the puerperium: a systematic review and meta-analysis. *J Matern.* 2014;28(3):245-253. Available from: <https://doi.org/10.3109/14767058.2014.913130>.
3. Pomp E, Lenselink A, Rosendaal F, Doggen C. Pregnancy, the postpartum period and prothrombotic defects: risk of venous thrombosis in the MEGA study. *J Thromb.* 2008;6(4):632-637. Available from: <https://doi.org/10.1111/j.1538-7836.2008.02921.x>.
4. Kane E, Calderwood C, Dobbie R, Morris C, Roman E, Greer I. A population-based study of venous thrombosis in pregnancy in Scotland 1980–2005. *Eur J Obstet Gynecol Reprod Biol.* 2013;169(2):223-229. Available from: <https://doi.org/10.1016/j.ejogrb.2013.03.024>.
5. James A, Tapson V, Goldhaber S. Thrombosis during pregnancy and the postpartum period. *Am J Obstet Gynecol.* 2005;193(1):216-219. Available from: <https://doi.org/10.1016/j.ajog.2004.11.037>.
6. Cantwell R, Clutton-Brock T, Cooper G, et al. Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer: 2006-2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG.* 2011;118 Suppl 1:1-203. Available from: <https://doi.org/10.1111/j.1471-0528.2010.02847.x>.
7. Heit J, Kobbervig C, James A, Petterson T, Bailey K, Melton L. Trends in the Incidence of Venous Thromboembolism during Pregnancy or Postpartum: A 30-Year Population-Based Study. *Ann Intern Med.* 2005;143(10):697. Available from: <https://doi.org/10.7326/0003-4819-143-10-200511150-00006>.
8. Jacobsen A, Skjeldestad F, Sandset P. Incidence and risk patterns of venous thromboembolism in pregnancy and puerperium—a register-based case-control study. *AJOG.* 2008;198(2):233.e1-233.e7. Available from: <https://doi.org/10.1016/j.ajog.2007.08.041>.
9. Kamel H, Navi B, Sriram N, Hovsepian D, Devereux R, Elkind M. Risk of a Thrombotic Event after the 6-Week Postpartum Period. *N Engl J Med.* 2014;370(14):1307-1315. Available from: <https://dx.doi.org/10.1056%2FNEJMoa1311485>.
10. Kline J, Richardson D, Than M, Penaloza A, Roy P. Systematic Review and Meta-analysis of Pregnant Patients Investigated for Suspected Pulmonary Embolism in the Emergency Department. *Acad Emerg Med.* 2014;21(9):949-959. Available from: <https://doi.org/10.1111/acem.12471>.

11. Chatur S, Islam S, Moore L, Sandhu R, Sheldon R, Kaul P. Incidence of Syncope During Pregnancy: Temporal Trends and Outcomes. *J Am Heart Assoc*, 2019;8(10), e011608. Available from: <https://doi.org/10.1161/JAHA.118.011608>.
12. Soma-Pillay P, Nelson-Piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr*. 2016;27(2):89-94. Available from: <https://dx.doi.org/10.5830%2FCVJA-2016-021>.
13. Boeldt D, Bird I. Vascular adaptation in pregnancy and endothelial dysfunction in preeclampsia. *J Endocrinol*. 2017;232(1):R27-R44. Available from: <https://doi.org/10.1530/joe-16-0340>.
14. Bates S, Greer I, Middeldorp S, Veenstra D, Prabalos A, Vandvik P. VTE, Thrombophilia, Antithrombotic Therapy, and Pregnancy. *Chest*. 2012;141(2):e691S-e736S. Available from: <https://doi.org/10.1378/chest.11-2300>.
15. Bibas M, Biava G, Antinori A. HIV-Associated Venous Thromboembolism. *Mediterr J Hematol Infect Dis*. 2011;3(1):e2011030. <https://dx.doi.org/10.4084%2FMJHID.2011.030>.
16. Di Prima FA, Valenti O, Hyseni E, et al. Antiphospholipid Syndrome during pregnancy: the state of the art. *J Prenat Med*. 2011;5(2):41-53. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3279165/pdf/prenatal-05-0041.pdf>. [Accessed 12th January 2022].
17. Blokhin I, Lentz S. Mechanisms of thrombosis in obesity. *Curr Opin Hematol*. 2013;20(5):437-444. Available from: <https://doi.org/10.1097/moh.0b013e3283634443>.
18. Kevane B, Donnelly J, D'Alton M, Cooley S, Preston RJ, Ní Ainle F. Risk factors for pregnancy-associated venous thromboembolism: a review. *J Perinat Med*. 2014;42(4):417-425. Available from: <https://doi.org/10.1515/jpm-2013-0207>.
19. Righini M, Le Gal G, Bounameaux H. Venous thromboembolism diagnosis: unresolved issues. *Thromb Haemost*. 2015;113(06):1184-1192. Available from: <https://doi.org/10.1160/th14-06-0530>.
20. Chan W, Ray J, Murray S, Coady G, Coates G, Ginsberg J. Suspected Pulmonary Embolism in Pregnancy. *Arch Intern Med*. 2002;162(10):1170. Available from: <http://doi.org/10.1001/archinte.162.10.1170>.
21. Wan T, Skeith L, Karovitch A, Rodger M, Le Gal G. Guidance for the diagnosis of pulmonary embolism during pregnancy: Consensus and controversies. *Thromb Res*. 2017;157:23-28. Available from: <https://doi.org/10.1016/j.thromres.2017.06.025>.
22. Chan W, Rey E, Kent N, Chan W, Kent N, Rey E et al. Venous Thromboembolism and Antithrombotic Therapy in Pregnancy. *J Obstet Gynaecol Can*. 2014;36(6):527-553. Available from: [https://doi.org/10.1016/s1701-2163\(15\)30569-7](https://doi.org/10.1016/s1701-2163(15)30569-7).
23. Royal College of Obstetricians & Gynaecologists. Thromboembolic disease in pregnancy and the puerperium: acute management, Green-top Guideline No. 37b.

- [homepage in the internet]. c2015 [updated 2015 April] Available from: <https://www.rcog.org.uk/globalassets/documents/guidelines/gtg-37b.pdf>. [Accessed 13th January 2022].
24. O'Connor C, Moriarty J, Walsh J, Murray J, Coulter-Smith S, Boyd W. The application of a clinical risk stratification score may reduce unnecessary investigations for pulmonary embolism in pregnancy. *J Matern Fetal Neonatal Med*. 2011;24(12):1461-1464. Available from: <https://doi.org/10.3109/14767058.2011.614652>.
 25. Cutts B, Tran H, Merriman E, Nandurkar D, Soo G, DasGupta D et al. The utility of the Wells clinical prediction model and ventilation-perfusion scanning for pulmonary embolism diagnosis in pregnancy. *Blood Coag Fibrinolysis*. 2014;25(4):375-378. Available from: <https://doi.org/10.1097/mbc.0000000000000054>.
 26. Chan W, Chunilal S, Lee A, Crowther M, Rodger M, Ginsberg J. A Red Blood Cell Agglutination d-Dimer Test to Exclude Deep Venous Thrombosis in Pregnancy. *Ann Intern Med*. 2007;147(3):165. Available from: <https://doi.org/10.7326/0003-4819-147-3-200708070-00005>.
 27. van der Pol L, Tromeur C, Bistervels I, Ni Ainle F, van Bommel T, Bertoletti L et al. Pregnancy-Adapted YEARS Algorithm for Diagnosis of Suspected Pulmonary Embolism. *N Engl J Med*. 2019;380(12):1139-1149. Available from: <http://doi.org/10.1056/NEJMoa1813865>.
 28. Linnemann B, Scholz U, Rott H, Halimeh S, Zotz R, Gerhardt A et al. Treatment of pregnancy-associated venous thromboembolism - position paper from the Working Group in Women's Health of the Society of Thrombosis and Haemostasis (GTH). *Vasa*. 2016;45(2):103-118. Available from: <https://doi.org/10.1024/0301-1526/a000504>.
 29. Righini M, Robert-Ebadi H, Elias A, Sanchez O, Le Moigne E, Schmidt J et al. Diagnosis of Pulmonary Embolism During Pregnancy: A Multicenter Prospective Management Outcome Study. *Ann Intern Med*. 2018;169(11):766-773. Available from: <https://doi.org/10.7326/m18-1670>.
 30. Goodacre S, Horspool K, Nelson-Piercy C, Knight M, Shephard N, Lecky F et al. The DiPEP study: an observational study of the diagnostic accuracy of clinical assessment, D-dimer and chest x-ray for suspected pulmonary embolism in pregnancy and postpartum. *BJOG*. 2019;126(3):383-392. Available from: <https://doi.org/10.1111/1471-0528.15286>.
 31. Dempster E, Gunatillake D, Kumarasena U. CTPA; can we say no to a raised d-dimer?. *Clin Radiol*. 2016;71:S11. Available from: <https://doi.org/10.1016/j.crad.2016.06.066>.
 32. Shahir K, Goodman L, Tali A, Thorsen K, Hellman R. Pulmonary Embolism in Pregnancy: CT Pulmonary Angiography Versus Perfusion Scanning. *AJR Am J Roentgenol*. 2010;195(3):W214-W220. Available from: <https://doi.org/10.2214/ajr.09.3506>.

33. U-King-Im J, Freeman S, Boylan T, Cheow H. Quality of CT pulmonary angiography for suspected pulmonary embolus in pregnancy. *Eur Radiol.* 2008;18(12):2709-2715. Available from: <https://doi.org/10.1007/s00330-008-1100-0>.
34. Sun S, Diaconescu M, Zhe T, Mesurolle B, Semionov A. Outcomes of Multidetector Computed Tomography Pulmonary Angiography in Pregnant and Postpartum Women With Suspected Pulmonary Embolism. *Can Assoc Radiol J.* 2021;72(3):512-518. Available from: <https://doi.org/10.1177/0846537119899552>.
35. Hohl C, Wildberger J, Suss C, Thomas C, Mühlenbruch G, Schmidt T et al. Radiation dose reduction to breast and thyroid during MDCT: effectiveness of an in-plane bismuth shield. *Acta Radiol.* 2006;47(6):562-567. Available from: <http://doi.org/10.1080/02841850600702150>.
36. Burton K, Park A, Fralick M, Ray J. Risk of early-onset breast cancer among women exposed to thoracic computed tomography in pregnancy or early postpartum. *J Thromb.* 2018;16(5):876-885. Available from: <http://doi.org/10.1111/jth.13980>.
37. Hill D, Preston-Martin S, Ross R, Bernstein L. Medical Radiation, Family History of Cancer, and Benign Breast Disease in Relation to Breast Cancer Risk in Young Women, USA. *Cancer causes control.* 2002;13(8):711-718. Available from: <http://doi.org/10.1023/a:1020201106117>.
38. Litmanovich D, Boiselle P, Bankier A, Kataoka M, Panykh O, Raptopoulos V. Dose Reduction in Computed Tomographic Angiography of Pregnant Patients With Suspected Acute Pulmonary Embolism. *JCAT.* 2009;33(6):961-966. Available from: <http://doi.org/10.1097/RCT.0b013e318198cd18>.
39. Jordan E, Godelman A, Levsky J, Zalta B, Haramati L. CT pulmonary angiography in pregnant and postpartum women: low yield, high dose. *Clin Imaging.* 2015;39(2):251-253. Available from: <http://doi.org/10.1016/j.clinimag.2014.11.006>.
40. Rotzinger D, Dunet V, Ilic V, Hugli O, Meuli R, Schmidt S. Pulmonary embolism during pregnancy: a 17-year single-center retrospective MDCT pulmonary angiography study. *Eur Radiol.* 2019;30(3):1780-1789. Available from: <http://doi.org/10.1007/s00330-019-06501-4>.
41. Konstantinides S, Torbicki A, Agnelli G, Danchin N, Fitzmaurice D, Galiè N et al. 2014 ESC Guidelines on the diagnosis and management of acute pulmonary embolism. *Eur.* 2014;72(11):997-1053. Available from: <http://doi.org/10.1093/eurheartj/ehu283>.
42. Leung A, Bull T, Jaeschke R, Lockwood C, Boiselle P, Hurwitz L et al. American Thoracic Society Documents: An Official American Thoracic Society/Society of Thoracic Radiology Clinical Practice Guideline: Evaluation of Suspected Pulmonary Embolism in Pregnancy. *Am J Respir Crit.* 2011;184(10):1200-8. Available from: <http://doi.org/10.1164/rccm.201108-1575ST>.

43. Balan K, Critchley M, Vedavathy K, Smith M, Vinjamuri S. The value of ventilation-perfusion imaging in pregnancy. *Br J Radiol.* 1997;70(832):338-340. Available from: <https://doi.org/10.1259/bjr.70.832.9166067>.
44. Pahade J, Litmanovich D, Pedrosa I, Romero J, Bankier A, Boiselle P. Imaging Pregnant Patients with Suspected Pulmonary Embolism: What the Radiologist Needs to Know. *RadioGraphics.* 2009;29(3):639-654. Available from: <http://doi.org/10.1148/rg.293085226>.
45. Tromeur C, van der Pol L, Le Roux P, Ende-Verhaar Y, Salaun P, Leroyer C et al. Computed tomography pulmonary angiography versus ventilation-perfusion lung scanning for diagnosing pulmonary embolism during pregnancy: a systematic review and meta-analysis. *Haematologica.* 2018;104(1):176-188. Available from: <http://doi.org/10.3324/haematol.2018.196121>.
46. Ahlers P, Said-Hartley M. A retrospective review of CT pulmonary angiogram confirmed pulmonary emboli in COVID-19 patients admitted to Groote Schuur Hospital, Cape Town. *S Afr J Rad.* 2022;26(1), a2280. Available from <https://dx.doi.org/10.4102/sajr.v26i1.2280>.

APPENDIX A: Data Collection Sheet

Pulmonary embolism in Obstetric HCU													
Participant Number													
Demographics													
DOB				Age			Gender			F			
Race	Black		White		Coloured		Indian		Asian				
History													
Pregnant?	Yes		No		Postpartum			Yes		No			
Multiple Gestation	Yes		No										
Gestational Age				Weeks Postpartum									
Method (GA)	LNMP		Early US		Mode of Delivery			CS		NVD			
	Late US		Clinical					Assisted					
Obstetric Hx	Gravidity			Parity				Miscarriages					
Pregnancy Complications present?				Yes					No				
If yes, which:	Anaemia		Antepartum Haemorrhage					Chorioamnionitis					
(Can select more than one)	IUGR		Placenta Praevia					Preterm Labour					
	Preterm ROM		Preeclampsia					Eclampsia					
	HELLP		PPH					PIH					
	PPCMO		Gestational DM					Puerperal Sepsis					
			Relook Laparotomy										
	ICU Admission		Reason:										
	Other		What other:										
Medical History													
Autoimmune Disease				Yes				No				Unknown	
Antiphospholipid Syndrome				Yes				No				Unknown	

Chronic Kidney Disease	Yes		No		Unknown	
Chronic Lung Disease	Yes		No		Unknown	
Diabetes	Yes		No		Unknown	
Epilepsy	Yes		No		Unknown	
Heart Failure	Yes		No		Unknown	
HIV	Yes		No		Unknown	
Hypertension	Yes		No		Unknown	
Prev. Tuberculosis	Yes		No		Unknown	
Valvular Heart Disease	Yes		No		Unknown	
Previous VTE?	Yes		No		Unknown	
Comments / Elaborate						

Examination									
BP		Pulse		Resp Rate					
Height		Weight		BMI					
MUAC		Saturation							
CVS									
Resp									
Modified YEARS Algorithm Score: _____									
Pregnant:	Yes		No						
Clinical signs of DVT:	Yes		No						
Haemoptysis:	Yes		No						
PE most likely dx:	Yes		No						
D-Dimer	< 0.5 mg/L		0.5 – 0.99 mg/L		> 1 mg/L				
Well's Score for PE Score: _____									
Clinical signs and symptoms of DVT:				Yes (+3)		No (0)			
PE is #1 likely diagnosis OR equally likely:				Yes (+3)		No (0)			
Heart rate >100:				Yes (+1.5)		No (0)			
Immobilization at least 3 days OR surgery in the previous 4 weeks:				Yes (+1.5)		No (0)			
Previous, objectively diagnosed PE or DVT:				Yes (+1.5)		No (0)			
Haemoptysis:				Yes (+1)		No (0)			
Malignancy:				Yes (+1)		No (0)			
ECG Feat.	Right atrial enlargement (P pulmonale)		Sinus tachycardia		Clockwise rotation				
	S ₁ Q ₃ T ₃ Pattern		Dominant R wave V1		Atrial tachyarrhythmias				
	Complete or incomplete RBBB		Right ventricular strain						
	Non-specific ST segment and T wave changes (including ST elevation and depression)								
Laboratory Data (within 24h of CTPA)									
FBC									
U&E									
D-Dimer		CRP		PCT					
Radiological Data									
CTPA	PE		No PE		Inconc.				
Distribution of PE	Proximal (Pulmonary Trunk/Pulmonary Artery)								

		Lobar (Interlobar Artery)				
		Segmental				
		Subsegmental				
Other CT Findings						
Pleural Effusion		Pneumothorax			GGO	
Consolidation		Cavitation			Bronchiectasis	
Atelectasis		Nodules			Masses	
Side:		Left		Right		
Comments (include location for each of the above findings)						



27/01/2022

Dear Prof. Chauke,

I wish to request permission to conduct research with patients from the Department of Obstetrics and Gynaecology, for the purpose of my MMed, titled A Review of the use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre.

The research will be supervised by Dr Shastra Bhoora (Dept Obstetrics & Gynaecology), Dr Jarrod Zamparini (Dept Internal Medicine), and Dr Halvani Moodley (Dept Radiology).

The participants will include admissions to the Obstetric High Care unit at Charlotte Maxeke Johannesburg Academic Hospital for the period January to December 2019. Access to patient files is requested.

Please find attached the protocol for the research project.

Your approval will be much appreciated.

Sincerely,

A handwritten signature in black ink, appearing to read "Wilhelm", written over a horizontal line.

Dr Wilhelm Herbst
Registrar: Internal Medicine



27/01/2022

Dear Prof. Ramaema,

I wish to request permission to conduct research with patients from the Department of Radiology, for the purpose of my MMed, titled A Review of the use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre.

The research will be supervised by Dr Halvani Moodley (Dept Radiology), Dr Jarrod Zamparini (Dept Internal Medicine), and Dr Shastra Bhoora (Dept Obstetrics & Gynaecology).

The participants will include admissions to the Obstetric High Care unit at Charlotte Maxeke Johannesburg Academic Hospital for the period January to December 2019, who received CT scans during this period. Access to PACS to review CT scans of study participants is requested.

Please find attached the protocol for the research project.

Your approval will be much appreciated.

Sincerely,

A handwritten signature in black ink, appearing to read "Wilhelm Herbst", written over a horizontal line.

Dr Wilhelm Herbst
Registrar: Internal Medicine



27/01/2022

Dear Prof. Chakane,

I wish to request permission to conduct research at Charlotte Maxeke Johannesburg Academic Hospital for the purpose of my MMed, titled A Review of the use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre.

The research will be supervised by Dr Shastra Bhoora (Dept Obstetrics & Gynaecology), Dr Jarrod Zamparini (Dept Internal Medicine), and Dr Halvani Moodley (Dept Radiology).

The participants will include admissions to the Obstetric High Care unit at CMJAH for the period January to December 2019. Access to patient files is requested.

Please find attached the protocol for the research project.

Your approval will be much appreciated.

Sincerely,

A handwritten signature in black ink, appearing to read "Wilhelm", written over a horizontal line.

Dr Wilhelm Herbst
Registrar: Internal Medicine

CHAPTER 2: MANUSCRIPT

This manuscript complies with the author guidelines for the journal *Obstetric Medicine* and has been formatted as such.

2.1 TITLE PAGE

A Submission to the journal Obstetric Medicine

Full Title: A Review of the Use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre

Short Title: CTPA in Pregnant and Postpartum Patients

Authors:

1. Wilhelm Herbst, MBChB (Pret) [**corresponding author**]
Department of Internal Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg
Postal: PO box 36381 Menlo Park Pretoria, 0181.
Email: wherbst.herbst@gmail.com
Tel: 079 885 9694
2. Shastra Bhoora, MBBCh (Wits), FCOG (SA), MMed (Wits), Cert Crit Care (SA), MSc (Wits)
Department of Obstetrics and Gynaecology, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand, Johannesburg.
Department of Obstetrics and Gynaecology, Baylor College of Medicine, Houston.
3. Halvani Moodley, MBChB(Pret), FC Rad Diag (SA), MMed Rad Diag (WITS), Dip Ped Rad (UCT), Department of Radiology, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand.
4. Ashesh Ishwarlal Ranchod, MBBCh (Wits), FCRAD (SA), EDiPNR Honorary lecturer, Department of Radiology, School of Medicine, Faculty of Health Sciences, University of the Witwatersrand. National Radiology Services Incorporated (NRS Inc).
5. Tracy Westgarth-Taylor, MBChB, FCRAD (Diag) (SA), Netcare Park Lane Hospital.
6. Jarrod Zamparini, MBChB (Pret), MMed (Wits), FCP (SA)
Obstetric Internal Medicine Unit, Charlotte Maxeke Johannesburg Academic Hospital and Department of Internal Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg

Declaration of Interests: The authors declare no conflicting interests

Funding: This research received no specific grant or funding

Informed Consent: Retrospective

Ethical Approval: The Wits Human Research Ethics Committee (Medical) approved this study (Clearance Certificate No. M220255)

Guarantor: WH

Authorship: WH and JZ conceived the study. SB assisted with data identification and retrieval. WH gained ethics approval, conducted the literature review, designed the data collection tool, collected raw data and wrote the first draft of the manuscript. HM, TWT and AR reviewed and reported all CT images. All authors contributed to the final manuscript.

Acknowledgements: We would like to thank all authors for their contribution.

Keywords: Pregnancy, VTE, CTPA

2.2 ABSTRACT

Background: Pulmonary embolism (PE), is a common cause of maternal death during pregnancy and the puerperium yet data on its prevalence in middle-income countries is lacking.

Methods: We examined the medical records and CTPAs of 67 women in an obstetric high care unit during pregnancy and the puerperium. We aimed to assess the efficacy of CTPA in diagnosing PE, evaluate associated clinical features, and determine the prevalence of alternative CT findings.

Results: CTPA detected PE in 11 women (16.42%) and alternative CT findings in 46 (68.6%). Women with PE had a significantly lower systolic BP than those without PE ($p = 0.001$). Multiple gestation, preterm rupture of membranes, and gestational diabetes were linked to higher PE prevalence.

Conclusions: This study, set in an upper middle-income country, demonstrated a higher CTPA yield for PE than in the international literature, emphasizing context-specific assessments.

2.3 INTRODUCTION

Venous thromboembolism (VTE), including pulmonary embolism (PE), is the most common cause of maternal death during pregnancy and the puerperium in developed countries.¹ The risk for venous thromboembolism (VTE) is significantly increased during pregnancy and the postpartum period, as these patients are in a physiological state of hypercoagulability, are prone to venous stasis and may have superimposed endothelial damage.²

Women have a 5-fold increased risk of developing VTE during pregnancy, as compared to their non-pregnant counterparts.³ Globally, the incidence of PE in pregnant or postpartum women ranges from 1-2 per 1,000 to 3 per 10,000 deliveries^{2,4}, and one death in every 100,000 deliveries.⁵ In South Africa PE accounted for 40 maternal deaths in 2022, with a maternal mortality rate of 4.1 per 100,000 live births.⁶

More than half of VTE cases in pregnancy occur before 20 weeks' gestation,⁴ and 80% of VTE cases in the postpartum period occur within the first 3 weeks following delivery.⁷

Despite the increased risk of VTE in pregnancy, detection rates using CTPA (computed tomography pulmonary angiography) are lower in pregnant than non-pregnant women. A 2015 study found that 2% of pregnant women had a positive CTPA compared to 16% of non-pregnant women.⁸ A 17-year retrospective study conducted at a single centre in Switzerland, published in 2020, observed that CTPA rarely indicated PE in pregnancy, with an alternative diagnosis found in 30% of pregnant women investigated.⁹ Similarly, a 2023 study including 86 pregnant and 354 non-pregnant women investigated for PE by either CTPA or V/Q scan, noted positive scan rates of 3.5% and 8.8% in pregnant and non-pregnant women respectively.¹⁰

Risk Factors

Pregnancy is a hypercoagulable physiological state owing to an increased concentration of certain clotting factors as well as a decrease in endogenous anticoagulants such as antithrombin and protein S¹¹. In addition, factors such as dehydration, secondary to nausea and vomiting in early pregnancy, immobility and iliac compression by the gravid uterus in late pregnancy, and endothelial injury from delivery trauma and venous distension^{11,12} also play a role in the increased risk for VTE. The

clinical presentation of VTE may also be masked by “symptoms” of pregnancy such as pedal oedema, tachypnoea and tachycardia.¹³ A number of postpartum factors are also associated with an increased risk of VTE, such as surgical delivery, particularly in the emergency setting or after prolonged labour, postpartum haemorrhage, thrombophilia, pre-eclampsia, and puerperal sepsis.¹⁴

Clinical predictors

The Wells’ criteria, which factors in tachycardia and lower limb oedema, is not particularly useful in pregnancy, given that these manifestations are frequently seen.¹⁵ This complicates the differentiation of PE from other potential diagnoses in these women.¹⁵ The recommendations outlined by the Society of Obstetricians and Gynaecologists of Canada (SOGC)¹⁶ and the Royal College of Obstetricians and Gynaecologists (RCOG)¹⁷, advise against clinical predictors to assess VTE in pregnant women.

D-dimer

In non-pregnant women PE can be safely excluded in the presence of a normal D-dimer in addition to a low pre-test probability. It must conversely be interpreted with caution in those with a high pre-test probability, as a falsely negative result can miss a diagnosis of PE.¹⁸ The American Thoracic Society (ATS)¹⁹, SOGC¹⁶, and RCOG¹⁷ recommendations state that a D-dimer should not be used in pregnancy to exclude PE as most studies investigating the use of D-dimer in diagnosing PE, have not included pregnant women as participants.²⁰

Van der Pol et al found that the pregnancy-adapted YEARS algorithm safely excluded acute PE in pregnant women with suspected PE, allowing CTPA to be avoided in 32 to 65% of women.²¹ A South African study found that the pregnancy-adapted YEARS algorithm successfully risk-stratified women in a single institution and would have reduced the number of CTPAs by 25.7% without missing any PE.²²

CT Pulmonary Angiography

Accepted standards outlined by the Royal College of Radiologists state that CTPA should detect PE in 15.4 - 37.4% and alternative diagnoses in up to 56% in the general population.²³

Most studies have disregarded pregnant women when investigating the diagnosis of PE.¹⁸ Certain physiological factors, such as increased circulatory volume and altered cardiac output, may lead to lower pulmonary artery opacification influencing the quality of CTPA.²⁴ Insufficient opacification of the pulmonary vasculature during CTPA is described twice as often among pregnant women compared to those who are not pregnant.²⁵ Normal CTPA imaging in pregnancy, has been found to have a high negative predictive value.²⁶

In this study we investigated the use of CTPA to diagnose PE in pregnant and postpartum women admitted to a high care setting in South Africa. We aimed to evaluate the prevalence of PE and alternative diagnoses as well as compare the clinical features of women with and without PE.

2.4 METHODS

We performed a cross-sectional analysis at the Charlotte Maxeke Johannesburg Academic Hospital, an academic centre in Johannesburg, South Africa. Pregnant and postpartum women admitted to the Obstetric High Care Unit (OHCU) between 1 January and 31 December 2019, who underwent CTPA for the suspicion of PE were consecutively enrolled in this study. We excluded women who had a PE diagnosed by a method other than CTPA, such as V/Q scan, due to limited availability in our setting. Women with imaging performed elsewhere, women with known PE diagnosed prior to admission, and women who were not pregnant or beyond 6 weeks post-delivery were also excluded.

The names and hospital numbers of all women admitted to the OHCU for the study period were collected from the ward admission register and cross-referenced with the hospital picture archiving and communicating system (PACS) to determine which women underwent CTPA. The included participant details were anonymised and allocated a unique study number. The CTPA images were reported on by two consultant radiologists (AR and TWT) who were blinded from each other's findings. The discrepant readings were reviewed by a third consultant radiologist (HM), also blinded from the findings of the other readers.

A final diagnosis was considered as agreement by two or more readers. Those studies with inconclusive findings or failing to achieve consensus among at least two readers, were interpreted as not having PE for the purposes of the study. The hospital files for the relevant participants were collected from hospital archives and clinical data was extracted. Data was captured in Microsoft Excel 16.78 (Microsoft Corporation, Redmond, USA) and statistical analysis was performed using Prism 10 (GraphPad Software Inc, La Jolla, CA, USA). Given the non-parametric nature of the data, the Mann-Whitney U test was employed for continuous variables, while the Fisher's exact test was utilized for categorical variables to determine significance.

2.5 RESULTS

A total of 732 women were admitted during the period under review, CTPAs were performed on 71 patients, of which 3 were not pregnant nor in the postpartum period and 1 had missing images, excluding them from the study, leaving a total of 67 women in the final analysis (Figure 1). The PE and non-PE groups were of similar age and race. The majority of the cohort were postpartum (82% in the PE group and 87.5% in the no PE group), and caesarean section was the most common mode of delivery in keeping with local trends. Most of the women were booked for antenatal care, and were multiparous. Table 1 provides demographic and pregnancy related data of the cohort.

Eleven women (16.4%) had a confirmed PE; 2 (18%) were pregnant and 9 (82%) postpartum and the majority (n=10, 90.9%) had peripheral PEs. Two women (18.2%) had a single embolism and 9 (81.8%) had multiple emboli.. The gestational age at delivery (in postpartum women) or time of scan (in pregnant women) was not significantly different ($p = 0.259$) in the non-PE group (median 34.5 weeks, IQR 30 – 38), compared to those with PE (median 30 weeks, IQR 26 – 38). Women with PE were more likely to have multiple gestation (27%) than without PE (4%) ($p = 0.028$). There was a larger proportion of women with alternative CT findings among those in the group with PE (n=9, 81.8%) compared to those without PE (n=37, 66.1%), yet the difference was not significant ($p=0.481$) (Table 2).

The difference in the technical adequacy of the CTPAs between the two groups was not significant ($p = 0.556$). This was assessed by measuring the opacification of the

pulmonary artery (Hounsfield units), between the two groups (median 237HU, IQR 216-277 versus median 247HU, IQR 209-285 in the PE and non-PE groups respectively). There was no difference in rate of motion artifact between the two groups ($p = 0.492$).

There was a significant difference in systolic blood pressure (sBP) between the two groups ($p = 0.001$) with lower sBP in the PE group (median 119mmHg, IQR 110 – 124) than in the non-PE group (median 134mmHg, IQR 120.3 - 146). There were no differences in other clinical features including the body mass index (BMI) or mid upper arm circumference (MUAC) between the two groups. Despite most women being tachycardic, there were no differences in electrocardiogram (ECG) findings between the two groups. It is not our standard protocol to analyse D-dimer routinely for the suspicion of PE in pregnancy, therefore only 27 women (40%) had D-dimers available. There was however no difference in D-dimers between the two groups ($p = 0.219$).

Biochemical inflammatory markers were available in a similar proportion of women, showing no difference between the groups, reflected by a median C-reactive protein (CRP) of 112mg/L in the PE group (10/11) and 100mg/L in the non-PE group (51/56) and a median procalcitonin (PCT) of 0.37ug/L in the PE group (8/11) and 0.31ug/L in the non-PE group. A diagnosis of puerperal sepsis was made in 36.4% in the PE group compared to 62.5% in the non-PE group, however the difference was not significant ($p = 0.18$). There were no other differences demonstrated in biochemistry between the two groups.

No women died in the non-PE group, while one woman died in the PE group (9%), however it is unclear if the death was directly caused by PE. Most women had a pregnancy-related complication, with the exception of 2 women (3.6%) in the non-PE group.

There were 2 women with HIV (18.2%) in the PE group, one of whom had a CD4 (CD4 = 464 cells/mm³) and HIV viral load (VL = 764 copies/ml) available, whilst there were 17 women with HIV in the non-PE group (30.4%) where 15 had a CD4 result (median 344.5 cells/mm³, IQR 269-457.75) and all had a viral load ($n = 17$, median 217 copies/ml, IQR 0 – 2472.25).

Gestational diabetes was significantly more prevalent ($p = 0.042$) among those with PE ($n = 1$, 9.1%) compared to those without PE ($n = 2$, 3.6%), as was premature rupture of membranes (PROM) ($p = 0.025$) comparing the PE group ($n = 2$, 18.2%) to the non-PE group ($n = 0$, 0%). There were no other complications or comorbidities with a significant difference between the groups (Table 3).

The Wells' score showed no correlation to PE, as all 67 (100%) women had a Wells' score of at least 4.5 (high risk for PE). When analysing the utility of the YEARS score²⁷, 7 women in the PE group and 20 women in the non-PE group had a D-dimer result available allowing the YEARS Score to be calculated. All 7 (100%) women in the PE group had a positive YEARS score, while 17 (85%) women in the non-PE group had a positive YEARS score. The difference was not significant ($p = 0.545$) (Figure 2).

2.6 DISCUSSION

In our study CTPA detected PE in 16.42% and alternative CT findings in 68.6% of our cohort, which is higher than reported in international literature. A large meta-analysis and systematic review of seventeen studies, including 25,339 women, found a 12.4% positivity rate for VTE in nonpregnant women, compared with 4.1% in pregnant women.²⁸ The high positivity rate in our cohort may be attributed to only including women admitted to a high care unit, assigning them to a higher risk category than other pregnant and postpartum women.

No clinical or biochemical biomarkers predicted PE in our cohort, similar to an Australian study evaluating the use of CTPA and V/Q in pregnancy and the puerperium.²⁹ The use of algorithms such as the modified YEARS and Wells' scores were limited in our cohort and would have been of no utility in reducing the number of CT scans done. However given our limited sample size, future studies are needed to further assess their utility in our setting. This is in keeping with recommendations from RCOG to avoid the use of clinical prediction tools in assessing VTE in pregnancy.¹⁷

Despite international guidelines considering maternal age over 35 years a risk factor for PE there was no significant difference in the age of the PE and the non-PE groups in our cohort.³⁰ There is evidence from the UK³¹ and USA³² that Black, Asian and minority ethnicities (BAME) have a much higher maternal mortality rate (MMR), with

the highest among Black women. There is evidence that PE is more prevalent in black ethnicities,³³ with a higher disease severity compared to other ethnicities.³⁴ The majority of our cohort were of Black ethnicity, making it difficult to compare outcomes related to ethnicity and to assess whether disparities exist.. This emphasises the importance of the high risk in this group, and the need for further research in a South African context

Caesarean delivery rate in the South African private sector was 73.2% in 2015³⁵, and increased from 15.1% in 2006³⁶ to 31.1% in 2022⁶ in the public sector. Caesarean delivery is not done on maternal request in the South African public sector. Compared to the UK, the rate of caesarean delivery was 29%³¹ in Wales, 31%³¹ in England and 35%³¹ in Scotland between 2019 - 2021. South African data shows that pregnant women who die from PE are more likely to have had a caesarean delivery.³⁷ In our cohort all women in the PE group were delivered by caesarean section, although this was for obstetric indications and not due to PE. While Caesarean delivery is a known risk factor for the development of PE³⁸, this was not demonstrated in our cohort and mode of delivery was not a risk factor for PE.

The most significant obstetric complication in our cohort was PROM with a higher prevalence in those with PE (n = 2, 18.2%) compared to those without PE (n = 0, 0%) (p = 0.025). Both women with PROM in the PE group were HIV positive, developed puerperal sepsis and had undergone caesarean delivery by gestational age of 28 and 35 weeks respectively. Only one of these women had a CD4 (CD4 = 464 cells/mm³) and HIV viral load (VL = 764 copies/ml) available, and neither had a history of previous tuberculosis. PE was diagnosed after delivery in both women. The development of PE in these women was likely multifactorial however prolonged bed rest prescribed for the treatment of PROM was associated with an increased risk for PE in a study by *Kovacevich et al* (16.5 cases per 1000 compared to 0.8 per 1000 among pregnant women for whom this therapy was not prescribed).³⁹

Systolic BP was significantly lower (p = 0.001) in the PE group, as was seen in the DiPEP study, however there were no differences in diastolic BP and heart rate. Out of keeping with the lower systolic BP in the PE group is that majority of these were peripheral PEs. However 81.8% of the women with PE had multiple emboli which may have contributed to the lower systolic BP. Hypotension associated with PE is attributed

to acute right ventricular dilatation compromising left ventricular filling and reducing cardiac output. Right ventricular dysfunction seen on echocardiography has been shown to correlate with pulmonary arterial obstruction of >30%, which is expected to occur with a large central PE⁴⁰. Echocardiography was not available to assess for right ventricular strain, however there were no ECG features to suggest this.

The high prevalence of alternative findings on CTPA may be explained by the hospital's protocol for investigating PE, where CTPA is warranted if the chest X-ray is abnormal. Despite there being no significant difference in prevalence of alternative CT findings between women with PE and women without PE, the high burden of alternative findings on CTPA in women with PE could signify underlying pathology contributing to development of PE. The high prevalence of pregnancy related complications in our cohort is due to it consisting only of women in a high care setting.

Multiple gestation was a risk factor for PE in our cohort ($p = 0.028$) in keeping with guidelines from RCOG¹⁷. Gestational diabetes has not been proven to be associated with an increased risk for thrombosis⁴¹, however was associated with PE in our cohort ($p = 0.042$). Despite HIV⁴², antiphospholipid syndrome⁴³, and obesity⁴⁴ known to be associated with increased risk for PE, none were significant risk factors in our cohort.

The strengths of this analysis are driven from the racial representation in our cohort, largely consisting of Black African women, an underrepresented group in other studies, which underpins the need and potential for further research. Additionally, the data investigated were largely complete, all the CTPAs in our cohort contained adequate quality images, and were reviewed by consultant radiologists. Yet, the findings of this analysis must be interpreted in the context of some limitations. The first being, the small sample size. Considering our study was performed at a single centre including only women admitted to a high care setting, this in itself might be a major contributory factor to the high PE burden in our population, as these women were all in a higher risk category by nature of their condition. Another limitation is that chest radiographs were not reviewed, as this could have contributed to analysing the pre-test probability for PE.

2.7 CONCLUSION

PE remains a significant contributor to morbidity during pregnancy and the puerperium. While clinical and laboratory features were not reliable to exclude PE in our cohort, it is justifiable to consider CTPA when suspecting PE in this population, considering the higher detection rate compared to international literature. CTPA has a good yield for diagnosing PE and unmasking alternative diagnoses in a middle-income country setting. The presence of a low systolic blood pressure, multiple gestation, PROM, and gestational diabetes should raise the suspicion for PE in a high care setting. Prospective studies evaluating the use of d-dimer and scoring systems to predict PE in our population are warranted.

2.8 DECLARATIONS:

A Submission to the journal Obstetric Medicine

Full Title: A Review of the Use of CT Pulmonary Angiography in Pregnant and Postpartum Patients at an Academic Centre

Short Title: CTPA in Pregnant and Postpartum Patients

Authors:

1. Wilhelm Herbst, MBChB (Pret) [***corresponding author***]
Department of Internal Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg
Postal: PO box 36381 Menlo Park Pretoria, 0181.
Email: wherbst.herbst@gmail.com
Tel: 079 885 9694
2. Shastra Bhoola, MBBCH (Wits), FCOG (SA), MMed (Wits), Cert Crit Care (SA), MSc (Wits)
Department of Obstetrics and Gynaecology, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand, Johannesburg.
Department of Obstetrics and Gynaecology, Baylor College of Medicine, Houston.
3. Halvani Moodley, MBChB(Pret), FC Rad Diag (SA), MMed Rad Diag (WITS), Dip Ped Rad (UCT), Department of Radiology, Charlotte Maxeke Johannesburg Academic Hospital, University of the Witwatersrand.
4. Ashesh Ranchod
5. Tracy Westgarth-Taylor
6. Jarrod Zamparini, MBChB (Pret), MMed (Wits), FCP (SA)
Obstetric Internal Medicine Unit, Charlotte Maxeke Johannesburg Academic Hospital and Department of Internal Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg

Declaration of Interests: The authors declare no conflicting interests

Funding: This research received no specific grant or funding

Informed Consent: Retrospective

Ethical Approval: The Wits Human Research Ethics Committee (Medical) approved this study (Clearance Certificate No. M220255)

Guarantor: WH

Authorship: WH and JZ conceived the study. SB assisted with data identification and retrieval WH gained ethics approval, conducted the literature review, designed the data collection tool, collected raw data and wrote the first draft of the manuscript. HM, TWT and AR reviewed and reported all CT images. All authors contributed to the final manuscript.

Acknowledgements: We would like to thank all authors for their contribution.

Keywords: Pregnancy, VTE, CTPA

2.9 TABLES AND FIGURES:




Figure 1: Study population




Figure 2: D-dimer values

Table 1: Demographics and pregnancy details

Demographics	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
Age (years), median, (IQR)	29 (26 – 35)	30 (26 – 33,75)	0.977
18 – 25	2 (18.18%)	12 (21.43%)	0.992
26 - 30	4 (36.36%)	18 (32.14%)	
31 - 35	3 (27.27%)	16 (28.57%)	
>35	2 (18.18%)	10 (17.86%)	
Race	PE, n (%)	No PE, n (%)	p-value
Black African	11 (100%)	55 (98%)	>0.999
White	0 (0%)	1 (2%)	
Pregnancy Details	PE, n (%)	No PE, n (%)	p-value
Postpartum	9 (82%)	49 (87.5%)	0.635
Pregnant	2 (18%)	7 (12.5%)	
Multiple gestation	3 (27%)	2 (4%)	0.028
Booked for antenatal care	8 (73%)	45 (80%)	0.686
Not booked for antenatal care	3 (27%)	11 (20%)	
Gestational age at delivery or time of scan (weeks), median (IQR)	30 (26 – 38)	34,5 (30 – 38)	0.259
Caesarean section	9 (82%)	41 (73.2%)	0.334
Normal vaginal delivery	0 (0%)	8 (14.3%)	
Gravidity (number of pregnancies), median (IQR)	3 (1 – 3)	2 (1 – 3)	0.971
Parity (number of live births) , median (IQR)	2 (1 -3)	2 (1 – 3)	0.714
Previous miscarriages (number), median (IQR)	0 (0 – 0)	0 (0 – 0)	0.841

Table 2: Alternative CT findings

CT findings	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
Pleural effusion	4 (36.4%)	14 (25%)	0.469
Ground glass opacification	6 (54.5%)	16 (28.6%)	0.157
Consolidation	3 (27.3%)	11 (19.6%)	0.686
Cavitation	0 (0%)	1 (1.8%)	>0.999
Bronchiectasis	0 (0%)	2 (3.6%)	>0.999
Atelectasis	8 (72.7%)	24 (42.9%)	0.101
Nodules	0 (0%)	5 (8.9%)	0.582

Table 3 YEARS score

YEARS score	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
YEARS positive	7 (100%)	17 (85%)	0.545
YEARS negative	0 (0%)	3 (15%)	
YEARS not calculated	4	36	

Table 4: Pregnancy complications and comorbidities

Pregnancy complications	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
No complications	0 (0%)	2 (3.6%)	>0,999
Obstetric Haemorrhage	5 (45.5%)	19 (33.9%)	0,505
Ante-partum haemorrhage	2 (18.2%)	6 (10.7%)	0,609
Post-partum haemorrhage	3 (27.3%)	13 (23.2%)	0,7159
Hypertensive Disorders	7 (63.6%)	42 (75%)	0,469
Pregnancy induced hypertension	1 (9.1%)	12 (21.4%)	0,677
Pre-Eclampsia	3 (27.3%)	18 (32.1%)	>0,999
Eclampsia	1 (9.1%)	9 (16.1%)	>0,999
HELLP	2 (18.2%)	3 (5.4%)	0,186
Obstetric Complications	4 (36.36%)	6 (10.71%)	0,051
Placenta praevia	1 (9.1%)	2 (3.6%)	0,421
Preterm labour	0 (0%)	2 (3.6%)	>0,999
Other	10 (90.1%)	42 (75%)	0,4329
Post-partum cardiomyopathy	2 (18.2%)	13 (23.2%)	>0,999
Anaemia	8 (72.7%)	29 (51.8%)	0,321
Relook laparotomy	3 (27.2%)	8 (14.3%)	0,371
ICU	5 (45.5%)	14 (25%)	0,27
Comorbidities	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
Autoimmune disease	1 (9.1%)	3 (5.4%)	0.521
Anti-phospholipid syndrome	0 (0%)	2 (3.6%)	>0.999
Chronic kidney disease	0 (0%)	4 (7.1%)	>0.999
Diabetes mellitus	1 (9.1%)	1 (1.8%)	0.304
Heart failure	0 (0%)	8 (14.3%)	0.335
Hypertension	2 (18.2%)	9 (16.1%)	>0.999
Previous Tuberculosis	0 (0%)	2 (3.6%)	>0.999
Valvular heart disease	0 (0%)	5 (8.9%)	0.582
Previous VTE	0 (0%)	4 (7.1%)	>0.999
HIV	2 (18.2%)	17 (30.4%)	0.722

2.10 REFERENCES:

1. MBRRACE-UK Update: Key messages from the UK and Ireland Confidential Enquiries into Maternal Death and Morbidity 2017. *The Obstetrician & Gynaecologist*. 2018;20(1):75-79. doi:10.1111/tog.12466.
2. Meng K, Hu X, Peng X, et al. Incidence of venous thromboembolism during pregnancy and the puerperium: A systematic review and meta-analysis. *J Matern -Fetal Neonatal Med*. 2015;28(3). doi:10.3109/14767058.2014.913130.
3. Pomp ER, Lenselink AM, Rosendaal FR, et al. Pregnancy, the postpartum period and prothrombotic defects: Risk of venous thrombosis in the MEGA study. *J Thromb Haemost*. 2008;6(4). doi:10.1111/j.1538-7836.2008.02921.x.
4. James AH, Tapson VF, Goldhaber SZ. Thrombosis during pregnancy and the postpartum period. *Am J Obstet Gynecol*. 2005;193(1). doi:10.1016/j.ajog.2004.11.037.
5. Cantwell R, Clutton-Brock T, Cooper G, et al. Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer: 2006-2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG*. 2011;118 Suppl. doi:10.1111/j.1471-0528.2010.02847.x.
6. National Committee For Confidential Enquiry Into Maternal Deaths (NCCEMD) Saving Mothers Fact Sheet For 2022 Maternal Deaths And Mortality Rates 2022. Report, South Africa, 2022.
7. Jacobsen AF, Skjeldestad FE, Sandset PM. Incidence and risk patterns of venous thromboembolism in pregnancy and puerperium-a register-based case-control study. *Am J Obstet Gynecol*. 2008;198(2). doi:10.1016/j.ajog.2007.08.041.
8. Jordan EJ, Godelman A, Levsky JM, et al. CT pulmonary angiography in pregnant and postpartum women: Low yield, high dose. *Clin Imaging*. 2015;39(2). doi:10.1016/j.clinimag.2014.11.006.
9. Rotzinger DC, Dunet V, Ilic V, et al. Pulmonary embolism during pregnancy: a 17-year single-center retrospective MDCT pulmonary angiography study. *Eur Radiol*. 2020;30(3). doi:10.1007/s00330-019-06501-4.
10. Ehrlich R, Lowe S. An audit of CTPA and V/Q scan for investigation of pulmonary embolism in pregnancy. *Obstet Med*. Published online 2023. doi:10.1177/1753495X231197563.
11. Soma-Pillay P, Nelson-Piercy C, Tolppanen H, et al. Physiological changes in pregnancy. *Cardiovasc J Afr*. 2016;27(2). doi:10.5830/CVJA-2016-021.
12. Boeldt DS, Bird IM. Vascular adaptation in pregnancy and endothelial dysfunction in preeclampsia. *J Endocrinol*. 2017;232(1). doi:10.1530/JOE-16-0340.
13. Chatur S, Islam S, Moore LE, et al. Incidence of Syncope During Pregnancy: Temporal Trends and Outcomes. *J Am Heart Assoc*. 2019;8(10). doi:10.1161/JAHA.118.011608.
14. Bates SM, Greer A, Middeldorp S, et al. VTE, thrombophilia, antithrombotic therapy, and pregnancy - Antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2012;141(2 SUPPL.). doi:10.1378/chest.11-2300.
15. Wan T, Skeith L, Karovitch A, et al. Guidance for the diagnosis of pulmonary embolism during pregnancy: Consensus and controversies. *Thromb Res*. 2017;157. doi:10.1016/j.thromres.2017.06.025.
16. Chan WS, Rey E, Kent NE, et al. Venous Thromboembolism and Antithrombotic Therapy in Pregnancy. *J Obstet Gynaecol Can*. 2014;36(6). doi:10.1016/S1701-2163(15)30569-7.

17. Royal College Obstetricians and Gynaecologists. Thromboembolic disease in pregnancy and the puerperium: Acute Management. Green-top Guideline No. 37b. RCOG Press. Report, Royal College Obstetricians and Gynaecologists, UK, 2015.
18. Chan WS, Ray JG, Murray S, et al. Suspected Pulmonary Embolism in Pregnancy. *Arch Intern Med*. 2002;162(10). doi:10.1001/archinte.162.10.1170.
19. Leung AN, Bull TM, Jaeschke R, et al. An official American Thoracic Society/Society of Thoracic Radiology clinical practice guideline: Evaluation of suspected pulmonary embolism in pregnancy. *Am J Respir Crit Care Med*. 2011;184(10):1200-1208. doi:10.1164/rccm.201108-1575ST.
20. Chan WS, Chunilal S, Lee A, et al. A red blood cell agglutination D-dimer test to exclude deep venous thrombosis in pregnancy. *Ann Intern Med*. 2007;147(3). doi:10.7326/0003-4819-147-3-200708070-00005.
21. van der Pol LM, Tromeur C, Bistervels IM, et al. Pregnancy-Adapted YEARS Algorithm for Diagnosis of Suspected Pulmonary Embolism. *J Vasc Surg Venous Lymphat Disord*. 2019;7(5). doi:10.1016/j.jvsv.2019.06.004.
22. Potgieter R, Becker P, Suleman F. *S Afr J Radiol*. Published online 2022. doi:10.4102/sajr.
23. The Royal College of Radiologists. Appropriateness of Usage of Computed Tomography Pulmonary Angiography (CTPA) Investigation of Suspected Pulmonary Embolism Descriptor: Background. Accessed November 23, 2023. <https://www.rcr.ac.uk/career-development/audit-quality-improvement/auditlive-radiology/appropriateness-of-usage-of-computed-tomography-pulmonary-angiography-ctpa-investigation-of-suspected-pulmonary-embolism/>.
24. Shahir K, Goodman LR, Tali A, et al. Pulmonary embolism in pregnancy: CT pulmonary angiography versus perfusion scanning. *AJR Am J Roentgenol*. 2010;195(3). doi:10.2214/AJR.09.3506.
25. U-King-Im JM, Freeman SJ, Boylan T, et al. Quality of CT pulmonary angiography for suspected pulmonary embolus in pregnancy. *Eur Radiol*. 2008;18(12). doi:10.1007/s00330-008-1100-0.
26. Sun S, Diaconescu M, Zhe T, et al. Outcomes of Multidetector Computed Tomography Pulmonary Angiography in Pregnant and Postpartum Women With Suspected Pulmonary Embolism. *Can Assoc Radiol J*. 2021;72(3). doi:10.1177/0846537119899552.
27. Duffy J, Berger FH, Cheng I, et al. Implementation of the YEARS algorithm to optimise pulmonary embolism diagnostic workup in the emergency department. *BMJ Open Qual*. 2023;12(2). doi:10.1136/bmjopen-2022-002119.
28. Kline JA, Richardson DM, Than MP, et al. Systematic review and meta-analysis of pregnant patients investigated for suspected pulmonary embolism in the emergency department. *J Acad Emerg Med*. 2014;21(9). doi:10.1111/acem.12471.
29. Scott K, Rutherford N, Fagermo N, et al. Use of imaging for investigation of suspected pulmonary embolism during pregnancy and the postpartum period. *Obstet Med*. 2011;4(1):20-23. doi:10.1258/om.2010.100065.
30. Dado CD, Levinson AT, Bourjeily G. Pregnancy and Pulmonary Embolism. *Clin Chest Med*. 2018;39(3):525-537. doi:10.1016/j.ccm.2018.04.007.
31. Knight M, Bunch K, Felker A, et al. Saving Lives, Improving Mothers' Care Core Report – Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2019-21. Report, University of Oxford, UK, 2023.

32. Roth GA, Fleszar LG, Bryant AS, et al. Trends in State-Level Maternal Mortality by Racial and Ethnic Group in the United States. *JAMA*. 2023;330(1):52-61. doi:10.1001/jama.2023.9043.
33. Zakai NA, McClure LA. Racial differences in venous thromboembolism. *J Thromb Haemost*. 2011;9(10):1877-1882. doi:10.1111/j.1538-7836.2011.04443.x.
34. Phillips AR, Reitz KM, Myers S, et al. Association between black race, clinical severity, and management of acute pulmonary embolism: A retrospective cohort study. *J Am Heart Assoc*. 2021;10(17). doi:10.1161/JAHA.121.021818.
35. Solanki G, Fawcus S, Daviaud E. A cross sectional analytic study of modes of delivery and caesarean section rates in a private health insured South African population. *PLoS One*. 2019;14(6). doi:10.1371/journal.pone.0219020.
36. Solanki GC, Cornell JE, Daviaud E, et al. Caesarean section rates in South Africa: A case study of the health systems challenges for the proposed National Health Insurance. *S Afr Med J*. 2020;110(8):747-750. doi:10.7196/SAMJ.2020.v110i8.14699.
37. Gebhardt GS, Fawcus S, Moodley J, et al. Maternal death and caesarean section in South Africa: Results from the 2011 - 2013 saving mothers report of the national committee for confidential enquiries into maternal deaths. *S Afr Med J*. 2015;105(4):287-291. doi:10.7196/SAMJ.9351.
38. Blondon M, Casini A, Hoppe KK, et al. Risks of Venous Thromboembolism After Cesarean Sections: A Meta-Analysis. In: *Chest*. Vol 150. Elsevier B.V.; 2016:572-596. doi:10.1016/j.chest.2016.05.021.
39. Kovacevich GJ, Gaich SA, Lavin JP, et al. The prevalence of thromboembolic events among women with extended bed rest prescribed as part of the treatment for premature labor or preterm premature rupture of membranes. *Am J Obstet Gynecol*. 2000;182(5):1089-1092. doi:10.1067/mob.2000.105405.
40. Morrone D, Morone V. Acute pulmonary embolism: Focus on the clinical picture. *Korean Circ J*. 2018;48(5):365-381. doi:10.4070/kcj.2017.0314.
41. Kevane B, Donnelly J, D'Alton M, et al. Risk factors for pregnancy-associated venous thromboembolism: A review. *J Perinat Med*. 2014;42(4). doi:10.1515/jpm-2013-0207.
42. Bibas M, Biava G, Antinori A. HIV-associated venous thromboembolism. *Mediterr J Hematol Infect Dis*. 2011;3(1). doi:10.4084/MJHID.2011.030.
43. Di Prima FAF, Valenti O, Hyseni E, et al. Antiphospholipid Syndrome during pregnancy: the state of the art. *J Prenat Med*. 2011;5(2).
44. Blokhin IO, Lentz SR. Mechanisms of thrombosis in obesity. *Curr Opin Hematol*. 2013;20(5). doi:10.1097/MOH.0b013e3283634443.


APPENDIX A: ETHICAL CLEARANCE - HUMAN RESEARCH ETHICS COMMITTEE
(MEDICAL)



R14/49 Dr Wilhelm Herbst

HUMAN RESEARCH ETHICS COMMITTEE (MEDICAL)

CLEARANCE CERTIFICATE NO. M220255

NAME: Dr Wilhelm Herbst
(Principal Investigator)
DEPARTMENT: Internal Medicine
Charlotte Maxeke Johannesburg Academic Hospital
PROJECT TITLE: A Review of the Use of CT Pulmonary Angiography in
Pregnant and Postpartum Patients at an Academic Centre
DATE CONSIDERED: 25/02/2022
DECISION: Approved unconditionally
CONDITIONS:
SUPERVISOR: Dr J. Zamparini, Dr H. Moodley and Dr S. Bhoora
APPROVED BY: 
Dr CB Penny, Chairperson, HREC (Medical)
DATE OF APPROVAL: 20/05/2022

This clearance certificate is valid for 5 years from date of approval. Extension may be applied for.

DECLARATION OF INVESTIGATORS

To be completed in duplicate and **ONE COPY** returned to the Research Office Secretary on the Third Floor, Faculty of Health Sciences, Phillip Tobias Building, 29 Princess of Wales Terrace, Parktown, 2193, University of the Witwatersrand. I/we fully understand the conditions under which I am/we are authorized to carry out the above-mentioned research and I/we undertake to ensure compliance with these conditions. Should any departure be contemplated, from the research protocol as approved, I/we undertake to resubmit the application to the Committee. I **agree to submit a yearly progress report**. The date for annual re-certification will be one year after the date of convened meeting where the study was initially reviewed. In this case, the study was initially reviewed in **February** and will therefore be due in the month of **February** each year. Unreported changes to the application may invalidate the clearance given by the HREC (Medical).



Principal Investigator Signature

22/05/2022

Date

PLEASE QUOTE THE PROTOCOL NUMBER IN ALL ENQUIRIES

APPENDIX B: ETHICAL CLEARANCE - CHARLOTTE MAXEKE JOHANNESBURG ACADEMIC HOSPITAL



CHARLOTTE MAXEKE JOHANNESBURG ACADEMIC HOSPITAL UNIVERSITY OF THE WITWATERSRAND

Cluster Research Committee
7 Jubilee Rd, Parktown 2196, Johannesburg
Enquiries: Office of the CEO
Ms. T. Ndlovu; Tel: 011 488 4211 Fax 011 488 3753
Ms M. Khumalo Tel: 011 717 2536
Email: Mandisa.Khumalo@wits.ac.za

Dear Dr Wilhelm Herbst

The Charlotte Maxeke Johannesburg Academic Hospital (CMJAH) Research Cluster Committee has reviewed your request to research "A Review of the Use of CT Pulmonary Angiography in Pregnant and Postpartum Patient at an Academic Centre".

I am pleased to inform you that you have been granted to research Charlotte Maxeke Johannesburg Academic Hospital provided that;

1. Charlotte Maxeke Johannesburg Academic Hospital will not anyway incur or inherit costs as result of the said study.
2. Your study shall not disrupt services at the study sites.
3. Strict confidentiality shall be observed at all times.
4. Informed consent shall be solicited from patients participating in your study.

The data collection to be started only after obtaining the Human Research Ethics Committee (HREC) unconditional approval.

Kindly note that it is a requirement at CMJAH that for unconditional approval to be granted, all pending documents highlighted below are to be submitted as soon as they are available or within 3 months, and if not applicable, kindly elaborate. Failure to comply with the above-stated requirements may result in the withdrawal of this research's approval at CMJAH.

Head(s) of Department(s) Approval Letter	Provided
Human Research Ethics Committee Approval	Provided
The National Health Research Database Registration	Provided
Data Collection Sheets	Provided
Informed Consent Documents	N/A

Kindly forward this office with the results of your study on completion of the research.

I wish you every success with your research.

Yours sincerely,

Ms G. Bogoshi
Chief Executive Officer

CMJA Hospital Research Checklist

Requirement	Yes	No	N/A
HOD Approval Letter	Yes		
HREC Approval	Yes		
NHRD Approval	Yes		
Postgraduate Approval	Yes - Pending		
Co-investigators	Yes		
Collaborators	Yes		

Study team and collaborators

Organisation	Investigator	Qualifications	Contact details
Principal Investigator	Dr Wilhelm Herbst	MBChB (Pret); Dip HIV (CMSA)	wherbst.herbst@gmail.com; 079 885 9694
Study Leads			
Co-investigators	Dr Jarrod Zamparini	MBChB (Pret); FCP (SA); MMed Int Med (Wits)	jarrod.zamparini@wits.ac.za; 0722781158
Collaborators			

Manuscript Submission Guidelines:

Manuscript Submission Guidelines: *Obstetric Medicine*

This Journal is a member of the Committee on Publication Ethics.

This Journal recommends that authors follow the Recommendations for the Conduct, Reporting, Editing, and Publication of Scholarly Work in Medical Journals formulated by the International Committee of Medical Journal Editors (ICMJE).

Please read the guidelines below then visit the Journal's submission site <http://mc.manuscriptcentral.com/obmed> to upload your manuscript. Please note that manuscripts not conforming to these guidelines may be returned.

Only manuscripts of sufficient quality that meet the aims and scope of *Obstetric Medicine* will be reviewed.

There are no fees payable to submit or publish in this journal.

As part of the submission process you will be required to warrant that you are submitting your original work, that you have the rights in the work, and that you have obtained and can supply all necessary permissions for the reproduction of any copyright works not owned by you, that you are submitting the work for first publication in the Journal and that it is not being considered for publication elsewhere and has not already been published elsewhere. Please see our guidelines on prior publication and note that *Obstetric Medicine* will consider submissions of papers that have been posted on preprint servers; please alert the Editorial Office when submitting (contact details are at the end of these guidelines) and include the DOI for the preprint in the designated field in the manuscript submission system. Authors should not post an updated version of their paper on the preprint server while it is being peer reviewed for possible publication in the Journal. If the article is accepted for publication, the author may re-use their work according to the Journal's author archiving policy. If your paper is accepted, you must include a link on your preprint to the final version of your paper.

1. What do we publish?

1.1 Aims & Scope

1.2 Article types

- 1.3 Writing your paper
 - 1.4 Inclusive Language
- 2. Editorial policies
 - 2.1 Peer review policy
 - 2.2 Declarations
 - 2.3 Authorship
 - 2.4 Acknowledgements
 - 2.5 Funding
 - 2.6 Declaration of conflicting interests
 - 2.7 Research ethics and patient consent
 - 2.8 Clinical trials
 - 2.9 Reporting guidelines
 - 2.10 Data
- 3. Publishing policies
 - 3.1 Publication ethics
 - 3.2 Contributor's publishing agreement
 - 3.3 Open access and author archiving
- 4. Preparing your manuscript
 - 4.1 Formatting
 - 4.2 Artwork, figures and other graphics
 - 4.3 Supplementary material
 - 4.4 Reference style
 - 4.5 English language editing services
- 5. Submitting your manuscript
 - 5.1 ORCID
 - 5.2 Information required for completing your submission
 - 5.3 Permissions
- 6. On acceptance and publication
 - 6.1 Sage Production
 - 6.2 Online First publication
 - 6.3 Access to your published article
 - 6.4 Promoting your article
- 7. Further information

1. What do we publish?

Submissions may be made under the following categories:

1. Original research articles
2. Systematic reviews
3. Case reports
4. Letters to the Editor

Commissioned articles: The following may be published following invitation or agreement by the Editors, prior to submission.

1. Editorials
2. Topic Reviews
3. Journal Club

1.1 Aims & Scope

Before submitting your manuscript to *Obstetric Medicine*, please ensure you have read the Aims & Scope.

1.2 Article Types

When assessing the length of your contribution, allow 250 words for each table, figure or group of eight references, since all of these will contribute to its total length.

****All papers (apart from Letters to the Editor) should be accompanied by a covering letter, title page and summary****

Covering

letter

The covering letter is important. To help the Editors in their preliminary evaluation, please indicate why you think the paper suitable for publication. If your paper should be considered for fast-track publication, please explain why.

Title page: All the following must be completed before submission

The first page should contain the full title of the manuscript, a short title, the author(s) name(s) and affiliation(s), and the name, postal and email addresses of the author for correspondence, as well as a full list of declarations (Authorship, Ethical approval, Funding, Guarantor, Declaration of Interests, Acknowledgements) as well as appropriate Keywords.

The title should be concise and informative, accurately indicating the content of the article. The short title should be no more than six words long.

Summary/Abstract

A summary of **no more than 150 words** must accompany all submissions. The summary should state the main purposes of the study, the basic procedures used and the most important conclusions drawn; numerical data should be included. Original Research Articles should include a structured summary, namely: Background, Methods, Results, Conclusion. Letters to the Editor do not require an abstract.

Editorials

Editorials will be commissioned by the Editors. Maximum length: 1,000 words, with up to 10 references.

Review Articles

Suggestions for review articles are welcomed by the Editors and the subject matter should be outlined to the Editors, if possible before writing the article. Maximum length: 3,000 words, with up to five tables or illustrations.

Original Research Articles

These are expected to contribute to the advancement of knowledge in the field of maternal medicine. Maximum length: 3,000 words, with up to five tables or illustrations. Structured summaries are required, namely: Background, Methods, Results, Conclusion.

Case Reports

These should be prepared in a narrative style and comprise a summary; a short introduction stating the reasons for reporting the case; the case report including history, investigations and treatment; and a discussion referring to the relevant literature. Maximum length: 1,000 words, with up to two tables or two tables or illustrations.

Letters to the Editors

These should relate to articles published recently in the journal. Maximum length: 300 words plus up to five references.

1.3 Writing your paper

The Sage Author Gateway has some general advice and on how to get published, plus links to further resources. Sage Author Services also offers authors a variety of ways to improve and enhance their article including English language editing, plagiarism detection, and video abstract and infographic preparation.

1.3.1 Make your article discoverable

When writing up your paper, think about how you can make it discoverable. The title, keywords and abstract are key to ensuring readers find your article through search engines such as Google. For information and guidance on how best to title your article, write your abstract and select your keywords, have a look at this page on the Gateway: [How to Help Readers Find Your Article Online](#).

[Back to top](#)

1.4 Inclusive Language

In our journal articles we use the term ‘woman’ and ‘women’s health’. However it is important to acknowledge that it is not only people who identify as women for whom it is necessary to access women’s health and reproductive services in order to maintain their health and reproductive well-being.

Inclusive language and representation recognises diversity and conveys respect for all people, irrespective of gender. We would encourage all contributors to avoid non-inclusive terminology where appropriate, such as “he,” “she” or “he/she” and instead, seek gender neutrality by using plural nouns, such as “they”, “individuals”, “patients” or “subjects”. New data should ideally specify sex and gender, avoiding gender-neutral terminology when all participants are cis-gendered, to avoid creating a false sense of inclusivity. The editorial staff would be happy to provide guidance on appropriate terminology.

2. Editorial policies

2.1 Peer review policy

Sage does not permit the use of author-suggested (recommended) reviewers at any stage of the submission process, be that through the web-based submission system or other communication. Reviewers should be experts in their fields and should be able

to provide an objective assessment of the manuscript. Our policy is that reviewers should not be assigned to a paper if:

- The reviewer is based at the same institution as any of the co-authors
- The reviewer is based at the funding body of the paper
- The author has recommended the reviewer
- The reviewer has provided a personal (e.g. Gmail/Yahoo/Hotmail) email account and an institutional email account cannot be found after performing a basic Google search (name, department and institution).

2.2 Declarations

Please see: Obstetric Medicine Declarations Policy Document

The following statements should be uploaded under the file type 'Declarations' for every manuscript submitted. These will be published at the end of every paper accepted for publication.

Please note that consent forms should not be uploaded.

Please read the Declaration guidelines for authors (available here carefully before submitting your Declarations document.

DECLARATIONS

1. Conflicting interests (section 2.6)
2. Funding (section 2.5)
3. Informed consent (section 2.7)
4. Ethical approval (section 2.7)
5. Guarantor (section 2.10)
6. Contributorship (section 2.11)
7. Acknowledgements (section 2.4)

Please see the below example of a completed declarations section:

DECLARATIONS

Conflicting interests: MS is an employee of XXX. BF has received grants from XXX.

Funding: This work was supported by the Medical Research Council [grant number XXX]

Informed consent: Written informed consent was obtained from the patient(s) for their anonymized information to be published in this article.

Ethical approval: The ethics committee of XXXX approved this study (REC number: XXXX)

Guarantor: BF

Contributorship: BF and NP researched literature and conceived the study. MS was involved in protocol development, gaining ethical approval, patient recruitment and data analysis. BF wrote the first draft of the manuscript. All authors reviewed and edited the manuscript and approved the final version of the manuscript

Acknowledgements: We would like to thank XXX XXXX for his assistance and guidance in this research.

2.3 Authorship

Papers should only be submitted for consideration once consent is given by all contributing authors. Those submitting papers should carefully check that all those whose work contributed to the paper are acknowledged as contributing authors.

The list of authors should include all those who can legitimately claim authorship. This is all those who:

1. Made a substantial contribution to the concept or design of the work; or acquisition, analysis or interpretation of data,
2. Drafted the article or revised it critically for important intellectual content,
3. Approved the version to be published,
4. Each author should have participated sufficiently in the work to take public responsibility for appropriate portions of the content.

Authors should meet the conditions of all of the points above. When a large, multicentre group has conducted the work, the group should identify the individuals who accept direct responsibility for the manuscript. These individuals should fully meet the criteria for authorship.

Acquisition of funding, collection of data, or general supervision of the research group alone does not constitute authorship, although all contributors who do not meet the

criteria for authorship should be listed in the Acknowledgments section. Please refer to the International Committee of Medical Journal Editors (ICMJE) authorship guidelines for more information on authorship.

Please note that AI chatbots, for example ChatGPT, should not be listed as authors. For more information see the policy on Use of ChatGPT and generative AI tools.

2.4 Acknowledgements

All contributors who do not meet the criteria for authorship should be listed in an Acknowledgements section. Examples of those who might be acknowledged include a person who provided purely technical help, or a department chair who provided only general support.

2.4.1 *Third party submissions*

Where an individual who is not listed as an author submits a manuscript on behalf of the author(s), a statement must be included in the Acknowledgements section of the manuscript and in the accompanying cover letter. The statements must:

- Disclose this type of editorial assistance – including the individual's name, company and level of input
- Identify any entities that paid for this assistance
- Confirm that the listed authors have authorized the submission of their manuscript via third party and approved any statements or declarations, e.g. conflicting interests, funding, etc.

Where appropriate, Sage reserves the right to deny consideration to manuscripts submitted by a third party rather than by the authors themselves.

2.4.2 *Writing assistance*

Individuals who provided writing assistance, e.g. from a specialist communications company, do not qualify as authors and so should be included in the Acknowledgements section. Authors must disclose any writing assistance – including the individual's name, company and level of input – and identify the entity that paid for this assistance”).

It is not necessary to disclose use of language polishing services.

Any acknowledgements should appear first at the end of your article prior to your Declaration of Conflicting Interests (if applicable), any notes and your References.

2.5 Funding

Obstetric Medicine requires all authors to acknowledge their funding in a consistent fashion under a separate heading. Please visit the Funding Acknowledgements page on the Sage Journal Author Gateway to confirm the format of the acknowledgment text in the event of funding, or state that: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

2.6 Declaration of conflicting interests

It is the policy of *Obstetric Medicine* to require a declaration of conflicting interests from all authors enabling a statement to be carried within the paginated pages of all published articles.

Please ensure that a 'Declaration of Conflicting Interests' statement is included at the end of your manuscript, after any acknowledgements and prior to the references. If no conflict exists, please state that 'The Author(s) declare(s) that there is no conflict of interest'. For guidance on conflict of interest statements, please see the ICMJE recommendations [here](#).

2.7 Research ethics and patient consent

Medical research involving human subjects must be conducted according to the World Medical Association Declaration of Helsinki.

Submitted manuscripts should conform to the ICMJE Recommendations for the Conduct, Reporting, Editing, and Publication of Scholarly Work in Medical Journals, and all papers reporting animal and/or human studies must state in the methods section that the relevant Ethics Committee or Institutional Review Board provided (or waived) approval. Please ensure that you have provided the full name and institution of the review committee, in addition to the approval number.

For research articles, authors are also required to state in the methods section whether participants provided informed consent and whether the consent was written or verbal.

Information on informed consent to report individual cases or case series should be included in the manuscript text. A statement is required regarding whether written informed consent for patient information and images to be published was provided by the patient(s) or a legally authorized representative.

Please also refer to the ICMJE Recommendations for the Protection of Research Participants.

All research involving animals submitted for publication must be approved by an ethics committee with oversight of the facility in which the studies were conducted. The Journal has adopted the ARRIVE guidelines.

2.8 Clinical trials

Obstetric Medicine conforms to the ICMJE requirement that clinical trials are registered in a WHO-approved public trials registry at or before the time of first patient enrolment as a condition of consideration for publication. The trial registry name and URL, and registration number must be included at the end of the abstract.

2.9 Reporting guidelines

The relevant EQUATOR Network reporting guidelines should be followed depending on the type of study. For example, all randomized controlled trials submitted for publication should include a completed CONSORT flow chart as a cited figure and the completed CONSORT checklist should be uploaded with your submission as a supplementary file. Systematic reviews and meta-analyses should include the completed PRISMA flow chart as a cited figure and the completed PRISMA checklist should be uploaded with your submission as a supplementary file. The EQUATOR wizard can help you identify the appropriate guideline.

Other resources can be found at NLM's Research Reporting Guidelines and Initiatives.

2.10 Data

Sage acknowledges the importance of research data availability as an integral part of the research and verification process for academic journal articles.

Obstetric Medicine requests all authors submitting any primary data used in their research articles alongside their article submissions to be published in the online version of the journal, or provide detailed information in their articles on how the data can be obtained. This information should include links to third-party data repositories or detailed contact information for third-party data sources. Data available only on an author-maintained website will need to be loaded onto either the journal's platform or a third-party platform to ensure continuing accessibility. Examples of data types include but are not limited to statistical data files, replication code, text files, audio files, images, videos, appendices, and additional charts and graphs necessary to understand the original research. The editor may consider limited embargoes on proprietary data. The editor can also grant exceptions for data that cannot legally or ethically be released. All data submitted should comply with Institutional or Ethical Review Board requirements and applicable government regulations. For further information, please contact the editorial office at jovie.mcmillan@sagepub.co.uk

[Back to top](#)

3. Publishing Policies

3.1 Publication ethics

Sage is committed to upholding the integrity of the academic record. We encourage authors to refer to the Committee on Publication Ethics' International Standards for Authors and view the Publication Ethics page on the Sage Author Gateway.

3.1.1 Plagiarism

Obstetric Medicine and Sage take issues of copyright infringement, plagiarism or other breaches of best practice in publication very seriously. We seek to protect the rights of our authors and we always investigate claims of plagiarism or misuse of published articles. Equally, we seek to protect the reputation of the journal against malpractice. Submitted articles may be checked with duplication-checking software. Where an article, for example, is found to have plagiarised other work or included third-party copyright material without permission or with insufficient acknowledgement, or where

the authorship of the article is contested, we reserve the right to take action including, but not limited to: publishing an erratum or corrigendum (correction); retracting the article; taking up the matter with the head of department or dean of the author's institution and/or relevant academic bodies or societies; or taking appropriate legal action.

3.1.2 Prior publication

If material has been previously published it is not generally acceptable for publication in a Sage journal. However, there are certain circumstances where previously published material can be considered for publication. Please refer to the guidance on the Sage Author Gateway or if in doubt, contact the Editor at the address given below.

3.2 Contributor's publishing agreement

Before publication, Sage requires the author as the rights holder to sign a Journal Contributor's Publishing Agreement. Sage's Journal Contributor's Publishing Agreement is an exclusive licence agreement which means that the author retains copyright in the work but grants Sage the sole and exclusive right and licence to publish for the full legal term of copyright. Exceptions may exist where an assignment of copyright is required or preferred by a proprietor other than Sage. In this case copyright in the work will be assigned from the author to the society. For more information please visit the Sage Author Gateway.

3.3 Open access and author archiving

Obstetric Medicine offers optional open access publishing via the Sage Choice programme. For more information on Open Access publishing options at Sage please visit Sage Open Access. For information on funding body compliance, and depositing your article in repositories, please visit Sage's Author Archiving and Re-Use Guidelines and Publishing Policies.

[Back to top](#)

4. Preparing your manuscript for submission

4.1 Formatting

Preferred formats for the text and tables of your manuscript are Word DOC, RTF, XLS. LaTeX files are also accepted. The text should be double-spaced throughout and with a minimum of 3cm for left and right hand margins and 5cm at head and foot. Text should be standard 10 or 12 point. Word and (La)Tex templates are available on the Manuscript Submission Guidelines page of our Author Gateway.

Please see [here](#) for journal specific preferences for formatting your submission.

4.2 Artwork, figures and other graphics

For guidance on the preparation of illustrations, pictures and graphs in electronic format, please visit Sage's Manuscript Submission Guidelines.

Figures supplied in colour will appear in colour online regardless of whether or not these illustrations are reproduced in colour in the printed version. For specifically requested colour reproduction in print, you will receive information regarding the costs from Sage after receipt of your accepted article.

Tables

Tables must be prepared using the Table feature of the word processor. Tables should not duplicate information given in the text, should be numbered in the order in which they are mentioned in the text, and should be given a brief title.

Figures

All figures should be numbered in the order in which they are mentioned in the text. All figures must be accompanied by a figure legend. If figures are supplied in separate files, the figure legends must all be listed at the end of the main text file.

Line drawings should be produced electronically and clearly labelled using a sans serif font such as Arial. Graphs may be supplied as Excel spreadsheets (one per sheet). Other line drawings should be supplied in a suitable vector graphic file format (e.g. .eps)

All photographic images should be submitted in camera-ready form (i.e. with all extraneous areas removed), and where necessary, magnification should be shown using a scale marker. Photographic images must be supplied at high resolution, preferably 600 dpi. Images supplied at less than 300 dpi are unsuitable for print and will delay publication. The preferred file format is .tif.

4.3 Supplementary material

This journal is able to host additional materials online (e.g. datasets, podcasts, videos, images etc) alongside the full-text of the article. For more information please refer to our guidelines on submitting supplementary files.

4.4 Reference style

Obstetric Medicine adheres to the Sage Vancouver reference style. View the Sage Vancouver guidelines to ensure your manuscript conforms to this reference style.

If you use EndNote to manage references, you can download the the Sage Vancouver EndNote output file.

4.5 English language editing services

Authors seeking assistance with English language editing, translation, or figure and manuscript formatting to fit the journal's specifications should consider using Sage Language Services. Visit Sage Language Services on our Journal Author Gateway for further information.

Back to top

5. Submitting your manuscript

Obstetric Medicine is hosted on Sage Track, a web based online submission and peer review system powered by ScholarOne™ Manuscripts. Visit <http://mc.manuscriptcentral.com/obmed> to login and submit your article online.

IMPORTANT: Please check whether you already have an account in the system before trying to create a new one. If you have reviewed or authored for the journal in the past year it is likely that you will have had an account created. For further guidance on submitting your manuscript online please visit ScholarOne Online Help.

If you encounter any difficulties during the online submission process, please contact ObMededitorial@sagepub.co.uk for further assistance

5.1 ORCID

As part of our commitment to ensuring an ethical, transparent and fair peer review process Sage is a supporting member of ORCID, the Open Researcher and Contributor ID. ORCID provides a unique and persistent digital identifier that distinguishes researchers from every other researcher, even those who share the same name, and, through integration in key research workflows such as manuscript and grant submission, supports automated linkages between researchers and their professional activities, ensuring that their work is recognized.

The collection of ORCID iDs from corresponding authors is now part of the submission process of this journal. If you already have an ORCID ID you will be asked to associate that to your submission during the online submission process. We also strongly encourage all co-authors to link their ORCID ID to their accounts in our online peer review platforms. It takes seconds to do: click the link when prompted, sign into your ORCID account and our systems are automatically updated. Your ORCID iD will become part of your accepted publication's metadata, making your work attributable to you and only you. Your ORCID iD is published with your article so that fellow researchers reading your work can link to your ORCID profile and from there link to your other publications.

If you do not already have an ORCID iD please follow this link to create one or visit our ORCID homepage to learn more.

5.2 Information required for completing your submission

You will be asked to provide contact details and academic affiliations for all co-authors via the submission system and identify who is to be the corresponding author. These details must match what appears on your manuscript. At this stage please ensure you have included all the required statements and declarations and uploaded any additional supplementary files (including reporting guidelines where relevant).

5.3 Permissions

Please also ensure that you have obtained any necessary permission from copyright holders for reproducing any illustrations, tables, figures or lengthy quotations previously published elsewhere. For further information including guidance on fair dealing for criticism and review, please see the Copyright and Permissions page on the Sage Author Gateway.

[Back to top](#)

6. On acceptance and publication

6.1 Sage Production

Your Sage Production Editor will keep you informed as to your article's progress throughout the production process. Proofs will be sent by PDF to the corresponding author and should be returned promptly. Authors are reminded to check their proofs carefully to confirm that all author information, including names, affiliations, sequence and contact details are correct, and that Funding and Conflict of Interest statements, if any, are accurate.

6.2 Online First publication

Online First allows final articles (completed and approved articles awaiting assignment to a future issue) to be published online prior to their inclusion in a journal issue, which significantly reduces the lead time between submission and publication. Visit the Sage Journals help page for more details, including how to cite Online First articles.

6.3 Access to your published article

Sage provides authors with online access to their final article.

6.4 Promoting your article

Publication is not the end of the process! You can help disseminate your paper and ensure it is as widely read and cited as possible. The Sage Author Gateway has numerous resources to help you promote your work. Visit the Promote Your Article page on the Gateway for tips and advice.

[Back to top](#)

7. Further information

Any correspondence, queries or additional requests for information on the manuscript submission process should be sent to the *Obstetric Medicine* editorial office as follows:

Emily Kent

Publishing Editor

Sage Publications Ltd

1 Oliver's Yard, 55 City Road

London, EC1Y 1SP, UK

emily.kent@sagepub.co.uk

Similarity draft.pdf

by Wilhelm Herbst

Submission date: 18-Dec-2023 03:10PM (UTC+0200)

Submission ID: 2261876683

File name: Similarity_draft.pdf (400.93K)

Word count: 3929

Character count: 19036

2.2 ABSTRACT

Background: ⁴ Pulmonary embolism (PE), is a common cause of maternal death during pregnancy and the puerperium yet data on its prevalence in middle-income countries is lacking.

Methods: We examined the medical records and CTPAs of 67 women in an obstetric high care unit during pregnancy and the puerperium. ¹ We aimed to assess the efficacy of CTPA in diagnosing PE, evaluate associated clinical features, and determine the prevalence of alternative CT findings.

Results: CTPA detected PE in 11 women (16.42%) and alternative CT findings in 46 (68.6%). Women with PE had a significantly lower systolic BP than those without PE ($p = 0.001$). Multiple gestation, preterm rupture of membranes, and gestational diabetes were linked to higher PE prevalence.

Conclusions: This study, set in an upper middle-income country, demonstrated a higher CTPA yield for PE than in the international literature, emphasizing context-specific assessments.

2.3 ⁴ INTRODUCTION

Venous thromboembolism (VTE), including pulmonary embolism (PE), is the most common cause of maternal death during pregnancy and the puerperium in developed countries.¹ The risk for venous thromboembolism (VTE) is significantly increased during pregnancy and the postpartum period, as these patients are in a physiological state of hypercoagulability, are prone to venous stasis and may have superimposed endothelial damage.²

Women have a 5-fold increased risk of developing VTE during pregnancy, as compared to their non-pregnant counterparts.³ Globally, the incidence of PE in pregnant or postpartum women ranges from 1-2 per 1,000 to 3 per 10,000 deliveries^{2,4}, and one death in every 100,000 deliveries.⁵ In South Africa PE accounted for 40 maternal deaths in 2022, with a maternal mortality rate of 4.1 per 100,000 live births.⁶

More than half of VTE cases in pregnancy occur before 20 weeks' gestation,⁴ and 80% of VTE cases in the postpartum period occur within the first 3 weeks following delivery.⁷

Despite the increased risk of VTE in pregnancy, detection rates using CTPA (computed tomography pulmonary angiography) are lower in pregnant than non-pregnant women. A 2015 study found that 2% of pregnant women had a positive CTPA compared to 16% of non-pregnant women.⁸ A 17-year retrospective study conducted at a single centre in Switzerland, published in 2020, observed that CTPA rarely indicated PE in pregnancy, with an alternative diagnosis found in 30% of pregnant women investigated.⁹ Similarly, a 2023 study including 86 pregnant and 354 non-pregnant women investigated for PE by either CTPA or V/Q scan, noted positive scan rates of 3.5% and 8.8% in pregnant and non-pregnant women respectively.¹⁰

Risk Factors

Pregnancy is a hypercoagulable physiological state owing to an increased concentration of certain clotting factors as well as a decrease in endogenous anticoagulants such as antithrombin and protein S¹¹. In addition, factors such as dehydration, secondary to nausea and vomiting in early pregnancy, immobility and iliac compression by the gravid uterus in late pregnancy, and endothelial injury from

delivery trauma and venous distension^{11,12} also play a role in the increased risk for VTE. The clinical presentation of VTE may also be masked by “symptoms” of pregnancy such as pedal oedema, tachypnoea and tachycardia.¹³ A number of postpartum factors are also associated with an increased risk of VTE, such as surgical delivery, particularly in the emergency setting or after prolonged labour, postpartum haemorrhage, thrombophilia, pre-eclampsia, and puerperal sepsis.¹⁴

Clinical predictors

The Wells’ criteria, which factors in tachycardia and lower limb oedema, is not particularly useful in pregnancy, given that these manifestations are frequently seen.¹⁵ This complicates the differentiation of PE from other potential diagnoses in these women.¹⁵ The recommendations outlined by the Society of Obstetricians and Gynaecologists of Canada (SOGC)¹⁶ and the Royal College of Obstetricians and Gynaecologists (RCOG)¹⁷, advise against clinical predictors to assess VTE in pregnant women.

D-dimer

In non-pregnant women PE can be safely excluded in the presence of a normal D-dimer in addition to a low pre-test probability. It must conversely be interpreted with caution in those with a high pre-test probability, as a falsely negative result can miss a diagnosis of PE.¹⁸ The American Thoracic Society (ATS)¹⁹, SOGC¹⁶, and RCOG¹⁷ recommendations state that a D-dimer should not be used in pregnancy to exclude PE as most studies investigating the use of D-dimer in diagnosing PE, have not included pregnant women as participants.²⁰

⁶ Van der Pol *et al* found that the pregnancy-adapted YEARS algorithm safely excluded acute PE in pregnant women with suspected PE, allowing CTPA to be avoided in 32 to 65% of women.²¹ A South African study found that the pregnancy-adapted YEARS algorithm successfully risk-stratified women in a single institution and would have reduced the number of CTPAs by 25.7% without missing any PE.²²

CT Pulmonary Angiography

Accepted standards outlined by the Royal College of Radiologists state that CTPA should detect PE in 15.4 - 37.4% and alternative diagnoses in up to 56% in the general population.²³

Most studies have disregarded pregnant women when investigating the diagnosis of PE.¹⁸ Certain physiological factors, such as increased circulatory volume and altered cardiac output, may lead to lower pulmonary artery opacification influencing the quality of CTPA.²⁴ Insufficient opacification of the pulmonary vasculature during CTPA is described twice as often among pregnant women compared to those who are not pregnant.²⁵ Normal CTPA imaging in pregnancy, has been found to have a high negative predictive value.²⁶

In this study we investigated ¹ the use of CTPA to diagnose PE in pregnant and postpartum ¹ women admitted to a high care setting in South Africa. We aimed to evaluate ¹ the prevalence of PE and alternative diagnoses as well as compare the clinical features of women with and without PE.

2.4 METHODS

We performed a cross-sectional analysis ¹¹ at the Charlotte Maxeke Johannesburg Academic Hospital, an academic centre in Johannesburg, South Africa. Pregnant and postpartum women admitted to the Obstetric High Care Unit (OHCU) between 1 January and 31 December 2019, who underwent CTPA for the suspicion of PE were consecutively enrolled in this study. We excluded women who had a PE diagnosed by a method other than CTPA, such as V/Q scan, due to limited availability in our setting. Women with imaging performed elsewhere, women with known PE diagnosed prior to admission, and women who were not pregnant or beyond 6 weeks post-delivery were also excluded.

The names and hospital numbers of all women admitted to the OHCU for the study period were collected from the ward admission register and cross-referenced with the hospital picture archiving and communicating system (PACS) to determine which women underwent CTPA. The included participant details were anonymised and

allocated a unique study number. The CTPA images were reported on by two consultant radiologists (AR and TWT) who were blinded from each other's findings. The discrepant readings were reviewed by a third consultant radiologist (HM), also blinded from the findings of the other readers.

A final diagnosis was considered as agreement by two or more readers. Those studies with inconclusive findings or failing to achieve consensus among at least two readers, were interpreted as not having PE for the purposes of the study. The hospital files for the relevant participants were collected from hospital archives and clinical data was extracted. Data was captured in Microsoft Excel 16.78 (Microsoft Corporation, Redmond, USA) and statistical analysis was performed using Prism 10 (GraphPad Software Inc, La Jolla, CA, USA). Given the non-parametric nature of the data, the Mann-Whitney U test was employed for continuous variables, while the Fisher's exact test was utilized for categorical variables to determine significance.

2.5 RESULTS

A total of 732 women were admitted during the period under review, CTPAs were performed on 71 patients, of which 3 were not pregnant nor in the postpartum period and 1 had missing images, excluding them from the study, leaving a total of 67 women in the final analysis (Figure 1). The PE and non-PE groups were of similar age and race. The majority of the cohort were postpartum (82% in the PE group and 87.5% in the no PE group), and caesarean section was the most common mode of delivery in keeping with local trends. Most of the women were booked for antenatal care, and were multiparous. Table 1 provides demographic and pregnancy related data of the cohort.

Eleven women (16.4%) had a confirmed PE; 2 (18%) were pregnant and 9 (82%) postpartum and the majority (n=10, 90.9%) had peripheral PEs. Two women (18.2%) had a single embolism and 9 (81.8%) had multiple emboli.. The gestational age at delivery (in postpartum women) or time of scan (in pregnant women) was not significantly different ($p = 0.259$) in the non-PE group (median 34.5 weeks, IQR 30 – 38), compared to those with PE (median 30 weeks, IQR 26 – 38). Women with PE were more likely to have multiple gestation (27%) than without PE (4%) ($p = 0.028$).

There was a larger proportion of women with alternative CT findings among those in the group with PE (n=9, 81.8%) compared to those without PE (n=37, 66.1%), yet the difference was not significant (p=0.481) (Table 2).

The difference in the technical adequacy of the CTPAs ³ between the two groups was not significant (p = 0.556). This was assessed by measuring the opacification of the pulmonary artery (Hounsfield units), between the two groups (median 237HU, IQR 216-277 versus ¹² median 247HU, IQR 209-285 in the PE and non-PE groups respectively). ¹² There was no difference in rate of motion artifact between the two groups (p = 0.492).

There was a ³ significant difference in systolic blood pressure (sBP) between the two groups (p = 0.001) with lower sBP in the PE group (median 119mmHg, IQR 110 – 124) than in the non-PE group (median 134mmHg, IQR 120.3 - 146). There were no differences in other clinical features including the ¹⁰ body mass index (BMI) or mid upper arm circumference (MUAC) between the two groups. Despite most women being tachycardic, there were no differences in electrocardiogram (ECG) findings between the two groups. It is not our standard protocol to analyse D-dimer routinely for the suspicion of PE in pregnancy, therefore only 27 women (40%) had D-dimers available. There was however no difference in D-dimers between the two groups (p = 0.219).

Biochemical inflammatory markers were available in a similar proportion of women, showing no difference between the groups, reflected by a median C-reactive protein (CRP) of 112mg/L in the PE group (10/11) and 100mg/L in the non-PE group (51/56) and a median procalcitonin (PCT) of 0.37ug/L in the PE group (8/11) and 0.31ug/L in the non-PE group. A diagnosis of puerperal sepsis was made in 36.4% in the PE group compared to 62.5% in the non-PE group, however the difference was not significant (p = 0.18). There were no other differences demonstrated in biochemistry between the two groups.

No women died in the non-PE group, while one woman died in the PE group (9%), however it is unclear if the death was directly caused by PE. Most women had a pregnancy-related complication, with the exception of 2 women (3.6%) in the non-PE group.

There were 2 women with HIV (18.2%) in the PE group, one of whom had a CD4 (CD4 = 464 cells/mm³) and HIV viral load (VL = 764 copies/ml) available, whilst there were 17 women with HIV in the non-PE group (30.4%) where 15 had a CD4 result (median 344.5 cells/mm³, IQR 269-457.75) and all had a viral load (n = 17, median 217 copies/ml, IQR 0 – 2472.25).

Gestational diabetes was significantly more prevalent (p = 0.042) among those with PE (n = 1, 9.1%) compared to those without PE (n = 2, 3.6%), as was premature rupture of membranes (PROM) (p = 0.025) comparing the PE group (n = 2, 18.2%) to the non-PE group (n = 0, 0%). There were no other complications or comorbidities with a significant difference between the groups (Table 3).

The Wells' score showed no correlation to PE, as all 67 (100%) women had a Wells' score of at least 4.5 (high risk for PE). When analysing the utility of the YEARS score²⁷, 7 women in the PE group and 20 women in the non-PE group had a D-dimer result available allowing the YEARS Score to be calculated. All 7 (100%) women in the PE group had a positive YEARS score, while 17 (85%) women in the non-PE group had a positive YEARS score. The difference was not significant (p = 0.545) (Figure 2).

2.6 DISCUSSION

In our study CTPA detected PE in 16.42% and alternative CT findings in 68.6% of our cohort, which is higher than reported in international literature. A large meta-analysis and systematic review of seventeen studies, including 25,339 women, found a 12.4% positivity rate for VTE in nonpregnant women, compared with 4.1% in pregnant women.²⁸ The high positivity rate in our cohort may be attributed to only including women admitted to a high care unit, assigning them to a higher risk category than other pregnant and postpartum women.

No clinical or biochemical biomarkers predicted PE in our cohort, similar to an Australian study evaluating the use of CTPA and V/Q in pregnancy and the puerperium.²⁹ The use of algorithms such as the modified YEARS and Wells' scores were limited in our cohort and would have been of no utility in reducing the number of

CT scans done. However given our limited sample size, future studies are needed to further assess their utility in our setting. This is in keeping with recommendations from RCOG to avoid the use of clinical prediction tools in assessing VTE in pregnancy.¹⁷

Despite international guidelines considering maternal age over 35 years a risk factor for PE¹⁵ there was no significant difference in the age of the PE and the non-PE groups in our cohort.³⁰ There is evidence from the UK³¹ and USA³² that Black, Asian and minority ethnicities (BAME) have a much higher maternal mortality rate (MMR), with the highest among Black women. There is evidence that PE is more prevalent in black ethnicities,³³ with a higher disease severity compared to other ethnicities.³⁴ The majority of our cohort were of Black ethnicity, making it difficult to compare outcomes related to ethnicity and to assess whether disparities exist.. This emphasises the importance of the high risk in this group, and the need for further research in a South African context

Caesarean delivery rate in the South African private sector was 73.2% in 2015³⁵, and increased from 15.1% in 2006³⁶ to 31.1% in 2022⁶ in the public sector. Caesarean delivery is not done on maternal request in the South African public sector. Compared to the UK, the rate of caesarean delivery was 29%³¹ in Wales, 31%³¹ in England and 35%³¹ in Scotland between 2019 - 2021. South African data shows that pregnant women who die from PE are more likely to have had a caesarean delivery.³⁷ In our cohort all women in the PE group were delivered by caesarean section, although this was for obstetric indications and not due to PE. While Caesarean delivery is a known risk factor for the development of PE³⁸, this was not demonstrated in our cohort and mode of delivery was not a risk factor for PE.

The most significant obstetric complication in our cohort was PROM with a higher prevalence in those with PE (n = 2, 18.2%) compared to those without PE (n = 0, 0%) (p = 0.025). Both women with PROM in the PE group were HIV positive, developed puerperal sepsis and had undergone caesarean delivery by gestational age of 28 and 35 weeks respectively. Only one of these women had a CD4 (CD4 = 464 cells/mm³) and HIV viral load (VL = 764 copies/ml) available, and neither had a history of previous tuberculosis. PE was diagnosed after delivery in both women. The development of PE in these women was likely multifactorial however prolonged bed rest prescribed for the

2 treatment of PROM was associated with an increased risk for PE in a study by Kovacevich *et al* (16.5 cases per 1000 compared to 0.8 per 1000 among pregnant women for whom this therapy was not prescribed).³⁹

3 Systolic BP was significantly lower ($p = 0.001$) in the PE group, as was seen in the DiPEP study, however there were no differences in diastolic BP and heart rate. Out of keeping with the lower systolic BP in the PE group is that majority of these were peripheral PEs. However 81.8% of the women with PE had multiple emboli which may have contributed to the lower systolic BP. Hypotension associated with PE is attributed to acute right ventricular dilatation compromising left ventricular filling and reducing cardiac output. Right ventricular dysfunction seen on echocardiography has been shown to correlate with pulmonary arterial obstruction of >30%, which is expected to occur with a large central PE⁴⁰. Echocardiography was not available to assess for right ventricular strain, however there were no ECG features to suggest this.

The high prevalence of alternative findings on CTPA may be explained by the hospital's protocol for investigating PE, where CTPA is warranted if the chest X-ray is abnormal. Despite there being no significant difference in prevalence of alternative CT findings between women with PE and women without PE, the high burden of alternative findings on CTPA in women with PE could signify underlying pathology contributing to development of PE. The high prevalence of pregnancy related complications in our cohort is due to it consisting only of women in a high care setting.

Multiple gestation was a risk factor for PE in our cohort ($p = 0.028$) in keeping with guidelines from RCOG¹⁷. Gestational diabetes has not been proven to be associated with an increased risk for thrombosis⁴¹, however was associated with PE in our cohort ($p = 0.042$). Despite HIV⁴², antiphospholipid syndrome⁴³, and obesity⁴⁴ known to be associated with increased risk for PE, none were significant risk factors in our cohort.

The strengths of this analysis are driven from the racial representation in our cohort, largely consisting of Black African women, an underrepresented group in other studies, which underpins the need and potential for further research. Additionally, the data investigated were largely complete, all the CTPAs in our cohort contained adequate quality images, and were reviewed by consultant radiologists. Yet, the findings of this

analysis must be interpreted in the context of some limitations. The first being, the small sample size. Considering our study was performed at a single centre including only women admitted to a high care setting, this in itself might be a major contributory factor to the high PE burden in our population, as these women were all in a higher risk category by nature of their condition. Another limitation is that chest radiographs were not reviewed, as this could have contributed to analysing the pre-test probability for PE.

2.7 CONCLUSION

PE remains a significant contributor to morbidity during pregnancy and the puerperium. While clinical and laboratory features were not reliable to exclude PE in our cohort, it is justifiable to consider CTPA when suspecting PE in this population, considering the higher detection rate compared to international literature. CTPA has a good yield for diagnosing PE and unmasking alternative diagnoses in a middle-income country setting. The presence of a low systolic blood pressure, multiple gestation, PROM, and gestational diabetes should raise the suspicion for PE in a high care setting. Prospective studies evaluating the use of d-dimer and scoring systems to predict PE in our population are warranted.

2.8 TABLES AND FIGURES:




Figure 1: Study population




Figure 2: D-dimer values

Table 1: Demographics and pregnancy details

Demographics	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
Age (years), median, (IQR)	29 (26 – 35)	30 (26 – 33,75)	0.977
18 – 25	2 (18.18%)	12 (21.43%)	0.992
26 - 30	4 (36.36%)	18 (32.14%)	
31 - 35	3 (27.27%)	16 (28.57%)	
>35	2 (18.18%)	10 (17.86%)	
Race	PE, n (%)	No PE, n (%)	p-value
Black African	11 (100%)	55 (98%)	>0.999
White	0 (0%)	1 (2%)	
Pregnancy Details	PE, n (%)	No PE, n (%)	p-value
Postpartum	9 (82%)	49 (87.5%)	0.635
Pregnant	2 (18%)	7 (12.5%)	
Multiple gestation	3 (27%)	2 (4%)	0.028
Booked for antenatal care	8 (73%)	45 (80%)	0.686
Not booked for antenatal care	3 (27%)	11 (20%)	
Gestational age at delivery or time of scan (weeks), median (IQR)	30 (26 – 38)	34,5 (30 – 38)	0.259
Caesarean section	9 (82%)	41 (73.2%)	0.334
Normal vaginal delivery	0 (0%)	8 (14.3%)	
Gravidity (number of pregnancies), median (IQR)	3 (1 – 3)	2 (1 – 3)	0.971
Parity (number of live births), median (IQR)	2 (1 – 3)	2 (1 – 3)	0.714
Previous miscarriages (number), median (IQR)	0 (0 – 0)	0 (0 – 0)	0.841

Table 2: Alternative CT findings

CT findings	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
Pleural effusion	4 (36.4%)	14 (25%)	0.469
Ground glass opacification	6 (54.5%)	16 (28.6%)	0.157
Consolidation	3 (27.3%)	11 (19.6%)	0.686
Cavitation	0 (0%)	1 (1.8%)	>0.999
Bronchiectasis	0 (0%)	2 (3.6%)	>0.999
Atelectasis	8 (72.7%)	24 (42.9%)	0.101
Nodules	0 (0%)	5 (8.9%)	0.582

Table 3: YEARS score

YEARS score	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
YEARS positive	7 (100%)	17 (85%)	0.545
YEARS negative	0 (0%)	3 (15%)	
YEARS not calculated	4	36	

Table 4: Pregnancy complications and comorbidities

Pregnancy complications	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
No complications	0 (0%)	2 (3.6%)	>0,999
Obstetric Haemorrhage	5 (45,5%)	19 (33.9%)	0,505
Ante-partum haemorrhage	2 (18.2%)	6 (10.7%)	0,609
Post-partum haemorrhage	3 (27.3%)	13 (23.2%)	0,7159
Hypertensive Disorders	7 (63.6%)	42 (75%)	0,469
Pregnancy induced hypertension	1 (9.1%)	12 (21.4%)	0,677
Pre-Eclampsia	3 (27.3%)	18 (32.1%)	>0,999
Eclampsia	1 (9.1%)	9 (16.1%)	>0,999
HELLP	2 (18.2%)	3 (5.4%)	0,186
Obstetric Complications	4 (36.36%)	6 (10.71%)	0,051
Placenta praevia	1 (9.1%)	2 (3.6%)	0,421
Preterm labour	0 (0%)	2 (3.6%)	>0,999
Other	10 (90.1%)	42 (75%)	0,4329
Post-partum cardiomyopathy	2 (18.2%)	13 (23.2%)	>0,999
Anaemia	8 (72.7%)	29 (51.8%)	0,321
Relook laparotomy	3 (27.2%)	8 (14.3%)	0,371
ICU	5 (45.5%)	14 (25%)	0,27
Comorbidities	PE (n = 11), n (%)	No PE (n = 56), n (%)	p-value
Autoimmune disease	1 (9.1%)	3 (5.4%)	0.521
Anti-phospholipid syndrome	0 (0%)	2 (3.6%)	>0.999
Chronic kidney disease	0 (0%)	4 (7.1%)	>0.999
Diabetes mellitus	1 (9.1%)	1 (1.8%)	0.304
Heart failure	0 (0%)	8 (14.3%)	0.335
Hypertension	2 (18.2%)	9 (16.1%)	>0.999
Previous Tuberculosis	0 (0%)	2 (3.6%)	>0.999
Valvular heart disease	0 (0%)	5 (8.9%)	0.582
Previous VTE	0 (0%)	4 (7.1%)	>0.999
HIV	2 (18.2%)	17 (30.4%)	0.722

Similarity draft.pdf

ORIGINALITY REPORT

7%

SIMILARITY INDEX

4%

INTERNET SOURCES

4%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

David C. Rotzinger, Vincent Dunet, Vesna Ilic, Olivier W. Hugli, Reto A. Meuli, Sabine Schmidt. "Pulmonary embolism during pregnancy: a 17-year single-center retrospective MDCT pulmonary angiography study", European Radiology, 2019

Publication

1%

2

"Abstracts from the 30th Annual Meeting of the Society of General Internal Medicine", Journal of General Internal Medicine, 2007

Publication

1%

3

Hathwar, Pranam. "A Prospective, Double Blind Comparative Clinical Study of Effect of 0.5% and 0.75% Isobaric Ropivacaine for Spinal Anaesthesia in Patients Undergoing Elective Lower Limb Orthopaedic Surgeries", Rajiv Gandhi University of Health Sciences (India), 2023

Publication

1%

4

[ashpublications.org](https://www.ashpublications.org)

Internet Source

1%

5	Groesbeck P Parham, Mulindi H Mwanahamuntu, Vikrant V Sahasrabuddhe, Andrew O Westfall et al. "Implementation of cervical cancer prevention services for HIV-infected women in Zambia: measuring program effectiveness", HIV Therapy, 2010 Publication	<1 %
6	www.one.be Internet Source	<1 %
7	www.sid.ir Internet Source	<1 %
8	www.mdpi.com Internet Source	<1 %
9	Submitted to Drexel University Student Paper	<1 %
10	journals.plos.org Internet Source	<1 %
11	Nqoba Tsabedze, Dineo Mpanya, Claude Bailly, Samantha Nel et al. "Clinical characteristics and one-year all-cause mortality outcomes in Africans with dilated cardiomyopathy", International Journal of Cardiology, 2023 Publication	<1 %
12	Ho Young Hwang, Kyung-Hwan Kim, Ki-Bong Kim, Hyuk Ahn. "Propensity Score Matching	<1 %

Analysis of Mechanical Versus Bioprosthetic Tricuspid Valve Replacements", The Annals of Thoracic Surgery, 2014

Publication

13

open.library.ubc.ca

Internet Source

<1 %

14

pure.uva.nl

Internet Source

<1 %

15

www.researchgate.net

Internet Source

<1 %

16

coek.info

Internet Source

<1 %

Exclude quotes On

Exclude matches < 8 words

Exclude bibliography On