

6. REFERENCES

1. UNAIDS. December 2005. AIDS Epidemic Update 2005. Available from: http://www.unaids.org/epi/2005/doc/EPIupdate2005_pdf_en/epi-update2005_en.pdf [Accessed 17 February 2006].
2. Department of Health. National HIV and syphilis sero-prevalence survey in South Africa 2004. Pretoria, South Africa: Department of Health; 2005.
3. Centers for Disease Control. 1994 Revised Classification System for Human Immunodeficiency Virus Infection in Children Less than 13 Years of Age. MMWR. 1994;43(RR-13):1-10.
4. Consensus Workshop. Early diagnosis of HIV infection in infants: Siena, Italy. J Acquir Immune Defic Syndr. 1992;5(11):1169-78.
5. Borkowsky W, Krasinski K, Pollack H, Hoover W, Kaul A, Ilmet-Moore T. Early diagnosis of human immunodeficiency virus infection in children less than 6 months of age: comparison of polymerase chain reaction, culture, and plasma antigen capture techniques. J Infect Dis. 1992 Sep;166(3):616-9.
6. De Rossi A, Amadori A, Chieco-Bianchi L, Giacchino C, Zacchello F, Buchbinder A, et al. Polymerase chain reaction and in-vitro antibody production for early diagnosis of paediatric HIV infection. Lancet. 1988 Jul 30;2(8605):278.
7. De Rossi A, Ades AE, Mammano F, Del Mistro A, Amadori A, Giacchino C, et al. Antigen detection, virus culture, polymerase chain reaction, and in vitro antibody production in the diagnosis of vertically transmitted HIV-1 infection. Aids. 1991 Jan;5(1):15-20.
8. Laure F, Courgnaud V, Rouzioux C, Blanche S, Veber F, Burgard M, et al. Detection of HIV1 DNA in infants and children by means of the polymerase chain reaction. Lancet. 1988 Sep 3;2(8610):538-41.

9. Young KK, Peter JB, Winters RE. Detection of HIV DNA in peripheral blood by the polymerase chain reaction: a study of clinical applicability and performance. *Aids*. 1990 May;4(5):389-91.
10. Krivine A, Yakudima A, Le May M, Pena-Cruz V, Huang AS, McIntosh K. A comparative study of virus isolation, polymerase chain reaction, and antigen detection in children of mothers infected with human immunodeficiency virus. *J Pediatr*. 1990 Mar;116(3):372-6.
11. Petru A, Dunphy MG, Azimi P, Janner D, Gallo D, Hanson C, et al. Reliability of polymerase chain reaction in the detection of human immunodeficiency virus infection in children. *Pediatr Infect Dis J*. 1992 Jan;11(1):30-3.
12. Rogers MF, Ou CY, Rayfield M, Thomas PA, Schoenbaum EE, Abrams E, et al. Use of the polymerase chain reaction for early detection of the proviral sequences of human immunodeficiency virus in infants born to seropositive mothers. New York City Collaborative Study of Maternal HIV Transmission and Montefiore Medical Center HIV Perinatal Transmission Study Group. *N Engl J Med*. 1989 Jun 22;320(25):1649-54.
13. Poussin K, Harter-Chaput A, Sayada C, Brechot C. Evaluation of the Amplicor HIV1 DNA diagnosis test. *Res Virol*. 1994 Nov-Dec;145(6):393-5.
14. Kovacs A, Xu J, Rasheed S, Li XL, Kogan T, Lee M, et al. Comparison of a rapid nonisotopic polymerase chain reaction assay with four commonly used methods for the early diagnosis of human immunodeficiency virus type 1 infection in neonates and children. *Pediatr Infect Dis J*. 1995 Nov;14(11):948-54.
15. Fransen K, van Kerckhoven I, Piot P, van der Groen G. Evaluation and comparison of the Amplicor HIV-1 PCR kit with an 'in-house' nested PCR. *Clin Diagn Virol*. 1995 Dec;4(4):311-9.
16. Bremer JW, Lew JF, Cooper E, Hillyer GV, Pitt J, Handelsman E, et al. Diagnosis of infection with human immunodeficiency virus type 1 by a DNA polymerase chain reaction assay among infants enrolled in the Women and Infants' Transmission Study. *J Pediatr*. 1996 Aug;129(2):198-207.

17. Cassol S, Salas T, Arella M, Neumann P, Schechter MT, O'Shaughnessy M. Use of dried blood spot specimens in the detection of human immunodeficiency virus type 1 by the polymerase chain reaction. *J Clin Microbiol.* 1991 Apr;29(4):667-71.
18. Cassol S, Salas T, Gill MJ, Montpetit M, Rudnik J, Sy CT, et al. Stability of dried blood spot specimens for detection of human immunodeficiency virus DNA by polymerase chain reaction. *J Clin Microbiol.* 1992 Dec;30(12):3039-42.
19. Cassol SA, Lapointe N, Salas T, Hankins C, Arella M, Fauvel M, et al. Diagnosis of vertical HIV-1 transmission using the polymerase chain reaction and dried blood spot specimens. *J Acquir Immune Defic Syndr.* 1992;5(2):113-9.
20. Comeau AM, Harris JA, McIntosh K, Weiblen BJ, Hoff R, Grady GF. Polymerase chain reaction in detecting HIV infection among seropositive infants: relation to clinical status and age and to results of other assays. *J Acquir Immune Defic Syndr.* 1992;5(3):271-8.
21. Comeau AM, Hsu HW, Schwerzler M, Mushinsky G, Grady GF. Detection of HIV in specimens from newborn screening programs. *N Engl J Med.* 1992 Jun 18;326(25):1703.
22. Yourno J, Conroy J. A novel polymerase chain reaction method for detection of human immunodeficiency virus in dried blood spots on filter paper. *J Clin Microbiol.* 1992 Nov;30(11):2887-92.
23. Amadori A, de Rossi A, Giaquinto C, Faulkner-Valle G, Zacchello F, Chieco-Bianchi L. In-vitro production of HIV-specific antibody in children at risk of AIDS. *Lancet.* 1988 Apr 16;1(8590):852-4.
24. Weiblen BJ, Lee FK, Cooper ER, Landesman SH, McIntosh K, Harris JA, et al. Early diagnosis of HIV infection in infants by detection of IgA HIV antibodies. *Lancet.* 1990 Apr 28;335(8696):988-90.
25. Landesman S, Weiblen B, Mendez H, Willoughby A, Goedert JJ, Rubinstein A, et al. Clinical utility of HIV-IgA immunoblot assay in the early diagnosis of perinatal HIV infection. *Jama.* 1991 Dec 25;266(24):3443-6.

26. Martin NL, Levy JA, Legg H, Weintrub PS, Cowan MJ, Wara DW. Detection of infection with human immunodeficiency virus (HIV) type 1 in infants by an anti-HIV immunoglobulin A assay using recombinant proteins. *J Pediatr*. 1991 Mar;118(3):354-8.
27. Quinn TC, Kline RL, Halsey N, Hutton N, Ruff A, Butz A, et al. Early diagnosis of perinatal HIV infection by detection of viral-specific IgA antibodies. *Jama*. 1991 Dec 25;266(24):3439-42.
28. Burgard M, Mayaux MJ, Blanche S, Ferroni A, Guihard-Moscato ML, Allemon MC, et al. The use of viral culture and p24 antigen testing to diagnose human immunodeficiency virus infection in neonates. The HIV Infection in Newborns French Collaborative Study Group. *N Engl J Med*. 1992 Oct 22;327(17):1192-7.
29. Simonds RJ, Brown TM, Thea DM, Orloff SL, Steketee RW, Lee FK, et al. Sensitivity and specificity of a qualitative RNA detection assay to diagnose HIV infection in young infants. Perinatal AIDS Collaborative Transmission Study. *Aids*. 1998 Aug 20;12(12):1545-9.
30. Delamare C, Burgard M, Mayaux MJ, Blanche S, Doussin A, Ivanoff S, et al. HIV-1 RNA detection in plasma for the diagnosis of infection in neonates. The French Pediatric HIV Infection Study Group. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1997 Jun 1;15(2):121-5.
31. Young NL, Shaffer N, Chaowanachan T, Chotpitayasunondh T, Vanparapar N, Mock PA, et al. Early diagnosis of HIV-1-infected infants in Thailand using RNA and DNA PCR assays sensitive to non-B subtypes. *J Acquir Immune Defic Syndr*. 2000 Aug 15;24(5):401-7.
32. Steketee RW, Abrams EJ, Thea DM, Brown TM, Lambert G, Orloff S, et al. Early detection of perinatal human immunodeficiency virus (HIV) type 1 infection using HIV RNA amplification and detection. New York City Perinatal HIV Transmission Collaborative Study. *J Infect Dis*. 1997 Mar;175(3):707-11.

33. Reisler RB, Thea DM, Pliner V, Green T, Lee F, Nesheim S, et al. Early detection of reverse transcriptase activity in plasma of neonates infected with HIV-1: a comparative analysis with RNA-based and DNA-based testing using polymerase chain reaction. *J Acquir Immune Defic Syndr*. 2001 Jan 1;26(1):93-102.
34. Nielsen K, Bryson YJ. Diagnosis of HIV infection in children. *Pediatr Clin North Am*. 2000 Feb;47(1):39-63.
35. Owens DK, Holodniy M, McDonald TW, Scott J, Sonnad S. A meta-analytic evaluation of the polymerase chain reaction for the diagnosis of HIV infection in infants. *Jama*. 1996 May 1;275(17):1342-8.
36. Mussi-Pinhata MM, Cervi MC, Isaac ML. Predicting perinatal HIV infection in young Brazilian infants: How accurate are signs, symptoms and immunological abnormalities? *Int J STD AIDS*. 2001 Mar;12(3):197-203.
37. Obimbo EM, Mbori-Ngacha DA, Ochieng JO, Richardson BA, Otieno PA, Bosire R, et al. Predictors of early mortality in a cohort of human immunodeficiency virus type 1-infected African children. *Pediatr Infect Dis J*. 2004 Jun;23(6):536-43.
38. Horwood C, Liebeschuetz S, Blaauw D, Cassol S, Qazi S. Diagnosis of paediatric HIV infection in a primary health care setting with a clinical algorithm. *Bull World Health Organ*. 2003;81(12):858-66.
39. Zijenah LS, Katzenstein DA, Nathoo KJ, Rusakaniko S, Tobaiwa O, Gwanzura C, et al. T lymphocytes among HIV-infected and -uninfected infants: CD4/CD8 ratio as a potential tool in diagnosis of infection in infants under the age of 2 years. *J Transl Med*. 2005 Feb 1;3(1):6.
40. Moodley D, Coovadia HM, Bobat RA, Madurai S, Sullivan JL. The relationship between maternal-infant antibody levels and vertical transmission of HIV-1 infection. *J Trop Pediatr*. 1997 Apr;43(2):75-9.
41. Madurai S, Moodley D, Coovadia HM, Gopaul W, Smith AN, York DF. Infant-maternal HIV-specific immunoglobulin G1 antibody ratios as an indicator of vertical transmission. *Aids*. 1997 Jul 15;11(9):1191-3.

42. Chantry CJ, Cooper ER, Pelton SI, Zorilla C, Hillyer GV, Diaz C. Seroreversion in human immunodeficiency virus-exposed but uninfected infants. *Pediatr Infect Dis J*. 1995 May;14(5):382-7.
43. Moodley D, Bobat RA, Coutsooudis A, Coovadia HM. Predicting perinatal human immunodeficiency virus infection by antibody patterns. *Pediatr Infect Dis J*. 1995 Oct;14(10):850-2.
44. Palasanthiran P, Robertson P, Ziegler JB, Graham GG. Decay of transplacental human immunodeficiency virus type 1 antibodies in neonates and infants. *J Infect Dis*. 1994 Dec;170(6):1593-6.
45. Newell ML, Loveday C, Dunn D, Kaye S, Tedder R, Peckham C, et al. Use of polymerase chain reaction and quantitative antibody tests in children born to human immunodeficiency virus-1-infected mothers. *J Med Virol*. 1995 Dec;47(4):330-5.
46. Candal D, Bulterys M, Abrams EJ, Steketee RW, Parekh BS. Efficacy of a less-sensitive enzyme immunoassay (3A11-LS) for early diagnosis of human immunodeficiency virus Type 1 infection in infants. *Clin Diagn Lab Immunol*. 2001 Nov;8(6):1282-5.
47. Sirinavin S, Atamasirikul K. Semiquantitative human immunodeficiency virus antibody tests in diagnosis of vertical infection. *Pediatr Infect Dis J*. 2000 Dec;19(12):1153-7.
48. Sherman GG. Quantitative enzyme-linked immunosorbent assays for diagnosis of human immunodeficiency virus infection in infancy. *Pediatr Infect Dis J*. 2001 Apr;20(4):462-3.
49. Johnson S, Nelson G, Violari A, Gray G. Early antibody testing to diagnose HIV infection in children. XIII International AIDS Conference. Durban, South Africa, 9-14 July 2000, abstract MoPeB2199.

50. Dunn DT, Brandt CD, Krivine A, Cassol SA, Roques P, Borkowsky W, et al. The sensitivity of HIV-1 DNA polymerase chain reaction in the neonatal period and the relative contributions of intra-uterine and intra-partum transmission. *Aids*. 1995 Sep;9(9):F7-11.
51. Benjamin DK, Jr., Miller WC, Fiscus SA, Benjamin DK, Morse M, Valentine M, et al. Rational testing of the HIV-exposed infant. *Pediatrics*. 2001 Jul;108(1):E3.
52. Wilfert C, Beck DT, Fleischman AR, Mofenson LM, Pantell RH, Schonberg SK, et al. American Academy of Pediatrics. Evaluation and Medical Treatment of the HIV-Exposed Infant. *Pediatrics*. 1997 Jun;99(6):909-18.
53. Webber LM, Cotton C, Stevens W. Diagnosis of Perinatal HIV-1 infection in South Africa. *S Afr J HIV Med*. 2001:21-6.
54. Gauteng Health Department. Management of HIV infection in children. Guidelines for clinics and hospitals. Pretoria, South Africa; November 2001.
55. African Network for the Care of Children affected by HIV/AIDS. Regional workshop report on early diagnosis and care of HIV infected children. Workshop report, 2003, Kampala, Uganda.
56. McCoy D. 7 March 2003. Interim Findings on the national PMTCT Pilot Sites: Lessons and Recommendations. Available from: <http://www.hst.org.za/pubs/pmtct/pmtctinterim.htm> [Accessed 5 December 2005].
57. Park M. Baby HIV success rate false due to incorrect info. *The Star*. 2004 June 10.
58. Doherty T, Besser M, Donohue S, Kamoga N, Stoops N, Williamson L, et al. September 2003. An Evaluation of the Prevention of Mother-To-Child Transmission (PMTCT) of HIV Initiative in South Africa. Lessons and Key recommendations. Available from: www.hst.org.za/uploads/files/pmtct_national.pdf. [Accessed 28 December 2005].

59. Sherman GG, Stevens WS, Stevens G, Galpin JS. Diagnosis of human immunodeficiency virus infection in perinatally exposed orphaned infants in a resource-poor setting. *Pediatr Infect Dis J*. 2000 Oct;19(10):1014-5.
60. Newell ML, Coovadia H, Cortina-Borja M, Rollins N, Gaillard P, Dabis F. Mortality of infected and uninfected infants born to HIV-infected mothers in Africa: a pooled analysis. *Lancet*. 2004 Oct 2;364(9441):1236-43.
61. Department of Health. Integrated Management of Childhood Illness: South African Edition. DOH, Pretoria, South Africa; October 2003.
62. Jones SA, Sherman GG, Coovadia AH. Can clinical algorithms deliver an accurate diagnosis of HIV infection in infancy? *Bull World Health Organ*. 2005 Jul;83(7):559-60.
63. O'Shea S, Mullen J, Tong CY. Pediatric diagnosis of human immunodeficiency virus type 1 infection: the problem of false negative DNA polymerase chain reaction results. *Pediatr Infect Dis J*. 2003 May;22(5):476-7.
64. Nesheim S, Palumbo P, Sullivan K, Lee F, Vink P, Abrams E, et al. Quantitative RNA testing for diagnosis of HIV-infected infants. *J Acquir Immune Defic Syndr*. 2003 Feb 1;32(2):192-5.
65. Schupbach J, Boni J, Bisset LR, Tomasik Z, Fischer M, Gunthard HF, et al. HIV-1 p24 antigen is a significant inverse correlate of CD4 T-cell change in patients with suppressed viremia under long-term antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2003 Jul 1;33(3):292-9.
66. Patton JC, Sherman GG, Coovadia AH, Stevens WS, Meyers TM. Ultrasensitive human immunodeficiency virus type 1 p24 antigen assay modified for use on dried whole-blood spots as a reliable, affordable test for infant diagnosis. *Clin Vaccine Immunol*. 2006 Jan;13(1):152-5.
67. De Baets AJ, Edidi BS, Kasali MJ, Beelaert G, Schrooten W, Litzroth A, et al. Pediatric human immunodeficiency virus screening in an African district hospital. *Clin Diagn Lab Immunol*. 2005 Jan;12(1):86-92.

68. Guay LA, Musoke P, Fleming T, Bagenda D, Allen M, Nakabiito C, et al. Intrapartum and neonatal single-dose nevirapine compared with zidovudine for prevention of mother-to-child transmission of HIV-1 in Kampala, Uganda: HIVNET 012 randomised trial. *Lancet*. 1999 Sep 4;354(9181):795-802.
69. Department of Health. 2005. Guidelines for the management of HIV-infected children. Available from: <http://www.doh.gov.za/docs/factsheets/guidelines/artguide04-f.html> [Accessed 10 January 2006].
70. Jones SA, Sherman GG, Varga CA. Exploring socio-economic conditions and poor follow-up rates of HIV-exposed infants in Johannesburg, South Africa. *AIDS Care*. 2005 May;17(4):466-70.
71. Loubser S, Balfe P, Sherman G, Hammer S, Kuhn L, Morris L. Decay of K103N mutants in cellular DNA and plasma RNA after single-dose nevirapine to reduce mother-to-child HIV transmission. *Aids*. 2006 (in press).
72. Varga CA, Sherman GG, Maphosa J, Jones SA. Psychosocial consequences of early diagnosis of HIV status in vertically exposed infants in Johannesburg, South Africa. *Health Care Women Int*. 2005 May;26(5):387-97.
73. Varga C, Sherman G, Jones S. HIV Disclosure in the Context of Vertical Transmission: HIV-Positive Mothers in Johannesburg, South Africa. *AIDS Care* 2006 (in press).
74. Green J. New ray of hope for 1000 AIDS orphans. *The Star*. 2003 December 8.
75. Stevens W, Sherman GG, Cotton M, Gerntholtz L, Webber L. Revised guidelines for diagnosis of perinatal HIV-1 infection in South Africa. *S Afr J HIV Med*. 2006 (in press).
76. Fischer A, Lejczak C, Lambert C, Servais J, Makombe N, Rusine J, et al. Simple DNA extraction method for dried blood spots and comparison of two PCR assays for diagnosis of vertical human immunodeficiency virus type 1 transmission in Rwanda. *J Clin Microbiol*. 2004 Jan;42(1):16-20.

77. Luo W, Yang H, Rathbun K, Pau CP, Ou CY. Detection of human immunodeficiency virus type 1 DNA in dried blood spots by a duplex real-time PCR assay. *J Clin Microbiol.* 2005 Apr;43(4):1851-7.
78. Li CC, Seidel KD, Coombs RW, Frenkel LM. Detection and quantification of human immunodeficiency virus type 1 p24 antigen in dried whole blood and plasma on filter paper stored under various conditions. *J Clin Microbiol.* 2005 Aug;43(8):3901-5.
79. Rouet F, Ekouevi DK, Chaix ML, Burgard M, Inwoley A, Tony TD, et al. Transfer and evaluation of an automated, low-cost real-time reverse transcription-PCR test for diagnosis and monitoring of human immunodeficiency virus type 1 infection in a West African resource-limited setting. *J Clin Microbiol.* 2005 Jun;43(6):2709-17.
80. Zijenah LS, Tobaiwa O, Rusakaniko S, Nathoo KJ, Nhembe M, Matibe P, et al. Signal-boosted qualitative ultrasensitive p24 antigen assay for diagnosis of subtype C HIV-1 infection in infants under the age of 2 years. *J Acquir Immune Defic Syndr.* 2005 Aug 1;39(4):391-4.
81. Dineva MA, Candotti D, Fletcher-Brown F, Allain JP, Lee H. Simultaneous visual detection of multiple viral amplicons by dipstick assay. *J Clin Microbiol.* 2005 Aug;43(8):4015-21.
82. Lujan-Zilbermann J, Bitar W, Knapp K, Flynn P. Human immunodeficiency virus type 1 RNA polymerase chain reaction reasonably excludes infection in exposed infants. *Pediatr Infect Dis J.* 2003 Jan;22(1):97-9.
83. Lambert JS, Harris DR, Stiehler ER, Moye J, Jr., Fowler MG, Meyer WA, 3rd, et al. Performance characteristics of HIV-1 culture and HIV-1 DNA and RNA amplification assays for early diagnosis of perinatal HIV-1 infection. *J Acquir Immune Defic Syndr.* 2003 Dec 15;34(5):512-9.