

https://dspace.org/

# A review of the clinical characteristics, management and outcomes of neonates exposed to meconium stained amniotic fluid at Chris Hani Baragwanath Academic Hospital

**Ali-Dikole, Masida Linda**

**2022**
https://hdl.handle.net/10539/35467

Downloaded from a DSpace repository. https://dspace.org/

# A Review of Neonates Exposed to Meconium Stained Amniotic Fluid at a Tertiary Level, Public Hospital in South Africa

Masida Linda Ali-Dikole[1†*], Alison van Kwawegen[1], Reenu Thomas[1]

[1]Department of Paediatrics, Chris Hani Baragwanath Academic Hospital and School of Clinical Medicine, University of the Witwatersrand.

*** Correspondence:**
M. Linda Ali-Dikole
ladikole@gmail.com

**Keywords: meconium stained amniotic fluid[1], meconium aspiration syndrome[2], incidence[3], resuscitation[4], hypoxic ischaemic encehalopathy[5], ventilation[6], mortality[7]**

**Abstract**

**Objective:** This study aimed to determine the incidence, clinical characteristics, management and outcomes of neonates born through meconium-stained amniotic fluid (MSAF) and those admitted with meconium aspiration syndrome (MAS).

**Study design:** This was a retrospective study conducted in a tertiary hospital in Johannesburg, South Africa. All neonates born through MSAF, and those admitted with MAS, from January to June 2018 were included.

**Results:** Incidence of neonates born through MSAF was 44/1000 live births. The median birth weight was 3125 grams and median gestational age was 39 weeks. Forty-eight percent required resuscitation and 55.5% required admission. The incidence of MAS was 10/1000 live births, with 32% requiring invasive ventilation. Mortality in neonates exposed to MSAF was 2.5%, and 9.3% in those with MAS.

**Conclusion:** The high observed incidence of MSAF and MAS, with associated mortality, necessitate effective intrapartum monitoring and appropriate postnatal management of infants exposed to MSAF to improve outcomes.