

## REFERENCES

Bakewell A, Higgins R, Edmunds M. 2002. Quality of life in peritoneal dialysis patients: decline over time and association with clinical outcomes. *Kidney international*. 61: 239-248.

Borg G. A.V. 1982. Psychosocial bases of perceived exertion. *Medicine & Science in Sport & Exercise*. 14 (5): 377 – 381

Cappy S, Jablonka J, Schroeder E.T. 1999. The effects of exercise during hemodialysis on physical performance and nutrition assessment. *Journal of Renal Nutrition*. 9: 63-70

Cheema B.S.B, Smith B.C.F, Singh M.A.F. 2005. A rationale for intradialytic exercise training as standard clinical practice in ESRD. *American Journal of Kidney Diseases*. 45: 912-916

Carney R.M, Templeton B, Hong B.A, Harter H.R, Hagberg J.M, Schechtman K.B, Goldberg A.P. 1987. Exercise reduces depression and increases performance of pleasant activities in hemodialysis patients. *Nephron*. 47: 194-198

Cleary J, Drennan J, 2005. Quality of life of patients on haemodialysis for end-stage renal disease. *Journal of Advanced Nursing*. 51 (6): 577-586

Daugirdas J.T, Blake P.G, Ing T.S. 2001. *Handbook of Dialysis*, Third Edition. Lippincott Williams & Wilkins, Philadelphia.

Daul A.E, Schäfers R.F, Daul K, Philipp T. 2004. Exercise during haemodialysis. *Clinical Nephrology*. 61: S26-S30

Diaz-Buxo J, Lowrie E, Lew N, Zhang H, Lazarus J. 2000. Quality-of-life evaluation using Short Form 36: comparison in hemodialysis and peritoneal dialysis patients. *American Journal of Kidney Diseases*. 35: 293-300

Diesel W, Emms M, Knight BK, Noakes T.D, Swanepoel C.R, van Zyl Smit R, Kaschula R.O, Sinclair-Smith C.C. 1993. Morphologic features of the myopathy associated with chronic renal failure. *American Journal of Kidney Diseases*. 22: 677-684

Dinenno F.A, Jones P.P, Seals D.R, Tanaka H. 1999. Limb blood flow and vascular conductance are related with age in healthy humans. *Circulation*. 100: 164-170

Endotec International. 2005. Venous Anti-Stasis Slipper (VASS) - Technical manual.

Fleming P, Fitzgerald P, Devitt A, Rice J, Murray P. 2000. The effect of the position of the limb on venous impulse foot pumps. *The Journal of Bone and Joint Surgery* 82-B: 433-434

Franks P.J, Posnett J. European Wound Management Association (EWMA): Position Document. Page 8: <<http://www.proguide.net/pdf/english.pdf>>

Fresenius Medical Care-Dialysis Information Guide, 2004. <<http://www.fmc-ag.com>>

Goldberg A.P, Geltman E.M, Gavin J.R, Carney R.M, Hagberg J.M, Delmez J.A, Naumovich A, Oldfield M.H, Harter H.R. 1986. Exercise training reduces coronary risk and effectively rehabilitates hemodialysis patients. *Nephron*. 42: 311-316

Gotch F.A, Sargent J.A. 1985. A mechanistic analysis of the National Cooperative Dialysis Study (NCDS). *Kidney International*. 28:526-534

Green J, 2005. Renal failure sufferer denied state treatment. *The Star Newspaper*, January 31, page 3.

Horl M. P, Horl W. H. 2002. Hemodialysis-associated hypertension: pathophysiology and therapy. *American Journal of Kidney Diseases*. 39: 227-244

Hruska K. A, Teitelbaum S.L. 1995. Renal Osteodystrophy. *New England Journal of Medicine*. 333:166-174

Hsieh R-L, Lee W-C, Chang C-H. 2006. Maximal cardiovascular fitness and its correlates in ambulatory hemodialysis patients. *American Journal of Kidney Diseases*. 48: 21-27

Hughes J, Jelsma J, Maclean E, Darder M, Tinise X. 2004. The health-related quality of life of people living with HIV / AIDS. *Disability and Rehabilitation*. 26(6):371-376

Jelsma J, Maclean E, Hughes J, Tinise X, Darder M. 2005. An investigation into the health-related quality of life of individuals living with HIV who are receiving HAART. *AIDS Care*. 17(5): 579-588

Johansen K.L, Sakkas G.K, Doyle J, Shubert T, Dudley R.A. 2003. Exercise counselling practices among nephrologists caring for patients on dialysis. *American Journal of Kidney Diseases*. 41: 171-178

Kong C.H, Tattersall J.E, Greenwood R.N, Farrington K. 1999. The effect of exercise during hemodialysis on solute removal. *Nephrology Dialysis Transplant*. 14: 2927-2931

Konstantinidou E, Koukouvou G, Kouidi E, Deligiannis A, Tourkantonis A. 2002. Exercise training in patients with end-stage renal disease on hemodialysis: comparison of three rehabilitation programs. *Journal of Rehabilitation Medicine*. 34: 40-45

Kouidi E. 2004. Health related quality of life in end-stage renal disease patients: The effects of renal rehabilitation. *Clinical Nephrology*. 61: S60-S71

Kouidi E, Grekas D, Deligiannis, Tourkantonis A. 2004. Outcomes of long-term exercise training in dialysis patients: comparison of two training programs. *Clinical Nephrology*. 61: S31-S38

Kouidi E, Iacovides A, Iordanidis P, Vassiliou S, Deligiannis A, Ierodiakonou C, Tourkantonis A. 1997. Exercise renal rehabilitation program (ERRP): psychological effects. *Nephron* 77: 152-158

Llach F. 1993. Chronic Renal Failure. Papper's Clinical Nephrology, Third Edition. Little, Brown and Company: 67-107

Lok P. 1996. Stressors, coping mechanisms and quality of life among dialysis patients in Australia. *Journal of Advanced Nursing*. 23 (5): 873-888

Miller B.W, Cress C.L, Johnson M.E, Nichols D.H, Schnitzler M.A. 2002. Exercise during hemodialysis decreases the use of antihypertensive medications. *American Journal of Kidney Diseases*. 39: 828-833

Molakeng S. 2002. Health Systems Trust. "15000 wait for donated organs" <[http://www.hst.co.za/news/item/php?nws\\_id=20020821](http://www.hst.co.za/news/item/php?nws_id=20020821)>

Moore G.E, Painter P.L, Brinker K.R, Stray-Gundersen J.S, Mitchell J. H. 1998. Cardiovascular response to submaximal stationary cycling during hemodialysis. *American Journal of Kidney Diseases*. 31: 631-637

Morgan AP. 1973. Long-Term hemodialysis, Second Edition. Grune and Stratton, Inc. 40-64

Morris R.J, Woodcock J.P. 2004. Evidence-Based Compression: Prevention of Stasis and Deep Vein Thrombosis. *Annals of Surgery* 239 (2):162-171

Moug S.J, Grant S, Creed G, Boulton Jones M. 2004. Exercise during haemodialysis: West Scotland pilot study. *Scottish Medical Journal*. 49: 14-17

Neal C.R, Resnikoff E, Unger A.M. 1981. Treatment of dialysis-related muscle cramps with hypertonic dextrose. *Archives of Internal Medicine*. 141: 171-173

Nephrology Channel.<[http:// www.nephrologychannel.com/crf/](http://www.nephrologychannel.com/crf/)>

O'Connor A.S, Leon J.B, Sehgal A.R. 2002. The relative predictive ability of four different measures of hemodialysis dose. *American Journal of Kidney Diseases*. 40: 1289-1294

Oh-Park M, Fast A, Gopal S, Lynn R, Frei G, Drenth R, Zohman L. 2002. Exercise for the dialyzed: Aerobic and strength training during hemodialysis. *American Journal of Physical Medicine and Rehabilitation*. 81: 814-821

Paganini E.P. 1994. In search of an optimal hemaocrit level in dialysis patients: rehabilitation and quality-of-life implications. *American Journal of Kidney Diseases*. 24 (1): S10-S16

Painter P and Johansen K. 2006. Improving Physical Functioning: time to Be a Part of Routine Care. *American Journal of Kidney Diseases*. 48 (1): 167-170

Painter P, Carlson L, Carey S, Paul S, Myll J. 2000**a**. Low-functioning haemodialysis patients improve with exercise training. *American Journal of Kidney Diseases*. 36: 600-608

Painter P, Carlson L, Carey S, Paul S, Myll J. 2000**b**. Physical functioning and health-related quality-of-life changes with exercise training in haemodialysis patients. *American Journal of Kidney Diseases*. 35: 482-492

Painter P, Nelson-Worel J, Hill M, Thornbery D, Shelp W, Harrington A, Weinstein A. 1986. Effects of exercise training during hemodialysis. *Nephron*. 43: 87-92

Parsons T.L, Toffelmire E.B, King-Van Vlack. 2006. Exercise Training During Hemodialysis Improves Dialysis Efficacy and Physical Performance. *Archives of Physical Medicine and Rehabilitation* 87: 680-687

Parsons T.L, Toffelmire E.B, King-Van Vlack. 2004. The effect of an exercise program during haemodialysis on dialysis efficacy, blood pressure and quality of life in end stage renal disease (ESRD) patients. *Clinical Nephrology*. 61: 261-274

Rabin R, de Charro F. 2001. EQ-5D: a measure of health status from the EuroQol Group. *The Finnish Medical Society Duodecim, Ann Med*. 33: 337-343

Rousseau P. 1993. Immobility in the aged. *Archives of Family Medicine*. 2 (2): 169-177

South African Nephrology Report 2001.  
<[http:// www.doh.gov.za/mts/reports/nephrology](http://www.doh.gov.za/mts/reports/nephrology)>

Suet-Ching W. L. 2001. The quality of life of Hong Kong dialysis patients. *Journal of Advanced Nursing*. 35: 218-227

University of Virginia, Health System.  
<[http://www.healthsystem.virginia.edu/uvahealth/adult\\_urology/endstage.cfm](http://www.healthsystem.virginia.edu/uvahealth/adult_urology/endstage.cfm)>

Wadhwa. Chronic Renal Failure.  
<[uhmcwww.sunysb.edu/internalmed/nephro/webpages/Part\\_G.htm](http://uhmcwww.sunysb.edu/internalmed/nephro/webpages/Part_G.htm) - 113k->

Zawada E. T. 2001. Initiation of dialysis. From: Daugirdas JT, Blake PG, Ing TS, editors. *Handbook of dialysis*. 3<sup>rd</sup> ed. Philadelphia: Lippincott Williams & Wilkins; pg. 3-11