

Generating functions for giant graviton bound states

Warren Carlson,^b Robert de Mello Koch^{a,b} and Minkyoo Kim^{b,c}

^a*School of Science, Huzhou University,
Huzhou 313000, China*

^b*School of Physics and Mandelstam Institute for Theoretical Physics,
University of the Witwatersrand,
Wits, 2050, South Africa*

^c*Center for Quantum Spacetime (CQUeST), Sogang University Seoul,
121-742, Seoul, South Korea*

E-mail: Warren.Carlson@wits.ac.za, robert@zjhu.edu.cn,
mimkim80@gmail.com

ABSTRACT: We construct generating functions for operators dual to systems of giant gravitons with open strings attached. These operators have a bare dimension of order N so that the usual methods used to solve the planar limit are not applicable. The generating functions are given as integrals over auxiliary variables, which implement symmetrization and antisymmetrization of the indices of the fields from which the operator is composed. Operators of a good scaling dimension (eigenstates of the dilatation operator) are known as Gauss graph operators. Our generating functions give a simple construction of the Gauss graph operators which were previously obtained using a Fourier transform on a double coset. The new description provides a natural starting point for a systematic $\frac{1}{N}$ expansion for these operators as well as the action of the dilatation operator on them, in terms of a saddle point evaluation of their integral representation.

KEYWORDS: AdS-CFT Correspondence, Brane Dynamics in Gauge Theories, Gauge-Gravity Correspondence

ARXIV EPRINT: [2212.06731](https://arxiv.org/abs/2212.06731)

Contents

1	Introduction	1
2	Projectors and generating functions	4
2.1	Young symmetrizers	5
2.2	Bound states of two AdS giants	7
2.3	Excited bound states of two AdS giants	10
2.4	Bound states of two sphere giants	21
2.5	Excited bound states of two sphere giants	22
3	Action of the dilatation operator	24
3.1	Open string boundaries	25
3.2	Gauss graph operators	27
4	Discussion and conclusions	33
A	Check of the generating function for a bound state of two AdS giants	36
A.1	Operators constructed from the generating function	36
A.2	Comparison to restricted Schur polynomial and Gauss graph operators	39
B	Check of the generating function for a bound state of two sphere giants	44
B.1	Operators constructed from the generating function	45
B.2	Comparison to restricted Schur polynomial and Gauss graph operators	47
C	A Gauss graph identity	50
D	Correlation functions	53

1 Introduction

The AdS/CFT correspondence has generated significant progress in understanding both quantum gravity in negatively curved spacetimes, as well as strongly coupled Yang-Mills theory [1–3]. For observables with a dimension that is fixed as we take $N \rightarrow \infty$ integrability was discovered [4, 5] giving unprecedented control, at both weak and strong coupling, over the planar limit of the theory. For these observables, a complete matching with the dual string is evident (see, for example, [6]) and we now know an exquisite matching between string states in the gravity Hilbert space and single trace gauge invariant operators in super Yang-Mills theory.

There are many observables whose bare dimensions scale with a power of N as we take $N \rightarrow \infty$, that are of interest in holography. These operators are very heavy states

and they probe non-perturbative aspects of the dual string theory. Nice examples include giant gravitons [7–9] which are dual to operators constructed with order N fields [10–12], and new spacetime geometries which are dual to operators constructed using order N^2 fields [13]. The construction of the large N expansion for these operators is not straight forward. We don’t have a nice analogue of the statement that Feynman diagrams become ribbon graphs and that the genus of the graph determines the dependence on N . The basic problem is that the number of ribbon graphs of a given genus is now dependent on N so that the N dependence is no longer determined by the genus of the ribbon graph being summed. The combinatorics is much more involved and there does not seem to be a nice subset of diagrams that, if summed, produce the leading effect at large N .

Without a better understanding of how to perform a systematic approximation, one could try to sum everything, producing results that are exact to all orders in $\frac{1}{N}$. Remarkably, the paper [11] showed that this can actually be done in practice, for the free theory. A key insight of [11] was to show that the combinatorics of ribbon graphs can be reformulated using symmetric groups and their characters. This initiated an active area of research, opening new directions that are still being developed. The original paper [11] showed that Schur polynomials provide a basis for the local gauge invariant operators, that they diagonalize the free field two point function and that the free field two point function can be explicitly computed, to all orders in $\frac{1}{N}$. In addition, all extremal correlators were also explicitly computed, again to all orders in $\frac{1}{N}$. Schur polynomials are labelled by Young diagrams, which may have columns with a maximum length of N as well as rows that are arbitrarily long. In the same way that there is a beautiful matching between the Hilbert space of strings and single trace operators in the planar limit, [11] argued for a matching between long columns and sphere giant gravitons, as well as between long rows and AdS giant gravitons, generalizing and completing the observations of [10] which related single sphere giant gravitons to subdeterminant operators. See also [14, 15]. These results were quickly generalized to multi-matrix models in a series of papers [16–20], as well as to other gauge groups [21–29].

Giant gravitons branes can be excited by attaching open strings to them. The gauge theory operators dual to these states are described using decorated Young diagrams [30, 31]. The spin chain language that worked so well for closed strings in the planar limit has to be modified when describing the open strings attached to giant gravitons. The new effect that needs to be included is that single fields can hop between the giant graviton and onto the string, so that we are forced to consider the dynamics of a spin chain defined on a dynamical lattice. A creative solution to this problem was put in place in [32, 33] for strings attached to single sphere giant gravitons, and then extended to more general brane bound states in [34, 35]. By now, a compelling physical picture is evident [36–41].

As we have just discussed, combinatorial approaches based on permutation groups and their characters, compute correlation functions exactly to all order in $\frac{1}{N}$. This is a valuable source of information and has stimulated a lot of progress, but it hasn’t given any guidance into how to develop a systematic $\frac{1}{N}$ expansion to the physics of excited giant graviton systems. Since we don’t expect any except the most symmetric problems to be exactly solvable, this is an obstacle to progress. One attempt at a simplification has been to use a displaced corners approximation [42, 43]. The idea is that the physics of the

open strings should simplify in the limit in which the giant gravitons are well separated in spacetime. This corresponds to a limit in which the lengths of distinct rows (or columns) of the Young diagram differ by an amount $\sim N$. In this limit one can simplify the description of the operators considered, at large N [42, 43], and some mixing problems can be solved exactly [44, 45]. There is a nice harmony between these results and another complimentary approach, based a coherent state description of the giant gravitons [46–49]. However, these small steps do not really improve the situation and a systematic $\frac{1}{N}$ expansion of the giant plus open string physics, in which, for example, we see the splitting and joining of open strings, remains to be developed.

An important recent development for the case of maximal giants was reported in the series of papers [50–52]. The maximal giant graviton is described by a determinant. The idea of [50–52] is to write the determinant as a fermionic integral. The rewriting replaces the symmetric group projector onto the representation given by a column of length N , written in terms of symmetric group characters, with an integral over some new auxiliary Grassmann fields. By writing the symmetric group projector as an integral one opens the possibility of performing a saddle point analysis which can provide the starting point for a systematic $\frac{1}{N}$ expansion. Similar integrals have been constructed for the projectors onto representations labelled by a single column or row, with an arbitrary length [53, 54]. Another development [55–58], which is closely related, has shown that many of the results from the combinatorial approach can be reproduced by writing suitable generating functions of coherent states as a matrix integral. In some cases these matrix integrals are related to the well known Harish-Chandra-Itzykson-Zuber integral and can be computed using localization. In this approach, non-trivial combinatorial information is repackaged in the manipulations of the coherent state generating function. This is a promising development, relevant to systematic $\frac{1}{N}$ expansions for heavy operators.

The central question motivating us in this paper, is the systematic $\frac{1}{N}$ expansion for giant graviton systems. Physically this expansion is naturally described in terms of splitting and joining interactions of open strings. Towards this end, we develop integral representations for generating functions of Schur polynomials labelled by Young diagrams that have two rows (corresponding to a boundstate of two AdS giants) or two columns (corresponding to a boundstate of two sphere giants). Our approach to the $\frac{1}{N}$ expansion entails developing a saddle point expansion of these integral representations [59]. Apart from this application, we will see that the new representations provide a dramatically simpler description of excited giant graviton systems. To demonstrate this, we explain how to recover known results in what follows. The class of problems we focus on is rather rich because a bound state of two giant gravitons is described by interesting dynamics. The world volume theory of p branes is a $U(p)$ gauge theory [60], so in this situation with $p = 2$ the effective Hamiltonian coming from the dilatation operator should reproduce a $U(2)$ non-Abelian gauge theory [30, 61]. To verify this expectation, we must necessarily correctly describe the splitting and joining of open strings, suggesting that this two giant bound state system is a good testing ground for the methods we are trying to develop.

Constructing the operators dual to a bound state of two giant gravitons requires symmetric group projectors labelled by Young diagrams that have two columns or two rows.

The key new progress reported in this paper starts with the construction of an integral representation for these projectors. To motivate the construction, in section 2 we describe how Schur polynomials are constructed using Young projectors. It is the Young projectors that can be given an integral representation. Since there is a direct connection between Schur polynomials and characters of the symmetric group, the same integral representations also provide new formulas for character generating functions. A simple generalization beyond the case of a single matrix produces integral representations for operators that are dual to excited giant graviton bound states. A detailed comparison of the operators constructed show that they correspond to the Gauss graph operators discovered in [44, 45]. We further develop this connection in section 3, where we evaluate the action of the dilatation operator on the excited giant graviton bound states. We discuss our results and suggest how they might be extended in section 4. The appendices collect technical details needed to perform the computations discussed in the body of the paper.

2 Projectors and generating functions

Half BPS operators in $\mathcal{N} = 4$ super Yang-Mills theory, dual to a system of giant graviton branes, are given by Schur polynomials $\chi_R(Z)$. The label R is a Young diagram with at most N rows. Each long column¹ in the Young diagram corresponds to a sphere giant graviton and each long row to an AdS giant graviton.² An innovative framework for thinking about the construction of these Schur polynomials and their correlation functions, was developed in [62] where the role of projection operators was stressed and an elegant diagrammatic formalism was established. The relevant projection operators used to construct an operator composed of n Z fields can be written in terms characters of the symmetric group S_n , as well as permutations acting on $V^{\otimes n}$ where V is the vector space that the matrix Z acts on. These characters and permutations are determined by the Young diagram R of the Schur polynomial as well as the number n of fields in the operator. The parameter N does not enter transparently in the construction.³ Consequently, simplifications as $N \rightarrow \infty$ are not immediate and it is not clear how one would pursue a systematic $\frac{1}{N}$ expansion. To go beyond the half BPS operators, we need to go beyond the single matrix Z sector. The simplest way to do this is by constructing operators which are systematically small deformations of the half BPS operators. These are nice systems to study as we expect that they will be simpler than the generic system that might be considered. In the dual gravity description, the giant graviton system is excited by attaching open strings to the giant gravitons. These generalizations also have a nice description in terms of projectors which proves useful when evaluating correlation functions and matrix elements of the dilatation operator, but again it is not obvious how a systematic large N expansion can be developed.

¹By a “long column” of a Young diagram R , we mean a column that has order N boxes in it. The number of boxes in each column is bounded by N . Similarly, a “long row” refers to a row with order N boxes. Row lengths are unbounded.

²Recall that a giant graviton is a spherical S^3 brane that carries a D3-brane dipole moment and the same quantum numbers as a point graviton. A sphere giant graviton is one for which the world volume $S^3 \subset S^5$, while an AdS giant graviton is one for which $S^3 \subset AdS_5$.

³ N does appear as a cut off on the number of rows of R .

In this section we briefly review the projection operator construction and its relation to Young symmetrizers. The language of Young symmetrizers gives a construction of the projectors which employs simple symmetrization and antisymmetrization operations on the indices of the Z fields appearing in the operator. Since these operations can be realized by introducing extra fermionic and bosonic auxiliary fields, we are able to construct an integral representation for the projection operators. More precisely, the integral representation gives a generating function for all Schur polynomials with two rows or two columns. This novel generating function provides a starting point for a systematic saddle point approximation. The generating function can also be interpreted as a generating function for characters, suggesting that one might be able to generate a systematic $\frac{1}{N}$ expansion of the characters themselves. There are natural ways to generalize the generating functions of Schur polynomials, to generating functions describing the Schur polynomials with strings attached. Remarkably, as we will demonstrate, the operators constructed in this way have an immediate and direct connection to other constructions of excited giant graviton operators based on restricted Schur polynomials and Gauss graph operators. We will develop this connection in the section 3 where we evaluate the action of the dilatation operator on the excited giant graviton states.

2.1 Young symmetrizers

Consider the construction of a Schur polynomial χ_R which is a function of a single complex adjoint scalar Z . The Young diagram R labelling the Schur polynomial specifies the system of D-branes as well as an irreducible representation R of the symmetric group S_n where n is the number of boxes in the diagram R , which is also the number of Z fields used to construct the operator. A simple formula for the Schur polynomial employs a projection operators P_R as follows [62]

$$\chi_R(Z) = \text{Tr}_{V^{\otimes n}}(P_R Z^{\otimes n}) \quad (2.1)$$

Here Z acts on the vector space V and both P_R and $Z^{\otimes n}$ are linear operators mapping $V^{\otimes n}$ to itself. The image of the projector corresponds simultaneously to a representation of the symmetric group S_n and a representation of the unitary group $U(N)$, both labelled by the same Young diagram⁴ R . A concrete expression for the projection operator is given by⁵

$$P_R = \frac{1}{n!} \sum_{\sigma \in S_n} \chi_R(\sigma) \sigma \quad (2.2)$$

where the matrix elements of the permutation σ are given by

$$\sigma_{j_1 j_2 \dots j_n}^{i_1 i_2 \dots i_n} = \delta_{j_{\sigma(1)}}^{i_1} \delta_{j_{\sigma(2)}}^{i_2} \dots \delta_{j_{\sigma(n)}}^{i_n} \quad (2.3)$$

so that

$$\text{Tr}(\sigma Z^{\otimes n}) = Z_{i_{\sigma(1)}}^{i_1} Z_{i_{\sigma(2)}}^{i_2} \dots Z_{i_{\sigma(n)}}^{i_n} \quad (2.4)$$

⁴The projector projects to a subspace of dimension $d_R \text{Dim}_R$, where d_R is the dimension of symmetric group representation R and Dim_R is the dimension of $U(N)$ representation R . States in this subspace carry both an S_n state label and a $U(N)$ state label.

⁵As written P_R is proportional to a projection operator. For a correctly normalized projection operator obeying $P_R P_S = \delta_{RS} P_R$ the right hand side of (2.2) must be multiplied by d_R .

Thus, given the characters of S_n a formula for P_R easily follows. These formulas can be extended to study operators constructed using more than one matrix, making it possible to consider perturbations of these half BPS operators. The characters that are needed are less well known and the combinatorics quickly becomes complicated. Nevertheless, at the level of two point functions we obtain a complete generalization of the Schur polynomial results i.e. we are able to construct a complete basis for the local operators, this basis takes finite N effects into account and this basis diagonalizes the free field theory two point function [19].

An alternative (but completely equivalent) approach uses the Young symmetrizer associated to R to construct the projector P_R . A Young symmetrizer is an element of the group algebra of the symmetric group. It is an endomorphism of the vector space $V^{\otimes n}$ which uses the action (2.4) of S_n on $V^{\otimes n}$ given by permuting indices. We illustrate the Young projectors making use of a specific example. Consider a Young diagram R that has two rows, with three boxes in the first row and two in the second. Each box corresponds to a field. The boxes in R are labelled, with a unique integer, as follows

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array} \quad (2.5)$$

There is no rule for the placement of integers in boxes, except that each box should be assigned a unique integer. Each box corresponds to the indices of a field in $Z^{\otimes 5}$ as shown below

$$Z_{j_1}^{i_1} Z_{j_2}^{i_2} Z_{j_3}^{i_3} Z_{j_4}^{i_4} Z_{j_5}^{i_5} \quad (2.6)$$

For every labelled Young diagram R we can define a “row group” for each row and a “column group” for each column. Each row group permutes the indices in the row, in all possible ways. Thus, for the example given in (2.5) the two row groups are

$$\mathcal{G}_{\text{row1}} = \{1, (12), (13), (23), (123), (132)\} \quad \mathcal{G}_{\text{row2}} = \{1, (45)\} \quad (2.7)$$

where we have written the permutations in cycle notation. The product of these groups is denoted $\mathcal{G}_{\text{row}}$. For the example we consider here $\mathcal{G}_{\text{row}}$ is of order 12. Similarly, the column groups are given by

$$\mathcal{G}_{\text{col1}} = \{1, (14)\} \quad \mathcal{G}_{\text{col2}} = \{1, (25)\} \quad (2.8)$$

and their product, denoted $\mathcal{G}_{\text{col}}$, has order 4. The Young symmetrizer is now given by

$$Y_{\begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \end{array}} = \sum_{\sigma_1 \in \mathcal{G}_{\text{row}}} \sum_{\sigma_2 \in \mathcal{G}_{\text{col}}} \text{sgn}(\sigma_2) \sigma_1 \sigma_2 \quad (2.9)$$

where $\text{sgn}(\sigma)$ is the parity of σ . A cycle of length 2 is called a transposition. Any permutation can be written as a product of transpositions. A particular permutation is even or odd if it can be expressed using an even or an odd number of transpositions. For an even permutation $\text{sgn}(\sigma) = 1$ and for an odd permutation $\text{sgn}(\sigma) = -1$. When applied to the tensor product (2.6) we find the σ_2 factor above antisymmetrizes over the pairs i_1, i_4 and i_2, i_5 of indices (i.e. the indices appearing in the columns of (2.5)) while the σ_1 factor

symmetrizes over the i_1, i_2, i_3 indices and over the i_4, i_5 indices (i.e. the indices associated to the rows of R). After a simple sum we find that

$$\begin{aligned} \text{Tr}(Y_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \end{array}} Z^{\otimes 5}) &= \text{Tr}(Z^5) + 2 \text{Tr}(Z^2) \text{Tr}(Z)^3 - 4 \text{Tr}(Z)^2 \text{Tr}(Z^3) + 3 \text{Tr}(Z) \text{Tr}(Z^2)^2 \\ &\quad - 6 \text{Tr}(Z) \text{Tr}(Z^4) + 4 \text{Tr}(Z^2) \text{Tr}(Z^3) \end{aligned} \quad (2.10)$$

The Schur polynomial corresponding to this Young diagram is given by

$$\begin{aligned} \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \end{array}}(Z) &= \frac{1}{5!} \left(5 \text{Tr}(Z^5) + 10 \text{Tr}(Z^2) \text{Tr}(Z)^3 - 20 \text{Tr}(Z)^2 \text{Tr}(Z^3) + 15 \text{Tr}(Z) \text{Tr}(Z^2)^2 \right. \\ &\quad \left. - 30 \text{Tr}(Z) \text{Tr}(Z^4) + 20 \text{Tr}(Z^2) \text{Tr}(Z^3) \right) \end{aligned} \quad (2.11)$$

which illustrates the general formula

$$\chi_R(Z) = \frac{d_R}{n!} \text{Tr} \left(Y_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \end{array}} Z^{\otimes 5} \right) \quad (2.12)$$

Thus, up to an unimportant overall constant, we can construct Schur polynomials by using Young projectors. This observation can be used to give an integral representation of the Young projectors needed to construct the two giant graviton bound states.

2.2 Bound states of two AdS giants

A bound state of two AdS giants, one of momentum p_1 and a second of momentum p_2 with $p_2 \leq p_1$, corresponds to a Schur polynomial labelled by a Young diagram with two rows. There are p_1 boxes in the first row and p_2 boxes in the second. Denote this Young diagram by listing row lengths, i.e. $R = (p_1, p_2)$. To describe a system of two giant gravitons, both p_1 and p_2 are of order N . The Young projector must symmetrize over all indices in row 1 and all indices in row 2, as well as antisymmetrize the pair of indices belonging to each column. The symmetrization to be performed represents the bulk of the work. Fortunately, this symmetrization of indices is easily accomplished with the help of a Gaussian integral. Indeed, a simple application of Wick's theorem implies that

$$\frac{1}{\pi^N} \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\phi}^a \phi_a} \bar{\phi}^{i_1} \bar{\phi}^{i_2} \dots \bar{\phi}^{i_k} \phi_{j_1} \phi_{j_2} \dots \phi_{j_k} = \sum_{\sigma \in S_k} \delta_{j_{\sigma(1)}}^{i_1} \delta_{j_{\sigma(2)}}^{i_2} \dots \delta_{j_{\sigma(k)}}^{i_k} \quad (2.13)$$

$$d^N \phi \equiv \prod_{a=1}^N d\phi_a \quad d^N \bar{\phi} \equiv \prod_{a=1}^N d\bar{\phi}^a \quad (2.14)$$

To construct the generating function, we perform the anti-symmetrization in each column by hand and use $(\varphi_i, \bar{\varphi}^i)$ to symmetrize indices in the first row and $(\phi_i, \bar{\phi}^i)$ to symmetrize indices in the second row. The two giant graviton bound state generating function can then be written as follows (as usual, repeated indices are summed)

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2) &= \sum_{p_1=p_2}^{\infty} \sum_{p_2=1}^{\infty} t_2^{2p_2} t_1^{p_1-p_2} \frac{(p_1+p_2)!}{d_{(p_1,p_2)} p_2! (p_1-p_2)!} \chi_{(p_1,p_2)}(Z) \\ &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \\ &\quad \times e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i + t_1 \bar{\varphi}^i Z_i^j \varphi_j + t_2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l} \end{aligned} \quad (2.15)$$

The coefficient of the Schur polynomial in the first line above combines factors arising from power expanding the exponential on the second line above (so that $t_2^{2p_2}$ comes with $1/p_2!$ and $t_1^{p_1-p_2}$ comes with $1/(p_1-p_2)!$), as well as from the coefficient in (2.12). In appendix A we check this result by performing the integral to second order in t_1 and t_2 and show that we reproduce the correct formulas for the relevant Schur polynomials.

This generating function can also be interpreted as a generating function for characters of the symmetric group $S_{p_1+p_2}$ in the representation labelled by Young diagram $R = (p_1, p_2)$. Permutations with a given cycle structure all belong to the same conjugacy class and the character is a function on these classes. We follow the standard notation $(1^{n_1}, 2^{n_2}, \dots, (p_1+p_2)^{n_{p_1+p_2}})$ to denote the cycle structure of the permutation which has n_k k -cycles. In what follows we use the shorthand $\{i^{n_i}\}$ for this cycle structure. As usual we have

$$\sum_{k=1}^{p_1+p_2} kn_k = p_1 + p_2 \quad (2.16)$$

for permutations $\sigma \in S_{p_1+p_2}$. Denote the symmetric group $S_{p_1+p_2}$ character, of a permutation with cycle structure $\{i^{n_i}\}$ in irreducible representation $R = (p_1, p_2)$ by $\chi_{(p_1, p_2)}(\{i^{n_i}\})$. The number of permutations with this cycle structure is given by

$$\frac{(p_1 + p_2)!}{1^{n_1} n_1! 2^{n_2} n_2! 3^{n_3} n_3! \dots (p_1 + p_2)^{n_{p_1+p_2}} n_{(p_1+p_2)}!} \quad (2.17)$$

We can then write the generating function for characters of the irreducible representations $R = (p_1, p_2)$ as follows

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2) = & \sum_{p_1=p_2}^{\infty} \sum_{p_2=1}^{\infty} \sum_{\{n_i\}} \frac{t_2^{2p_2} t_1^{p_1-p_2}}{1^{n_1} n_1! 2^{n_2} n_2! \dots (p_1 + p_2)^{n_{p_1+p_2}} n_{(p_1+p_2)}!} \frac{(p_1 + 1)!}{(p_1 - p_2 + 1)!} \\ & \times \chi_{(p_1, p_2)}(\{i^{n_i}\}) \text{Tr}(Z)^{n_1} \text{Tr}(Z^2)^{n_2} \dots \text{Tr}(Z^{p_1+p_2})^{n_{p_1+p_2}} \end{aligned} \quad (2.18)$$

where $\sum_{\{n_i\}}$ denotes a sum over the conjugacy classes of $S_{p_1+p_2}$. As a simple illustration of this formula, expanding to linear order in t_1 and t_2^2 , we find

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2) = & 1 + t_1 \chi_{\square}((\cdot)) \text{Tr}(Z) + t_2^2 \chi_{\square\square}((\cdot)(\cdot)) \text{Tr}(Z)^2 + t_2^2 \chi_{\square\square}((\cdot\cdot)) \text{Tr}(Z^2) + \dots \\ = & 1 + t_1 \text{Tr}(Z) + t_2^2 \text{Tr}(Z)^2 - t_2^2 \text{Tr}(Z^2) + \dots \end{aligned} \quad (2.19)$$

reproducing the correct characters for S_1 and S_2

$$\chi_{\square}((\cdot)) = 1 \quad \chi_{\square\square}((\cdot)(\cdot)) = 1 \quad \chi_{\square\square}((\cdot\cdot)) = -1 \quad (2.20)$$

This integral representation is a nice repackaging of the computation of characters using Young symmetrizers. Since the computation of the power series expansion of $Z_{\text{AdS}}(t_1, t_2)$ involves evaluating moments of a Gaussian integral, there is no obstacle to computing characters for any reasonable value of p_1 and p_2 . Evaluating $Z_{\text{AdS}}(t_1, t_2)$ explicitly is frustrated by the fact that the term with coefficient t_2^2 is quartic in the integration variables.

By using the simple identity

$$\begin{aligned} \frac{1}{\pi} \int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 e^{-(x_1+ix_2)(x_1-ix_2)+A(x_1+ix_2)+B(x_1-ix_2)} &\equiv \frac{1}{\pi} \int dx \int d\bar{x} e^{-x\bar{x}+Ax+B\bar{x}} \\ &= e^{AB} \end{aligned} \quad (2.21)$$

we find

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2) &= \frac{1}{\pi^{2N+2}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \int dx \int d\bar{x} \int dy \int d\bar{y} \\ &\quad \times e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i - x\bar{x} - y\bar{y} + t_1 \bar{\varphi}^i Z_i^j \varphi_j + t_2 x \bar{\varphi}^i Z_i^k \varphi_k + t_2 \bar{x} \bar{\phi}^j Z_j^l \phi_l + t_2 y \bar{\varphi}^i Z_i^l \phi_l - t_2 \bar{y} \bar{\phi}^j Z_j^k \varphi_k} \end{aligned}$$

Notice that after this rewriting the integral over both $\varphi_i, \bar{\varphi}^i$ and $\phi_i, \bar{\phi}^i$ are Gaussian so that we can do these integrals exactly. The result is

$$Z_{\text{AdS}}(t_1, t_2) = \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} \frac{e^{-x\bar{x}-y\bar{y}}}{\det M} \quad (2.22)$$

where

$$M = \mathbf{1}_2 \otimes \mathbf{1}_N - m \otimes Z \quad m = \begin{bmatrix} t_1 + t_2 x & t_2 y \\ -t_2 \bar{y} & t_2 \bar{x} \end{bmatrix} \quad (2.23)$$

and $\mathbf{1}_k$ is the $k \times k$ identity matrix. The eigenvalues λ_a with $a = 1, 2$ of m are given by

$$\lambda_{1,2} = \frac{1}{2} \left(t_1 + t_2(x + \bar{x}) \pm \sqrt{(t_1 + t_2(x + \bar{x}))^2 - 4(\bar{x}t_1 t_2 + t_2^2(x\bar{x} + y\bar{y}))} \right) \quad (2.24)$$

Denote the eigenvalues of Z_j^i by z_i . A straight forward computation now shows that

$$\begin{aligned} \det M &= \prod_{i=1}^N \frac{1}{(1 - \lambda_1 z_i)(1 - \lambda_2 z_i)} \\ &= \exp \left(\sum_{n=1}^{\infty} \frac{\lambda_1^n + \lambda_2^n}{n} \text{Tr}(Z^n) \right) \\ &= \exp \left(\sum_{n=1}^{\infty} f_n \text{Tr}(Z^n) \right) \end{aligned} \quad (2.25)$$

where f_n is the following polynomial

$$\begin{aligned} f_n &= \sum_{r=0}^{\text{int}(\frac{n}{2})} \frac{(-\beta)^r}{r!} \frac{\alpha^{n-2r}}{(n-2r)!} (n-r-1)! \\ \alpha &= t_1 + t_2(x + \bar{x}) \quad \beta = t_1 t_2 \bar{x} + t_2^2(x\bar{x} + y\bar{y}) \end{aligned} \quad (2.26)$$

where $\text{int}(\frac{n}{2})$ is an instruction to take the integer part of $\frac{n}{2}$. Thus, we have

$$Z_{\text{AdS}}(t_1, t_2) = \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}+\sum_{n=1}^{\infty} f_n \text{Tr}(Z^n)} \quad (2.27)$$

This implies the following formula for the character

$$\chi_{(p_1, p_2)}(\{i^{n_i}\}) = [t_1^{p_1-p_2} t_2^{2p_2}] \left(\frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}} \frac{(p_1-p_2+1)!}{(p_1+1)!} \prod_{i=1}^{p_1+p_2} (f_{n_i})^{n_i} \right) \quad (2.28)$$

where $[x^n]p(x)$ stands for the coefficient of x^n in the power series expansion of the polynomial $p(x)$. This integral formula for characters of the symmetric group is a new result. Lets test this for the case that $p_2 = 0$, in which we case we know that

$$\chi_{(p_1,0)}(\{i^{n_i}\}) = 1 \quad (2.29)$$

When $t_2 = 0$ we easily find $f_n = t_1^n/n$ and it is possible to compute (2.27) explicitly

$$\begin{aligned} Z_{\text{AdS}}(t_1, 0) &= e^{\sum_{n=1}^{\infty} \frac{t_1^n}{n} \text{Tr}(Z^n)} \\ &= \sum_{p_1=0}^{\infty} \sum_{\{n_i\}} t_1^{p_1} \frac{1}{n_1!} \frac{1}{2^{n_2} n_2!} \cdots \frac{1}{2^{n_{p_1}} n_{p_1}!} \text{Tr}(Z)^{n_1} \text{Tr}(Z^2)^{n_2} \cdots \text{Tr}(Z^{p_1})^{n_{p_1}} \\ &= \sum_{p_1=0}^{\infty} \sum_{\{n_i\}} t_1^{p_1} \frac{1}{n_1!} \frac{1}{2^{n_2} n_2!} \cdots \frac{1}{2^{n_{p_1}} n_{p_1}!} \chi_{(p_1,0)}(\{i^{n_i}\}) \text{Tr}(Z)^{n_1} \text{Tr}(Z^2)^{n_2} \cdots \text{Tr}(Z^{p_1})^{n_{p_1}} \end{aligned}$$

which immediately implies (2.29). As a second example, consider $p_1 + p_2 = 6$ and choose a character from the conjugacy class (2^3) . We easily find

$$\frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}} (f_2)^3 = t_1^6 - 6t_1^4 t_2^2 + 60t_1^2 t_2^4 - 72t_2^6 \quad (2.30)$$

Thus we find

$$\begin{aligned} \chi_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}}([2^3]) &= [t_2^6] \left(\frac{1}{24} \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}} (f_2)^3 \right) = -3 \\ \chi_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}}([2^3]) &= [t_2^4 t_1^2] \left(\frac{1}{20} \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}} (f_2)^3 \right) = 3 \\ \chi_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}}([2^3]) &= [t_2^2 t_1^4] \left(\frac{1}{6} \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}} (f_2)^3 \right) = -1 \\ \chi_{\begin{array}{|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array}}([2^3]) &= [t_1^6] \left(\frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{-x\bar{x}-y\bar{y}} (f_2)^3 \right) = 1 \end{aligned} \quad (2.31)$$

which are indeed the correct results. This demonstrates that the computation of characters of S_n is reduced to evaluating moments of a Gaussian integral.

An interesting extension of these ideas would be to consider characters for $R = (p_1, p_2)$ with both $p_1 - p_2$ and p_2 of order N . The relevant character can be extracted from $Z_{\text{AdS}}(t_1, t_2)$ by performing the following contour integrals

$$\oint_{C_\infty} \frac{dt_1}{t_1^{1+p_1-p_2}} \oint_{C_\epsilon} \frac{dt_2}{t_2^{1+2p_2}} Z_{\text{AdS}}(t_1, t_2) \quad (2.32)$$

Employing a saddle point evaluation would give a $\frac{1}{N}$ expansion of the characters. The leading term in this expansion should agree with the results of [63].

2.3 Excited bound states of two AdS giants

To extend the analysis of the previous subsection we consider excited giant graviton bound-states obtained by attaching open strings to the brane system. Open strings are described by words composed of $O(\sqrt{N})$ letters [64]. The letters can be any of the scalar or fermion

fields of $\mathcal{N} = 4$ super Yang-Mills theory, the field strength tensor or covariant derivatives of them. Attach m_{11} open strings described by the words $W_{A_1}, W_{A_2}, \dots, W_{A_{m_{11}}}$ to the first brane (of momentum p_1). Since indices of the first brane are symmetrized by the $\bar{\varphi}^i$, φ_i fields attaching these strings is achieved by including the factor

$$\prod_{r=1}^{m_{11}} \bar{\varphi}^{i_r} (W_{A_r})_{i_r}^{j_r} \varphi_{j_r} \quad (2.33)$$

in the integrand of the generating function. We also attach m_{22} open strings, described by the words $W_{B_1}, W_{B_2}, \dots, W_{B_{m_{22}}}$, to the second brane (of momentum p_2). Since these fields are in the second row of the Young diagram describing the brane system, we must anti-symmetrize these words with the Z field that lives in the same column but in the first row. Thus, attaching these strings is achieved by including the factor

$$\prod_{s=1}^{m_{22}} (\bar{\varphi}^{i_s} Z_{i_s}^{j_s} \varphi_{j_s} \bar{\phi}^{k_s} (W_{B_s})_{k_s}^{l_s} \phi_{l_s} - \bar{\varphi}^{i_s} Z_{i_s}^{j_s} \phi_{j_s} \bar{\phi}^{k_s} (W_{B_s})_{k_s}^{l_s} \varphi_{l_s}) \quad (2.34)$$

in the integrand. We can also attach strings that stretch between the two branes. To do this, we start with the product of a factor which attaches a string to the first brane and a factor which attaches a string to the second brane. From this we obtain a pair of strings stretched between the branes by permuting column indices, thereby swapping the endpoints of the two strings. There is a non-trivial fact [30] about the Hilbert space of states on a two giant graviton brane bound state that this construction must respect. The world volume theory of a system of p branes is a $U(p)$ gauge theory. One consequence of this gauge symmetry is the Gauss Law, which states that for a giant graviton brane, which has a compact world volume, the total charge on the brane vanishes. In the open string language, this forces the number of strings leaving any particular brane to be equal to the number of strings terminating on the brane. We implement the Gauss Law constraint by adding open strings in pairs, one stretching from 1 to 2 and one from 2 to 1. Describe the m_{12} open strings stretching from the first to the second brane using the words $W_{C_1}, W_{C_2}, \dots, W_{C_{m_{12}}}$ and describe the m_{21} ($= m_{12}$) open strings stretching from the second to the first brane using the words $W_{D_1}, W_{D_2}, \dots, W_{D_{m_{21}}}$. Thus the generating function for this class of excited bound states, using an obvious dot product notation, is

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2, \{W\}) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi} \\ &\times e^{t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2^2 \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - t_2^2 \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi} \prod_{r=1}^{m_{11}} \bar{\varphi} \cdot W_{A_r} \cdot \varphi \\ &\times \prod_{s=1}^{m_{22}} (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi) \\ &\times \prod_{t=1}^{m_{12}} (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{D_t} \cdot \varphi \bar{\varphi} \cdot W_{C_t} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{D_t} \cdot \varphi \bar{\varphi} W_{C_t} \varphi) \end{aligned} \quad (2.35)$$

A graphical representation of this generating function is given in figure 1. The dependence on t_1 and t_2 continues to tell us about the brane bound state with no open strings attached,




Figure 1. This figure gives a graphical representation of the excited brane bound states described by the generating function (2.35). The node on the left represents the brane with momentum p_1 while the node on the right is for the brane of momentum p_2 .

i.e. the coefficient of the term $t_1^{p_1-p_2} t_2^{2p_2}$ is a two giant bound state for which the first brane has momentum p_1 and the second momentum p_2 . To simplify our discussion, we focus on the $\text{su}(2)$ sector, in which case the open string words are constructed using two complex adjoint scalars, Z and Y . Further, the last and first letter of the word are both Y s. By acting on these operators with the dilatation operator, the dynamics of the open string words is described by a Hamiltonian for a spin chain, with non-trivial boundary conditions. In section 3 we consider this Hamiltonian.

Operators dual to a giant graviton system with strings attached were constructed in [31]. For operators constructed using n Z fields and q open string words, the construction of [31] starts with a chain of subgroups $S_n \subset S_{n+1} \subset S_{n+2} \subset \dots \subset S_{n+q}$. Using this chain of subgroups, we can define a restriction of a representation R of S_{n+q} to a representation r of S_n , by giving the representation obtained as one successively restricts to each subgroup in the chain. The sequence of representations of the subgroups specify where the open strings are attached, and the representation r defines the giant graviton bound state. All of this information can be summarized by decorating the Young diagram label of the operator. For example, a two AdS giant graviton bound state with four strings attached to the first giant graviton is given by the operator

$$\chi_{\begin{array}{cccc|ccc} & & & & & 4 & 3 & 2 & 1 \\ & & & & & & & & \end{array}} \quad (2.36)$$

The chain of representations used to define the operator are obtained by dropping the numbered boxes in order. For the example above, we have the following 5 representations defining the restriction from the original Young diagram R to the Young diagram⁶ R'''' defining the giant bound state

$$R \equiv \begin{array}{cccc|ccc} & & & & & 4 & 3 & 2 & 1 \\ & & & & & & & & \end{array} \rightarrow \begin{array}{cccc|ccc} & & & & & 4 & 3 & 2 \\ & & & & & & & \end{array} \rightarrow \begin{array}{cccc|ccc} & & & & & 4 & 3 \\ & & & & & & \end{array} \rightarrow \begin{array}{cccc|ccc} & & & & & 4 \\ & & & & & & \end{array} \rightarrow \begin{array}{cccc|ccc} & & & & & & & & \end{array} \equiv R''''$$

⁶The number of primes appearing on R'''' tells us how many open string boxes were dropped from R .

Using this set of restrictions the restricted Schur polynomial of [31] is given by

$$\chi_{R,R'''}(Z, W^{(i)}) = \frac{1}{n!} \sum_{\sigma \in S_{14}} \text{Tr}_{R'''}(\Gamma_R(\sigma)) Z_{i_{\sigma(14)}}^{i_{14}} \cdots Z_{i_{\sigma(5)}}^{i_5} (W_4)_{i_{\sigma(4)}}^{i_4} (W_3)_{i_{\sigma(3)}}^{i_3} (W_2)_{i_{\sigma(2)}}^{i_2} (W_1)_{i_{\sigma(1)}}^{i_1} \quad (2.37)$$

In [31], $\text{Tr}_{R'''}(\Gamma_R(\sigma))$ is called a restricted character because the trace over R to define the character has been restricted to the subspace R''' . We could also attach two strings to each giant graviton brane

$$\chi \begin{array}{|c|c|c|c|c|c|c|c|} \hline & & & & & & 3 & 1 \\ \hline & & & & 4 & 2 & & \\ \hline \end{array} \quad (2.38)$$

by making use of the following restrictions

$$R \equiv \begin{array}{|c|c|c|c|c|c|c|c|} \hline & & & & & & 3 & 1 \\ \hline & & & & 4 & 2 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|c|c|} \hline & & & & & & 3 \\ \hline & & & & 4 & 2 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|c|} \hline & & & & & 3 \\ \hline & & & & 4 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|c|} \hline & & & & 4 \\ \hline & & & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline \end{array} \equiv R''' \quad (2.39)$$

To stretch the strings between the branes we use a different restriction for the row and column index of the restricted character. The row restriction specifies the open string start points while the column restriction specifies the end points.

What is the relation between the operators of [31] and those constructed above? In appendix A we explicitly evaluate the form of the operators constructed in [31] and compare them to the operators which follow from the generating function (2.35). Simple examples immediately confirm that

1. In the case that $m_{22} = 0$, $m_{12} = 0 = m_{21}$ with m_{11} arbitrary, the generating function (2.35) precisely reproduces the restricted Schur polynomials constructed in [31].
2. In the case that $m_{11} = 0$, $m_{12} = 0 = m_{21}$ with m_{22} arbitrary, the generating function (2.35) again precisely reproduces the restricted Schur polynomials constructed in [31].

This agreement is not an accident and we can prove that, for the cases described in 1. and 2. above, the two constructions agree. A central result of [31], is a rule for the derivative of a restricted Schur polynomial with respect to an open string. The idea behind the proof is to show that the operators constructed from the generating function (2.35) obey this derivative rule.

To state the derivative rule of [31] we need to review a few facts. When strings are added to the giant system, the order in which strings are added matters.⁷ Bearing in mind how the operator was constructed, this is not surprising: changing the order in which open strings are attached changes the set of restrictions used to define the restricted character and the restricted character depends sensitively on which subspace is traced. If we differentiate with respect to the last open string added, the open string together with its box is removed from the restricted Schur polynomial and the remaining polynomial is multiplied by the factor c of the box removed. Recall that the factor of a box in row i and column j is $c = N - i + j$.

⁷For the specific case that we attach all the open strings to a single giant, as in (2.36) it turns out that attaching the strings in different orders gives the same operator. In this case and only this case the order in which strings are attached does not matter.

We can extend this result to operators corresponding to a bound state of two giant gravitons with all strings attached to the second brane. In this case the derivative rule is

$$\frac{\partial}{\partial (W_{B_1})_a} \chi_{\begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline \hline \hline \end{array}} = (N + p_2 + q - 2) \chi_{\begin{array}{|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|c|} \hline \hline \hline \end{array}} \quad (2.42)$$

Note that the coefficient on the right hand side is the coefficient of (2.40) with p_1 replaced by p_2 minus one. The extra minus one is because the open string now occupies the second row. To derive this rule for the operators derived from the generating function, consider

$$\begin{aligned}
& \frac{\partial}{\partial (W_{B_1})_a} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} \\
& \times \prod_{s=1}^q (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi) \\
& = \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot \varphi) \\
& \quad \times \prod_{s=2}^q (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi) \\
& = (N + p_2 + q - 1) \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} \\
& \quad \times \bar{\varphi} \cdot Z \cdot \varphi \prod_{s=2}^q (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi) \\
& \quad - \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \left(\frac{e^{S_0}}{\pi^{2N}} \right) \\
& \quad \times \varphi_a \frac{\partial}{\partial \phi_a} \left[\bar{\varphi} \cdot Z \cdot \phi e^{S_1} \prod_{s=2}^q (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi) \right] \quad (2.43)
\end{aligned}$$

Using the obvious facts

$$\varphi_a \frac{\partial}{\partial \phi_a} \left[\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi \right] = 0 \quad (2.44)$$

$$\varphi_a \frac{\partial}{\partial \varphi_a} \left[t_2^2 (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi) \right] = 0 \quad (2.45)$$

we find

$$\begin{aligned}
& \frac{\partial}{\partial (W_{B_1})_a^a} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} \prod_{s=1}^q (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi) \\
&= (N + p_2 + q - 2) \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} \\
&\quad \times \bar{\varphi} \cdot Z \cdot \varphi \prod_{s=2}^q (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_s} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_s} \cdot \varphi)
\end{aligned} \tag{2.46}$$

The extra factor of $\bar{\varphi} \cdot Z \cdot \varphi$ is simply adding one box to the first row of the Young diagram labelling the operator. The open string W_{B_1} was located in a column of length 2, with the

other box in the column occupied by a Z field. After we differentiate with respect to W_{B_1} the second box in the column is left over and this is precisely the extra factor $\bar{\varphi} \cdot Z \cdot \varphi$. This reproduces the correct derivative rule in complete detail. As above, since there is no term in the operators considered that is independent of the open string word W_{B_1} , the equality of these derivatives implies the equality of the operators. Thus, as long as we attach strings to a specific giant, the generating function we have written down is exactly reproducing the restricted Schur polynomials constructed in [31].

This agreement between the operators obtained from the generating function (2.35) and those constructed in [31] does not hold in general. Indeed, it is simple to verify that

3. In the case that $m_{12} = 0 = m_{21}$ with both $m_{11} \neq 0$ and $m_{22} \neq 0$ but otherwise arbitrary, the operators defined by the generating function (2.35) do not agree with the restricted Schur polynomials constructed in [31].

As a simple example, the operator with $p_1 = 1$ and $p_2 = 0$, with a single string attached to each row, is given by

$$\begin{aligned} & \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} \bar{\varphi} \cdot W_A \cdot \varphi (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_B \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_B \cdot \varphi) \\ &= \text{Tr}(W_A) \text{Tr}(W_B) \text{Tr}(Z) + \text{Tr}(ZW_A) \text{Tr}(W_B) - \text{Tr}(W_A) \text{Tr}(W_B Z) - \text{Tr}(W_A Z W_B) \end{aligned} \quad (2.47)$$

while the construction of [31] gives

$$\begin{aligned} \chi_{\begin{smallmatrix} \square & A \\ B & \end{smallmatrix}} &= \text{Tr}(W_A) \text{Tr}(W_B) \text{Tr}(Z) + \frac{1}{2} \text{Tr}(W_A W_B) \text{Tr}(Z) - \text{Tr}(W_A) \text{Tr}(W_B Z) \\ &+ \frac{1}{2} \text{Tr}(ZW_A) \text{Tr}(W_B) - \frac{1}{2} \text{Tr}(W_A Z W_B) - \frac{1}{2} \text{Tr}(W_A W_B Z) \end{aligned} \quad (2.48)$$

This proves that the restricted Schur construction of [31] and the construction using the generating function (2.35) are not the same. There is a simple and general way to see that these two sets of operators are different. The operators produced using the generating function (2.35) treats all open strings with the same orientation identically. Thus, we can permute any of the strings of the same letter⁸ and this is a symmetry of the operator. This symmetry is not present in the restricted Schur polynomial construction of [31].

Describing open string excitations of the giant graviton system in terms of open strings has the advantage that we can make contact with a worldsheet description of the open string [32]. Another possibility is that we excite of the half BPS giant graviton bound state by including a second field Y in the description of the operator.⁹ This description is naturally related to the restricted Schur polynomial construction developed in [19, 20]. Consider operators constructed using n Z fields and m Y fields. The restricted Schur polynomial $\chi_{R,(r,s)\alpha\beta}(Z, Y)$ construction of [19, 20] uses a representation $R \vdash m+n$ to organize

⁸By this we mean that we can permute the W_{A_i} 's among themselves, the W_{B_i} 's among themselves and so on.

⁹We can include the complete set of fields in $\mathcal{N} = 4$ super Yang Mills theory. This restriction to a single field Y is for simplicity only.

the complete set of fields, as well as a representation $r \vdash n$ for the Z fields and a representation $s \vdash m$ for the Y fields. In addition, there are multiplicity labels α, β needed¹⁰ because in general there is more than one way in which the $S_n \times S_m$ representation (r, s) can be obtained after restricting the S_{n+m} representation R to the $S_n \times S_m$ subgroup. This description of the local operators does not immediately have a visible open string interpretation. By diagonalizing the one loop dilatation operator, the studies [42–45] discovered an alternative basis, known as the Gauss graph basis. The operators of this basis, called Gauss graph operators, are eigenstates of the one loop dilatation operator. These eigenstates are labelled by graphs, which provide a diagrammatic picture of the open strings stretching between the different giant graviton branes. The complete basis of operators manifestly obey the Gauss Law of the brane world volume theory, motivating the terminology “Gauss graph basis”. We will now argue that it is the Gauss graph operators that correspond to the operators constructed from our generating function. Towards this end, consider the generating function

$$\begin{aligned}
 Z_{\text{AdS}}(t_1, t_2, Y) = & \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi} \\
 & \times e^{t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2^2 \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - t_2^2 \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi} (\bar{\varphi} \cdot Y \cdot \varphi)^{m_{11}} \\
 & \times (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi)^{m_{22}} \\
 & \times (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \varphi)^{m_{12}} \quad (2.49)
 \end{aligned}$$

which is obtained by replacing $W_* \rightarrow Y$ in (2.35). The bound states described by this generating function have m_{11} strings attached to the brane corresponding to the first row, m_{22} strings attached to the brane corresponding to the second row, m_{12} strings stretching from the first to the second branes and $m_{21} = m_{12}$ strings stretched from the second to the first. The Gauss Law is clearly manifest in this description since the above expression is only defined for $m_{12} = m_{21}$. In addition, the construction is symmetric under swaps of any strings that have the same orientation. This is a symmetry that the Gauss graph operators enjoy and this agreement of the symmetry enjoyed by the two constructions is a key piece of evidence suggesting that the operators constructed from the generating function (2.49) are the Gauss graph operators. The Gauss graph operators with all strings attached to a single brane agree with the restricted Schur polynomials [43, 45] with all open strings attached in a given row. Consequently, the results we have obtained above already establish that restricted Schur polynomials and Gauss graph operators coincide when the open strings are all attached to a single brane, i.e. when there are no strings stretching between branes.

For a detailed comparison of operators that involve stretched strings, study the simpler expression obtained for $m_{11} = 0 = m_{22}$ and $m_{12} = 1 = m_{21}$

$$\begin{aligned}
 G_{\text{AdS}}(t_1, t_2) = & \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} \\
 & \times (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \varphi) \quad (2.50)
 \end{aligned}$$

¹⁰To define the relevant restricted character, we need one multiplicity label for the row index and one for the column index, that we restrict to.

The dependence on t_1 and t_2 continues to keep track of the momentum carried by each brane, exactly as above. A simple computation proves that

$$\begin{aligned}
 G_{\text{AdS}}(t_1, t_2) = & \text{Tr}(Z)\text{Tr}(Y^2) - \text{Tr}(ZY)\text{Tr}(Y) + t_1 (\text{Tr}(Z)^2\text{Tr}(Y^2) + \text{Tr}(Z^2)\text{Tr}(Y^2) + \text{Tr}(Z)\text{Tr}(ZY^2) \\
 & + \text{Tr}(Z^2Y^2) - \text{Tr}(ZY)\text{Tr}(Z)\text{Tr}(Y) - \text{Tr}(Z^2Y)\text{Tr}(Y) - \text{Tr}(ZY)^2 - \text{Tr}(ZYZY)) \\
 & + t_2 (\text{Tr}(Z)^3\text{Tr}(Y^2) + 2\text{Tr}(Z^2)\text{Tr}(Y^2Z) - \text{Tr}(Z)^2\text{Tr}(Y)\text{Tr}(YZ) + 2\text{Tr}(Z)^2\text{Tr}(Y^2Z) \\
 & - \text{Tr}(YZ)\text{Tr}(YZ^2) - \text{Tr}(Y^2)\text{Tr}(Z^3) - 2\text{Tr}(Z)\text{Tr}(YZ)^2 - 2\text{Tr}(Z)\text{Tr}(Y)\text{Tr}(YZ^2) \\
 & - 2\text{Tr}(Z)\text{Tr}(Y^2Z^2) + \text{Tr}(Z)\text{Tr}(YZYZ) + 2\text{Tr}(Y)\text{Tr}(YZ)\text{Tr}(Z^2) + \text{Tr}(Y)\text{Tr}(YZ^3) \\
 & - 2\text{Tr}(Y^2Z^3) + 2\text{Tr}(YZYZ^2)) + \dots
 \end{aligned} \tag{2.51}$$

In the appendices we prove that the Gauss graph operators are given by

$$\begin{aligned}
 \left(\square, \bullet \right) &= \text{Tr}(Z)\text{Tr}(Y^2) - \text{Tr}(ZY)\text{Tr}(Y) \\
 \left(\square\square, \bullet \right) &= \text{Tr}(Z)^2\text{Tr}(Y^2) + \text{Tr}(Z^2)\text{Tr}(Y^2) + \text{Tr}(Z)\text{Tr}(ZY^2) + \text{Tr}(Z^2Y^2) \\
 &\quad - \text{Tr}(ZY)\text{Tr}(Z)\text{Tr}(Y) - \text{Tr}(Z^2Y)\text{Tr}(Y) - \text{Tr}(ZY)^2 - \text{Tr}(ZYZY) \\
 \left(\square\square\square, \bullet \right) &= \text{Tr}(Z)^3\text{Tr}(Y^2) + 2\text{Tr}(Z^2)\text{Tr}(Y^2Z) - \text{Tr}(Z)^2\text{Tr}(Y)\text{Tr}(YZ) + 2\text{Tr}(Z)^2\text{Tr}(Y^2Z) \\
 &\quad - \text{Tr}(YZ)\text{Tr}(YZ^2) - \text{Tr}(Y^2)\text{Tr}(Z^3) - 2\text{Tr}(Z)\text{Tr}(YZ)^2 - 2\text{Tr}(Z)\text{Tr}(Y)\text{Tr}(YZ^2) \\
 &\quad - 2\text{Tr}(Z)\text{Tr}(Y^2Z^2) + \text{Tr}(Z)\text{Tr}(YZYZ) + 2\text{Tr}(Y)\text{Tr}(YZ)\text{Tr}(Z^2) + \text{Tr}(Y)\text{Tr}(YZ^3) \\
 &\quad - 2\text{Tr}(Y^2Z^3) + 2\text{Tr}(YZYZ^2)
 \end{aligned} \tag{2.52}$$

in complete agreement with (2.51).

With this encouraging evidence in hand, we would now like to prove that the operators generated by (2.49) are indeed identical to the Gauss graph operators. First in appendix C we prove that, at large N , operators derived from the generating function for arbitrary m_{11} and m_{22} agree with the Gauss graph operators. Thus, all that is left is to consider operators with non-zero m_{12} . The proof introduces a graph changing operator: acting on a given Gauss graph, we can easily generate any other Gauss graph operator. Then, given the equality between our operators and the Gauss graph operators when all strings are attached to a definite brane (i.e. there are no stretched string states), and the fact that we can generate any boundstate with any number of strings stretched between the giants with the graph changing operator, the proof is complete.

For the proof that follows, it is necessary to briefly review the construction of the Gauss graph operators. The restricted Schur polynomial $\chi_{R,(r,s)\alpha\beta}(Z, Y)$ constructed using n Z fields and m Y fields is labelled by three Young diagrams $R \vdash m+n$, $r \vdash n$ and $s \vdash m$ and two multiplicity labels α, β . The Gauss graph operators $O_{R,r}(\sigma)$ have the same R and r labels, but the s label and the multiplicity labels are replaced by a permutation σ . If we think about each individual Y field as an open string, to specify a state of the Y fields we need to specify a configuration of the open strings. This can be achieved with a permutation, as illustrated in figure 2. This is the permutation σ labelling the Gauss graph operator. From the figure it is clear that swapping endpoints of open strings terminating on the brane and swapping endpoints of open strings originating from the brane is a symmetry




Figure 2. Any open string configuration can be mapped to a labelled graph as shown. The two bold horizontal lines in the figure on the right are identified. Rows of R form nodes in the graph. For the graph shown, R has three rows. There are 7 open strings, so the operator is constructed using 7 Y fields. The graph determines a permutation, so each open string configuration maps to a permutation. For the graph shown the permutation, in cycle notation, is $\sigma = (24)(536)$. A nice example is the configuration in which all open strings loop back to the brane they start from, described by the identity permutation.

of the open string configuration. To correctly enumerate distinct physical states we must divide out this symmetry so that each configuration corresponds to an element of a double coset. For the configuration shown in figure 2 the relevant double coset is $H \setminus S_7 / H$ where H is the group of symmetries for the end points, given by $H = S_3 \times S_2 \times S_2$.

The transformation to the Gauss graph basis is given by

$$O_{R,r}(\sigma) = \sum_{s \vdash m} \sum_{\alpha, \beta} C_{\alpha\beta}^s(\sigma) \chi_{R,(r,s)\alpha\beta} \quad (2.53)$$

where

$$C_{\alpha\beta}^s(\tau) = |H| \sqrt{\frac{d_s}{m!}} \sum_{k,l=1}^{d_s} \Gamma_{kl}^{(s)}(\tau) B_{k\alpha}^{s \rightarrow 1_H} B_{l\beta}^{s \rightarrow 1_H} \quad (2.54)$$

This last formula makes use of branching coefficients, which are defined as follows

$$\frac{1}{|H|} \sum_{\gamma \in H} \Gamma_{ik}^{(s)}(\gamma) = \sum_{\mu} B_{i\mu}^{s \rightarrow 1_H} B_{k\mu}^{s \rightarrow 1_H} \quad (2.55)$$

where the index μ on the right hand side is a multiplicity label.

As we have just reviewed, the Gauss graph operator is labelled by a permutation σ that specifies how the open strings are stretched between the giant gravitons. We can easily

rewrite the formula (2.49) in terms of σ as follows ($m = m_{11} + m_{12} + m_{21} + m_{22}$)

$$\begin{aligned}
 Z_{\text{AdS}}(t_1, t_2, Y, \sigma) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi} \\
 &\quad \times e^{t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2^2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l} \prod_{i=1}^{m_{11}+m_{12}} \bar{\varphi}^{b_i} Y_{b_i}^{a_i} \varphi_{a_{\sigma(i)}} \\
 &\quad \times \prod_{j=m_{11}+m_{12}+1}^m (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi}^{b_j} Y_{b_j}^{a_j} \phi_{a_{\sigma(j)}} - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi}^{b_j} Y_{b_j}^{a_j} \varphi_{a_{\sigma(j)}})
 \end{aligned}$$

It is clear that σ specifies the row to which the upper indices of the Y fields belong i.e. σ specifies how the Y excitations connect to the giants. The graph changing operator, acting on a system with a total of m Y excitations, has the form

$$\hat{G}(\sigma) = Y_{j_{\sigma(1)}}^{i_1} \cdots Y_{j_{\sigma(m)}}^{i_m} \frac{\partial}{\partial Y_{j_1}^{i_1}} \cdots \frac{\partial}{\partial Y_{j_m}^{i_m}} \quad (2.56)$$

It is simple to verify that

$$\hat{G}(\sigma_1) Z_{\text{AdS}}(t_1, t_2, Y, \sigma_2) = Z_{\text{AdS}}(t_1, t_2, Y, \sigma_1 \sigma_2) \quad (2.57)$$

We will now argue that this graph changing operator has exactly the same action on the Gauss graph operators. When R has only two rows we have no multiplicity labels so that, up to a normalization factor of $\frac{|H|}{m!}$ which we drop, we have¹¹

$$\begin{aligned}
 O(\sigma) &= \sum_{s \vdash m} \sqrt{d_s} \Gamma_{jk}^{(s)}(\sigma) B_j^{s \rightarrow 1_H} B_k^{s \rightarrow 1_H} O_{R,(r,s)} \\
 &= \frac{1}{\prod_i m_i!} \sum_{s \vdash m} \sum_{\alpha \in \prod_i S_{m_i}} \sqrt{d_s} \Gamma_{jk}^{(s)}(\sigma) \Gamma_{kj}^{(s)}(\alpha) O_{R,(r,s)} \\
 &= \frac{1}{\prod_i m_i!} \sum_{s \vdash m} \sum_{\alpha \in \prod_i S_{m_i}} \sqrt{d_s} \chi_s(\sigma \alpha) O_{R,(r,s)} \quad (2.58)
 \end{aligned}$$

where $\chi_s(\sigma)$ is the character of $\sigma \in S_m$ in irrep s . Now, again up to a normalization that plays no role in this proof, we have

$$O_{R,(r,s)} = \sum_{\rho \in S_{n+m}} \sum_{\sigma_1 \in S_n} \sum_{\sigma_2 \in S_m} \frac{1}{\sqrt{d_s}} \chi_r(\sigma_1) \chi_s(\sigma_2) \chi_R(\sigma_1 \circ \sigma_2) \text{Tr}(\rho Z^{\otimes n} Y^{\otimes m}) \quad (2.59)$$

In writing this formula we have explicitly used the crucial fact that restricted Schur polynomials with only two rows carry no multiplicity labels.¹² Using this formula, as well as the character orthogonality identity, which says

$$\sum_{s \vdash m} \chi_s(g) \chi_s(h) = |[g]| \delta_{[g]}[h] \quad (2.61)$$

¹¹For a detailed discussion and derivation of this formula, see [45].

¹²In the case that there are no multiplicities we can write the projection operator, which projects to the (r, s) irreducible representation of $S_n \times S_m$, within the carrier space of R as

$$P_{R \rightarrow (r,s)} = \frac{d_r d_s}{n! m!} \sum_{\sigma_1 \in S_n} \sum_{\sigma_2 \in S_m} \chi_r(\sigma_1) \chi_s(\sigma_2) \Gamma_R(\sigma_1 \circ \sigma_2) \quad (2.60)$$

we have

$$O(\sigma) = \sum_{\rho \in S_{n+m}} \sum_{\sigma_1 \in S_n} \sum_{\alpha \in S_{m_1} \times S_{m_2}} \chi_r(\sigma_1) \chi_R(\sigma_1 \circ \sigma \alpha \rho) \text{Tr}(\rho Z^{\otimes n} Y^{\otimes m}) \quad (2.62)$$

Using this last formula it is simple to verify that

$$\hat{G}(\sigma_1)O(\sigma_2) = O(\sigma_1\sigma_2) \quad (2.63)$$

In appendix C we argue that

$$\sum_{p_1, p_2} t_1^{p_1-p_2} t_2^{p_2} O_{(p_1, p_2)}(\mathbf{1}) = Z_{\text{AdS}}(t_1, t_2, Y, \mathbf{1}) \quad (2.64)$$

and since the graph changing operator acts in the same way on both sides of this equation, the proof is complete.

2.4 Bound states of two sphere giants

A bound state of two sphere giants, one of momentum p_1 and a second of momentum p_2 with $p_2 \leq p_1$ corresponds to a Schur polynomial labeled by a Young diagram with two columns. There are p_1 boxes in the first column and p_2 boxes in the second. To describe giant gravitons, both p_1 and p_2 must be order N . The Young symmetrizer must antisymmetrize over all indices in column 1, and all indices in column 2, as well as symmetrize the pair of indices belonging to each row. In contrast to the AdS giants, it is now the antisymmetrization to be performed that represents the bulk of the work. This antisymmetrization of indices can again be accomplished with the help of a Gaussian integral, but now we integrate over Grassmann variables. Again it is a simple application of Wick's theorem to learn that

$$\int d^N \bar{\xi} \int d^N \xi e^{-\bar{\xi}^a \xi_a} \bar{\xi}^{i_1} \xi_{j_1} \bar{\xi}^{i_2} \xi_{j_2} \cdots \bar{\xi}^{i_k} \xi_{j_k} = \sum_{\sigma \in S_k} \text{sgn}(\sigma) \delta_{j_{\sigma(1)}}^{i_1} \delta_{j_{\sigma(2)}}^{i_2} \cdots \delta_{j_{\sigma(k)}}^{i_k} \quad (2.65)$$

$$d^N \xi \equiv d\xi_1 d\xi_2 \cdots d\xi_N \quad d^N \bar{\xi} \equiv d\bar{\xi}^N \cdots d\bar{\xi}^2 d\bar{\xi}^1 \quad (2.66)$$

To write down the generating function for Young diagrams $R = (2^{p_2}, 1^{p_1-p_2})$ that have two columns, one of length p_1 and one of length p_2 , we perform the symmetrization in each row by hand and use $(\psi_i, \bar{\psi}^i)$ to antisymmetrize indices in the first column and $(\xi_i, \bar{\xi}^i)$ to perform the antisymmetrization of indices in the second column. The two giant graviton boundstate generating function can then be written as follows

$$\begin{aligned} Z_S(t_1, t_2) &= \sum_{p_1 \geq p_2} \sum_{p_2=1}^{\infty} t_2^{p_2} t_1^{p_1-p_2} \frac{(p_1 + p_2)!}{d_{(2^{p_2}, 1^{p_1-p_2})} p_2! (p_1 - p_2)!} \chi_{(2^{p_2}, 1^{p_1-p_2})}(Z) \\ &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi \\ &\quad \times e^{-\bar{\psi}^i \psi_i - \bar{\xi}^i \xi_i - t_1 \bar{\psi}^i Z_i^j \psi_j + t_2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l} \end{aligned} \quad (2.67)$$

In appendix A we check this result by performing the integral to second order in t_1 and t_2 and show that we reproduce the correct formulas for the relevant Schur polynomials.

As was the case as for the AdS giants, this generating function can also be interpreted as a generating function for characters of the symmetric group $S_{p_1+p_2}$ in the representation labelled by Young diagram $R = (2^{p_2}, 1^{p_1-p_2})$. The generating function is

$$Z_S(t_1, t_2) = \sum_{p_1=p_2}^{\infty} \sum_{p_2=1}^{\infty} \sum_{\{n_i\}} \frac{t_2^{2p_2} t_1^{p_1-p_2}}{1^{n_1} n_1! 2^{n_2} n_2! \dots (p_1+p_2)^{n_{p_1+p_2}} n_{(p_1+p_2)}!} \frac{(p_1+1)!}{(p_1-p_2+1)(p_1-p_2)!} \times \chi_{(2^{p_2}, 1^{p_1-p_2})}(\{i^{n_i}\}) \text{Tr}(Z)^{n_1} \text{Tr}(Z^2)^{n_2} \dots \text{Tr}(Z^{p_1+p_2})^{n_{p_1+p_2}} \quad (2.68)$$

where $\sum_{\{n_i\}}$ denotes a sum over the conjugacy classes of $S_{p_1+p_2}$. To build some intuition for this formula, expand to linear order in t_1 and t_2 to find

$$\begin{aligned} Z_S(t_1, t_2) &= 1 + t_1 \chi_{\square}((\cdot)) \text{Tr}(Z) + t_2^2 \chi_{\square\square}((\cdot)(\cdot)) \text{Tr}(Z)^2 + t_2^2 \chi_{\square\square}((\cdot\cdot)) \text{Tr}(Z^2) + \dots \\ &= 1 + t_1 \text{Tr}(Z) + t_2^2 \text{Tr}(Z)^2 + t_2^2 \text{Tr}(Z^2) + \dots \end{aligned} \quad (2.69)$$

reproducing the correct characters for S_1 and S_2

$$\chi_{\square}((\cdot)) = 1 \quad \chi_{\square\square}((\cdot)(\cdot)) = 1 \quad \chi_{\square\square}((\cdot\cdot)) = 1 \quad (2.70)$$

Following our discussion for characters with long rows, we could easily use the result (2.68) to develop a formula for characters in representations labelled by Young diagrams with two columns.

2.5 Excited bound states of two sphere giants

In this section we extend the results obtained above, constructing generating functions for operators dual to excited sphere giant gravitons. Again consider a boundstate of two sphere giant gravitons, the first with momentum p_1 and the second with momentum p_2 . The boundstate is excited by attaching open strings. The generating function for this class of bound states is

$$\begin{aligned} Z_S(t_1, t_2, \{W\}) &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{-\bar{\psi} \cdot \psi - \bar{\xi} \cdot \xi} \\ &\times e^{-t_1 \bar{\psi} \cdot Z \cdot \psi + t_2^2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l} \prod_{r=1}^{m_{11}} \bar{\psi} \cdot W_{A_r} \cdot \psi \\ &\times \prod_{s=1}^{m_{22}} (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot W_{B_s} \cdot \xi - \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot W_{B_s} \cdot \psi) \\ &\times \prod_{t=1}^{m_{12}} (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot W_{D_t} \cdot \psi \bar{\psi} \cdot W_{C_t} \cdot \xi + \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot W_{D_t} \cdot \psi \bar{\psi} \cdot W_{C_t} \cdot \psi) \end{aligned} \quad (2.71)$$

The signs in the final two lines would both have been positive if we had kept all barred variables to the left with $\bar{\psi}$'s before $\bar{\xi}$'s and all unbarred variables to the right with ψ 's before ξ 's. The dependence on t_1 and t_2 again tells us about the quantum numbers of the brane boundstate with no open strings attached: the coefficient of the term $t_1^{p_1-p_2} t_2^{2p_2}$ describes the case that the first brane has momentum p_1 and the second momentum p_2 . We again focus mainly on the $\text{su}(2)$ sector in which case the open string words are constructed

using two complex adjoint scalars, Z and Y , with the last and first letter of the word both Y s. The Gauss Law is again respected because the open strings are added to the bound state in pairs, with one string (W_{C_t}) stretching from brane 1 to brane 2 and another (W_{D_t}) stretching from 2 to 1. In a discussion which closely parallels the discussion for the AdS giant bound state, we can again prove that the operators with strings attached only to brane 1 or only to brane 2 agree with the construction given in [31]. The proof again proceeds by verifying that derivatives with respect to open strings words agree with the rule derived in [31]. We will sketch the case that strings are attached to the giant of momentum p_1 since it is interesting to see the role of the Grassman nature of the integration variables. Recall that in the case of the AdS giant (a long row), the factor of the box containing W_{A_1} is $N + (q - 1)$, since each time we move along the Young diagram, to the right, we add 1 to the factor. In the case of the sphere giant (a long column), the factor of the box containing W_{A_1} is $N - (q - 1)$, since each time we move down the Young diagram we subtract 1 from the factor. It is the change in the sign of the $q - 1$ contribution to the factor that we want to demonstrate. Towards this end, consider ($S_0 = -\bar{\psi} \cdot \psi - \bar{\xi} \cdot \xi$, $S_1 = -t_1 \bar{\psi} \cdot Z \cdot \psi + t_2^2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l$)

$$\begin{aligned}
 & \frac{\partial}{\partial(W_{A_1})_a} \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{S_0} e^{S_1} \prod_{r=1}^q \bar{\psi} \cdot W_{A_r} \cdot \psi \\
 &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{S_0} e^{S_1} \bar{\psi} \cdot \psi \prod_{r=2}^q \bar{\psi} \cdot W_{A_r} \cdot \psi \\
 &= - \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi \left(\bar{\psi}^i \frac{\partial}{\partial \bar{\psi}^i} e^{S_0} \right) e^{S_1} \prod_{r=2}^q \bar{\psi} \cdot W_{A_r} \cdot \psi \\
 &= \left(N - \bar{\psi}^i \frac{\partial}{\partial \bar{\psi}^i} \right) \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{S_0 + S_1} \prod_{r=2}^q \bar{\psi} \cdot W_{A_r} \cdot \psi \quad (2.72)
 \end{aligned}$$

where the negative sign in the expression in brackets on the last line above is exactly what is required to get the correct factor. This sign comes from the Grassman nature of $\bar{\psi}^i$ since

$$\frac{\partial}{\partial \bar{\psi}^i} \bar{\psi}^i = N - \bar{\psi}^i \frac{\partial}{\partial \bar{\psi}^i} \quad (2.73)$$

Again, exactly as we found for the excited AdS giant bound states, by constructing and comparing operators we learn that the restricted Schur construction of [31] is in general not the same as the construction using the generating function (2.71).

The operators constructed using the generating function (2.71) are again symmetric under swapping strings of the same orientation. This is a strong hint that they are closely related to the Gauss graph operators. This expectation turns out to be correct. The

relevant generating function is

$$\begin{aligned}
Z_S(t_1, t_2, Y) = & \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{-\bar{\psi} \cdot \psi - \bar{\xi} \cdot \xi} \\
& \times e^{-t_1 \bar{\psi} \cdot Z \cdot \psi + t_2^2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l} \prod_{r=1}^{m_{11}} \bar{\psi} \cdot Y \cdot \psi \\
& \times \prod_{s=1}^{m_{22}} (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot Y \cdot \xi - \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot Y \cdot \psi) \\
& \times \prod_{t=1}^{m_{12}} (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot Y \cdot \psi \bar{\psi} \cdot Y \cdot \xi + \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot Y \cdot \psi \bar{\psi} Y \psi) \quad (2.74)
\end{aligned}$$

The bound states described by this generating function have m_{11} strings attached to the brane corresponding to the first column, m_{22} strings attached to the brane corresponding to the second column, m_{12} strings stretching from the first to the second branes and $m_{21} = m_{12}$ strings stretched from the second to the first. The construction is manifestly symmetric under swaps of any strings that have the same orientation. In appendix B we give detailed checks of the construction of operators coming from (2.74), as well as detailed examples demonstrating their equality to the Gauss graph operators. It is possible to generalize the proof given in appendix C to the case of operators labelled by Young diagrams with long columns, and then to repeat the argument that worked for the AdS giant graviton, making use of the graph changing operator. The details are an obvious generalization of the discussion for the AdS gaint graviton.

3 Action of the dilatation operator

In the previous section we have obtained an integral representation of the operators dual to excited giant graviton bound states. Our goal in this section is study the action of the dilatation operator on these integral representations. We will restrict to the SU(2) sector, which corresponds to studying open string words constructed using two complex matrices Y and Z . The one loop dilatation operator in the SU(2) sector (up to overall normalization) is [66]

$$D = -g_{YM}^2 \text{Tr} ([Z, Y][\partial_Z, \partial_Y]) \quad (3.1)$$

When the open strings are described using words, one can derive a Hamiltonian [32, 33] for particles (representing the Z fields) hopping on a lattice (defined by the Y fields). The Z fields can hop between the open string world sheet and the giant graviton bound state. A key physical question is the description of this hopping, relevant for the dynamics of the string endpoints. We take this up in section 3.1. For the description in which the open strings are described using the Y field, we have argued that the operators we construct are the Gauss graph operators. The Gauss graph operators are eigenstates of the one loop dilatation operator at large N . We will define an integral representation for the action of the dilatation operator D on the generating function (2.49) in section 3.2. This integral representation is a natural starting point for a systematic $\frac{1}{N}$ expansion.

3.1 Open string boundaries

Our goal is to work out the action of the dilatation operator, in the large N limit. Concretely, this amounts to constructing a lattice Hamiltonian for each of the four types of open strings (attaching to brane 1 or 2, and stretching from 1 to 2 or from 2 to 1) we can consider. What is of most interest is the boundary conditions for the strings. There are two derivatives in D , so it is possible that the two derivatives will act on different open strings. However, these effects are subleading at large N and we ignore them.¹³ In this case, we are not losing any information by simplifying the discussion to a single (for W_A and W_B) or pair (for W_C and W_D) of open strings.

The number of fields in the giant graviton is $O(N)$, the planar approximation fails and to obtain the correct large N limit we need to sum all possible contractions of fields in the two giant gravitons. The number of fields in each open string is $\sim L$ with $L \sim O(\sqrt{N})$. If we take $\frac{L^2}{N} \ll 1$, then at large N it is accurate to contract the open string words planarly [31].

We restrict to the $\text{su}(2)$ sector which amounts to constructing all open string words W_* from the pair Z, Y of matrices. To obtain the world sheet description of the open strings, following [33], we interpret the Y fields in W_* as a “lattice” which can be populated by inserting excitations (in this case Z ’s) into the lattice (gaps between the Y ’s). We will refer to the Z ’s as impurities. For a word containing L Y fields, there are $L - 1$ sites in the lattice

$$W_* = Y Z^{n_1} Y Z^{n_2} Y \dots Y Z^{n_{L-1}} Y \quad (3.2)$$

The $\{n_i\}$ are lattice site occupation numbers. The one loop dilatation operator preserves the number of Y ’s (the lattice is not dynamical) and allows impurities to hop between neighbouring sites (Z ’s and Y ’s can swap their order in W_*). This interpretation maps the problem of determining the anomalous dimensions of operators in the super Yang-Mills theory into the dynamics of a Cuntz oscillator chain [33]. The bulk interactions are described by the Hamiltonian

$$H_{\text{bulk}} = 2\lambda \sum_{l=1}^L \hat{a}_l^\dagger \hat{a}_l - \lambda \sum_{l=1}^{L-1} (\hat{a}_l^\dagger \hat{a}_{l+1} + \hat{a}_l \hat{a}_{l+1}^\dagger), \quad (3.3)$$

where $\lambda = g_{YM}^2 N$ is the ’t Hooft coupling and the Cuntz oscillators obey

$$\hat{a}_i \hat{a}_i^\dagger = I, \quad \hat{a}_i^\dagger \hat{a}_i = I - |0\rangle\langle 0|.$$

To obtain the full Hamiltonian, we need to include boundary interactions arising from the string/brane system interaction. This interaction, which introduces sources and sinks for the impurities at the boundaries of the lattice, was first derived in [33] for a string attached to a single sphere giant graviton, and then derived in [34, 35] for strings attached to an arbitrary bound state. One such interaction allows a Z to hop from the first or last site of the string onto the giant, or from the giant into the first or last site of the string. In this process

¹³Here we are simply alluding to the fact that at large N string interactions are suppressed and hence open string words do not mix. In the closed string sector the analogous statement is that in the planar limit, different trace structures do not mix.

the string exchanges momentum with the giant graviton, so that the giant graviton is exerting a force on the string. There is also a boundary interaction in which a Z belonging to the giant “kisses” the first (or last) Y in the open string word. In this case, no momentum is exchanged, and the string does not feel a force. The logic developed in [34] derives the interaction for the “hop off” process, in which a Z hops off the string and onto the giant. Requiring that the Hamiltonian is hermitian then fixes the “hop on” interaction. The momentum conserving boundary interaction follows by expressing the kiss as a hop on followed by a hop off.

The hop off boundary term for open string W_* is given by [34, 35]

$$H_{\text{hop off}} = \lambda \sqrt{\frac{c}{N}} \hat{A}^\dagger \hat{a}_1 + \lambda \sqrt{\frac{c}{N}} \hat{A}^\dagger \hat{a}_L \quad (3.4)$$

where c is the factor of the box occupied by the open string word W_* , $\hat{a}_1$ and $\hat{a}_L$ annihilate a Z from the last and first sites of the open string word W_* and $\hat{A}^\dagger$ adds a Z to the giant graviton i.e. it adds a box in the Young diagram label of the operator. The key step in the derivation of this hop off boundary term is an identity obeyed by the restricted Schur polynomials. When a string hops out of the first (or last) site we obtain an open string word of the form ZW_* (or W_*Z). The identity expresses the restricted Schur polynomial with words ZW_* or W_*Z in terms of restricted Schur polynomials with word W_* . This identity is derived by writing the sum over σ in (2.37) as a sum over a subgroup and its cosets and again using known results about the eigenvalues of the Jucys-Murphy elements. This is how the factor of the box appears in (3.4). In terms of the generating function the derivation of this identity is a trivial exercise. Indeed, the identity of [34] is given by ($S_0 = -\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi$ and $S_1 = t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2^2 (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi)$)

$$\begin{aligned} 0 &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{\partial}{\partial \bar{\varphi}^j} \left(e^{S_0 + S_1} \bar{\varphi}^i W_i^j \right) \\ &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{S_0 + S_1} \left(\text{Tr}(W) - \bar{\varphi} \cdot W \cdot \varphi + t_1 \bar{\varphi} \cdot W \cdot Z \cdot \varphi \right. \\ &\quad \left. + t_2^2 (\bar{\varphi} \cdot W \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - \bar{\varphi} \cdot W \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi) \right) \end{aligned} \quad (3.5)$$

This is identical to the identities given in appendix C.1 of [35]. The term proportional to $\text{Tr}(W)$ describes gravitational radiation, whereby the open string W is emitted as a closed string $\text{Tr}(W)$ from the excited giant bound state. It represents a subleading contribution at large N . The term proportional to $\bar{\varphi} \cdot W \cdot \varphi$ produces restricted Schur polynomials constructed with open string word W . The remaining terms produce restricted Schur polynomials constructed with open string word $W \cdot Z$. The generating function provides a remarkably simple derivation of this formula.

The identity above describes hopping out of the last site of the open string word. To obtain the identity needed for hopping out of the first site we consider

$$0 = \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{\partial}{\partial \varphi_i} \left(e^{S_0 + S_1} W_i^j \varphi_j \right) \quad (3.6)$$

The identities needed for the sphere giant are equally easy to derive.

3.2 Gauss graph operators

We have already argued that when each open string word is replaced by a single letter, the operators we generate are the Gauss graph operators. From the results obtained in [45] we know that these are eigenstates of D , in the large N limit. In [43, 45] the simplifications of large N were implemented by exploiting simplifications in the action of the symmetric group. This analysis, known as the distant corners approximation, is clumsy. In particular, it is not obvious exactly what has been discarded in this approximation and, consequently, it is not easy to systematically correct and improve the approximation. It is interesting to reconsider this computation using the generating function where we should have better control over the large N expansion. Indeed, a systematic large N expansion should follow from a saddle point approximation of the integral representations we have obtained. To carry out this analysis we need to apply D to the operators defined by (2.49). In this subsection we study some illustrative examples. Return to (2.49) which we write as

$$Z_{\text{AdS}}(t_1, t_2, Y) = \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi} e^{t_1 \Phi_1 + t_2^2 \Phi_2} \Phi_3^{m_{11}} \Phi_4^{m_{22}} \Phi_5^{m_{12}}$$

where we have introduced the new variables

$$\begin{aligned} \Phi_1 &= \bar{\varphi} \cdot Z \cdot \varphi \\ \Phi_2 &= \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi \\ \Phi_3 &= \bar{\varphi} \cdot Y \cdot \varphi \\ \Phi_4 &= \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi \\ \Phi_5 &= \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \varphi \end{aligned} \quad (3.7)$$

Our operators are constructed using a large number of Z fields (order N) and very few Y fields (order 1). It proves helpful to remove the Y fields, as much as possible, from the manipulations that follow. With this in mind, introduce sources for the fields φ , $\bar{\varphi}$, ϕ and $\bar{\phi}$ and write

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2, Y) &= \hat{\Phi}_3^{m_{11}} \hat{\Phi}_4^{m_{22}} \hat{\Phi}_5^{m_{12}} \left(\frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \times \right. \\ &\quad \left. \times e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi} e^{t_1 \Phi_1 + t_2^2 \Phi_2} e^{\bar{j}^\varphi \cdot \varphi + \bar{\varphi} \cdot j^\varphi + \bar{j}^\phi \cdot \phi + \bar{\phi} \cdot j^\phi} \right) \Big|_{j=0} \\ &\equiv \hat{\Phi}_3^{m_{11}} \hat{\Phi}_4^{m_{22}} \hat{\Phi}_5^{m_{12}} Z_{\text{AdS, reduced}}(t_1, t_2) \Big|_{j=0} \end{aligned} \quad (3.8)$$

where $|_{j=0}$ is an instruction to set all sources to zero and

$$\begin{aligned} \hat{\Phi}_3 &= \frac{\partial}{\partial j^\varphi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\varphi} \\ \hat{\Phi}_4 &= \frac{\partial}{\partial j^\varphi} \cdot Z \cdot \frac{\partial}{\partial \bar{j}^\varphi} \frac{\partial}{\partial j^\phi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\phi} - \frac{\partial}{\partial j^\varphi} \cdot Z \cdot \frac{\partial}{\partial \bar{j}^\phi} \frac{\partial}{\partial j^\phi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\varphi} \\ \hat{\Phi}_5 &= \frac{\partial}{\partial j^\varphi} \cdot Z \cdot \frac{\partial}{\partial \bar{j}^\varphi} \frac{\partial}{\partial j^\phi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\varphi} \frac{\partial}{\partial j^\varphi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\phi} \\ &\quad - \frac{\partial}{\partial j^\varphi} \cdot Z \cdot \frac{\partial}{\partial \bar{j}^\phi} \frac{\partial}{\partial j^\phi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\varphi} \frac{\partial}{\partial j^\varphi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\phi} \end{aligned} \quad (3.9)$$

The integration is not a Gaussian integral because Φ_2 introduces terms that are quartic in the integration variables. To proceed perform a Hubbard-Stratanovich like transformation, introducing two new complex fields, and rewrite $Z_{\text{AdS, reduced}}(t_1, t_2)$ as follows

$$Z_{\text{AdS, reduced}}(t_1, t_2) = \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \int dx \int d\bar{x} \int dy \int d\bar{y} \\ \times e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi - x\bar{x} - y\bar{y} + t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2 x \bar{\varphi} \cdot Z \cdot \varphi + t_2 \bar{x} \bar{\phi} \cdot Z \cdot \phi + t_2 y \bar{\phi} \cdot Z \cdot \phi - t_2 \bar{y} \bar{\phi} \cdot Z \cdot \varphi + \bar{j}^\varphi \cdot \varphi + \bar{\varphi} \cdot j^\varphi + \bar{j}^\phi \cdot \phi + \bar{\phi} \cdot j^\phi}$$

We can streamline this expression as follows

$$Z_{\text{AdS, reduced}}(t_1, t_2) = \frac{1}{\pi^{2N+2}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \int dx \int d\bar{x} \int dy \int d\bar{y} \\ \times e^{-v \cdot M \cdot \bar{v} - x\bar{x} - y\bar{y} + v \cdot \bar{j} + j \cdot \bar{v}} \quad (3.10)$$

where

$$v_i = \begin{bmatrix} \varphi_i & \phi_i \end{bmatrix} \quad \bar{v}^i = \begin{bmatrix} \bar{\varphi}^i \\ \bar{\phi}^i \end{bmatrix} \quad j_i = \begin{bmatrix} j_i^\varphi & j_i^\phi \end{bmatrix} \quad \bar{j}^i = \begin{bmatrix} \bar{j}^{\varphi i} \\ \bar{j}^{\phi i} \end{bmatrix} \\ M = \begin{bmatrix} \mathbf{1}_N - t_1 Z - t_2 x Z & -t_2 y Z \\ t_2 \bar{y} Z & \mathbf{1}_N - t_2 \bar{x} Z \end{bmatrix} = \mathbf{1}_2 \otimes \mathbf{1}_N - \begin{bmatrix} t_1 + t_2 x & t_2 y \\ -t_2 \bar{y} & t_2 \bar{x} \end{bmatrix} \otimes Z \\ \equiv \mathbf{1}_2 \otimes \mathbf{1}_N - \mathcal{M} \otimes Z \quad (3.11)$$

The integrals over φ , $\bar{\varphi}$, ϕ and $\bar{\phi}$ are Gaussian and can be done exactly. The result is

$$Z_{\text{AdS, reduced}}(t_1, t_2) = \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} \frac{e^{j \cdot (M^{-1}) \cdot \bar{j} - x\bar{x} - y\bar{y}}}{\det M} \quad (3.12)$$

The eigenvalues of Z are z_i . The eigenvalues of $\mathcal{M}$ are given in (2.24). In terms of these eigenvalues we can write

$$\frac{1}{\det(M)} = \prod_{a=1}^2 \prod_{i=1}^N \frac{1}{1 - \lambda_a z_i} \quad (3.13)$$

The Cauchy-Littlewood identity [67] is

$$\prod_{i=1}^L \prod_{j=1}^M \frac{1}{1 - x_i y_j} = \sum_{\Lambda} \chi_{\Lambda}(x_1, \dots, x_L) \chi_{\Lambda}(y_1, \dots, y_M) \quad (3.14)$$

where the sum on the right hand side only includes Young diagrams Λ with less than the smaller of L, M rows, and χ_{Λ} is a Schur polynomial. Applying this formula to our problem we find

$$\frac{1}{\det(M)} = \sum_{\Lambda} \chi_{\Lambda}(\mathcal{M}) \chi_{\Lambda}(Z) \quad (3.15)$$

so that

$$Z_{\text{AdS, reduced}}(t_1, t_2) = \sum_{\Lambda} f_{\Lambda}(j, \bar{j}) \chi_{\Lambda}(Z) \quad (3.16)$$

$$f_{\Lambda}(j, \bar{j}) = \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} e^{j \cdot (M^{-1}) \cdot \bar{j} - x\bar{x} - y\bar{y}} \chi_{\Lambda}(\mathcal{M}) \quad (3.17)$$

The sum over Λ is only over Young diagrams with at most 2 rows.

We can perform a convincing check of (3.16) after we set $j = \bar{j} = 0$. When the sources vanish we know that the generating function produces the Schur polynomials with two rows. Further, the Schur polynomial labelled by a Young diagram with r_1 boxes in the first row and r_2 boxes in the second is multiplied by a monomial $t_1^{r_1-r_2}t_2^{r_2}$. Using the usual formula for the Schur polynomials, we find

$$\chi_{\square}(\mathcal{M}) = t_1 + t_2x + t_2\bar{x} \quad (3.18)$$

$$\chi_{\square\square}(\mathcal{M}) = t_1^2 + 2t_1t_2x + t_2^2x^2 + t_1t_2\bar{x} + t_2^2x\bar{x} + t_2^2\bar{x}^2 - t_2^2y\bar{y} \quad (3.19)$$

$$\chi_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}(\mathcal{M}) = t_1t_2\bar{x} + t_2^2x\bar{x} + t_2^2y\bar{y} \quad (3.20)$$

$$\begin{aligned} \chi_{\square\square\square}(\mathcal{M}) = & t_1^3 + 3t_1^2t_2x + 3t_1t_2^2x^2 + t_2^3x^3 + t_1^2t_2\bar{x} + 2t_1t_2^2x\bar{x} + t_2^3x^2\bar{x} \\ & + t_1t_2^2\bar{x}^2 + t_2^3x\bar{x}^2 + t_2^3\bar{x}^3 - 2t_1t_2^2y\bar{y} - 2t_2^3xy\bar{y} - 2t_2^3\bar{x}y\bar{y} \end{aligned} \quad (3.21)$$

$$\begin{aligned} \chi_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}}(\mathcal{M}) = & t_1^2t_2\bar{x} + 2t_1t_2^2x\bar{x} + t_2^3x^2\bar{x} + t_1t_2^2\bar{x}^2 + t_2^3x\bar{x}^2 + t_1t_2^2y\bar{y} \\ & + t_2^3xy\bar{y} + t_2^3\bar{x}y\bar{y} \end{aligned} \quad (3.22)$$

We then easily obtain

$$f_{\square} = t_1 \quad f_{\square\square} = t_1^2 \quad f_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} = 2t_2^2 \quad f_{\square\square\square} = t_1^3 \quad f_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} = 3t_1t_2^2 \quad (3.23)$$

which are the correct results. We have also tested that the integer coefficient of each monomial is correct.

If we work in a basis in which Z is diagonal, after a little work we can rewrite (3.12) as

$$\begin{aligned} Z_{\text{AdS, reduced}}(t_1, t_2) = & \frac{1}{\pi^2} \int dx \int d\bar{x} \int dy \int d\bar{y} \exp \left[-x\bar{x} - y\bar{y} + \sum_{n=1}^{\infty} f_n \text{Tr}(Z^n) \right. \\ & + j_{\varphi} \cdot \left(\sum_{n=0}^{\infty} X^n (1 - t_2\bar{x}Z) \right) \cdot \bar{j}_{\varphi} - j_{\varphi} \cdot \left(\sum_{n=0}^{\infty} X^n t_2yZ \right) \cdot \bar{j}_{\phi} + j_{\phi} \cdot \left(\sum_{n=0}^{\infty} X^n t_2\bar{y}Z \right) \cdot \bar{j}_{\varphi} \\ & \left. + j_{\phi} \cdot \left(\sum_{n=0}^{\infty} X^n (1 - t_2\bar{x}Z - t_1Z) \right) \cdot \bar{j}_{\phi} \right] \end{aligned} \quad (3.24)$$

where f_n has been defined in (2.26) and

$$X = (t_1 + t_2(x + \bar{x}))Z - (t_1t_2\bar{x} + t_2^2(x\bar{x} + y\bar{y}))Z^2 \quad (3.25)$$

This formula is exact. We are now ready to apply the operators defined in (3.9) to this expression. We evaluate the result at $j = \bar{j} = 0$. One derivative acting produces a term linear in the sources. To get a non-zero result we need to differentiate this again, with respect to the source that appears. Consequently, a pair of derivatives must act before we get a non-zero result. The combinatorics is therefore identical to the combinatorics from Wick's theorem and we only need to know what the basic pairings are. It is easy to read off from the above expression what each possible pairing gives.

It is instructive to consider the case that $t_2 = 0$. This corresponds to a single giant system, which we know is a $\frac{1}{2}$ -BPS operator. A simple computation shows that

$$Z_{\text{AdS, reduced}}(t_1, 0) = \exp \left(\sum_{n=1}^{\infty} f_n \text{Tr}(Z^n) + j_{\varphi} \cdot \left(\sum_{n=0}^{\infty} t_1^n Z^n \right) \cdot \bar{j}_{\varphi} + j_{\phi} \cdot \bar{j}_{\phi} \right) \quad (3.26)$$

For two strings attached to the giant graviton, the generating function is

$$\begin{aligned}
 Z_{\text{AdS}}(t_1, 0, Y) &= \left(\frac{\partial}{\partial j^\varphi} \cdot Y \cdot \frac{\partial}{\partial \bar{j}^\varphi} \right)^2 Z_{\text{AdS, reduced}}(t_1, 0) \\
 &= e^{\sum_{n=1}^{\infty} \frac{t_1^n}{n} \text{Tr}(Z^n)} \sum_{n_1, n_2=1}^{\infty} t_1^{n_1+n_2} \left(\text{Tr}(Y Z^{n_1}) \text{Tr}(Y Z^{n_2}) + \text{Tr}(Y Z^{n_1} Y Z^{n_2}) \right) \\
 &= \sum_{q=1}^{\infty} \frac{t_1^q}{q!} O_q^{(2)}
 \end{aligned} \tag{3.27}$$

For k strings attached to the giant graviton, the generating function is

$$\begin{aligned}
 Z_{\text{AdS}}(t_1, 0, Y) &= e^{\sum_{n=1}^{\infty} \frac{t_1^n}{n} \text{Tr}(Z^n)} \sum_{n_1, \dots, n_k=1}^{\infty} t_1^{n_1+n_2+\dots+n_k} \\
 &\quad \times \sum_{\sigma \in S_k} (Y Z^{n_1})_{i_{\sigma(1)}}^{i_1} (Y Z^{n_2})_{i_{\sigma(2)}}^{i_2} \dots (Y Z^{n_k})_{i_{\sigma(k)}}^{i_k} \\
 &= \sum_{q=1}^{\infty} \frac{t_1^q}{q!} O_q^{(k)}
 \end{aligned} \tag{3.28}$$

This gives a very explicit description of the Gauss graph operator, for a single giant graviton with k strings attached. It is straight forward to verify that

$$\begin{aligned}
 O_q^{(k)} &= \sum_{\sigma \in S_{k+q}} Z_{i_{\sigma(1)}}^{i_1} \dots Z_{i_{\sigma(q)}}^{i_q} Y_{i_{\sigma(q+1)}}^{i_{q+1}} \dots Y_{i_{\sigma(q+k)}}^{i_{q+k}} \\
 &= k! q! \chi_{(k+q), (k)(q)}(Z, Y)
 \end{aligned} \tag{3.29}$$

where $\chi_{(k+q), (k)(q)}(Z, Y)$ is a restricted Schur polynomial. The corresponding projection operator $P_{(k+q), (k)(q)}$ needed to define the restricted trace is trivial since the carrier space of representation $(k+q)$ is one dimensional. The results of [68] then imply that

$$D O_q^{(k)} = 0 \tag{3.30}$$

to all orders in $\frac{1}{N}$.

As an example of a non-trivial eigenstate of the dilatation operator D , consider a pair of stretched strings, i.e. $m_{11} = 0 = m_{22}$ and $m_{12} = 1 = m_{21}$. To obtain a manageable problem, consider the coefficient of t_1^n in the expansion of (3.8), which is given by

$$\begin{aligned}
 \mathcal{O} &= \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \dots Z_{i_{\sigma(n+1)}}^{i_{n+1}} (Y^2)_{i_{\sigma(n+2)}}^{i_{n+2}} \\
 &\quad - \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \dots Z_{i_{\sigma(n)}}^{i_n} Y_{i_{\sigma(n+1)}}^{i_{n+1}} (ZY)_{i_{\sigma(n+2)}}^{i_{n+2}} \\
 &= \mathcal{O}_1 - \mathcal{O}_2
 \end{aligned} \tag{3.31}$$

We work at the leading order in a large N expansion and we take n to be of order N . In this limit $\mathcal{O}$ simplifies. Recall that in any conformal field theory, we have the operator/state correspondence, which associates a state to every operator. Thus associated to the two

operators above, we have the states $|\mathcal{O}_1\rangle$ and $|\mathcal{O}_2\rangle$. Norms of states map to correlation functions of operators

$$\langle \mathcal{O}_1 | \mathcal{O}_1 \rangle = \langle \mathcal{O}_1 \mathcal{O}_1^\dagger \rangle \quad \langle \mathcal{O}_2 | \mathcal{O}_2 \rangle = \langle \mathcal{O}_2 \mathcal{O}_2^\dagger \rangle \quad (3.32)$$

In appendix D we show that

$$\frac{\langle \mathcal{O}_2 | \mathcal{O}_2 \rangle}{\langle \mathcal{O}_1 | \mathcal{O}_1 \rangle} = O(N^{-1}) \quad (3.33)$$

so that at large N we can simplify

$$\mathcal{O} = \mathcal{O}_1 + O(N^{-1}) = \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} (Y^2)_{i_{\sigma(n+2)}}^{i_{n+2}} \quad (3.34)$$

It is interesting to trace this simplification back to the generating function $Z_{\text{AdS}}(t_1, t_2, Y)$. This simplification amounts to dropping the second term in

$$\Phi_5 = \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \phi \quad (3.35)$$

which is intuitively appealing. Indeed the factor $\bar{\phi} \cdot Y \cdot \varphi \bar{\varphi} \cdot Y \cdot \phi$ has the very natural interpretation that the two Y fields have endpoints on different branes. Acting on $\mathcal{O}$ with D given in (3.1) we obtain

$$D\mathcal{O} = -g_{YM}^2(n+1) \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(2)}}^{i_2} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} \left(Y_{i_{\sigma(n+2)}}^{i_1} [Z, Y]_{i_{\sigma(1)}}^{i_{n+2}} + Y_i^{i_{n+2}} [Z, Y]_{i_{\sigma(1)}}^i \delta_{i_{\sigma(n+2)}}^{i_1} \right. \\ \left. - [Z, Y]_j^{i_1} Y_{i_{\sigma(n+2)}}^j \delta_{i_{\sigma(1)}}^{i_{n+2}} - Y_{i_{\sigma(1)}}^{i_{n+2}} [Z, Y]_{i_{\sigma(n+2)}}^{i_1} \right) \quad (3.36)$$

To simplify this expression we need a few identities. The most basic of these follow by simply changing variables in the sum. For example

$$\sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(2)}}^{i_2} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} Y_{i_{\sigma(n+2)}}^{i_1} [Z, Y]_{i_{\sigma(1)}}^{i_{n+2}} = \sum_{\rho \in S_{n+2}} Z_{i_{\rho(2)}}^{i_2} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} Y_{i_{\rho(1)}}^{i_1} [Z, Y]_{i_{\rho(n+2)}}^{i_{n+2}} \\ = \sum_{\psi \in S_{n+2}} Z_{i_{\psi(1)}}^{i_1} \cdots Z_{i_{\psi(n)}}^{i_n} Y_{i_{\psi(n+1)}}^{i_{n+1}} [Z, Y]_{i_{\psi(n+2)}}^{i_{n+2}}$$

where for the first equality $\sigma = \rho(1, n+2)$ and for the second $\rho = (1, n+1)\psi(1, n+1)$. Another useful trick is to write sums in terms of cosets of a subgroup. For example, examine the sum

$$\mathcal{S} = \sum_{\sigma \in S_{n+3}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} Y_{i_{\sigma(n+2)}}^{i_{n+2}} Y_{i_{\sigma(n+3)}}^{i_{n+3}} \quad (3.37)$$

Consider the subgroup S_{n+2} defined by $\rho(n+3) = n+3$ for $\rho \in S_{n+2}$, i.e. the subgroup leaves $n+3$ inert. Now rewrite the above sum in terms of the right cosets, labelled by the integer $i = 1, 2, \dots, n+3$ defined by

$$\sigma = \rho(i, n+3) \quad \sigma \in S_{n+3} \quad \rho \in S_{n+2} \quad (3.38)$$

The sum over σ is equivalent to separately summing over ρ and i . After rewriting we find

$$\mathcal{S} = \sum_{\rho \in S_{n+2}} Z_{i_{\rho(1)}}^{i_1} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} Y_{i_{\rho(n+2)}}^{i_{n+2}} \text{Tr}(Y) + \sum_{\rho \in S_{n+2}} Z_{i_{\rho(1)}}^{i_1} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} (Y^2)_{i_{\rho(n+2)}}^{i_{n+2}} \\ + (n+1) \sum_{\rho \in S_{n+2}} Z_{i_{\rho(1)}}^{i_1} \cdots Y_{i_{\rho(n+1)}}^{i_{n+1}} (ZY)_{i_{\rho(n+2)}}^{i_{n+2}} \quad (3.39)$$

If we use left cosets we find

$$\begin{aligned} \mathcal{S} = & \sum_{\rho \in S_{n+2}} Z_{i_{\rho(1)}}^{i_1} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} Y_{i_{\rho(n+2)}}^{i_{n+2}} \text{Tr}(Y) + \sum_{\rho \in S_{n+2}} Z_{i_{\rho(1)}}^{i_1} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} (Y^2)_{i_{\sigma(n+2)}}^{i_{n+2}} \\ & + (n+1) \sum_{\rho \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Y_{i_{\sigma(n+1)}}^{i_{n+1}} (YZ)_{i_{\sigma(n+2)}}^{i_{n+2}} \end{aligned} \quad (3.40)$$

Comparing (3.39) and (3.40) we have

$$\sum_{\rho \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Y_{i_{\sigma(n+1)}}^{i_{n+1}} (ZY)_{i_{\sigma(n+2)}}^{i_{n+2}} = \sum_{\rho \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Y_{i_{\sigma(n+1)}}^{i_{n+1}} (YZ)_{i_{\sigma(n+2)}}^{i_{n+2}} \quad (3.41)$$

Making use of these identities we find

$$D\mathcal{O} = -g_{YM}^2(n+1) \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} (2YZY - Y^2Z - ZY^2)_{i_{\sigma(n+1)}}^{i_{n+1}} \delta_{i_{\sigma(n+2)}}^{i_{n+2}}$$

Next, introduce the S_{n+1} subgroup defined by $\rho(n+2) = n+2$. By making use of right cosets exactly as above, we find

$$D\mathcal{O} = -g_{YM}^2(N+n+1)(n+1) \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} (2YZY - Y^2Z - ZY^2)_{i_{\sigma(n+1)}}^{i_{n+1}}$$

The next identity we need is obtained by starting from the sum which defines $\mathcal{O}$

$$\mathcal{O} = \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} Z_{i_{\sigma(n+1)}}^{i_{n+1}} (Y^2)_{i_{\sigma(n+2)}}^{i_{n+2}} \quad (3.42)$$

and writing in terms of cosets of the subgroup S_{n+1} which leaves $n+2$ inert. We learn that

$$\mathcal{O} = \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} Z_{i_{\sigma(n+1)}}^{i_{n+1}} \text{Tr}(Y^2) + (n+1) \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} (ZY^2)_{i_{\sigma(n+1)}}^{i_{n+1}}$$

when using right cosets, and

$$\mathcal{O} = \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} Z_{i_{\sigma(n+1)}}^{i_{n+1}} \text{Tr}(Y^2) + (n+1) \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} (Y^2Z)_{i_{\sigma(n+1)}}^{i_{n+1}}$$

when using left cosets. Recalling that n is of order N , it is clear that the first term on the right hand side of these last two formulas can be dropped at large N . We then find

$$D\mathcal{O} = 2g_{YM}^2(N+n+1)\mathcal{O} + 2\mathcal{O}' \quad (3.43)$$

where

$$\mathcal{O}' = \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} (YZY)_{i_{\sigma(n+1)}}^{i_{n+1}} \quad (3.44)$$

In appendix D we show that

$$\frac{\langle \mathcal{O}' | \mathcal{O}' \rangle}{\langle \mathcal{O} | \mathcal{O} \rangle} = O(N^{-1}) \quad (3.45)$$

so that the $\mathcal{O}'$ contribution can be dropped at large N . Thus, in the end we have

$$D\mathcal{O} = 2g_{YM}^2(N+n+1)\mathcal{O} \quad (3.46)$$

proving that our operators are indeed eigenstates of D at large N . There are a few points worth stressing:

1. The above calculation has summed a lot more than just the planar diagrams.
2. The operator $\mathcal{O}$ is a sum of many different trace structures.
3. It is not the usual 't Hooft coupling $g_{YM}^2 N$ that appears, but rather $g_{YM}^2(N+n+1)$. This is also fixed in the large N limit so that we have a finite renormalization of the 't Hooft coupling. Notice that $N+n+1$ is the factor of the last box in the first row of the R label of the Gauss graph operator. The same renormalization is evident in the results of [61, 68], obtained using very different methods.
4. The above result is another nice check of the equivalence of the operators defined by the generating function and the Gauss graph operators.

4 Discussion and conclusions

The spectrum of type IIB string theory on $\text{AdS}_5 \times \text{S}^5$ includes giant graviton branes. These are $\frac{1}{2}$ -BPS solutions. Superposing a collection of giant graviton branes gives a $\frac{1}{2}$ -BPS bound state of giant gravitons. The CFT operators dual to these bound states are given by Schur polynomials [11]. Excitations of these bound states, which are in general not BPS states, are obtained by attaching open strings to the branes. CFT operators describing these excited giant graviton bound states, known as Gauss graph operators, were described in [44, 45]. Their construction employs group representation theory and a Fourier transform on a double coset. The construction is technical which obscures many details. For example, it is not easy to see how the description simplifies at large N or how systematic $\frac{1}{N}$ corrections can be computed. The discussion is general in the sense that it applies to arbitrary bound states of giant gravitons. An important assumption is that the number of open strings that are attached is limited so that we are considering a small deformation of the original $\frac{1}{2}$ -BPS operator.

The question of developing a systematic $\frac{1}{N}$ expansion for excited giant graviton systems is the central motivation for our study. A standard way of generating a $\frac{1}{N}$ expansion is through a saddle point evaluation of the path integral. Consequently, a promising starting point is an integral representations for the relevant operators. This philosophy was recently employed, in a very similar setting, in [55]. In this article we have obtained these integral representations, for operators dual to a boundstate of two sphere giant gravitons or two AdS giant gravitons. These representations provide a simple description of the relevant operators.

The starting point for the construction is the observation that Schur polynomials can be written using Young symmetrizers. These Young symmetrizers symmetrize and anti-symmetrize indices of matrix fields from which our operator is composed. Motivated by the results of [50–54] these symmetrization and anti-symmetrizations can be implemented as integrals over ordinary (for symmetrization) and Grassman (for anti-symmetrization) variables which carry a single $U(N)$ index. In this way one obtains an integral representation for Schur polynomials with two rows or two columns. One consequence of this representation is a simple integral formula for characters of the symmetric group, in irreps labelled by Young

diagrams with two rows, or two columns. There is a natural proposal for operators dual to excited giant graviton boundstates. One of our most important results has been the demonstration that this gives an integral representation for the Gauss graph operators themselves.

We have considered the action of the dilatation operator on these novel generating functions. In the case that the open strings are described using an open string word, a key physical question is the boundary condition imposed on the string endpoints. The dilatation operator translates into a world sheet Hamiltonian as usual. The open string endpoints are represented by sources and sinks in this Hamiltonian, allowing the brane and open string to exchange momentum, i.e. to exert a force on each other. The form of this interaction was first obtained in [32, 33] for a single brane, and in [34, 35] for a general brane boundstate. In the case of the general brane boundstate, the discussion makes use of identities obeyed by restricted characters and these identities were derived using sums over cosets of the group and properties of Jucys-Murphy elements. We have recovered these known results, but with much less effort: the restricted character identity follows from the statement that the integral of a total derivative vanishes! In the case that each open string is described by a single Y field, we are considering the Gauss graph operators which are eigenstates of D [44, 45]. By introducing a set of sources we have written down a set of differential operators (3.9) which create the open strings when applied to the generating function (3.24). In (3.24) there are only 4 integrals left to do and consequently, this is a promising starting point to generate a systematic $\frac{1}{N}$ expansion through a saddle point evaluation.

Our analysis has been specialized to the case of a boundstate of two giant gravitons. A natural extension is to boundstates of more giant gravitons. The extension is straight forward, if a little tedious. To sketch what is involved, we will give some of the details relevant to describe bound states of three AdS giant gravitons. Since there are three rows, and the indices belonging to each row must be symmetrized, the description will involve three sets of fields, $\varphi_i, \bar{\varphi}^i, \phi_i, \bar{\phi}^i$ as well as $\eta_i, \bar{\eta}^i$. There is also a new antisymmetrization, associated to the new columns of length three. As usual, we list row lengths so that $R = (p_1, p_2, p_3)$ denotes a Young diagram with p_1 boxes in the first row, p_2 in the second and p_3 in the third. The generating function for Schur polynomials with three rows is given by

$$\begin{aligned}
 Z_{\text{AdS}}(t_1, t_2, t_3) &= \sum_{p_1=p_2+p_3}^{\infty} \sum_{p_2=p_3}^{\infty} \sum_{p_3=1}^{\infty} \frac{t_3^{p_3} t_2^{p_2-p_3} t_1^{p_1-p_2} (p_1 + p_2 + p_3)!}{d_{(p_1, p_2, p_3)} p_3! (p_2 - p_3)! (p_1 - p_2)!} \chi_{(p_1, p_2, p_3)}(Z) \\
 &= \frac{1}{\pi^{3N}} \int d^N \varphi \int d^N \bar{\varphi} \int d^N \phi \int d^N \bar{\phi} \int d^N \eta \int d^N \bar{\eta} \\
 &\quad \times e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i - \bar{\eta}^i \eta_i + t_1 S_1 + t_2^2 S_2 + t_3^3 S_3}
 \end{aligned} \tag{4.1}$$

where

$$\begin{aligned}
 S_1 &= \bar{\varphi}^i Z_i^j \varphi_j \\
 S_2 &= \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l \\
 S_3 &= \bar{\varphi}^i \bar{\phi}^j \bar{\eta}^k (Z_i^l Z_j^m Z_k^n - Z_i^m Z_j^l Z_k^n - Z_i^n Z_j^m Z_k^l - Z_i^l Z_j^n Z_k^m \\
 &\quad + Z_i^n Z_j^l Z_k^m + Z_i^m Z_j^n Z_k^l) \varphi_l \phi_m \eta_n
 \end{aligned} \tag{4.2}$$

When considering excited bound states, we now have many different possibilities for the open strings that can be attached, since the open string can start on any brane and terminate on any brane. Factors that attach strings to brane 1, brane 2 and brane 3 respectively, are given by

$$\begin{aligned} & \bar{\varphi}^i Y_i^j \varphi_j \quad \bar{\varphi}^i \bar{\phi}^j (Z_i^k Y_j^l - Z_i^l Y_j^k) \varphi_k \phi_l \\ & \bar{\varphi}^i \bar{\phi}^j \bar{\eta}^k (Z_i^l Z_j^m Y_k^n - Z_i^m Z_j^l Y_k^n - Z_i^n Z_j^m Y_k^l - Z_i^l Z_j^n Y_k^m \\ & \quad + Z_i^n Z_j^l Y_k^m + Z_i^m Z_j^n Y_k^l) \varphi_l \phi_m \eta_n \end{aligned} \quad (4.3)$$

To describe strings that stretch between branes, we again use a product of the above factors and simply swap column indices. It is no longer necessary to attach pairs of strings. We can for example define a factor with three Y fields, corresponding to a configuration in which a Y stretches from the first to the second brane, another from the second to the third and the last from the third to the first.

We can also attach strings that stretch between the two branes. To do this, we start with the product of a factor which attaches a string to the first brane and a factor which attaches a string to the second brane. From this we obtain a pair of strings stretched between the branes by permuting column indices, thereby swapping the endpoints of the two strings. The resulting generating function is again consistent with the Gauss Law arising from the world volume theory of the giant gravitons. Further, it is manifestly invariant under swapping strings of the same orientation. This again suggests that these operators are identical to the corresponding Gauss graph operators, something that we have checked in some examples. It is equally easy to consider bound states with even more AdS giant gravitons, or to consider bound states of more than two sphere giant gravitons.

The world volume dynamics of a bound state of p giant gravitons is described in terms of the open string excitations of the branes [69]. At low energy this should become a non-Abelian Yang-Mills theory with gauge group $U(p)$ [60]. Attempts to construct this emergent Yang-Mills theory starting from a description of the giant graviton bound state in conformal field theory have been given in [30, 61, 70, 71]. Although these initial results are encouraging, they are frustrated by the technical difficulty of the combinatorics. Perhaps the simpler description of giant graviton bound states obtained in this paper will allow progress. Specifically, a systematic $\frac{1}{N}$ expansion, through a saddle point evaluation of (3.24), should lead to $\frac{1}{N}$ interactions which allow open strings to split and join. These interactions should then match those of the expected emergent Yang-Mills theory.

Acknowledgments

RdMK is supported by the South African Research Chairs initiative of the Department of Science and Technology and the National Research Foundation. MK is supported by the National Research Foundation of Korea Grants funded by the Korea government (MSIT), NRF-2020R1A6A1A03047877 (Center for Quantum Space Time) and NRF-2022R1I1A1A010. This work was initiated while RdMK was a participant of the KITP program “Integrability in String, Field, and Condensed Matter Theory.” This research

was also supported in part by the National Science Foundation under Grant No. NSF PHY-1748958. We thank David Berenstein and Adolfo Holguin for useful discussions.

A Check of the generating function for a bound state of two AdS giants

In this appendix we expand the generating function for the bound states of two AdS giant gravitons, in a power series in t_1 and t_2 . This allows us to test their correctness since, for a small number of fields, the construction of the relevant Schur polynomials is entirely straightforward.

A.1 Operators constructed from the generating function

No open strings attached. To start, consider the two AdS giant graviton bound state generating function, which is given by

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2) &= \sum_{p_1=p_2}^{\infty} \sum_{p_2=1}^{\infty} t_2^{2p_2} t_1^{p_1-p_2} \frac{(p_1+p_2)!}{d_{(p_1,p_2)} p_2! (p_1-p_2)!} \chi_{(p_1,p_2)}(Z) \\ &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \\ &\quad \times e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i + t_1 \bar{\varphi}^i Z_i^j \varphi_j + t_2^2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l} \end{aligned} \quad (\text{A.1})$$

where $d_{(p_1,p_2)}$ is the dimension of the symmetric group $S_{p_1+p_2}$ representation labelled by a Young diagram with p_1 boxes in the first row and p_2 boxes in the second row, $p_1 \geq p_2$. After a simple computation we have

$$\begin{aligned} d_{(p_1,p_2)} &= \frac{(p_1+p_2)!(p_1-p_2+1)}{(p_1+1)!p_2!} \\ \mathcal{M}_{p_1,p_2} &= \frac{(p_1+p_2)!}{d_{(p_1,p_2)} p_2! (p_1-p_2)!} = \frac{(p_1+1)!}{(p_1-p_2+1)!} \end{aligned} \quad (\text{A.2})$$

Power series expanding the integrand of (A.1) in the parameters t_1 and t_2 , we generate a sequence of polynomials in the variables $\bar{\varphi}^i$, φ_i , $\bar{\phi}^i$ and ϕ_i . The resulting integrals are all easily performed using (2.13) and the obvious analogous formula for integrals over $\bar{\varphi}^i$ and φ_i . The result is

$$\begin{aligned} Z(t_1, t_2) &= 1 + t_1 \text{Tr} Z + t_2^2 \left(\text{Tr}(Z)^2 - \text{Tr}(Z^2) \right) + t_1 t_2^2 \left(\text{Tr}(Z)^3 - \text{Tr}(Z^3) \right) \\ &\quad + \frac{t_1^2}{2} \left(\text{Tr}(Z)^2 + \text{Tr}(Z^2) \right) + \frac{t_2^4}{2} \left(\text{Tr}(Z)^4 + 3 \text{Tr}(Z^2)^2 - 4 \text{Tr} Z \text{Tr}(Z^3) \right) + \dots \end{aligned} \quad (\text{A.3})$$

In the first column below we show the value of $\mathcal{M}_{p_1,p_2}$, in the second we show the Schur polynomial $\chi_{(p_1,p_2)}(Z)$ and in the third we show the coefficient of $t_1^{p_1-p_2} t_2^{2p_2}$. The fact that the product of the first two entries on any row reproduces the third entry shows that the

result (A.3) has indeed confirmed the formula (A.1).

$$\begin{aligned}
 \mathcal{M}_{1,0}=1 \quad \chi_{\square} &= \text{Tr} Z & \text{Tr} Z \\
 \mathcal{M}_{1,1}=2 \quad \chi_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} &= \frac{1}{2}((\text{Tr} Z)^2 - \text{Tr} Z^2) & (\text{Tr}(Z)^2 - \text{Tr}(Z^2)) \\
 \mathcal{M}_{2,1}=3 \quad \chi_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} &= \frac{1}{3}((\text{Tr} Z)^3 - \text{Tr} Z^3) & (\text{Tr}(Z)^3 - \text{Tr}(Z^3)) \\
 \mathcal{M}_{2,0}=1 \quad \chi_{\square\square} &= \frac{1}{2}((\text{Tr} Z)^2 + \text{Tr} Z^2) & \frac{1}{2}(\text{Tr}(Z)^2 + \text{Tr}(Z^2)) \\
 \mathcal{M}_{2,2}=6 \quad \chi_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} &= \frac{((\text{Tr} Z)^4 - 4\text{Tr} Z \text{Tr} Z^3 + 3(\text{Tr} Z^2)^2)}{12} & \frac{1}{2}(\text{Tr}(Z)^4 + 3\text{Tr}(Z^2)^2 - 4\text{Tr} Z \text{Tr}(Z^3))
 \end{aligned} \tag{A.4}$$

This example nicely illustrates how the checks that follow will be performed. We will not however quote the explicit formulas for all of the operators we produce, since these become rather lengthy expressions that are not very illuminating.

Two open strings attached, to the giant of momentum p_1 . Next consider the generating function obtained when we attach two open strings to the giant graviton with momentum p_1 . We employ the description in which each open string is described with a word. The relevant generating function is

$$\begin{aligned}
 Z(t_1, t_2) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i + t_2^2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l} \\
 &\quad \times e^{t_1 \bar{\varphi}^i Z_i^j \varphi_j} \bar{\varphi}^{b_1} (W_{A_1})_{b_1}^{a_1} \varphi_{a_1} \bar{\varphi}^{b_2} (W_{A_2})_{b_2}^{a_2} \varphi_{a_2}
 \end{aligned} \tag{A.5}$$

Expanding to second order in t_1 and t_2^2 we obtain

$$Z(t_1, t_2) = a_0 + a_1 t_1 + a_2 t_2^2 + t_1^2 a_3 + t_1 t_2^2 a_4 + \frac{t_2^4}{2} a_5 + \dots \tag{A.6}$$

where,¹⁴ for example

$$\begin{aligned}
 a_0 &= \text{Tr}(W_{A_1}) \text{Tr}(W_{A_2}) + \text{Tr}(W_{A_1} W_{A_2}) \\
 a_1 &= \text{Tr}(Z) \text{Tr}(W_{A_1}) \text{Tr}(W_{A_2}) + \text{Tr}(Z) \text{Tr}(W_{A_1} W_{A_2}) + \text{Tr}(Z W_{A_1}) \text{Tr}(W_{A_2}) \\
 &\quad + \text{Tr}(W_{A_1}) \text{Tr}(Z W_{A_2}) + \text{Tr}(W_{A_1} W_{A_2} Z) + \text{Tr}(W_{A_1} Z W_{A_2})
 \end{aligned} \tag{A.7}$$

Two open strings attached, to the giant of momentum p_2 . Next consider the generating function obtained when we attach two open strings to the giant graviton with momentum p_2 . The relevant generating function is

$$\begin{aligned}
 Z(t_1, t_2) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i + t_2^2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l} \\
 &\quad \times e^{t_1 \bar{\varphi}^i Z_i^j \varphi_j} (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_1} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_1} \cdot \varphi) \\
 &\quad \times (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_{B_2} \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_{B_2} \cdot \varphi)
 \end{aligned} \tag{A.8}$$

¹⁴When writing Z_i^j the upper i index is a row index and the lower j index is a column index.

Expanding to second order in t_1 and t_2^2 we obtain

$$Z(t_1, t_2) = b_0 + b_1 t_1 + b_2 t_2^2 + t_1^2 b_3 + t_1 t_2^2 b_4 + \frac{t_2^4}{2} b_5 + \dots \quad (\text{A.9})$$

where, for example

$$\begin{aligned} b_0 &= \text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z)^2 + \text{Tr}(Z)^2\text{Tr}(W_{B_1}W_{B_2}) - \text{Tr}(W_{B_2})\text{Tr}(Z)\text{Tr}(W_{B_1}Z) \\ &\quad - \text{Tr}(W_{B_1})\text{Tr}(Z)\text{Tr}(W_{B_2}Z) + 2\text{Tr}(W_{B_1}Z)\text{Tr}(W_{B_2}Z) + \text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z^2) \\ &\quad + \text{Tr}(W_{B_1}W_{B_2})\text{Tr}(Z^2) - \text{Tr}(Z)\text{Tr}(W_{B_1}W_{B_2}Z) - \text{Tr}(W_{B_2})\text{Tr}(W_{B_1}Z^2) - \text{Tr}(W_{B_2}W_{B_1}Z^2) \\ &\quad - \text{Tr}(Z)\text{Tr}(W_{B_2}W_{B_1}Z) - \text{Tr}(W_{B_1})\text{Tr}(W_{B_2}Z^2) - \text{Tr}(W_{B_1}W_{B_2}Z^2) + 2\text{Tr}(W_{B_1}ZW_{B_2}Z) \\ b_1 &= \text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z)^3 + \text{Tr}(Z)^3\text{Tr}(W_{B_1}W_{B_2}) - \text{Tr}(W_{B_2})\text{Tr}(Z)^2\text{Tr}(W_{B_1}Z) \\ &\quad - \text{Tr}(W_{B_1})\text{Tr}(Z)^2\text{Tr}(W_{B_2}Z) + 2\text{Tr}(Z)\text{Tr}(W_{B_1}Z)\text{Tr}(W_{B_2}Z) + 3\text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z)\text{Tr}(Z^2) \\ &\quad + 3\text{Tr}(Z)\text{Tr}(W_{B_1}W_{B_2})\text{Tr}(Z^2) - \text{Tr}(W_{B_2})\text{Tr}(W_{B_1}Z)\text{Tr}(Z^2) - \text{Tr}(W_{B_1})\text{Tr}(W_{B_2}Z)\text{Tr}(Z^2) \\ &\quad - \text{Tr}(Z)^2\text{Tr}(W_{B_1}W_{B_2}Z) - \text{Tr}(Z^2)\text{Tr}(W_{B_1}W_{B_2}Z) - 2\text{Tr}(W_{B_2})\text{Tr}(Z)\text{Tr}(W_{B_1}Z^2) \\ &\quad + 2\text{Tr}(W_{B_2}Z)\text{Tr}(W_{B_1}Z^2) - \text{Tr}(Z)^2\text{Tr}(W_{B_2}W_{B_1}Z) - \text{Tr}(Z^2)\text{Tr}(W_{B_2}W_{B_1}Z) \\ &\quad - 2\text{Tr}(W_{B_1})\text{Tr}(Z)\text{Tr}(W_{B_2}Z^2) + 2\text{Tr}(W_{B_1}Z)\text{Tr}(W_{B_2}Z^2) + 2\text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z^3) \\ &\quad + 2\text{Tr}(W_{B_1}W_{B_2})\text{Tr}(Z^3) - 2\text{Tr}(Z)\text{Tr}(W_{B_1}W_{B_2}Z^2) + 2\text{Tr}(Z)\text{Tr}(W_{B_1}ZW_{B_2}Z) \\ &\quad - 2\text{Tr}(W_{B_2})\text{Tr}(W_{B_1}Z^3) - 2\text{Tr}(Z)\text{Tr}(W_{B_2}W_{B_1}Z^2) - 2\text{Tr}(W_{B_1})\text{Tr}(W_{B_2}Z^3) \\ &\quad - 2\text{Tr}(W_{B_1}W_{B_2}Z^3) + 2\text{Tr}(W_{B_1}ZW_{B_2}Z^2) - 2\text{Tr}(W_{B_2}W_{B_1}Z^3) \\ &\quad + 2\text{Tr}(W_{B_2}ZW_{B_1}Z^2) \end{aligned} \quad (\text{A.10})$$

A pair of strings attached, one to each giant. The generating function for two strings attached, one to each of the two giants in the bound state, again employing a description in which each open string is described with a word, is given by

$$\begin{aligned} Z(t_1, t_2) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i + t_2^2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l + t_1 \bar{\varphi}^i Z_i^j \varphi_j} \\ &\quad \times \bar{\varphi} \cdot W_A \cdot \varphi (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_B \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_B \cdot \varphi) \end{aligned} \quad (\text{A.11})$$

Again expanding to second order in t_1 and t_2 we obtain

$$Z(t_1, t_2) = c_0 + t_1 c_1 + t_2^2 c_2 + t_1^2 c_3 + t_1 t_2^2 c_4 + t_2^4 c_5 + \dots \quad (\text{A.12})$$

where, for example

$$\begin{aligned} c_0 &= \text{Tr}(W_A)\text{Tr}(W_B)\text{Tr}(Z) - \text{Tr}(W_B)\text{Tr}(W_A Z) + \text{Tr}(W_A)\text{Tr}(W_B Z) - \text{Tr}(W_A W_B Z) \\ c_1 &= \text{Tr}(W_A)\text{Tr}(W_B)\text{Tr}(Z)^2 - \text{Tr}(W_B)\text{Tr}(Z)\text{Tr}(W_A Z) + 2\text{Tr}(W_A)\text{Tr}(Z)\text{Tr}(W_B Z) \\ &\quad - \text{Tr}(W_A Z)\text{Tr}(W_B Z) + \text{Tr}(W_A)\text{Tr}(W_B)\text{Tr}(Z^2) - \text{Tr}(Z)\text{Tr}(W_A W_B Z) \\ &\quad - \text{Tr}(W_B)\text{Tr}(W_A Z^2) + 2\text{Tr}(W_A)\text{Tr}(W_B Z^2) - \text{Tr}(W_A W_B Z^2) - \text{Tr}(W_A Z W_B Z) \end{aligned} \quad (\text{A.13})$$

A pair of strings stretched between the giants. Finally, consider the generating function for two strings stretched between the two giants in the bound state, again employing a description in which each open string is described with a word

$$\begin{aligned} Z(t_1, t_2) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi}^i \varphi_i - \bar{\phi}^i \phi_i + t_2^2 \bar{\varphi}^i \bar{\phi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \varphi_k \phi_l + t_1 \bar{\varphi}^i Z_i^j \varphi_j} \\ &\quad \times (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_C \cdot \varphi \bar{\varphi} \cdot W_D \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_C \cdot \varphi \bar{\varphi} \cdot W_D \cdot \varphi) \end{aligned} \quad (\text{A.14})$$

Again expanding to second order in t_1 and t_2 we obtain

$$Z(t_1, t_2) = s_0 + t_1 s_1 + t_2^2 s_2 + t_1^2 s_3 + t_1 t_2^2 s_4 + t_2^4 s_5 + \dots \quad (\text{A.15})$$

where, for example

$$\begin{aligned} s_0 &= \text{Tr}(Z)\text{Tr}(W_C W_D) - \text{Tr}(W_D)\text{Tr}(W_C Z) - \text{Tr}(W_C W_D Z) + \text{Tr}(W_D W_C Z) \\ s_1 &= \text{Tr}(Z)^2 \text{Tr}(W_C W_D) - \text{Tr}(W_D)\text{Tr}(Z)\text{Tr}(W_C Z) - \text{Tr}(W_C Z)\text{Tr}(W_D Z) + \text{Tr}(W_C W_D)\text{Tr}(Z^2) \\ &\quad - \text{Tr}(Z)\text{Tr}(W_C W_D Z) - \text{Tr}(W_D)\text{Tr}(W_C Z^2) + 2\text{Tr}(Z)\text{Tr}(W_D W_C Z) - \text{Tr}(W_C W_D Z^2) \\ &\quad - \text{Tr}(W_C Z W_D Z) + 2\text{Tr}(W_D W_C Z^2) \end{aligned} \quad (\text{A.16})$$

A.2 Comparison to restricted Schur polynomial and Gauss graph operators

The formulas derived above, in terms of open string words W_* , are extremely useful. As written, they describe the operators obtained when each open string excitation is described using a word. These are the operators we use in section 3 to determine the lattice Hamiltonians describing the dynamics of open strings attached to giant gravitons. Alternatively, we can set all open string words equal to a single letter $W_* = Y$ and compare to the Gauss graph operators. It is this second exercise that we will carry out in this appendix.

To construct the Gauss graph operators, we need to construct the restricted Schur polynomials. There are a number of simplifications that arise when we consider restricted Schur polynomial operators labeled by a single row (this is the case for the coefficients a_0 , a_1 and a_3 in (A.6)). In this case the Gauss graph operators and the restricted Schur polynomial bases coincide. Further, since the representation R is one dimensional, the subspace we restrict to when defining the restricted characters is not a proper subspace and the restricted characters simply reduce to ordinary characters. In this case it is simple to check the following equalities

$$\begin{aligned} a_0 &= 2\chi_{\square\square}(\cdot, \square\square)(Z, Y) = \left(\cdot, \bullet \begin{array}{c} \curvearrowright \\ \curvearrowright \end{array} \right) \\ a_1 &= 2\chi_{\square\square\square}(\square, \square\square)(Z, Y) = \left(\square, \bullet \begin{array}{c} \curvearrowright \\ \curvearrowright \end{array} \right) \\ a_3 &= 2\chi_{\square\square\square\square}(\square\square, \square\square)(Z, Y) = \left(\square\square, \bullet \begin{array}{c} \curvearrowright \\ \curvearrowright \end{array} \right) \end{aligned}$$

The equality above is between (1) the coefficient a_* appearing in (A.7), (2) the restricted Schur polynomial with three Young diagram labels and (3) the Gauss graph operator which has both a Young diagram label and a graph label. The graph is the same in all three cases because in all three we have a single brane with two strings attached to it. The equalities above are explicit checks of the equality between Gauss graph operators with all strings attached to a single brane, and our generating function for excited bound states with all excitations associated to a given row. Here we are, of course, using the fact that these equalities are meant to be exact and not just large N identities.

The remaining terms in (A.6) should be compared to Gauss graph operators constructed from restricted Schur polynomials that are labelled by Young diagrams that have

two rows. The two row restricted Schur polynomial is still simpler than the general case, since with only two rows we do not need to introduce multiplicity labels. However, the representation R is no longer one dimensional and the subspace we restrict to when defining the restricted characters is usually a proper subspace. Thus, the restricted characters are not equal to ordinary characters, and they must be computed. Since we are attaching both strings to a single brane, it is again true that, for these operators, the Gauss graph operators and the restricted Schur polynomial bases coincide. This is again an exact statement. The restricted Schur polynomials that we need are given by

$$\begin{aligned}
 \chi_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) &= \frac{1}{4} \left(\text{Tr}(Z)^2 \text{Tr}(Y)^2 + \text{Tr}(Y^2) \text{Tr}(Z)^2 + 2 \text{Tr}(YZ) \text{Tr}(Z) \text{Tr}(Y) \right. \\
 &\quad \left. - \text{Tr}(Z^2) \text{Tr}(Y)^2 - \text{Tr}(Y^2) \text{Tr}(Z^2) - 2 \text{Tr}(YZ^2) \text{Tr}(Y) \right. \\
 &\quad \left. + 2 \text{Tr}(ZY^2) \text{Tr}(Z) - 2 \text{Tr}(Y^2 Z^2) \right) \\
 \chi_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & & \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) &= \frac{1}{6} \left(\text{Tr}(Z)^3 \text{Tr}(Y)^2 + 4 \text{Tr}(Z)^2 \text{Tr}(Y) \text{Tr}(ZY) + 4 \text{Tr}(Z)^2 \text{Tr}(ZY^2) \right. \\
 &\quad + 2 \text{Tr}(Z) \text{Tr}(Y) \text{Tr}(Z^2 Y) + \text{Tr}(Z)^3 \text{Tr}(Y^2) + 2 \text{Tr}(Z) \text{Tr}(ZY)^2 \\
 &\quad + 2 \text{Tr}(Z) \text{Tr}(Z^2 Y^2) + 2 \text{Tr}(Z) \text{Tr}(ZY ZY) - 2 \text{Tr}(YZ) \text{Tr}(Y) \text{Tr}(Z^2) \\
 &\quad - \text{Tr}(Y)^2 \text{Tr}(Z^3) - \text{Tr}(Y^2) \text{Tr}(Z^3) - 2 \text{Tr}(ZY) \text{Tr}(Z^2 Y) \\
 &\quad - 2 \text{Tr}(Z^2) \text{Tr}(ZY^2) - 4 \text{Tr}(Z^3 Y) \text{Tr}(Y) - 4 \text{Tr}(Y^2 Z^3) \\
 &\quad \left. - 2 \text{Tr}(Y ZY Z^2) \right) \\
 \chi_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) &= \frac{1}{24} \text{Tr}(Y)^2 \text{Tr}(Z)^4 + \frac{1}{24} \text{Tr}(Z)^4 \text{Tr}(Y^2) + \frac{1}{6} \text{Tr}(Y) \text{Tr}(Z)^3 \text{Tr}(YZ) \\
 &\quad + \frac{1}{12} \text{Tr}(Z)^2 \text{Tr}(YZ)^2 + \frac{1}{12} \text{Tr}(Y Z)^2 \text{Tr}(Z^2) + \frac{1}{8} \text{Tr}(Y)^2 \text{Tr}(Z^2)^2 \\
 &\quad + \frac{1}{8} \text{Tr}(Y^2) \text{Tr}(Z^2)^2 + \frac{1}{6} \text{Tr}(Z)^3 \text{Tr}(Y^2 Z) - \frac{1}{6} \text{Tr}(Z) \text{Tr}(YZ) \text{Tr}(Y Z^2) \\
 &\quad + \frac{1}{2} \text{Tr}(Y) \text{Tr}(Z^2) \text{Tr}(Y Z^2) + \frac{1}{6} \text{Tr}(Y Z^2)^2 - \frac{1}{6} \text{Tr}(Y)^2 \text{Tr}(Z) \text{Tr}(Z^3) \\
 &\quad - \frac{1}{6} \text{Tr}(Z) \text{Tr}(Y^2) \text{Tr}(Z^3) - \frac{1}{6} \text{Tr}(Y) \text{Tr}(YZ) \text{Tr}(Z^3) - \frac{1}{6} \text{Tr}(Y^2 Z) \text{Tr}(Z^3) \\
 &\quad + \frac{1}{2} \text{Tr}(Z^2) \text{Tr}(Y^2 Z^2) + \frac{1}{12} \text{Tr}(Z)^2 \text{Tr}(Y ZY Z) + \frac{1}{12} \text{Tr}(Z^2) \text{Tr}(Y ZY Z) \\
 &\quad - \frac{1}{2} \text{Tr}(Y) \text{Tr}(Z) \text{Tr}(Y Z^3) - \frac{1}{6} \text{Tr}(YZ) \text{Tr}(Y Z^3) - \frac{1}{2} \text{Tr}(Z) \text{Tr}(Y^2 Z^3) \\
 &\quad \left. - \frac{1}{6} \text{Tr}(Z) \text{Tr}(Y ZY Z^2) - \frac{1}{6} \text{Tr}(Y ZY Z^3) + \frac{1}{6} \text{Tr}(Y Z^2 Y Z^2) \right) \quad (\text{A.17})
 \end{aligned}$$

It is now a simple exercise to check the following equalities

$$\begin{aligned}
 a_2 &= 4 \chi_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \\ \hline \end{array} \right) \\
 a_4 &= 6 \chi_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & & \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \\ \hline \end{array} \right) \\
 a_5 &= 12 \chi_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \\ \hline \end{array} \right)
 \end{aligned}$$

after setting $W_1 = W_2 = Y$ in a_2 , a_4 and a_5 .

It is equally easy to test the results for strings attached to the second giant graviton, given in (A.9). In this case, again because all strings are attached to a single giant, there is an exact equality between the restricted Schur polynomials and the Gauss graph operators. We have tested the equalities

$$\begin{aligned}
 b_0 &= 4\chi_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})(Z, Y) = \left(\begin{smallmatrix} \square & \square \\ \bullet & \bullet \end{smallmatrix}, \text{graph} \right) \\
 b_1 &= 12\chi_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix})(Z, Y) = \left(\begin{smallmatrix} \square & \square & \square \\ \bullet & \bullet & \bullet \end{smallmatrix}, \text{graph} \right) \\
 b_2 &= 16\chi_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix})(Z, Y) = \left(\begin{smallmatrix} \square & \square & \square \\ \bullet & \bullet & \bullet \end{smallmatrix}, \text{graph} \right) \\
 b_3 &= 24\chi_{\begin{smallmatrix} \square & \square & \square & \square \\ \square & \square & \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square & \square & \square \\ \square & \square & \square & \square \end{smallmatrix})(Z, Y) = \left(\begin{smallmatrix} \square & \square & \square & \square \\ \bullet & \bullet & \bullet & \bullet \end{smallmatrix}, \text{graph} \right)
 \end{aligned}$$

after setting $W_1 = W_2 = Y$ in b_0 , b_1 , b_2 and b_3 .

Testing the results when there is a string attached to each giant graviton is also interesting. In this case we gave a proof that the two sets of operators agree at large N , but have not proved an exact statement. These results are given in (A.12). For this class of operators it is no longer true that the Gauss graph operators and the restricted Schur polynomials coincide. Recall that from (2.58) we have

$$O(\sigma) = \frac{1}{\prod_i m_i!} \sum_{s \vdash m} \sum_{\alpha \in \prod_i S_{m_i}} \sqrt{d_s} \chi_s(\sigma \alpha) O_{R,(r,s)} \quad (\text{A.18})$$

where $\chi_s(\sigma)$ is the character of $\sigma \in S_m$ in irrep s . Since we only have a single Y box in each row, the groups S_{m_i} are trivial we have the lovely simple formula

$$O(\sigma) = \sum_{s \vdash m} \sqrt{d_s} \chi_s(\sigma) O_{R,(r,s)} \quad (\text{A.19})$$

To obtain the state with a string attached to each of the two branes we need to choose σ equals the identity in (A.19). We then have $\chi_s(\mathbf{1}) = d_s$ and hence

$$O(\mathbf{1}) = O_{R,(r,\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix})} + O_{R,(r,\begin{smallmatrix} \square \\ \square \end{smallmatrix})} \quad (\text{A.20})$$

Using this formula, we can test the polynomials defined in (A.12). A subset of the restricted Schur polynomials that we need are

$$\begin{aligned}
 \chi_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) &= \frac{1}{2!} \left(\text{Tr}(Y)^2 \text{Tr}(Z) + \text{Tr}(Y^2) \text{Tr}(Z) - \text{Tr}(Y) \text{Tr}(YZ) - \text{Tr}(Y^2 Z) \right) \\
 \chi_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) &= \frac{1}{2!} \left(\text{Tr}(Y)^2 \text{Tr}(Z) - \text{Tr}(Y^2) \text{Tr}(Z) + \text{Tr}(Y) \text{Tr}(YZ) - \text{Tr}(Y^2 Z) \right) \\
 \chi_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}}(\begin{smallmatrix} \square & \square & \square \\ \square & \square & \square \end{smallmatrix}) &= \frac{1}{4} \left(\text{Tr}(Z^2) \text{Tr}(Y^2) + \text{Tr}(Z^2) \text{Tr}(Y)^2 + \text{Tr}(Y^2) \text{Tr}(Z)^2 \right. \\
 &\quad \left. - 2 \text{Tr}(YZYZ) + \text{Tr}(Y^2) \text{Tr}(Z^2) - 2 \text{Tr}(Y Z)^2 \right)
 \end{aligned}$$

$$\begin{aligned}
 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} &= \frac{1}{4} \left(\text{Tr}(Z)^2 \text{Tr}(Y)^2 + \text{Tr}(Z^2) \text{Tr}(Y)^2 - \text{Tr}(Y^2) \text{Tr}(Z)^2 \right. \\
 &\quad - 2\text{Tr}(Y^2 Z^2) - \text{Tr}(Y^2) \text{Tr}(Z^2) + 2\text{Tr}(Y Z^2) \text{Tr}(Y) \\
 &\quad \left. - 2\text{Tr}(Z Y^2) \text{Tr}(Z) + 2\text{Tr}(Y Z) \text{Tr}(Z) \text{Tr}(Y) \right) \quad (\text{A.21})
 \end{aligned}$$

We have tested the equalities

$$\begin{aligned}
 c_0 &= \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) + \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 c_1 &= \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) + \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 c_2 &= 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) + 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \quad (\text{A.22}) \\
 c_3 &= \frac{3}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) + \frac{3}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 c_4 &= 8\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) + 8\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right)
 \end{aligned}$$

after setting $W_1 = W_2 = Y$ in c_0, c_1, c_2, c_3 and c_4 . The fact that we obtain exact agreement is not expected, suggesting we should attempt to extend the proof given in appendix C to finite N .

We would now like to test the results for the stretched string operators, summarized in (A.16) above. Since the operators with a string on each brane are related to the stretched string operators by the action of the graph changing operator, the results (A.22) suggest we should again find exact agreement. To obtain the state with strings stretching between the two branes we need to choose $\sigma = (1, 2)$ in (A.19). Thus, in the end we have

$$O((1, 2)) = O_{R, (r, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array})} - O_{R, (r, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array})} \quad (\text{A.23})$$

Using this formula, we have argued for and tested the equalities

$$\begin{aligned}
 s_0 &= \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) - \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 s_1 &= \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) - \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 s_2 &= 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) - 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 s_3 &= \frac{3}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) - \frac{3}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right) \\
 s_4 &= 8\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) - 8\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \bullet \circlearrowleft \bullet \circlearrowleft \bullet \\ \hline \end{array} \right)
 \end{aligned}$$

after setting $W_1 = W_2 = Y$ in s_0, s_1, s_2, s_3 and s_4 .

Our final comparison is between operators for which the open string excitations are described using open string words. A simple example is given by attaching two strings to the first brane and setting $p_1 = p_2 = 1$. As above, each row of R is, of course, not a giant

graviton, but this simple example allows us to compare the construction of [31] with the construction developed above. From the above construction this operator is given as the coefficient of t_2^2 in the power series expansion of the expression

$$\int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi}}{\pi^{2N}} e^{t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2^2 \bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Z \cdot \phi - t_2^2 \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi} \prod_{r=1}^2 \bar{\varphi} \cdot W_{A_r} \cdot \varphi \quad (\text{A.24})$$

The result is

$$\begin{aligned} O_{11}(Z, W_{A_1}, W_{A_2}) = & \text{Tr}(Z)^2 \text{Tr}(W_{A_1}) \text{Tr}(W_{A_2}) - \text{Tr}(Z^2) \text{Tr}(W_{A_1}) \text{Tr}(W_{A_2}) + \text{Tr}(ZW_{A_1}) \text{Tr}(Z) \text{Tr}(W_{A_2}) \\ & + \text{Tr}(ZW_{A_2}) \text{Tr}(Z) \text{Tr}(W_{A_1}) + \text{Tr}(Z)^2 \text{Tr}(W_{A_1} W_{A_2}) - \text{Tr}(Z^2) \text{Tr}(W_{A_1} W_{A_2}) \\ & - \text{Tr}(Z^2 W_{A_1}) \text{Tr}(W_{A_2}) - \text{Tr}(Z^2 W_{A_2}) \text{Tr}(W_{A_1}) + \text{Tr}(ZW_{A_1} W_{A_2}) \text{Tr}(Z) \\ & + \text{Tr}(ZW_{A_2} W_{A_1}) \text{Tr}(Z) - \text{Tr}(Z^2 W_{A_1} W_{A_2}) - \text{Tr}(Z^2 W_{A_2} W_{A_1}) \end{aligned} \quad (\text{A.25})$$

We want to compare this to the construction given in [31], which makes use of projection operators. For the operator that we are studying, we define a representation R_1'' by removing two boxes from representation R , as follows

$$R = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array} \longrightarrow \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array} \longrightarrow R_1'' = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \quad (\text{A.26})$$

We use the notation R_1'' to match the notation of appendix C1 of [31]. The projection operator that acts within the carrier space of R and projects from R to R_1'' is denoted by $P_{R, R_1''}$. In terms of this projector, the giant bound state with two strings attached is given by

$$\chi_{R, R_1''}(Z, W_{A_1}, W_{A_2}) = \frac{1}{2} \sum_{\sigma \in S_4} Z_{i_{\sigma(1)}}^{i_1} Z_{i_{\sigma(2)}}^{i_2} (W_{A_1})_{i_{\sigma(3)}}^{i_3} (W_{A_2})_{i_{\sigma(4)}}^{i_4} \text{Tr}(P_{R, R_1''} \Gamma_R(\sigma)) \quad (\text{A.27})$$

Using the results of appendix C1 of [31] and the explicit expression (A.25) it is simple to verify that

$$O(Z, W_{A_1}, W_{A_2}) = 2\chi_{R, R_1''}(Z, W_{A_1}, W_{A_2}) \quad (\text{A.28})$$

so that our generating function has correctly reproduced the restricted polynomial constructed in [31]. In section 2.3 our arguments for the equality of restricted Schur polynomials and the operators obtained from the generating function made extensive use of the derivative rule proved in [31]. It is worth reviewing this rule with the formula (A.25) in hand. Towards this end, we employ a convenient graphical notation (for this example, when defining the restrictions, string W_{A_1} is removed first and then W_{A_2} is removed) as follows

$$\chi_{R, R_1''}(Z, W_{A_1}, W_{A_2}) = \chi_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \quad (\text{A.29})$$

More details about this notation are available in [31]. For this example, the derivative rule of [31] states that

$$\frac{\partial}{\partial (W_{A_1})_a^a} \chi_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = (N+2) \chi_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}} \quad \frac{\partial}{\partial (W_{A_2})_a^a} \chi_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = (N+1) \chi_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}} \quad (\text{A.30})$$

It is a simple matter to verify these formulas using (A.25).

Finally, consider the case that a pair of strings are stretched between the two giant gravitons. Giant graviton bound states with a single pair of stretched strings are generated by the integral

$$\int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi \frac{e^{S_0}}{\pi^{2N}} e^{S_1} (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot W_D \cdot \varphi \bar{\varphi} \cdot W_C \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot W_D \cdot \varphi \bar{\varphi} \cdot W_C \cdot \varphi)$$

The term independent of t_1 and t_2 in the power series expansion of integral is

$$O_{ss} = \text{Tr}(Z)\text{Tr}(W_C W_D) + \text{Tr}(Z W_C W_D) - \text{Tr}(Z W_D)\text{Tr}(W_C) - \text{Tr}(Z W_D W_C) \quad (\text{A.31})$$

which is in precise agreement with the construction of [31]. To see this, note that the restricted character needed to construct the restricted Schur polynomial is given by a single matrix element (we use the Young's orthonormal basis)

$$\chi_{ss}(\sigma) = \langle \begin{smallmatrix} 3 & 1 \\ 2 \end{smallmatrix} | \Gamma_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}}(\sigma) | \begin{smallmatrix} 3 & 2 \\ 1 \end{smallmatrix} \rangle \quad (\text{A.32})$$

The complete set of restricted characters are now easily evaluated

$$\begin{aligned} \chi_{ss}((1)(2)(3)) &= 0 & \chi_{ss}((12)) &= \frac{\sqrt{3}}{2} & \chi_{ss}((13)) &= -\frac{\sqrt{3}}{2} \\ \chi_{ss}((23)) &= 0 & \chi_{ss}((123)) &= -\frac{\sqrt{3}}{2} & \chi_{ss}((132)) &= \frac{\sqrt{3}}{2} \end{aligned} \quad (\text{A.33})$$

It is now trivial to verify that

$$\chi_{ss}(Z, W_C, W_D) \equiv \sum_{\sigma \in S_3} \chi_{ss}(\sigma) (W_D)_{i_{\sigma(1)}}^{i_1} (W_C)_{i_{\sigma(2)}}^{i_2} Z_{i_{\sigma(3)}}^{i_3} = \frac{\sqrt{3}}{2} O_{ss} \quad (\text{A.34})$$

It is again interesting to explore the derivative of this operator. The derivative rule of [31] states that the derivative with respect to string D , stretching between the two branes, vanishes.¹⁵ It is a simple exercise to verify that this is indeed the case

$$\frac{\partial}{\partial (W_D)_a^a} O_{ss} = 0 \quad (\text{A.35})$$

B Check of the generating function for a bound state of two sphere giants

In this appendix we expand the generating functions for the bound states of two sphere giant gravitons, in a power series in t_1 and t_2 , in order to test their correctness.

¹⁵The thing that is special about string D is that it occupies slot 1, i.e. its upper index is i_1 .

B.1 Operators constructed from the generating function

No open strings attached. The two sphere giant graviton bound state generating function is given by

$$\begin{aligned}
 Z(t_1, t_2) &= \sum_{p_1 \geq p_2}^{\infty} \sum_{p_2=1}^{\infty} t_2^{2p_2} t_1^{p_1-p_2} \frac{(p_1+p_2)!}{d_{(2^{p_2} 1^{p_1-p_2})} p_2! (p_1-p_2)!} \chi_{(2^{p_2}, 1^{p_1-p_2})}(Z) \\
 &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi \\
 &\quad \times e^{-\bar{\psi}^i \psi_i - \bar{\xi}^i \xi_i - t_1 \bar{\psi}^i Z_i^j \psi_j - t_2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l}
 \end{aligned} \tag{B.1}$$

where $d_{(2^{p_2} 1^{p_1-p_2})}$ is the dimension of the symmetric group $S_{p_1+p_2}$ representation labelled by a Young diagram with p_1 boxes in the first column and $p_2 \leq p_1$ boxes in the second column. After a simple computation we have

$$\begin{aligned}
 d_{(2^{p_2} 1^{p_1-p_2})} &= \frac{(p_1+p_2)!(p_1-p_2+1)}{(p_1+1)!p_2!} \\
 \mathcal{M}_{p_1, p_2} &= \frac{(p_1+p_2)!}{d_{(2^{p_2} 1^{p_1-p_2})} p_2! (p_1-p_2)!} = \frac{(p_1+1)!}{(p_1-p_2+1)!}
 \end{aligned} \tag{B.2}$$

Power series expanding this integral in the parameters t_1 and t_2^2 we find a sequence of polynomials in the variables $\bar{\psi}^i$, ψ_i , $\bar{\xi}^i$ and ξ_i . These are all easily performed using (2.65) and the obvious analogous formula for integrals over $\bar{\xi}^i$ and ξ_i . The result is

$$Z(t_1, t_2, \{W\}) = 1 + t_1 d_1 + t_2^2 d_2 + t_1^2 d_3 + t_1 t_2^2 d_4 + t_2^4 d_5 + \dots \tag{B.3}$$

where

$$\begin{aligned}
 d_1 &= \text{Tr}(Z) = \mathcal{M}_{1,0} \chi_{\square} \\
 d_2 &= \text{Tr}(Z)^2 + \text{Tr}(Z^2) = \mathcal{M}_{2,0} \chi_{\square\square} \\
 d_3 &= \text{Tr}(Z)^2 - \text{Tr}(Z^2) = \mathcal{M}_{1,1} \chi_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \\
 d_4 &= \text{Tr}(Z)^3 - \text{Tr}(Z^3) = \mathcal{M}_{2,1} \chi_{\begin{smallmatrix} \square & \square \\ \square & \end{smallmatrix}} \\
 d_5 &= \text{Tr}(Z)^4 + 3\text{Tr}(Z^2)^2 - 4\text{Tr}(Z)\text{Tr}(Z^3) = \mathcal{M}_{2,2} \chi_{\begin{smallmatrix} \square & \square & \square \\ \square & \square & \end{smallmatrix}}
 \end{aligned} \tag{B.4}$$

confirming (B.1).

Two open strings attached, to the giant of momentum p_1 . Next consider the generating function describing bound states of two sphere giant gravitons, with two strings attached to the first giant graviton. The generating function is given by

$$\begin{aligned}
 Z(t_1, t_2, \{W\}) &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{-\bar{\psi}^i \psi_i - \bar{\xi}^i \xi_i} \\
 &\quad \times e^{-t_1 \bar{\psi}^i Z_i^j \psi_j - t_2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l} \psi_{a_1}^{a_1} (W_{A_1})_{b_1}^{b_1} \bar{\psi}^{b_1} \psi_{a_2}^{a_2} (W_{A_2})_{b_2}^{a_2} \bar{\psi}^{b_2} \\
 &= f_0 + t_1 f_1 + t_2^2 f_2 + t_1^2 f_3 + t_1 t_2^2 f_4 + t_2^4 f_5 + \dots
 \end{aligned} \tag{B.5}$$

where, for example

$$\begin{aligned}
 f_0 &= \text{Tr}(W_{A_1})\text{Tr}(W_{A_2}) - \text{Tr}(W_{A_1}W_{A_2}) \\
 f_1 &= \text{Tr}(W_{A_1})\text{Tr}(W_{A_2})\text{Tr}(Z) - \text{Tr}(ZW_{A_1})\text{Tr}(W_{A_2}) - \text{Tr}(ZW_{A_2})\text{Tr}(W_{A_1}) \\
 &\quad - \text{Tr}(W_{A_1}W_{A_2})\text{Tr}(Z) + \text{Tr}(W_{A_1}W_{A_2}Z) + \text{Tr}(W_{A_2}W_{A_1}Z)
 \end{aligned} \tag{B.6}$$

Two open strings attached, to the giant of momentum p_2 . Next consider the generating function describing bound states of two sphere giant gravitons, with two strings attached to the first giant graviton. The generating function is given by

$$\begin{aligned}
 Z(t_1, t_2, \{W\}) &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{-t_1 \bar{\psi}^i Z_i^j \psi_j - t_2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l + Z_i^l Z_j^k) \psi_k \xi_l} \\
 &\quad \times e^{-\bar{\psi}^i \psi_i - \bar{\xi}^i \xi_i} (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot W_{B_1} \cdot \xi - \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot W_{B_1} \cdot \psi) \\
 &\quad \times (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot W_{B_2} \cdot \xi - \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot W_{B_2} \cdot \psi) \\
 &= g_0 + t_1 g_1 + t_2^2 g_2 + t_1^2 g_3 + t_1 t_2^2 g_4 + t_2^4 g_5 + \dots
 \end{aligned} \tag{B.7}$$

where, for example

$$\begin{aligned}
 g_0 &= \text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z)^2 - \text{Tr}(Z)^2\text{Tr}(W_{B_1}W_{B_2}) + \text{Tr}(W_{B_2})\text{Tr}(Z)\text{Tr}(W_{B_1}Z) \\
 &\quad + \text{Tr}(W_{B_1})\text{Tr}(Z)\text{Tr}(W_{B_2}Z) + 2\text{Tr}(W_{B_1}Z)\text{Tr}(W_{B_2}Z) - \text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z^2) \\
 &\quad + \text{Tr}(W_{B_1}W_{B_2})\text{Tr}(Z^2) - \text{Tr}(Z)\text{Tr}(W_{B_1}W_{B_2}Z) - \text{Tr}(W_{B_2})\text{Tr}(W_{B_1}Z^2) - \text{Tr}(Z)\text{Tr}(W_{B_2}W_{B_1}Z) \\
 &\quad - \text{Tr}(W_{B_1})\text{Tr}(W_{B_2}Z^2) + \text{Tr}(W_{B_1}W_{B_2}Z^2) - 2\text{Tr}(W_{B_1}ZW_{B_2}Z) + \text{Tr}(W_{B_2}W_{B_1}Z^2) \\
 g_1 &= \text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z)^3 - \text{Tr}(Z)^3\text{Tr}(W_{B_1}W_{B_2}) + \text{Tr}(W_{B_2})\text{Tr}(Z)^2\text{Tr}(W_{B_1}Z) \\
 &\quad + \text{Tr}(W_{B_1})\text{Tr}(Z)^2\text{Tr}(W_{B_2}Z) + 2\text{Tr}(Z)\text{Tr}(W_{B_1}Z)\text{Tr}(W_{B_2}Z) - 3\text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z)\text{Tr}(Z^2) \\
 &\quad + 3\text{Tr}(Z)\text{Tr}(W_{B_1}W_{B_2})\text{Tr}(Z^2) - \text{Tr}(W_{B_2})\text{Tr}(W_{B_1}Z)\text{Tr}(Z^2) - \text{Tr}(W_{B_1})\text{Tr}(W_{B_2}Z)\text{Tr}(Z^2) \\
 &\quad - \text{Tr}(Z)^2\text{Tr}(W_{B_1}W_{B_2}Z) + \text{Tr}(Z^2)\text{Tr}(W_{B_1}W_{B_2}Z) - 2\text{Tr}(W_{B_2})\text{Tr}(Z)\text{Tr}(W_{B_1}Z^2) \\
 &\quad - 2\text{Tr}(W_{B_2}Z)\text{Tr}(W_{B_1}Z^2) - \text{Tr}(Z)^2\text{Tr}(W_{B_2}W_{B_1}Z) + \text{Tr}(Z^2)\text{Tr}(W_{B_2}W_{B_1}Z) \\
 &\quad - 2\text{Tr}(W_{B_1})\text{Tr}(Z)\text{Tr}(W_{B_2}Z^2) - 2\text{Tr}(W_{B_1}Z)\text{Tr}(W_{B_2}Z^2) + 2\text{Tr}(W_{B_1})\text{Tr}(W_{B_2})\text{Tr}(Z^3) \\
 &\quad - 2\text{Tr}(W_{B_1}W_{B_2})\text{Tr}(Z^3) + 2\text{Tr}(Z)\text{Tr}(W_{B_1}W_{B_2}Z^2) - 2\text{Tr}(Z)\text{Tr}(W_{B_1}ZW_{B_2}Z) \\
 &\quad + 2\text{Tr}(W_{B_2})\text{Tr}(W_{B_1}Z^3) + 2\text{Tr}(Z)\text{Tr}(W_{B_2}W_{B_1}Z^2) + 2\text{Tr}(W_{B_1})\text{Tr}(W_{B_2}Z^3) \\
 &\quad - 2\text{Tr}(W_{B_1}W_{B_2}Z^3) + 2\text{Tr}(W_{B_1}ZW_{B_2}Z^2) - 2\text{Tr}(W_{B_2}W_{B_1}Z^3) + 2\text{Tr}(W_{B_2}ZW_{B_1}Z^2)
 \end{aligned} \tag{B.8}$$

A pair of strings, one attached to each giant. The generating function describing bound states of two sphere giant gravitons, with a string attached to each of the two giant gravitons, is given by

$$\begin{aligned}
 Z(t_1, t_2, \{W\}) &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{-\bar{\psi}^i \psi_i - \bar{\xi}^i \xi_i} \\
 &\quad \times e^{-t_1 \bar{\psi}^i Z_i^j \psi_j - t_2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \psi_k \xi_l} \bar{\psi} \cdot W_A \cdot \psi \\
 &\quad \times (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot W_B \cdot \xi - \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot W_B \cdot \psi) \\
 &= h_0 + t_1 h_1 + t_2^2 h_2 + t_1^2 h_3 + t_1 t_2^2 h_4 + t_2^4 h_5 + \dots
 \end{aligned} \tag{B.9}$$

where, for example

$$\begin{aligned}
 h_0 &= \text{Tr}(W_A)\text{Tr}(W_B)\text{Tr}(Z) + \text{Tr}(W_B)\text{Tr}(W_A Z) - \text{Tr}(W_A)\text{Tr}(W_B Z) - \text{Tr}(W_A W_B Z) \\
 h_1 &= \text{Tr}(W_A)\text{Tr}(W_B)\text{Tr}(Z)^2 + \text{Tr}(W_B)\text{Tr}(Z)\text{Tr}(W_A Z) - 2\text{Tr}(W_A)\text{Tr}(Z)\text{Tr}(W_B Z) \\
 &\quad - \text{Tr}(W_A Z)\text{Tr}(W_B Z) - \text{Tr}(W_A)\text{Tr}(W_B)\text{Tr}(Z^2) - \text{Tr}(Z)\text{Tr}(W_A \cdot W_B Z) + \text{Tr}(W_A W_B Z^2) \\
 &\quad - \text{Tr}(W_B)\text{Tr}(W_A Z^2) + 2\text{Tr}(W_A)\text{Tr}(W_B Z^2) + \text{Tr}(W_A Z W_B Z)
 \end{aligned} \tag{B.10}$$

A pair of strings stretched between the giants. Finally consider the generating function describing bound states of two sphere giant gravitons, with two strings stretched between the two giant gravitons. The generating function is given by

$$\begin{aligned}
 Z(t_1, t_2, \{W\}) &= \int d^N \bar{\psi} \int d^N \psi \int d^N \bar{\xi} \int d^N \xi e^{-\bar{\psi}^i \psi_i - \bar{\xi}^i \xi_i} \\
 &\quad \times e^{-t_1 \bar{\psi}^i Z_i^j \psi_j - t_2^2 \bar{\psi}^i \bar{\xi}^j (Z_i^k Z_j^l - Z_i^l Z_j^k) \psi_k \xi_l} \\
 &\quad \times (\bar{\psi} \cdot Z \cdot \psi \bar{\xi} \cdot W_C \cdot \psi \bar{\psi} \cdot W_D \cdot \xi - \bar{\psi} \cdot Z \cdot \xi \bar{\xi} \cdot W_C \cdot \psi \bar{\psi} W_D \psi) \\
 &= j_0 + t_1 j_1 + t_2^2 j_2 + t_1^2 j_3 + t_1 t_2^2 j_4 + t_2^4 j_5 + \dots
 \end{aligned} \tag{B.11}$$

where, for example

$$\begin{aligned}
 j_0 &= \text{Tr}(Z)\text{Tr}(W_C W_D) - \text{Tr}(W_D)\text{Tr}(W_C Z) + \text{Tr}(W_C W_D Z) - \text{Tr}(W_D W_C Z) \\
 j_1 &= \text{Tr}(Z)^2 \text{Tr}(W_C W_D) - \text{Tr}(W_D)\text{Tr}(Z)\text{Tr}(W_C Z) + \text{Tr}(Z)\text{Tr}(W_C W_D Z) - 2\text{Tr}(Z)\text{Tr}(W_D W_C Z) \\
 &\quad + \text{Tr}(W_C Z)\text{Tr}(W_D Z) - \text{Tr}(Z^2)\text{Tr}(W_C W_D) + \text{Tr}(W_D)\text{Tr}(W_C Z^2) - \text{Tr}(W_C W_D Z^2) \\
 &\quad - \text{Tr}(W_C Z W_D Z) + 2\text{Tr}(W_D W_C Z^2)
 \end{aligned} \tag{B.12}$$

B.2 Comparison to restricted Schur polynomial and Gauss graph operators

Our goal is again to compare the operators we have obtained above to the Gauss graph operators, after setting the open string words equal to Y . As for the case of the AdS giants, the simplest possibility to consider is when the Young diagram labelling the operator has a single column, corresponding to a single giant graviton. In this case, the restricted Schur polynomial basis and the Gauss graph basis coincide, and restricted characters reduce to ordinary characters. It is simple to verify that

$$\begin{aligned}
 f_0 &= 2\chi_{\begin{array}{|c|} \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array}}(Z, Y) = \left(\cdot, \begin{array}{c} \bullet \\ \curvearrowright \end{array} \right) \\
 f_1 &= 2\chi_{\begin{array}{|c|} \hline \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\square, \begin{array}{c} \bullet \\ \curvearrowright \end{array} \right) \\
 f_3 &= 2\chi_{\begin{array}{|c|} \hline \square \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}}(Z, Y) = \left(\begin{array}{c} \square \\ \square \end{array}, \begin{array}{c} \bullet \\ \curvearrowright \end{array} \right)
 \end{aligned}$$

The remaining terms in (B.5)) correspond to restricted Schur polynomials that are labelled by Young diagrams with two columns. Since we are attaching both strings to a single brane, it is still true that the Gauss graph operators and the restricted Schur

polynomial bases coincide. The restricted Schur polynomials that we need to evaluate are given by

$$\begin{aligned}
 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} &= \frac{1}{4} \left(\text{Tr}(Z)^2 \text{Tr}(Y)^2 + 2 \text{Tr}(Y^2 Z^2) - \text{Tr}(Y^2) \text{Tr}(Z^2) - 2 \text{Tr}(Y Z^2) \text{Tr}(Y) \right. \\
 &\quad \left. + 2 \text{Tr}(Z Y^2) \text{Tr}(Z) - \text{Tr}(Y^2) \text{Tr}(Z)^2 - 2 \text{Tr}(Y Z) \text{Tr}(Z) \text{Tr}(Y) + \text{Tr}(Z^2) \text{Tr}(Y)^2 \right) \\
 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} &= \frac{1}{6} \left(\text{Tr}(Z)^3 \text{Tr}(Y)^2 - 4 \text{Tr}(Z)^2 \text{Tr}(Y) \text{Tr}(Z Y) - \text{Tr}(Z)^3 \text{Tr}(Y^2) \right. \\
 &\quad - 2 \text{Tr}(Z^2) \text{Tr}(Z Y) \text{Tr}(Y) + 2 \text{Tr}(Z^2 Y) \text{Tr}(Z) \text{Tr}(Y) - \text{Tr}(Z^3) \text{Tr}(Y)^2 \\
 &\quad + 4 \text{Tr}(Y^2 Z) \text{Tr}(Z)^2 + 2 \text{Tr}(Z Y)^2 \text{Tr}(Z) + \text{Tr}(Y^2) \text{Tr}(Z^3) \\
 &\quad + 2 \text{Tr}(Z Y) \text{Tr}(Z^2 Y) - 2 \text{Tr}(Z Y Z Y) \text{Tr}(Z) + 2 \text{Tr}(Z^2) \text{Tr}(Z Y^2) \\
 &\quad \left. - 2 \text{Tr}(Z^2 Y^2) \text{Tr}(Z) + 4 \text{Tr}(Z^3 Y) \text{Tr}(Y) - 2 \text{Tr}(Z^2 Y Z Y) - 4 \text{Tr}(Z^3 Y^2) \right) \\
 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} &= \frac{1}{24} \text{Tr}(Y)^2 \text{Tr}(Z)^4 - \frac{1}{24} \text{Tr}(Z)^4 \text{Tr}(Y^2) - \frac{1}{6} \text{Tr}(Y) \text{Tr}(Z)^3 \text{Tr}(Y Z) + \frac{1}{12} \text{Tr}(Z)^2 \text{Tr}(Y Z)^2 \\
 &\quad - \frac{1}{12} \text{Tr}(Y Z)^2 \text{Tr}(Z^2) + \frac{1}{8} \text{Tr}(Y)^2 \text{Tr}(Z^2)^2 - \frac{1}{8} \text{Tr}(Y^2) \text{Tr}(Z^2)^2 + \frac{1}{6} \text{Tr}(Z)^3 \text{Tr}(Y^2 Z) \\
 &\quad + \frac{1}{6} \text{Tr}(Z) \text{Tr}(Y Z) \text{Tr}(Y Z^2) - \frac{1}{2} \text{Tr}(Y) \text{Tr}(Z^2) \text{Tr}(Y Z^2) + \frac{1}{6} \text{Tr}(Y Z^2)^2 \\
 &\quad - \frac{1}{6} \text{Tr}(Y)^2 \text{Tr}(Z) \text{Tr}(Z^3) + \frac{1}{6} \text{Tr}(Z) \text{Tr}(Y^2) \text{Tr}(Z^3) + \frac{1}{6} \text{Tr}(Y) \text{Tr}(Y Z) \text{Tr}(Z^3) \\
 &\quad - \frac{1}{6} \text{Tr}(Y^2 Z) \text{Tr}(Z^3) + \frac{1}{2} \text{Tr}(Z^2) \text{Tr}(Y^2 Z^2) - \frac{1}{12} \text{Tr}(Z)^2 \text{Tr}(Y Z Y Z) \\
 &\quad + \frac{1}{12} \text{Tr}(Z^2) \text{Tr}(Y Z Y Z) + \frac{1}{2} \text{Tr}(Y) \text{Tr}(Z) \text{Tr}(Y Z^3) - \frac{1}{6} \text{Tr}(Y Z) \text{Tr}(Y Z^3) \\
 &\quad - \frac{1}{2} \text{Tr}(Z) \text{Tr}(Y^2 Z^3) - \frac{1}{6} \text{Tr}(Z) \text{Tr}(Y Z Y Z^2) + \frac{1}{6} \text{Tr}(Y Z Y Z^3) - \frac{1}{6} \text{Tr}(Y Z^2 Y Z^2) \quad (\text{B.13})
 \end{aligned}$$

It is now a simple exercise to check the following equalities

$$\begin{aligned}
 f_2 &= 4 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \bullet \begin{array}{c} \curvearrowright \end{array} \bullet \begin{array}{c} \curvearrowright \end{array} \bullet \right) \\
 f_4 &= 6 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \bullet \begin{array}{c} \curvearrowright \end{array} \bullet \begin{array}{c} \curvearrowright \end{array} \bullet \right) \\
 f_5 &= 12 \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \bullet \begin{array}{c} \curvearrowright \end{array} \bullet \begin{array}{c} \curvearrowright \end{array} \bullet \right)
 \end{aligned}$$

after setting $W_1 = W_2 = Y$ in f_2 , f_4 and f_5 .

Using the results for strings attached to the second giant graviton, given in (B.7) we have checked that the identities

$$\begin{aligned}
 g_0 &= 4\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 g_1 &= 2\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 g_2 &= 16\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 g_3 &= 24\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right) (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right)
 \end{aligned}$$

are obeyed. To test the results when there is one string attached to each giant graviton, we again have to compare to a linear combination of restricted Schur polynomials. The discussion is parallel to the discussion for the AdS giants, so we simply quote the results. We have verified the identities

$$\begin{aligned}
 h_0 &= \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) + \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 h_1 &= 2\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) + 2\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 h_2 &= \frac{1}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) + \frac{1}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 h_3 &= 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) + 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 h_4 &= 8\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) + 8\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right)
 \end{aligned}$$

Note once again that we have a complete match, even at finite N . Finally, for the operators corresponding to the stretched string states we have verified the identities

$$\begin{aligned}
 j_0 &= \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) - \chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 j_1 &= 2\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) - 2\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right) \\
 j_2 &= \frac{1}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) - \frac{1}{2}\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} (Z, Y) = \left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \bullet \curvearrowright \bullet \right)
 \end{aligned}$$

$$\begin{aligned}
 j_3 &= 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} (Z, Y) - 3\chi_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{c} \bullet \leftarrow \bullet \\ \bullet \rightarrow \bullet \end{array} \right) \\
 j_4 &= 8\chi_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} (Z, Y) - 8\chi_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} (Z, Y) = \left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}, \begin{array}{c} \bullet \leftarrow \bullet \\ \bullet \rightarrow \bullet \end{array} \right)
 \end{aligned}$$

C A Gauss graph identity

In this appendix we will prove an identity for Gauss graph operators which do not involve any stretched strings. This corresponds to setting $\sigma = \mathbf{1}$ in $O_{R,r}(\sigma)$. We choose r to have p_1 boxes in the first row and p_2 boxes in the second and take both p_1 and p_2 to be $O(N)$. R has $p_1 + m_1$ boxes in the first row and $m_2 + p_2$ boxes in the second row so that m_1 of the boxes in the first row and m_2 in the second row correspond to Y 's. We take m_1 and m_2 to be $O(1)$. Recall formula (2.62) which, after setting σ to the identity, says

$$O_{R,r}(\mathbf{1}) = \sum_{\rho \in S_{n+m}} \sum_{\sigma_1 \in S_n} \sum_{\alpha \in S_{m_1} \times S_{m_2}} \chi_r(\sigma_1) \chi_R(\sigma_1 \circ \alpha \rho) \text{Tr}(\rho Z^{\otimes n} Y^{\otimes m}) \quad (\text{C.1})$$

There is a nice interpretation of this formula that follows after recalling that the projection operator onto representation q , within a vector space V , is given by

$$P_q = \frac{d_q}{|\mathcal{G}|} \sum_{\sigma \in \mathcal{G}} \chi_q(\sigma) \sigma \quad (\text{C.2})$$

Here σ is the action of group $\mathcal{G}$ on the space V , $\chi_q(\sigma)$ is the character of group element σ in representation q , d_q is the dimension of representation q and $|\mathcal{G}|$ is the order of the group $\mathcal{G}$. If there is more than one subspace of V that carries representation q , P_q will project to their direct sum. Using (m) to denote the trivial representation of S_m , labelled by a Young diagram with a single row of length m , we can write (up to an unimportant constant which we drop)

$$O_{R,r}(\mathbf{1}) = \sum_{\rho \in S_{n+m}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho) \text{Tr}(\rho Z^{\otimes n} \otimes Y^{\otimes m}) \quad (\text{C.3})$$

We are going to work out a formula for the derivative of this operator with respect to Y_a^a . This analysis is motivated by appendix C of [31] and we recommend that the reader consult this source for further details that we do not spell out here. It is useful to generalize the operators we consider by including a third field

$$O_{R,r}^{(3)}(\mathbf{1}) = \sum_{\rho \in S_{n+m}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho) \text{Tr}(\rho Z^{\otimes n} \otimes X^{\otimes m_2} \otimes Y^{\otimes m_1}) \quad (\text{C.4})$$

Describing the states in the carrier space of $R \vdash m+n$ using Young-Yammanouchi patterns, the indices of the fields Y correspond to boxes with labels from 1 to m_1 , the indices of the fields X to boxes with labels from $m_1 + 1$ to $m_1 + m_2 = m$ and Z to labels from $m + 1$ to $m + n$. The Y fields are in representation (m_1) and the X fields in representation

(m_2) . The Y fields populate boxes in the first row, while the X fields populate boxes in the second row. Now, consider the derivative

$$\frac{d}{dY_a^a} O_{R,r}^{(3)}(\mathbf{1}) = m_1 \sum_{\rho \in S_{n+m}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho) \text{Tr}(\rho Z^{\otimes n} \otimes X^{\otimes m_2} \otimes Y^{\otimes m_1-1} \otimes \mathbf{1}) \quad (\text{C.5})$$

where a is summed on the left hand side. Here we have chosen the index of $\mathbf{1}$ to correspond to the box in R labelled 1 and we have used the fact that we can differentiate any of the Y fields and then cycle the identity $\mathbf{1}$ into slot 1, possible because the Y fields are in representation (m_1) . Writing the indices more explicitly we have

$$\begin{aligned} \frac{d}{dY_a^a} O_{R,r}^{(3)}(\mathbf{1}) &= m_1 \sum_{\rho \in S_{n+m}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho) Z_{i_{\rho(n+m)}}^{i_{n+m}} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} X_{i_{\rho(m_1+m_2)}}^{i_{m_1+m_2}} \cdots X_{i_{\rho(m_1+1)}}^{i_{m_1+1}} \\ &\quad \times Y_{i_{\rho(m_1)}}^{i_{m_1}} \cdots Y_{i_{\rho(2)}}^{i_2} \delta_{i_{\rho(1)}}^{i_1} \end{aligned} \quad (\text{C.6})$$

To proceed, write the sum over S_{n+m} as a sum over the subgroup S_{n+m-1} which permutes $2, 3, \dots, m+n$ but leaves 1 inert. The result is

$$\begin{aligned} \frac{d}{dY_a^a} O_{R,r}^{(3)}(\mathbf{1}) &= m_1 \sum_{\rho \in S_{n+m-1}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho C) Z_{i_{\rho(n+m)}}^{i_{n+m}} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} X_{i_{\rho(m_1+m_2)}}^{i_{m_1+m_2}} \cdots X_{i_{\rho(m_1+1)}}^{i_{m_1+1}} \\ &\quad \times Y_{i_{\rho(m_1)}}^{i_{m_1}} \cdots Y_{i_{\rho(2)}}^{i_2} \end{aligned} \quad (\text{C.7})$$

where

$$C = N + \sum_{i=2}^{n+m} (1, i) = N + \hat{j} \quad (\text{C.8})$$

In this formula, $\hat{j}$ is a Jucys-Murphy element. Denote the representation obtained after dropping the box labelled 1 from R by R' . The box that is dropped is from the first row of R . Acting on this representation $\hat{j}$ gives the content of the dropped box and we have

$$\frac{d}{dY_a^a} O_{R,r}^{(3)}(\mathbf{1}) = m_1 c_{RR'} O_{R',r}(\mathbf{1}) \quad (\text{C.9})$$

This is an exact result. Now consider the derivative with respect to X . A simple calculation gives

$$\begin{aligned} \frac{d}{dX_a^a} O_{R,r}^{(3)}(\mathbf{1}) &= m_2 \sum_{\rho \in S_{n+m}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho) Z_{i_{\rho(n+m)}}^{i_{n+m}} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} X_{i_{\rho(m_1+m_2)}}^{i_{m_1+m_2}} \cdots \delta_{i_{\rho(m_1+1)}}^{i_{m_1+1}} \\ &\quad \times Y_{i_{\rho(m_1)}}^{i_{m_1}} \cdots Y_{i_{\rho(2)}}^{i_2} Y_{i_{\rho(1)}}^{i_1} \end{aligned} \quad (\text{C.10})$$

To proceed, we write the sum over S_{n+m} as a sum over the subgroup S_{n+m-1} which permutes $1, 2, \dots, m_1, m_1+2, \dots, m+n$ but leaves m_1+1 inert. The result is

$$\begin{aligned} \frac{d}{dX_a^a} O_{R,r}^{(3)}(\mathbf{1}) &= m_2 \sum_{\rho \in S_{n+m-1}} \chi_R(P_r P_{(m_1)} P_{(m_2)} \rho C) Z_{i_{\rho(n+m)}}^{i_{n+m}} \cdots Z_{i_{\rho(n+1)}}^{i_{n+1}} X_{i_{\rho(m_1+m_2)}}^{i_{m_1+m_2}} \cdots X_{i_{\rho(m_1+2)}}^{i_{m_1+2}} \\ &\quad \times Y_{i_{\rho(m_1)}}^{i_{m_1}} \cdots Y_{i_{\rho(1)}}^{i_1} \end{aligned} \quad (\text{C.11})$$

where

$$C = \kappa_1 + \kappa_2 \quad (C.12)$$

$$\kappa_1 = N + \sum_{i=m_1+2}^{n+m} (m_1 + 1, i) \quad \kappa_2 = \sum_{i=1}^{m_1} (m_1 + 1, i) \quad (C.13)$$

We will now argue that at large N , the terms collected in κ_2 can be ignored. Above we are summing transpositions (2-cycles). Transpositions square to the identity matrix, so that their eigenvalues square to 1. Consequently, the matrix elements of the transpositions are bounded, independent of N . In the term κ_2 we sum only $m_1 = O(1)$ terms, while in κ_1 we sum $m_2 + n = O(N)$ terms. In addition, the first term in κ_1 is N . This certainly bounds the eigenvalues of κ_2 to be $O(1)$ at most. One might worry that there are cancellations between terms summed in κ_1 , which might in the end imply that κ_1 itself has eigenvalues of order 1. This is not the case: it is clear $\kappa_1 = N + \hat{j}$. For the case that we consider here, $\hat{j}$ gives the content of the last box in the second row, which is $p_2 - 2$. Thus, κ_1 has all eigenvalues of order N and we can safely ignore κ_2 at large N . In this case we have

$$\frac{d}{dX_a^a} O_{R,r}^{(3)}(\mathbf{1}) = m_2 c_{RR'} O_{R',r}(\mathbf{1}) \quad (C.14)$$

In this formula the box dropped to obtain R' from R is the right most box in the second row. The formula we want to derive, is for the derivative of $O_{R,r}(\mathbf{1})$. This can be written as

$$\begin{aligned} \frac{d}{dY_a^a} O_{R,r}(\mathbf{1}) &= \frac{d}{dY_a^a} O_{R,r}^{(3)}(\mathbf{1})|_{X=Y} + \frac{d}{dX_a^a} O_{R,r}^{(3)}(\mathbf{1})|_{X=Y} \\ &= m_1 c_{RR'_1} O_{R'_1,r}(\mathbf{1}) + m_2 c_{RR'_2} O_{R'_2,r}(\mathbf{1}) \end{aligned} \quad (C.15)$$

where we have used the notation R'_i to denote the representation obtained by dropping a box from row i of R .

It is a much simpler exercise to show that this is the rule obeyed by operators obtained from generating function. The relevant generating function is

$$\begin{aligned} Z_{\text{AdS}}(t_1, t_2, Y) &= \frac{1}{\pi^{2N}} \int d^N \bar{\varphi} \int d^N \varphi \int d^N \bar{\phi} \int d^N \phi e^{-\bar{\varphi} \cdot \varphi - \bar{\phi} \cdot \phi} \\ &\quad \times e^{t_1 \bar{\varphi} \cdot Z \cdot \varphi + t_2^2 \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \phi - t_2^2 \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Z \cdot \varphi} (\bar{\varphi} \cdot Y \cdot \varphi)^{m_{11}} \\ &\quad \times (\bar{\varphi} \cdot Z \cdot \varphi \bar{\phi} \cdot Y \cdot \phi - \bar{\varphi} \cdot Z \cdot \phi \bar{\phi} \cdot Y \cdot \varphi)^{m_{22}} \end{aligned} \quad (C.16)$$

Setting m_{11} equal to m_1 and m_{22} equal to m_2 , and extracting the coefficient of $(t_2^2)^{p_2} t_1^{p_1 - p_2}$ gives the operator $O_{R,r}(\mathbf{1})$ with m_1 Y boxes in the first row, m_2 Y boxes in the second and r a Young diagram with p_1 boxes in the first row and p_2 boxes in the second row. Differentiating with respect to Y_a^a and using the arguments of section 2.3, we easily recover (C.15).

This proves that the derivative with respect to Y of the Gauss graph operator matches the derivative with respect to Y of the operators defined by our generating function. Since there are no contributions to either operator that is independent of Y , this proves their equality.

There is an obvious extension of these arguments to the case of Gauss graph operators for sphere giant gravitons.

D Correlation functions

In this section we evaluate the correlation functions that were used in section 3.2. Consider

$$\mathcal{O}_1 = \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} (Y^2)_{i_{\sigma(n+2)}}^{i_{n+2}} \quad (\text{D.1})$$

We would like to evaluate the correlator $\langle \mathcal{O}_1 \mathcal{O}_1^\dagger \rangle$. The Y fields only contract with $Y^\dagger$ fields. Note that

$$\langle (Y^2)_j^i (Y^{\dagger 2})_l^k \rangle = \delta_l^k \delta_j^i + N \delta_l^i \delta_j^k \quad (\text{D.2})$$

At large N we need only keep the second term on the right hand side. This gives

$$\langle (Y^2)_j^i (Y^{\dagger 2})_l^k \rangle = N \langle Y_j^i Y_l^k \rangle (1 + O(N^{-1})) \quad (\text{D.3})$$

We will not display the $O(N^{-1})$ correction in what follows. Using this formula we have (the operator $P_{(n+2)}$ is defined in (2.2) above)

$$\begin{aligned} \langle \mathcal{O}_1 \mathcal{O}_1^\dagger \rangle &= N \sum_{\sigma_1, \sigma_2 \in S_{n+2}} \langle Z_{i_{\sigma_1(1)}}^{i_1} \cdots Z_{i_{\sigma_1(n+1)}}^{i_{n+1}} Y_{i_{\sigma_1(n+2)}}^{i_{n+2}} Z_{j_{\sigma_2(1)}}^{\dagger j_1} \cdots Z_{j_{\sigma_2(n+1)}}^{\dagger j_{n+1}} Y_{j_{\sigma_2(n+2)}}^{\dagger j_{n+2}} \rangle \\ &= N \sum_{\sigma_1, \sigma_2 \in S_{n+2}} \sum_{\rho \in S_{n+1}} \text{Tr}(\sigma_1 \rho \sigma_2 \rho^{-1}) \\ &= N(n+1)! \sum_{\sigma_1, \sigma_2 \in S_{n+2}} \text{Tr}(\sigma_1 \sigma_2) \\ &= N(n+1)!(n+2)! \sum_{\sigma_1 \in S_{n+2}} \text{Tr}(\sigma_1) \\ &= N(n+1)!((n+2)!)^2 \text{Tr}(P_{(n+2)}) \\ &= N(n+1)!(n+2)! \prod_{i=0}^{n+1} (N+i) \end{aligned} \quad (\text{D.4})$$

Next, consider

$$\mathcal{O}_2 = \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} Y_{i_{\sigma(n+1)}}^{i_{n+1}} (ZY)_{i_{\sigma(n+2)}}^{i_{n+2}} \quad (\text{D.5})$$

Our previous strategy of performing the Y contractions by hand is not very useful when evaluating the $\langle \mathcal{O}_2 \mathcal{O}_2^\dagger \rangle$ correlator. A better approach is to start from the identity

$$\begin{aligned} \sum_{\sigma \in S_{n+3}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} Y_{i_{\sigma(n+2)}}^{i_{n+2}} Y_{i_{\sigma(n+3)}}^{i_{n+3}} &= \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} Y_{i_{\sigma(n+2)}}^{i_{n+2}} \text{Tr}(Y) \\ &\quad + \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} (Y^2)_{i_{\sigma(n+2)}}^{i_{n+2}} \\ &\quad + (n+1) \sum_{\sigma \in S_{n+2}} Z_{i_{\sigma(1)}}^{i_1} \cdots Y_{i_{\sigma(n+1)}}^{i_{n+1}} (ZY)_{i_{\sigma(n+2)}}^{i_{n+2}} \end{aligned}$$

which follows after summing, on the right hand side, over the subgroup and its cosets. The first two terms on the right hand side can be neglected at large N so that

$$\mathcal{O}_2 = \frac{1}{n+1} \sum_{\sigma \in S_{n+3}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n+1)}}^{i_{n+1}} Y_{i_{\sigma(n+2)}}^{i_{n+2}} Y_{i_{\sigma(n+3)}}^{i_{n+3}} \quad (\text{D.6})$$

A simple computation now gives

$$\langle \mathcal{O}_2 \mathcal{O}_2^\dagger \rangle = 2(n!)^2 (n+2)(n+3) \prod_{i=0}^{n+2} (N+i) \quad (\text{D.7})$$

and hence we have

$$\frac{\langle \mathcal{O}_2 \mathcal{O}_2^\dagger \rangle}{\langle \mathcal{O}_1 \mathcal{O}_1^\dagger \rangle} = \frac{2(n+3)(N+n+2)}{(n+1)^2 N} = O(N^{-1}) \quad (\text{D.8})$$

Finally, consider

$$\mathcal{O}' = \sum_{\sigma \in S_{n+1}} Z_{i_{\sigma(1)}}^{i_1} \cdots Z_{i_{\sigma(n)}}^{i_n} (YZY)_{i_{\sigma(n+1)}}^{i_{n+1}} \quad (\text{D.9})$$

A very similar argument to the one just given establishes that

$$\frac{\langle \mathcal{O}' \mathcal{O}'^\dagger \rangle}{\langle \mathcal{O}_1 \mathcal{O}_1^\dagger \rangle} = O(N^{-2}) \quad (\text{D.10})$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] J.M. Maldacena, *The large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231 [[hep-th/9711200](#)] [[INSPIRE](#)].
- [2] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](#)] [[INSPIRE](#)].
- [3] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](#)] [[INSPIRE](#)].
- [4] J.A. Minahan and K. Zarembo, *The Bethe ansatz for N = 4 super Yang-Mills*, *JHEP* **03** (2003) 013 [[hep-th/0212208](#)] [[INSPIRE](#)].
- [5] N. Beisert et al., *Review of AdS/CFT integrability: an overview*, *Lett. Math. Phys.* **99** (2012) 3 [[arXiv:1012.3982](#)] [[INSPIRE](#)].
- [6] N. Beisert, S. Frolov, M. Staudacher and A.A. Tseytlin, *Precision spectroscopy of AdS/CFT*, *JHEP* **10** (2003) 037 [[hep-th/0308117](#)] [[INSPIRE](#)].
- [7] J. McGreevy, L. Susskind and N. Toumbas, *Invasion of the giant gravitons from anti-de Sitter space*, *JHEP* **06** (2000) 008 [[hep-th/0003075](#)] [[INSPIRE](#)].
- [8] A. Hashimoto, S. Hirano and N. Itzhaki, *Large branes in AdS and their field theory dual*, *JHEP* **08** (2000) 051 [[hep-th/0008016](#)] [[INSPIRE](#)].
- [9] M.T. Grisaru, R.C. Myers and O. Tafjord, *SUSY and goliath*, *JHEP* **08** (2000) 040 [[hep-th/0008015](#)] [[INSPIRE](#)].
- [10] V. Balasubramanian, M. Berkooz, A. Naqvi and M.J. Strassler, *Giant gravitons in conformal field theory*, *JHEP* **04** (2002) 034 [[hep-th/0107119](#)] [[INSPIRE](#)].

- [11] S. Corley, A. Jevicki and S. Ramgoolam, *Exact correlators of giant gravitons from dual $N = 4$ SYM theory*, *Adv. Theor. Math. Phys.* **5** (2002) 809 [[hep-th/0111222](#)] [[INSPIRE](#)].
- [12] D. Berenstein, *A toy model for the AdS/CFT correspondence*, *JHEP* **07** (2004) 018 [[hep-th/0403110](#)] [[INSPIRE](#)].
- [13] H. Lin, O. Lunin and J.M. Maldacena, *Bubbling AdS space and 1/2 BPS geometries*, *JHEP* **10** (2004) 025 [[hep-th/0409174](#)] [[INSPIRE](#)].
- [14] O. Aharony, Y.E. Antebi, M. Berkooz and R. Fishman, *‘Holey sheets’: Pfaffians and subdeterminants as D-brane operators in large N gauge theories*, *JHEP* **12** (2002) 069 [[hep-th/0211152](#)] [[INSPIRE](#)].
- [15] D. Berenstein, *Shape and holography: studies of dual operators to giant gravitons*, *Nucl. Phys. B* **675** (2003) 179 [[hep-th/0306090](#)] [[INSPIRE](#)].
- [16] Y. Kimura and S. Ramgoolam, *Branes, anti-branes and Brauer algebras in gauge-gravity duality*, *JHEP* **11** (2007) 078 [[arXiv:0709.2158](#)] [[INSPIRE](#)].
- [17] T.W. Brown, P.J. Heslop and S. Ramgoolam, *Diagonal multi-matrix correlators and BPS operators in $N = 4$ SYM*, *JHEP* **02** (2008) 030 [[arXiv:0711.0176](#)] [[INSPIRE](#)].
- [18] T.W. Brown, P.J. Heslop and S. Ramgoolam, *Diagonal free field matrix correlators, global symmetries and giant gravitons*, *JHEP* **04** (2009) 089 [[arXiv:0806.1911](#)] [[INSPIRE](#)].
- [19] R. Bhattacharyya, S. Collins and R. de Mello Koch, *Exact multi-matrix correlators*, *JHEP* **03** (2008) 044 [[arXiv:0801.2061](#)] [[INSPIRE](#)].
- [20] R. Bhattacharyya, R. de Mello Koch and M. Stephanou, *Exact multi-restricted Schur polynomial correlators*, *JHEP* **06** (2008) 101 [[arXiv:0805.3025](#)] [[INSPIRE](#)].
- [21] P. Caputa, R. de Mello Koch and P. Diaz, *A basis for large operators in $N = 4$ SYM with orthogonal gauge group*, *JHEP* **03** (2013) 041 [[arXiv:1301.1560](#)] [[INSPIRE](#)].
- [22] P. Caputa, R. de Mello Koch and P. Diaz, *Operators, correlators and free fermions for $SO(N)$ and $Sp(N)$* , *JHEP* **06** (2013) 018 [[arXiv:1303.7252](#)] [[INSPIRE](#)].
- [23] G. Kemp, *$SO(N)$ restricted Schur polynomials*, *J. Math. Phys.* **56** (2015) 022302 [[arXiv:1405.7017](#)] [[INSPIRE](#)].
- [24] G. Kemp, *Restricted Schurs and correlators for $SO(N)$ and $Sp(N)$* , *JHEP* **08** (2014) 137 [[arXiv:1406.3854](#)] [[INSPIRE](#)].
- [25] J. Pasukonis and S. Ramgoolam, *Quivers as calculators: counting, correlators and Riemann surfaces*, *JHEP* **04** (2013) 094 [[arXiv:1301.1980](#)] [[INSPIRE](#)].
- [26] P. Mattioli and S. Ramgoolam, *Gauge invariants and correlators in flavoured quiver gauge theories*, *Nucl. Phys. B* **911** (2016) 638 [[arXiv:1603.04369](#)] [[INSPIRE](#)].
- [27] S. Ramgoolam, *Permutation invariant Gaussian matrix models*, *Nucl. Phys. B* **945** (2019) 114682 [[arXiv:1809.07559](#)] [[INSPIRE](#)].
- [28] G. Barnes, A. Padellaro and S. Ramgoolam, *Permutation invariant Gaussian two-matrix models*, *J. Phys. A* **55** (2022) 145202 [[arXiv:2104.03707](#)] [[INSPIRE](#)].
- [29] G. Barnes, A. Padellaro and S. Ramgoolam, *Permutation symmetry in large- N matrix quantum mechanics and partition algebras*, *Phys. Rev. D* **106** (2022) 106020 [[arXiv:2207.02166](#)] [[INSPIRE](#)].
- [30] V. Balasubramanian, D. Berenstein, B. Feng and M.-X. Huang, *D-branes in Yang-Mills theory and emergent gauge symmetry*, *JHEP* **03** (2005) 006 [[hep-th/0411205](#)] [[INSPIRE](#)].

- [31] R. de Mello Koch, J. Smolic and M. Smolic, *Giant gravitons — with strings attached (I)*, *JHEP* **06** (2007) 074 [[hep-th/0701066](#)] [[INSPIRE](#)].
- [32] D. Berenstein, D.H. Correa and S.E. Vazquez, *Quantizing open spin chains with variable length: an example from giant gravitons*, *Phys. Rev. Lett.* **95** (2005) 191601 [[hep-th/0502172](#)] [[INSPIRE](#)].
- [33] D. Berenstein, D.H. Correa and S.E. Vazquez, *A study of open strings ending on giant gravitons, spin chains and integrability*, *JHEP* **09** (2006) 065 [[hep-th/0604123](#)] [[INSPIRE](#)].
- [34] R. de Mello Koch, J. Smolic and M. Smolic, *Giant gravitons — with strings attached (II)*, *JHEP* **09** (2007) 049 [[hep-th/0701067](#)] [[INSPIRE](#)].
- [35] D. Bekker, R. de Mello Koch and M. Stephanou, *Giant gravitons — with strings attached (III)*, *JHEP* **02** (2008) 029 [[arXiv:0710.5372](#)] [[INSPIRE](#)].
- [36] D. Berenstein and E. Dzienkowski, *Giant gravitons and the emergence of geometric limits in beta-deformations of $N = 4$ SYM*, *JHEP* **01** (2015) 126 [[arXiv:1408.3620](#)] [[INSPIRE](#)].
- [37] H. Lin, *Relation between large dimension operators and oscillator algebra of Young diagrams*, *Int. J. Geom. Meth. Mod. Phys.* **12** (2015) 1550047 [[arXiv:1407.7815](#)] [[INSPIRE](#)].
- [38] R. de Mello Koch, N.H. Tahiridimbisoa and C. Mathwin, *Anomalous dimensions of heavy operators from magnon energies*, *JHEP* **03** (2016) 156 [[arXiv:1506.05224](#)] [[INSPIRE](#)].
- [39] D. Berenstein and A. Holguin, *Open giant magnons suspended between dual giant gravitons in $N = 4$ SYM*, *JHEP* **09** (2020) 019 [[arXiv:2006.08649](#)] [[INSPIRE](#)].
- [40] A. Holguin, *Giant gravitons intersecting at angles from integrable spin chains*, [arXiv:2111.05981](#) [[INSPIRE](#)].
- [41] D. Berenstein and A. Holguin, *String junctions suspended between giants*, *JHEP* **11** (2022) 085 [[arXiv:2202.11729](#)] [[INSPIRE](#)].
- [42] W. Carlson, R. de Mello Koch and H. Lin, *Nonplanar integrability*, *JHEP* **03** (2011) 105 [[arXiv:1101.5404](#)] [[INSPIRE](#)].
- [43] R. de Mello Koch, M. Dessein, D. Giataganas and C. Mathwin, *Giant graviton oscillators*, *JHEP* **10** (2011) 009 [[arXiv:1108.2761](#)] [[INSPIRE](#)].
- [44] R. de Mello Koch, G. Kemp and S. Smith, *From large N nonplanar anomalous dimensions to open spring theory*, *Phys. Lett. B* **711** (2012) 398 [[arXiv:1111.1058](#)] [[INSPIRE](#)].
- [45] R. de Mello Koch and S. Ramgoolam, *A double coset ansatz for integrability in AdS/CFT*, *JHEP* **06** (2012) 083 [[arXiv:1204.2153](#)] [[INSPIRE](#)].
- [46] D. Berenstein, *Giant gravitons: a collective coordinate approach*, *Phys. Rev. D* **87** (2013) 126009 [[arXiv:1301.3519](#)] [[INSPIRE](#)].
- [47] D. Berenstein and E. Dzienkowski, *Open spin chains for giant gravitons and relativity*, *JHEP* **08** (2013) 047 [[arXiv:1305.2394](#)] [[INSPIRE](#)].
- [48] D. Berenstein, *Sketches of emergent geometry in the gauge/gravity duality*, *Fortsch. Phys.* **62** (2014) 776 [[arXiv:1404.7052](#)] [[INSPIRE](#)].
- [49] D. Berenstein, *On the central charge extension of the $N = 4$ SYM spin chain*, *JHEP* **05** (2015) 129 [[arXiv:1411.5921](#)] [[INSPIRE](#)].
- [50] Y. Jiang, S. Komatsu and E. Vescovi, *Structure constants in $N = 4$ SYM at finite coupling as worldsheet g -function*, *JHEP* **07** (2020) 037 [[arXiv:1906.07733](#)] [[INSPIRE](#)].
- [51] Y. Jiang, S. Komatsu and E. Vescovi, *Exact three-point functions of determinant operators in planar $N = 4$ supersymmetric Yang-Mills theory*, *Phys. Rev. Lett.* **123** (2019) 191601 [[arXiv:1907.11242](#)] [[INSPIRE](#)].

- [52] P. Yang, Y. Jiang, S. Komatsu and J.-B. Wu, *D-branes and orbit average*, *SciPost Phys.* **12** (2022) 055 [[arXiv:2103.16580](#)] [[INSPIRE](#)].
- [53] G. Chen, R. de Mello Koch, M. Kim and H.J.R. Van Zyl, *Absorption of closed strings by giant gravitons*, *JHEP* **10** (2019) 133 [[arXiv:1908.03553](#)] [[INSPIRE](#)].
- [54] G. Chen, R. De Mello Koch, M. Kim and H.J.R. Van Zyl, *Structure constants of heavy operators in ABJM and ABJ theory*, *Phys. Rev. D* **100** (2019) 086019 [[arXiv:1909.03215](#)] [[INSPIRE](#)].
- [55] D. Berenstein and S. Wang, *BPS coherent states and localization*, *JHEP* **08** (2022) 164 [[arXiv:2203.15820](#)] [[INSPIRE](#)].
- [56] A. Holguin and S. Wang, *Giant gravitons, Harish-Chandra integrals, and BPS states in symplectic and orthogonal $N = 4$ SYM*, *JHEP* **10** (2022) 078 [[arXiv:2206.00020](#)] [[INSPIRE](#)].
- [57] H. Lin, *Coherent state excitations and string-added coherent states in gauge-gravity correspondence*, *Nucl. Phys. B* **986** (2023) 116066 [[arXiv:2206.06524](#)] [[INSPIRE](#)].
- [58] A. Holguin and W.W. Weng, *Orbit averaging coherent states: holographic three-point functions of AdS giant gravitons*, [arXiv:2211.03805](#) [[INSPIRE](#)].
- [59] W. Carlson, R. de Mello Koch, E. Gandote and M. Kim, work in progress.
- [60] E. Witten, *Bound states of strings and p-branes*, *Nucl. Phys. B* **460** (1996) 335 [[hep-th/9510135](#)] [[INSPIRE](#)].
- [61] R. de Mello Koch, J.-H. Huang, M. Kim and H.J.R. Van Zyl, *Emergent Yang-Mills theory*, *JHEP* **10** (2020) 100 [[arXiv:2005.02731](#)] [[INSPIRE](#)].
- [62] S. Corley and S. Ramgoolam, *Finite factorization equations and sum rules for BPS correlators in $N = 4$ SYM theory*, *Nucl. Phys. B* **641** (2002) 131 [[hep-th/0205221](#)] [[INSPIRE](#)].
- [63] A.M. Vershik and S.V. Kerov, *Asymptotic theory of characters of the symmetric group*, *Funkt. Anal. Pril.* **15** (1981) 15 [*Funct. Anal. Appl.* **15** (1981) 246].
- [64] D.E. Berenstein, J.M. Maldacena and H.S. Nastase, *Strings in flat space and pp waves from $N = 4$ super Yang-Mills*, *JHEP* **04** (2002) 013 [[hep-th/0202021](#)] [[INSPIRE](#)].
- [65] R. de Mello Koch and R. Gwyn, *Giant graviton correlators from dual $SU(N)$ super Yang-Mills theory*, *JHEP* **11** (2004) 081 [[hep-th/0410236](#)] [[INSPIRE](#)].
- [66] N. Beisert, C. Kristjansen and M. Staudacher, *The dilatation operator of conformal $N = 4$ super Yang-Mills theory*, *Nucl. Phys. B* **664** (2003) 131 [[hep-th/0303060](#)] [[INSPIRE](#)].
- [67] I.G. Macdonald, *Symmetric functions and Hall polynomials*, second edition, Oxford University Press (1995).
- [68] V. De Comarmond, R. de Mello Koch and K. Jefferies, *Surprisingly simple spectra*, *JHEP* **02** (2011) 006 [[arXiv:1012.3884](#)] [[INSPIRE](#)].
- [69] J. Dai, R.G. Leigh and J. Polchinski, *New connections between string theories*, *Mod. Phys. Lett. A* **4** (1989) 2073 [[INSPIRE](#)].
- [70] S. de Carvalho, R. de Mello Koch and M. Kim, *Central charges for the double coset*, *JHEP* **05** (2020) 007 [[arXiv:2001.10181](#)] [[INSPIRE](#)].
- [71] C. Su, *Interacting emergent Yang-Mills theory*, *JHEP* **05** (2022) 008 [[arXiv:2203.03215](#)] [[INSPIRE](#)].