



CRITICAL CARE

Glucocorticoids with or without fludrocortisone in septic shock: a narrative review from a biochemical and molecular perspective

Gladness D. Nethathe^{1,2,3,*} , Jeffrey Lipman^{3,4,5}, Ronald Anderson⁶, Peter J. Fuller⁷  and Charles Feldman¹

¹School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa, ²Department of Anaesthesia and Perioperative Medicine, Royal Brisbane and Women's Hospital, Brisbane, QLD, Australia, ³Academy of Critical Care, Faculty of Medicine, University of Queensland, Brisbane, QLD, Australia, ⁴Jamieson Trauma Institute and Intensive Care Services, Royal Brisbane and Women's Hospital, Butterfield Street, Herston, Brisbane, 4029, QLD, Australia, ⁵Nimes University Hospital, University of Montpellier, Nimes, France, ⁶Department of Immunology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa and ⁷Endocrinology Unit, Monash Health, Centre for Endocrinology and Metabolism, Hudson Institute of Medical Research, Melbourne, VIC, Australia

*Corresponding author. E-mail: gladness.nethathe@wits.ac.za

Summary

Two randomised controlled trials have reported a reduction in mortality when adjunctive hydrocortisone is administered in combination with fludrocortisone compared with placebo in septic shock. A third trial did not support this finding when hydrocortisone administered in combination with fludrocortisone was compared with hydrocortisone alone. The underlying mechanisms for this mortality benefit remain poorly understood. We review the clinical implications and potential mechanisms derived from laboratory and clinical data underlying the beneficial role of adjunctive fludrocortisone with hydrocortisone supplementation in septic shock. Factors including distinct biological effects of glucocorticoids and mineralocorticoids, tissue-specific and mineralocorticoid receptor-independent effects of mineralocorticoids, and differences in downstream signalling pathways between mineralocorticoid and glucocorticoid binding at the mineralocorticoid receptor could contribute to this interaction. Furthermore, pharmacokinetic and pharmacodynamic disparities exist between aldosterone and its synthetic counterpart fludrocortisone, potentially influencing their effects. Pending publication of well-designed, randomised controlled trials, a molecular perspective offers valuable insights and guidance to help inform clinical strategies.

Keywords: adrenal insufficiency; corticosteroid insufficiency; critical illness; glucocorticoids; mineralocorticoids; stress response

Editor's key points

- A reduction in mortality has been reported in septic shock when adjunctive hydrocortisone is administered in combination with fludrocortisone. The mechanisms underpinning this mortality benefit remain unclear.
- The authors explored potential mechanisms derived from laboratory and clinical data, elucidating the

distinct biological effects of glucocorticoids and mineralocorticoids, and the tissue-specific and receptor-independent effects.

- An understanding of the interplay between glucocorticoid and mineralocorticoid action in critical illness could help guide future clinical strategies.

Received: 7 August 2023; Accepted: 24 October 2023

© 2023 The Author(s). Published by Elsevier Ltd on behalf of British Journal of Anaesthesia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

For Permissions, please email: permissions@elsevier.com

Adrenal insufficiency in critically ill patients is characterised by hypothalamic–pituitary–adrenal axis derangements and involves dysfunction in cortisol handling and the renin–angiotensin–aldosterone system (RAAS). Current guide lines suggest adding corticosteroid therapy for patients with septic shock requiring ongoing vasopressor support.^{1,2} Although adjunctive hydrocortisone (i.e. cortisol) alone has not shown improvement in mortality in septic shock, two randomised controlled trials have demonstrated a mortality improvement when fludrocortisone is combined with hydrocortisone.^{3–5} However, mechanisms underlying the observed comparative effectiveness of adjunctive fludrocortisone with hydrocortisone over hydrocortisone alone remain poorly understood.⁶

The Ger-Inf-05 trial examined the combination of fludrocortisone and hydrocortisone in septic shock.^{3,4} A total of 300 patients were assigned to receive either the combination of a 50-mg i.v. bolus of hydrocortisone 6 hourly with 0.05 mg fludrocortisone daily administered through a nasogastric tube or placebo for 7 days.³ Patients were classified as responders or non-responders based on a corticotropin test.³ Combined low-dose hydrocortisone with fludrocortisone was observed to significantly reduce the risk of death in non-responders with septic shock (73 deaths [63%] in the placebo group, 60 deaths [53%] in the corticosteroid group [hazard ratio, 0.67; 95% confidence interval or CI, 0.47–0.95; $P=0.02$]). No significant mortality benefit was observed in responders (18 deaths [53%] in the placebo group and 22 deaths [61%] in the corticosteroid group, $P=0.96$). These results suggested that combining mineralocorticoid with glucocorticoid therapy improved mortality outcomes in septic shock.

Recent large, randomised, controlled trials examining the role of hydrocortisone in septic shock have shown divergent effects on mortality with fludrocortisone in combination with hydrocortisone treatment. *Corticosteroid Treatment and Intensive Insulin Therapy for Septic Shock in Adults* (COITSS) was a randomised, 2×2 factorial trial of 509 patients with septic shock and multiple organ dysfunction assigned randomly to one of four groups: continuous i.v. insulin infusion with hydrocortisone alone, continuous i.v. insulin infusion with hydrocortisone and fludrocortisone, conventional insulin therapy with hydrocortisone alone, or conventional insulin therapy with i.v. hydrocortisone and fludrocortisone.⁷ The primary objective was to assess the efficacy of intensive insulin therapy in patients with septic shock treated with hydrocortisone. The secondary objective was to evaluate the additive benefit of fludrocortisone administered for 7 days. Hydrocortisone was administered as a 50-mg i.v. bolus, followed by 50 mg i.v. 6 hourly for 7 days. A total of 245 patients received additional oral fludrocortisone at a dose of 0.05 mg daily. Overall, 105 of 245 patients treated with fludrocortisone died (42.9%) and 121 of 264 (45.8%) patients in the control group died (risk reduction, 0.94; 95% CI, 0.77–1.14; $P=0.50$). These findings provided evidence against intensive insulin therapy in patients receiving hydrocortisone for septic shock and indicated a lack of significant mortality benefit from the additional fludrocortisone. Notably, the COITSS trial was underpowered as the mortality in the control group was underestimated and the sample size chosen to detect a large effect size (12.5% risk difference).⁵ Furthermore, sample size calculation was based on the estimated effects of the insulin intervention, but not on the effects of the added fludrocortisone.⁷ Thus, the trial was not powered to find a mortality difference with respect to the comparison of hydrocortisone with hydrocortisone plus fludrocortisone.

In the *Recombinant Human Activated Protein C and Low Dose of Hydrocortisone and Fludrocortisone in Adult Septic Shock* (APROGCHSS) trial, 614 patients were randomised to receive a combination of a hydrocortisone 50 mg i.v. bolus 6 hourly and fludrocortisone 0.05 mg daily through the nasogastric tube for 7 days, whereas 627 patients received hydrocortisone and fludrocortisone placebos.⁴ The 90-day all-cause mortality at ICU discharge was observed to be lower in the hydrocortisone-plus-fludrocortisone group than among those who received placebo rather than hydrocortisone and fludrocortisone (35.4% vs 41.0%, $P=0.04$). The hydrocortisone-plus-fludrocortisone group, additionally, had more vasopressor-free and organ failure-free days.

Nonetheless, two randomised, placebo-controlled clinical trials revealed a survival advantage in septic shock through a combination of hydrocortisone and the mineralocorticoid fludrocortisone, compared with placebo. Another smaller trial ($n=509$ patients) contrasted hydrocortisone alone with hydrocortisone in conjunction with fludrocortisone. The findings did not exhibit a significant decrease in mortality with the addition of fludrocortisone but did show a 3% numerically lower mortality with combination therapy (relative risk, 0.94; 95% CI, 0.77–1.14; $P=0.50$). Thus, the unresolved potential value of combining fludrocortisone with hydrocortisone for septic shock treatment persists.

Apparently discrepant findings from these three large trials have led to divergent perspectives on the benefits of adjunctive hydrocortisone in combination with fludrocortisone in septic shock. However, distinctions and the limited power of these studies make direct comparisons difficult with respect to hydrocortisone and fludrocortisone.^{1,6,8} Cortisol binds to both the glucocorticoid (GR) and the mineralocorticoid receptor (MR). One perspective suggests that as cortisol binds to the MR, the administration of high doses of hydrocortisone (≥ 200 mg daily) offers an adequate level of mineralocorticoid activity.^{1,6} A dose of 240 mg of hydrocortisone confers a mineralocorticoid effect equivalent to 1.2 mg (1200 μ g) daily of fludrocortisone and thus additional fludrocortisone therapy is not required.⁹ An alternative perspective is that adjunctive hydrocortisone with fludrocortisone treatment is required in septic shock as although cortisol binds to the MR, the subsequent signalling pathways differ from those mediated by aldosterone (and by implication fludrocortisone).

Pending a well-designed, randomised, controlled trial on the role of fludrocortisone in combination with hydrocortisone in septic shock, a molecular perspective offers valuable guidance for clinical strategies.

Biological rationale for the combination of hydrocortisone with fludrocortisone in the management of septic shock

Sepsis, which is a life-threatening time-dependent organ dysfunction, arises from a dysregulated host response to pathogens.^{10,11} This response manifests as an overwhelming systemic inflammatory response or immune dysfunction, and involves pattern recognition receptors sensing pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs).^{10,11} The resultant injury occurs at the molecular, cellular, and organ levels and involves the breakdown of both the endothelial and epithelial barriers, and the microvasculature, compromising multiple circulatory beds.¹² Various organ systems, including the cardiovascular and pulmonary systems, renal system, the coagulation cascade, and the immune system become compromised and multiorgan

failure ensues.^{10,12} In the blood–brain barrier, it causes perivascular oedema, oxidative stress, leukoencephalopathy, and neurotransmitter alterations.¹² The nervous system additionally plays an anti-inflammatory role, particularly in the early stages.¹² With regard to the immune system, this interaction leads to an excessive innate immune response.^{10,12} The persistence of a severe and prolonged compensatory anti-inflammatory innate immune response results in a state of acquired immunodeficiency termed immunoparalysis.¹³ Corticosteroids, recognised for their immunomodulatory properties, constitute an integral component of the host response to sepsis.^{10,14}

Septic shock initiates a neuro-humoral response, which includes the activation of the RAAS.^{15,16} The RAAS is a multi-organ system, which regulates homeostasis, inflammation, apoptosis, and fibrosis.¹⁷ The classical RAAS pathway is considered a cardiovascular circulating hormonal system mediated by angiotensin II (AII), and the non-classical RAAS pathway, a local tissue-based system mediated by angiotensin-(1–7). These two systems can operate synergistically or independently.^{18–20} In contrast to the classical RAAS and its circulating mediators, the non-classical RAAS comprises tissue axes, which regulate renal, cardiovascular, and nervous systems.^{21,22}

Angiotensin-(1–7) is found in the heart, renal, and brain tissues, and in plasma.¹⁹ Relative contributions of the classical (systemic) vs the non-classical (intra-renal) RAAS to blood pressure regulation remain debatable; however, the crosstalk between the RAAS and aldosterone production is well established.²¹ As such, these two systems should not be seen in isolation.²³

Both glucocorticoids and mineralocorticoids are integral components of the stress response.^{10,14,24} Notably, steroidogenesis pathways also demonstrate overlap between mineralocorticoid and corticosteroid production and the synthesis of corticosterone, an aldosterone precursor, is exclusively triggered by corticotropin.²⁵ As such, adrenal corticosteroid insufficiency is likely to occur concurrently with mineralocorticoid dysfunction. Physiologically, daily adrenal production of cortisol oscillates between 5 and 11 mg m⁻² day⁻¹ with a mean of 7 mg m⁻² day⁻¹, which equates to 20–30 mg daily of hydrocortisone in adults.^{26,27} Production rates are highly variable between and within individuals.^{26,27} Peak mean cortisol levels have been documented at 400 nmol L⁻¹ with nadir levels of <50 nmol L⁻¹. In adrenal insufficiency, the traditional replacement dose from consensus guidelines is 15–25 mg of hydrocortisone daily.^{28,29} In septic shock, the currently accepted dose for relative adrenal insufficiency is 100–200 mg daily.¹

In contrast, aldosterone secretion is 200- to 500-fold lower at a rate of 50–200 µg daily, with plasma levels ranging 40–210 µg L⁻¹.^{30,31} Isolated serum aldosterone measurement is of limited clinical value and is often interpreted alongside plasma renin activity.³²

Aldosterone administration has been demonstrated to restore vascular endothelial alpha-1 receptor expression, improving sensitivity to catecholamines in septic shock models.^{14,33} Fludrocortisone, a synthetic mineralocorticoid, may be used as replacement therapy for this purpose. The usual starting dose of fludrocortisone for adults with primary adrenal cortical insufficiency is 100 µg daily with a dose range of 100–500 µg.^{34,35} Higher doses may be required in high salt-losing states, in paediatric patients because of relative renal resistance to mineralocorticoids, and during the last trimester

of pregnancy because of the counter-regulatory effects of high progesterone levels.^{29,36–38}

Mechanisms for the beneficial role of adjunctive use of fludrocortisone with hydrocortisone in septic shock

Reasons for the mortality improvement observed in the APROCCHSS and Ger-Inf-05 trials remain poorly understood.⁶ Explanatory mechanisms include the mineralocorticoid activity of fludrocortisone, although high doses of cortisol purportedly possess such activity.^{9,39} Furthermore, aldosterone has non-mineralocorticoid effects, which potentially play a role.⁴⁰ Although pharmacologically different, a similar implication can be extended to its synthetic counterpart, fludrocortisone.

The additional glucocorticoid effects of fludrocortisone should also be considered.^{41,42} However, although fludrocortisone is a potent glucocorticoid,⁴¹ its effects are largely mineralocorticoid at the low dosage used because of its significantly greater potency at the MR. Additionally, various genomic and non-genomic effects of fludrocortisone, and a yet unidentified mechanism resulting from the combination of hydrocortisone and fludrocortisone should also be considered.

Aldosterone has non-mineralocorticoid effects: classical mineralocorticoid receptor-mediated signalling vs alternate modes of signalling

Aldosterone has genomic and non-genomic actions (Fig. 1).⁴⁵ Genomic actions are the classic nuclear actions occurring over hours in epithelial and non-epithelial cells attributable to activation of the MR, a ligand-dependent transcription factor.⁴⁵ Aldosterone also displays rapid actions occurring over minutes. Termed non-genomic or rapid non-nuclear actions, these are independent of gene transcription and occur in classical MR targets and non-epithelial tissue and include actions such as the activation of sodium transport in target cells.^{46,47} Rapid actions are clinically relevant in the acute setting as a result of their role in increasing peripheral vascular resistance and blood pressure.⁴⁸

Notably, plasma aldosterone levels, not cortisol, rapidly change with posture as part of normal blood pressure postural control.^{49,50} This suggests rapid cellular responses mediated by the acute increase in aldosterone and a role in blood pressure control in the acute setting.⁴⁸ Non-genomic actions were thought to be mediated through membrane-bound MR acting through second messenger pathways.⁴⁶ The G protein-coupled receptor, GPR30, or the G protein-coupled oestrogen receptor-1 (GPER-1), has been implicated as the receptor that mediates some of the rapid actions of aldosterone on vascular smooth muscle.^{45,51–54} Experimentally, aldosterone has also been demonstrated to potentiate the constrictor effect of AII on vascular smooth muscle, which is mediated through the angiotensin receptor type 1 (AT1). Experimental evidence further suggests that GPR30 (GPER-1) has a high affinity for aldosterone but does not bind cortisol.⁴³

Clinically, aldosterone (and by implication fludrocortisone) thus has MR-independent actions that are relevant in the acute setting with an effect on vascular smooth muscle and blood pressure control. The clinical rationale for fludrocortisone in combination with hydrocortisone therapy in critical illness should thereby not solely relate to aldosterone or fludrocortisone activation of the renal epithelial MR.^{6,8}




Fig 1. Genomic and non-genomic actions of aldosterone. Key features distinguishing genomic and non-genomic actions of aldosterone. Aldosterone action; genomic and non-genomic effects of aldosterone and cortisol. Classical slow genomic actions mediated through the mineralocorticoid receptor (MR) result in gene transcription and the production of effector proteins. Fast actions mediated through a surface receptor, which interact with the classical genomic actions, have recently been described to be mediated through the G protein-coupled receptor GPR30 (GPER), possibly through a membrane-bound MR, or both. Aldosterone action through GPR30 (GPER) is currently understood to be specific for aldosterone. BP, blood pressure; c-SRC, non-receptor tyrosine kinase c-SRC protein; ERK1/2, extracellular-signal regulated kinase 1/2; JNK, c-Jun NH2-terminal kinase; PI3-K, Phosphatidylinositol 3-kinase. Reused with permission from AME Publishing Company.^{43,44}

The multifarious mineralocorticoid receptor

The MR and GR are members of the nuclear receptor superfamily, which encompasses seven subfamilies (sub-group 0 to sub-group 6) and 48 members.⁵⁵ They act as ligand-activated transcription factors and consist of four structurally distinct domains namely: the amino terminal, a central DNA-binding

domain, the hinge region, and the C-terminal ligand-binding domains.⁵⁶ Lipophilic ligands including steroids, retinoids, phospholipids, and as yet unidentified ligands (orphan members) regulate their activity.^{55,57} Steroid receptors belong to sub-group 3, which includes the GR, MR, androgen receptor, progesterone receptor, and the oestrogen receptors (ERs): ER α and ER β .⁵⁵ The MR and GR amino acid sequences exhibit

significant homology being 56% identical in the steroid-binding domain.⁵⁶ Distinctively, the MR has the longest amino terminal domain among steroid receptors, sharing less than 15% homology with the GR in this region.⁵⁶ Additionally, the MR has affinity for mineralocorticoids, endogenous glucocorticoids, and progesterone.^{56,58–60}

In view of higher circulating levels of cortisol and equal affinity of the MR for cortisol and aldosterone, a system for receptor selectivity exists in some tissues. This is mediated at a pre-receptor level in mineralocorticoid target tissues by the 11 β -hydroxysteroid dehydrogenase (11 β -HSD) enzyme system.⁵⁸ Two 11 β -HSD isoenzymes exist.^{61,62} First, 11 β -HSD type-1 (11 β -HSD1) is a bi-directional enzyme *in vitro* and an activating enzyme *in vivo* by converting receptor-inactive cortisone to cortisol.⁶² In contrast, 11 β -HSD type-2 (11 β -HSD2) is expressed in mineralocorticoid target tissues and is a deactivating enzyme through its conversion of cortisol to cortisone.^{50–52,63} The role of 11 β -HSD2 is to protect the MR from cortisol in mineralocorticoid target tissues, conferring specificity for aldosterone.^{63,64} The observation that both cortisol and aldosterone have a high affinity for the MR indicates a lack of selectivity of the C-terminal ligand-binding domain for aldosterone (or cortisol), but not necessarily comparative activation of the receptor.^{56,58–60} Notably, although the GR is activated by cortisol alone, not all glucocorticoids activate the MR. Synthetic glucocorticoids show minimal or no activity at the MR.^{65–70}

The observation that different ligands have variable activity at the MR, and that not all glucocorticoids activate the MR, is supportive of the rationale that the choice of corticosteroid supplementation potentially influences clinical outcomes and that cortisol and aldosterone (and by inference their synthetic counterparts) do not necessarily have the same effect as ligands at the MR.

Mineralocorticoid receptors are capable of divergent responses to different ligands

Glucocorticoid conformational mobility is another factor that confers mineralocorticoid action of glucocorticoids on the MR.^{71–74} The molecular skeleton of steroids consists of four carbon atom rings defined as rings A, B, C, and D.⁷⁵ Steroid ring A conveys glucocorticoid action and is rigid in *pure* glucocorticoids and flexible in aldosterone. The structure of steroid ring C is required for mineralocorticoid action and is rigid in aldosterone (and relatively flexible in *pure* glucocorticoids). The exception is 11-deoxycorticosterone and its synthetic derivative, delta-11,12-deoxycorticosterone, which has a flexible C ring (and thus should be a *pure* glucocorticoid), but is actually a specific mineralocorticoid with marked glucocorticoid activity.^{69,76} The lack of C-11 oxygenation of 11-deoxycorticosterone confers versatility.^{76,77}

The MR exhibits ligand- and tissue-specific dichotomy. Cortisol's actions on the MR, although primarily agonist, can be antagonistic depending on the underlying pathophysiology.^{78–81} The latter has been reported under certain experimental conditions in tissues in which 11 β -HSD2 is not expressed (heart and specific regions in the central nervous system).^{78–81} In conditions of tissue damage and in the presence of reactive oxygen species, cortisol acts as MR agonist.^{80–82} Thus, functions of the MR are diverse, tightly regulated by ligand-specifying mechanisms, and are tissue-dependent.^{61,83,84} Furthermore, MRs bound to aldosterone are demonstrably more resistant to proteolysis than MRs bound to

cortisol, a result of ligand-induced conformational changes within the receptor protein.⁸⁵ Notably, the MR bound to fludrocortisone, aldosterone's synthetic counterpart, is highly resistant to degradation.^{86–88} Thus, the MR is capable of divergent responses to different ligands.^{39,89} Furthermore, receptor–ligand interactions differ according to ligand, and mineralocorticoid supplementation may help optimise MR activation.

Downstream signalling pathways between aldosterone and cortisol binding at the mineralocorticoid receptor differ

Although the MR binds multiple ligands, unique mineralocorticoid and glucocorticoid effects are evident. Their main ligands, aldosterone and cortisol, perform diverse roles and are governed by different regulatory mechanisms.^{90–92} The effects of cortisol are influenced by various factors, including whether the signalling occurs through the GR, MR, or perhaps both.⁹³ Furthermore, therapeutic agents such as spironolactone, a competitive antagonist with weak agonist activity at the MR, are able to exert their protective effects not only through the exclusion of cortisol or aldosterone from the MR but potentially through the induction of protective factors.⁹⁴ Differences in differential downstream signalling pathways that have been observed when the MR is bound to aldosterone compared with cortisol are also likely to be cell- and tissue-specific.

Mineralocorticoid activity in the regulation of salt and water balance

Aldosterone release is the final step in the classical RAAS signalling system. In renal 11 β -HSD2-expressing cells, cortisol is metabolised to bio-inactive cortisone. This pre-receptor exclusion of cortisol in 11 β -HSD2-expressing tissues renders the MR aldosterone-selective, suggesting that cortisol and aldosterone may have similar or overlapping effects on the MR 11 β -HSD2-expressing tissues.⁹⁵ This implicates the utilisation of similar signalling pathways between cortisol and aldosterone, specifically with respect to sodium and potassium regulation. The process of filtered sodium (Na⁺) reabsorption in collecting duct cells begins with luminal Na⁺ crossing the apical plasma membrane through epithelial sodium channels (ENaC) driven by its electrochemical gradient.⁹⁶ Subsequently, intracellular Na⁺ is actively transported into the interstitial space by the basolateral Na⁺/K⁺-ATPase. Aldosterone-induced genes modulate ENaC subunit turnover, enhance ENaC synthesis, ENaC activity, and stimulate Na⁺/K⁺-ATPase activity.^{71,96–98} The interaction of aldosterone with the MR leads to induction or repression of additional aldosterone-regulated genes, including serum and glucocorticoid-induced kinase (SGK1), which also play a key role in the regulation of Na⁺ transport in distal kidney nephron epithelia.⁹⁹ SGK1 is the main regulator of ENaC.¹⁰⁰ Additionally, aldosterone has been found to have acute, non-mineralocorticoid effects on GPR30, which are not observed with cortisol.^{51,53}

Mineralocorticoid activity in the brain

Neuroinflammation, along with ischaemic injury, neuronal cell death, and neuroanatomy changes, are major factors associated with cerebral dysfunction secondary to sepsis and septic shock.¹⁰¹ Inflammation associated with sepsis and

septic shock has a direct impact on the structural integrity of the hippocampus and prefrontal cortex.^{102,103} No effective therapeutic interventions exist; however, there is evidence to suggest that elevated levels of glucocorticoids may exacerbate neural inflammation, and evidence to the contrary.^{102–105} The role of glucocorticoids, acting through both MR and GR, in modulating the immune response has been well studied.¹⁰ Although, interest in the potential immune effects of mineralocorticoids has only recently emerged, the general consensus is that there is a role for both GR and MR in modulating the neuroinflammation associated with sepsis and septic shock.^{10,102}

Compared with the GR, which is expressed ubiquitously in neurones and glial cells, the MR has a more restricted distribution, with the highest expression in limbic structures.^{102,106} Aldosterone-selective neurones are limited to areas (periventricular, brain stem nucleus tractus solitarius) that regulate haemodynamic, fluid, and electrolyte homeostasis.⁹⁰

Brain aldosterone levels are normally low because of the absence of aldosterone synthase and poor penetration across the blood–brain barrier caused by the presence of P-glycoprotein, an ATP-driven, drug efflux pump, which transports certain substrates across the cerebral vascular endothelium and into the circulation.^{106–108} In contrast, cortisol (or corticosterone in rodents) circulates at significantly higher levels (100–1000 fold), acting as the main ligand for brain MR and GR.¹⁰⁹ Although MR signalling in the brain has been demonstrated to induce a pro-inflammatory response, the results of MR activation in the brain can vary depending on the ligand.^{10,110}

Numerous studies show the diverse impacts of cortisol and aldosterone on brain MR.^{111,112} Despite this variability, experimental data indicate that both hormones, when interacting with the brain cortisol-responsive MR, can influence cellular activity in similar ways through both genomic and non-genomic pathways.¹¹³ Notably, these neurones also contain 11 β -HSD1, which regenerates cortisol. However, it is essential to consider that poor penetration of the blood–brain barrier by aldosterone may be a limiting factor in its effects on brain MR activation.

Indirect evidence of downstream signalling differences is seen in murine models, where corticosterone antagonises aldosterone-mediated hypertension at the brain MR.¹¹⁴ The brain shows widespread 11 β -HSD-1 expression, whereas 11 β -HSD-2 is limited to neurones in the fasciculus solitarius.^{61,70,106} Aldosterone interactions with these neurones regulate salt appetite.¹¹² The subsequent blood pressure response to dietary sodium is inhibited by spironolactone.¹¹⁵ The presence of 11 β -HSD-1 and lack of 11 β -HSD-2 imply significant roles for both cortisol and aldosterone in the brain, beyond salt regulation. In these cells, salt regulation is not the primary function, providing indirect evidence of the requirement for both cortisol and aldosterone in those tissues, although the specific role of aldosterone in these tissues is not well defined.

Our hypothesis is that in aldosterone-selective tissues, downstream signalling mechanisms are similar between the two ligands, hence the pre-receptor exclusion of cortisol to prevent MR activation by cortisol, which is ubiquitous. In non-aldosterone selective tissues, downstream signalling mechanisms activated by the MR differ between ligands, and pre-receptor modulation of access is therefore not required. These variable downstream actions resulting from aldosterone

and cortisol binding at the MR may extend to their synthetic counterparts.

Mineralocorticoid activity in the heart

Differences in aldosterone and corticosterone signalling have been described in murine cardiomyocytes.^{116,117} Both GR and MR are present in the heart. In a murine model, the activation of MR and GR resulted in both common and distinct responses, suggesting similarities and differences in signalling. In a study using isolated neonate cardiomyocyte cells *in vitro*, both aldosterone and corticosterone induced acceleration of spontaneous cardiomyocyte contractions with a chronotropic response to corticosteroids mediated by both MR and GR.^{116,117}

Increased chronotropy was dependent on low threshold T-type calcium channel expression, which was disrupted by GR antagonism, but not by MR antagonism. Aldosterone's chronotropic action was further found to be additive to forskolin's effect on intracellular cyclic adenosine monophosphate (cAMP) levels, an additive response selectively abolished by GR inhibition.¹¹⁸ Moreover, GR antagonism prevented cardiomyocyte hypertrophy induced by aldosterone, whereas MR antagonism (spironolactone) had limited effect. This suggests that, in murine cardiomyocytes, complete electrical remodeling and the maximal chronotropic response to corticosteroids require both GR and MR activation, with the GR alone involved in sensitising cells to aldosterone's chronotropic regulation through the cAMP pathway and in the hypertrophic response.¹¹⁷

Further evidence suggestive of signalling differences between cortisol and aldosterone in cardiomyocytes include well-established recognition of rapid non-genomic effects unique to aldosterone, not mimicked by cortisol.¹¹⁹ Acute non-genomic effects of aldosterone in cardiac and other tissues mimicked by classical mineralocorticoid agonists, fludrocortisone, and deoxycorticosterone, but not by cortisol, have been demonstrated.¹¹⁹ These effects are not blocked by spironolactone.¹¹⁹ In a murine cardiomyocyte model, aldosterone specifically increased [³H]leucine incorporation into protein, which was not observed with MR occupancy by corticosterone.¹²⁰

The observation that reduced GR signalling and increased MR signalling are associated with an increased risk of heart disease, whereas alterations that favour increased GR signalling and diminished MR signalling exhibit cardioprotective effects, provides further indirect evidence of differences in signalling between glucocorticoids and mineralocorticoids.¹²¹ GR and MR have complementary roles in regulating the hypothalamic–pituitary axis. The balance of glucocorticoid signalling through GR and MR has been demonstrated to play a role in disease.^{90,92,121} Increased GR signalling and decreased MR signalling is considered cardioprotective, with unopposed MR signalling, irrespective of ligand (cortisol or aldosterone), associated with accelerated cardiovascular disease in experimental models. These responses can also be viewed as being anti-inflammatory or pro-inflammatory, respectively.^{94,122}

Molecular interactions of the MR that differ by ligand have been described, providing compelling evidence of distinct MR conformation when bound to aldosterone vs cortisol.^{91,109,123–125}

Collectively, these findings support the view that downstream signalling pathways vary depending on whether the MR is bound to aldosterone or cortisol and emphasise the

clinical implications of unique signalling pathways for cortisol, aldosterone, and their synthetic analogues. Although these represent potential mechanisms and, in a sense, proof-of-concept, a link to specific responses or outputs remains to be fully elucidated in humans.

Aldosterone versus fludrocortisone mechanism of action: is there a difference?

Synthetic hormones structurally vary from physiological steroid hormones, leading to differing effects.^{126–128} Although they may mimic selected biological actions of endogenous hormones, they may not exert identical effects at a molecular level.^{90,126,127,129–133}

Pharmacokinetic differences between synthetic and physiological steroid hormones are evident in their binding to corticosteroid-binding globulin (CBG). The majority of cortisol (>90%) is bound to CBG, whereas synthetic glucocorticoids (prednisone, prednisolone, dexamethasone, fludrocortisone) have lower CBG affinity, leading to increased free fractions.^{67,134} This difference in bioavailability supports additive glucocorticoid activity as another mechanism explaining the mortality improvement in patients administered hydrocortisone in combination with fludrocortisone in septic shock. In contrast to fludrocortisone, aldosterone plays a minimal role in glycogen deposition, glycogen phosphorylation, glycogenolysis, insulin resistance, or sensitivity.⁷⁶ Thus, additive fludrocortisone may exacerbate hyperglycaemia compared with hydrocortisone alone.

Moreover, fludrocortisone is a synthetic steroid with potent mineralocorticoid and substantial glucocorticoid activity (Fig. 2a) which was developed from the addition of a fluorine molecule at the 9- α -position to hydrocortisone (Fig. 2b and c).^{76,87,136–139} It is a structural analogue of hydrocortisone with the same A ring, but is functionally similar to aldosterone.¹⁴⁰ Fluorination results in inert bonds, a prolonged half-life, improved bioavailability, and increased affinity for GR.^{86–88} Fludrocortisone has similar effects to hydrocortisone, but has marked effects on electrolyte balance and carbohydrate metabolism with 10-fold higher glucocorticoid activity than hydrocortisone.⁸⁶

Although human comparative pharmacokinetic and pharmacodynamic data for aldosterone and fludrocortisone dosing are limited, during the characterisation of the novel MR modulator, AZD9977, in a murine model, distinct mineralocorticoid effects of fludrocortisone compared with aldosterone were observed.^{76,133} These effects were subsequently extrapolated to humans using preclinical and clinical data modelling. The mechanism of this effect remains unexplained.

Comparison of aldosterone and fludrocortisone revealed distinct effects.¹²⁸ A 1-mg dose of intramuscular aldosterone exhibited an electrolyte regulating effect equivalent to a daily oral dose of 250 μ g of fludrocortisone.¹²⁸ However, fludrocortisone exerted a more prolonged effect. A 1-mg dose of intramuscular aldosterone showed a greater effect on eosinopenia than 1-mg dose of oral fludrocortisone, indicating that fludrocortisone's effect on immune cell function requires much higher doses than aldosterone.^{128,141} Notably, differences in the administration routes of aldosterone and fludrocortisone may have contributed to the observed effects. Fludrocortisone also exhibited pituitary adrenocorticotropic




Fig 2. Molecular structure of (a) aldosterone (DB04630), (b) fludrocortisone (DB00687), and (c) hydrocortisone (DB00741).¹³⁵ Molecular structures are depicted in a standard format. Distinguishing features between depicted steroids: fludrocortisone has a fluorine atom on C9 replacing hydrogen and aldosterone has an aldehyde group on C18. Bond line notation: double lines indicate a double bond, single arrows indicate a single bond, dashed lines indicate that the bond extends behind the surface plane, bold wedged lines indicate bonds protruding out of the surface plane. DB, DrugBank Accession number.

hormone (ACTH) inhibition, consistent with glucocorticoid activity, an effect not observed with aldosterone, providing further mechanisms whereby fludrocortisone may compound the hyperglycaemia and immune changes observed with hydrocortisone administration.^{3,4,128,142}

Thus, although both aldosterone and fludrocortisone are classified as mineralocorticoids, their pharmacological actions differ and may translate into distinct clinical effects. However, although both fludrocortisone and aldosterone have demonstrable non-genomic effects in murine models, evidence for this mechanism in humans as an explanation for the mortality improvement observed with adjunctive fludrocortisone with hydrocortisone in septic shock is minimal.¹⁴³

Role of adrenal age-related changes and mineralocorticoid supplementation

The traditional concept of compartmentalised enzymatic reactions within distinct zones of the adrenal cortex, known as functional zonation, has been challenged.¹⁴⁴ Adrenal cortex zones, namely the zona glomerulosa, zona fasciculata, and zona reticularis, were thought to have distinct roles in synthesising aldosterone, glucocorticoids, and androgens, respectively.^{145,146} The distribution of enzymes and receptors does not, however, suggest autonomous steroid production by each adrenal zone. This is particularly relevant for the glomerulosa, which seemingly lacks the complete suite of the necessary enzymes for aldosterone biosynthesis.^{147,148} Hence, the adrenal gland, seen as an integrated structure, diminishes the likelihood of diseases such as septic shock being restricted to a particular zone.¹⁴⁷

Furthermore, with progressing age (from about 30 yr), adrenal topographical changes occur, resulting in zona fasciculata enlargement, reduction in zona reticularis and glomerulosa cell masses, and the development of aldosterone-synthase expressing cell clusters at the border of the zona glomerulosa with the zona fasciculata.^{147,149,150} Such histological changes suggest reduced functional differentiation between zones. Of clinical relevance is that the reduction in the aldosterone-producing cell mass and observed decrease in aldosterone release with advancing age suggests that adjunctive mineralocorticoid supplementation in septic shock may be more effective in the older population.^{149–151} This aligns with the demographic profile of participants in the APROCCHSS trial, where the average age was 66 yr.⁴ Thus, adjunctive hydrocortisone and fludrocortisone therapy may hold greater promise for improving outcomes in septic shock in older patients.

Impact of bioavailability of orally administered fludrocortisone

Fludrocortisone differs from other corticosteroids in its 100% bioavailability and long half-life (18–36 h) and duration of action (1–2 days). It is dosed orally and is stable as a solid.¹⁴⁰ However, pharmacokinetic data on the bioavailability of oral fludrocortisone in critical illness is limited. In one-third of patients with septic shock, a single oral dose of 50 µg led to undetectable plasma levels in one study.¹⁵² The usual replacement dose of fludrocortisone in primary adrenal insufficiency is 100–200 µg, a dose significantly higher than the 50 µg administered in studies conducted in septic shock. More recently, an average of 100 µg fludrocortisone has been used in septic shock.⁵ Additionally, the different minera-

locorticoid effects of synthetic fludrocortisone compared with endogenous aldosterone suggested by preclinical data highlights that translation of the mineralocorticoid effects of fludrocortisone to those of aldosterone requires further exploration.^{133,153}

Clinical applications and suggestions for clinical practice

Which population is likely to benefit?

The decline of the aldosterone-producing cell mass with age and the consequent reduction in aldosterone release suggests that adjunctive mineralocorticoid supplementation in septic shock may be more efficacious in the older population. Based on the demographic profile of participants in the APROCCHSS trial, consideration of adjunctive mineralocorticoid therapy in septic shock may be more beneficial in the population group above 60 yr of age.

Dosage and mode of fludrocortisone administration

Current recommendations suggest adjunctive hydrocortisone at a non-tapered dose of 200 mg daily for 7 days if used in septic shock. If adjunctive fludrocortisone therapy is to be commenced in combination with hydrocortisone, we suggest a dose of 50 µg daily for 7 days without tapering. Notably, a recent international survey of clinician preferences for corticosteroid prescription in septic shock suggests a low preference for fludrocortisone.¹⁵⁴ Well-powered RCTs to investigate patient selection, optimal dosage, and treatment regimen are required before any firm recommendations can be made regarding fludrocortisone dosing.

Adverse effects of hydrocortisone and fludrocortisone use

Risks associated with adjunctive corticosteroid therapy in severe sepsis and septic shock include gastroduodenal bleeding, superinfection, neuromuscular weakness, hyperglycaemia, and hypernatraemia.¹⁴² Findings of a systematic review on corticosteroids in the treatment of severe sepsis and septic shock in adults suggested no significant differences in rates of gastroduodenal bleeding, superinfections, or neuromuscular weakness between treated and control patients. However, there were higher risks of hyperglycaemia (nine trials; 51.6% in the treatment group [363 out of 703] vs 46% in the control group [308 out of 670]; $P < 0.001$; $I^2 = 0\%$) and hypernatraemia (three trials; 31.4% in the treatment group [127 out of 404] vs 19.2% in the control group [77 out of 401]; $P < 0.001$; $I^2 = 0\%$) observed in the treated group.¹⁴² In the APROCCHSS trial, the use of hydrocortisone in combination with fludrocortisone significantly increased the risk of hyperglycaemia (relative risk, 1.07; 95% CI, 1.03–1.12; $P = 0.002$). There were no significant differences in the risks of gastroduodenal bleeding (relative risk, 0.88; 95% CI, 0.58–1.34; $P = 0.56$) or superinfection (relative risk, 1.09; 95% CI, 0.92–1.30; $P = 0.30$) when comparing hydrocortisone plus fludrocortisone with placebo. The relatively minimal adverse implications, along with the substantial potential benefit, would seem to favour adjunctive hydrocortisone and fludrocortisone administration in septic shock.

Conclusions

Human and laboratory data provide insights into potential mechanisms explaining the beneficial role of adjunctive

hydrocortisone with fludrocortisone in septic shock. Cortisol and aldosterone actions at the MR differ and by implication their synthetic counterparts, however, this has not yet translated into compelling clinical evidence of effect. Aldosterone exhibits mineralocorticoid and clinically relevant non-mineralocorticoid effects; thus, when considering potential beneficial mechanisms, the focus should not solely be on the activity of aldosterone or fludrocortisone at the MR. Signalling pathways activated by aldosterone on the MR are likely distinct from those of fludrocortisone. Consequently, specific effects of adjunctive fludrocortisone in combination with hydrocortisone in septic shock are still not well defined.

The precise targets of mineralocorticoid therapy within the context of sepsis require further elucidation. Additionally, the effect of 'pro-inflammatory' mineralocorticoid actions in relation to immunoparalysis in septic shock and on vascular reactivity requires further elucidation. Additionally, there is clearly a need for further investigation into the pharmacokinetics of orally administered fludrocortisone in critical illness.

Authors' contributions

Researched the scientific literature: GDN

Read and approved the final manuscript: all authors

Declaration of interest

The authors declare that they have no competing interests.

Funding

The National Research Foundation of South Africa, Pretoria, South Africa and The Discovery Foundation, Braamfontein, Johannesburg, South Africa (to GDN).

References

- Cohen J, Venkatesh B. Adjunctive corticosteroid treatment in septic shock. *Anesthesiol J Am Soc Anesthesiol* 2019; **131**: 410–9
- Evans L, Rhodes A, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Crit Care Med* 2021; **49**: e1063–143
- Annane D, Sébille V, Charpentier C, et al. Effect of treatment with low doses of hydrocortisone and fludrocortisone on mortality in patients with septic shock. *JAMA* 2002; **288**: 862–71
- Annane D, Renault A, Brun-Buisson C, et al. Hydrocortisone plus fludrocortisone for adults with septic shock. *N Engl J Med* 2018; **378**: 809–18
- Bosch NA, Teja B, Law AC, Pang B, Jafarzadeh SR, Walkey AJ. Comparative effectiveness of fludrocortisone and hydrocortisone vs hydrocortisone alone among patients with septic shock. *JAMA Intern Med* 2023; **183**: 451
- Venkatesh B, Finfer S, Myburgh J, ADRENAL investigators. Glucocorticoids with or without fludrocortisone in septic shock. *N Engl J Med* 2018; **379**: 893–6
- Annane D, Cariou A, Maxime V, et al. COIITSS Study Investigators, Corticosteroid treatment and intensive insulin therapy for septic shock in adults: a randomized controlled trial. *JAMA* 2010; **303**: 341–8
- Yamamoto R, Nahara I, Toyosaki M, Fukuda T, Masuda Y, Fujishima S. Hydrocortisone with fludrocortisone for septic shock: a systematic review and meta-analysis. *Acute Med Surg* 2020; **7**: e563
- Ramsahoye BH, Davies SV, El-Gaylani N, Sandeman D, Scanlon MF. The mineralocorticoid effects of high dose hydrocortisone. *BMJ* 1995; **310**: 656–7
- Heming N, Sivanandamoorthy S, Meng P, Bounab R, Annane D. Immune effects of corticosteroids in sepsis. *Front Immunol* 2018; **9**: 1736
- Berghe GV den, Téblick A, Langouche L, Gunst J. The hypothalamus-pituitary-adrenal axis in sepsis- and hyperinflammation-induced critical illness: gaps in current knowledge and future translational research directions. *eBioMedicine* 2022; **84**, 104284
- Gotts JE, Matthay MA. Sepsis: pathophysiology and clinical management. *BMJ* 2016; **353**: i1585
- Frazier WJ, Hall MW. Immunoparalysis and adverse outcomes from critical illness. *Pediatr Clin North Am* 2008; **55**: 647–68
- Ramanan M, Cohen J, Venkatesh B. Steroids and sepsis: the debate continues. *Int Anesthesiol Clin* 2019; **57**: 17–30
- Corrêa TD, Takala J, Jakob SM. Angiotensin II in septic shock. *Crit Care* 2015; **19**: 98
- Antonucci E, Gleeson PJ, Annoni F, et al. Angiotensin II in refractory septic shock. *Shock* 2017; **47**: 560–6
- Fountain JH, Lappin SL. *Physiology, renin angiotensin system*, StatPearls. Treasure Island, FL: StatPearls Publishing; 2020. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK470410/>. [Accessed 26 October 2022]
- Kaparianos A, Argyropoulou E. Local renin-angiotensin II systems, angiotensin-converting enzyme and its homologue ACE2: their potential role in the pathogenesis of chronic obstructive pulmonary diseases, pulmonary hypertension and acute respiratory distress syndrome. *Curr Med Chem* 2011; **18**: 3506–15
- Bitker L, Burrell LM. Classic and nonclassic renin-angiotensin systems in the critically ill. *Crit Care Clin* 2019; **35**: 213–27
- Mascolo A, Scavone C, Rafaniello C, et al. The role of renin-angiotensin-aldosterone system in the heart and lung: focus on COVID-19. *Front Pharmacol* 2021; **12**, 667254
- Zhuo JL, Ferrao FM, Zheng Y, Li XC. New frontiers in the intrarenal renin-angiotensin system: a critical review of classical and new paradigms. *Front Endocrinol* 2013; **4**: 166
- Santos RA. Angiotensin-(1–7). *Hypertension* 2014; **63**: 1138–47
- Briet M, Schiffrin EL. Vascular actions of aldosterone. *J Vasc Res* 2013; **50**: 89–99
- Tolstoy NS, Aized M, McMonagle MP, et al. Mineralocorticoid deficiency in hemorrhagic shock. *J Surg Res* 2013; **180**: 232–7
- Briegel J, Möhnle P, Keh D, et al. Corticotropin-stimulated steroid profiles to predict shock development and mortality in sepsis: from the HYPRESS study. *Crit Care* 2022; **26**: 343
- Seo KH. Perioperative glucocorticoid management based on current evidence. *Anesth Pain Med* 2021; **16**: 8–15
- Oprea A, Bonnet NCG, Pollé O, Lysy PA. Novel insights into glucocorticoid replacement therapy for pediatric and adult adrenal insufficiency. *Ther Adv Endocrinol Metab* 2019; **10**, 204201881882129

28. Caetano CM, Sliwiska A, Madhavan P, Grady J, Malchoff CD. Empiric determination of the daily glucocorticoid replacement dose in adrenal insufficiency. *J Endocr Soc* 2020; 4: bvaa145
29. Bornstein SR, Barthel A, Husebye ES, et al. Diagnosis and treatment of primary adrenal insufficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2016; 101: 364–89
30. Pearce D, Bhalla V, Funder JW. *Aldosterone and mineralocorticoid receptors: renal and extrarenal roles, brenner and rector's the kidney*. 11th ed. Elsevier; 2020. p. 335–56
31. Auchus RJ, Parker KL. *The adrenal glands, Textbook of endocrine physiology*. 6th Edn. New York: Oxford University Press; 2011
32. Amar L, Plouin P-F, Steichen O. Aldosterone-producing adenoma and other surgically correctable forms of primary aldosteronism. *Orphanet J Rare Dis* 2010; 5: 9
33. Black RE, Laxminarayan R, Temmerman M, Walker N. *Reproductive, maternal, newborn, and child health: disease control priorities, third edition (volume 2)*. Washington (DC): The International Bank for Reconstruction and Development/The World Bank; 2016 Apr 5. <https://doi.org/10.1596/978-1-4648-0348-2>. PMID: 27227205
34. Esposito D, Pasquali D, Johannsson G. Primary adrenal insufficiency: managing mineralocorticoid replacement therapy. *J Clin Endocrinol Metab* 2018; 103: 376–87
35. Rahman M, Anjum F. *Fludrocortisone, StatPearls*. Treasure Island, FL: StatPearls Publishing; 2022. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK564331/>. [Accessed 24 March 2022]
36. Bowden SA, Henry R. Pediatric adrenal insufficiency: diagnosis, management, and new therapies. *Int J Pediatr* 2018; 2018, 1739831
37. Albanese A, Hindmarsh P, Stanhope R. Management of hyponatraemia in patients with acute cerebral insults. *Arch Dis Child* 2001; 85: 246–51
38. Grossman A, Johannsson G, Quinkler M, Zelissen P. Therapy of endocrine disease: perspectives on the management of adrenal insufficiency: clinical insights from across Europe. *Eur J Endocrinol* 2013; 169: R165–75
39. Nethathe GD, Cohen J, Lipman J, Anderson R, Feldman C. Mineralocorticoid dysfunction during critical illness: a review of the evidence. *Anesthesiology* 2020; 133: 439–57
40. Williams JS, Williams GH. 50th anniversary of aldosterone. *J Clin Endocrinol Metab* 2003; 88: 2364–72
41. Grossmann C, Scholz T, Rochel M, et al. Transactivation via the human glucocorticoid and mineralocorticoid receptor by therapeutically used steroids in CV-1 cells: a comparison of their glucocorticoid and mineralocorticoid properties. *Eur J Endocrinol* 2004; 151: 397–406
42. Sutanto W, De Kloet ER. Mineralocorticoid receptor ligands: biochemical, pharmacological, and clinical aspects. *Med Res Rev* 1991; 11: 617–39
43. Gomez-Sanchez E, Gomez-Sanchez CE. The multifaceted mineralocorticoid receptor. *Compr Physiol* 2014; 4: 965–94
44. Nethathe GD, Lipman J, Anderson R, Feldman C. CIRMI—a new term for a concept worthy of further exploration: a narrative review. *Ann Transl Med* 2022; 10: 646
45. Hermidorff MM, Assis LVM de, Isoldi MC. Genomic and rapid effects of aldosterone: what we know and do not know thus far. *Heart Fail Rev* 2017; 22: 65–89
46. Connell JMC, Davies E. The new biology of aldosterone. *J Endocrinol* 2005; 186: 1–20
47. Wehling M. Specific, nongenomic actions of steroid hormones. *Annu Rev Physiol* 1997; 59: 365–93
48. Boldyreff B, Wehling M. Aldosterone: refreshing a slow hormone by swift action. *Physiology* 2004; 19: 97–100. [Accessed 30 May 2020]
49. Hucklebridge F, Mellins J, Evans P, Clow A. The awakening cortisol response: no evidence for an influence of body posture. *Life Sci* 2002; 71: 639–46
50. Vernikos-Danellis J, Winget CM. The importance of light, postural and social cues in the regulation of the plasma cortisol rhythm in man 1979. Available from: <http://ntrs.nasa.gov/search.jsp?R=19790063385>.
51. Wendler A, Wehling M. Is GPR30 the membrane aldosterone receptor postulated 20 years ago? *Hypertension* 2011; 57: e16. ; author reply e17
52. Mihailidou AS, Tzakos AG, Ashton AW. Non-genomic effects of aldosterone, vitamins and hormones. *Vitam Horm* 2019; 109: 133–49
53. Funder JW. GPR30, mineralocorticoid receptors, and the rapid vascular effects of aldosterone. *Hypertension* 2011; 57: 370–2
54. Feldman RD, Limbird LE. GPER (GPR30): a nongenomic receptor (GPCR) for steroid hormones with implications for cardiovascular disease and cancer. *Annu Rev Pharmacol Toxicol* 2017; 57: 567–84
55. Weikum ER, Liu X, Ortlund EA. The nuclear receptor superfamily: a structural perspective. *Protein Sci Publ Protein Soc* 2018; 27: 1876
56. Pippal JB, Fuller PJ. Structure–function relationships in the mineralocorticoid receptor. *J Mol Endocrinol* 2008; 41: 405–13
57. Eberwine J. Glucocorticoid and mineralocorticoid receptors as transcription factors. In: Siegel GJ, Agranoff BW, Albers RW, et al., editors. *Basic Neurochemistry: Molecular, Cellular and Medical Aspects*. 6th Edn. Philadelphia: Lippincott-Raven; 1999 Available from: <http://www.ncbi.nlm.nih.gov/books/NBK28050/>
58. Funder JW, Pearce PT, Smith R, Smith AI. Mineralocorticoid action: target tissue specificity is enzyme, not receptor, mediated. *Science* 1988; 242: 583–5
59. Funder JW. Target tissue specificity of mineralocorticoids. *Trends Endocrinol Metab* 1990; 1: 145–8
60. Arriza J, Weinberger C, Cerelli G, et al. Cloning of human mineralocorticoid receptor complementary DNA: structural and functional kinship with the glucocorticoid receptor. *Science* 1987; 237: 268–75
61. Gomez-Sanchez EP. The mammalian mineralocorticoid receptor: tying down a promiscuous receptor: mineralocorticoid receptor selectivity. *Exp Physiol* 2010; 95: 13–8
62. Diederich S, Eigendorff E, Burkhardt P, et al. 11 β -Hydroxysteroid dehydrogenase types 1 and 2: an important pharmacokinetic determinant for the activity of synthetic mineralo- and glucocorticoids. *J Clin Endocrinol Metab* 2002; 87: 5695–701
63. Venkatesh B, Cohen J, Hickman I, et al. Evidence of altered cortisol metabolism in critically ill patients: a prospective study. *Intensive Care Med* 2007; 33: 1746–53
64. Boonen E, Van den Berghe G. Cortisol metabolism in critical illness: implications for clinical care. *Curr Opin Endocrinol Diabetes Obes* 2014; 21: 185–92
65. Nicolaides NC, Chrousos G, Kino T. Glucocorticoid Receptor, Endotext. In: Feingold KR, Anawalt B, Boyce A, editors. *South Dartmouth (MA)*. MDTText.com, Inc.; 2000.

- Available from: <http://www.ncbi.nlm.nih.gov/books/NBK279171/>. [Accessed 21 October 2022]
66. Timmermans S, Souffriau J, Libert C. A general introduction to glucocorticoid biology. *Front Immunol* 2019; **10**: 1545
 67. Funder J. Experimental and clinical pharmacology: corticosteroids - mechanisms of action. *Aust Prescr* 1996; **16**: 41–4
 68. Fuller PJ, Lim-Tio SS, Brennan FE. Specificity in mineralocorticoid versus glucocorticoid action. *Kidney Int* 2000; **57**: 1256–64
 69. Brookes JC, Galigniana MD, Harker AH, Stoneham AM, Vinson GP. System among the corticosteroids: specificity and molecular dynamics. *J R Soc Interface* 2012; **9**: 43–53
 70. Seckl JR. 11β -Hydroxysteroid dehydrogenase in the brain: a novel regulator of glucocorticoid action? *Front Neuroendocrinol* 1997; **18**: 49–99
 71. Ngarmukos C, Grekin RJ. Nontraditional aspects of aldosterone physiology. *Am J Physiol Endocrinol Metab* 2001; **281**: E1122–7
 72. Annane D. Is there a mineralocorticoid deficiency in critically ill patients? How can it be diagnosed? Should it be treated?. In: *Evidence-Based Practice of Critical Care*. Philadelphia: Elsevier; 2010. p. 521–4
 73. Funder JW. Aldosterone action. *Annu Rev Physiol* 1993; **55**: 115–30
 74. Trapp T, Holsboer F. Ligand-induced conformational changes in the mineralocorticoid receptor analyzed by protease mapping. *Biochem Biophys Res Commun* 1995; **215**: 286–91
 75. Kirk DN, Marples BA. The structure and nomenclature of steroids. In: Makin HLJ, Gower DB, Kirk DN, editors. *Steroid Analysis*. Dordrecht: Springer Netherlands; 1995. p. 1–24
 76. Vinson GP. The mislabelling of deoxycorticosterone: making sense of corticosteroid structure and function. *J Endocrinol* 2011; **211**: 3–16
 77. Galigniana MD, Piwien Pilipuk G, Kanelakis KC, Burton G, Lantos CP. Molecular mechanism of activation and nuclear translocation of the mineralocorticoid receptor upon binding of pregnanosteroids. *Mol Cell Endocrinol* 2004; **217**: 167–79
 78. Good DW, George T, Watts BA. Aldosterone inhibits HCO_3^- absorption via a nongenomic pathway in medullary thick ascending limb. *Am J Physiol Renal Physiol* 2002; **283**: F699–706
 79. Qin W, Rudolph AE, Bond BR, et al. Transgenic model of aldosterone-driven cardiac hypertrophy and heart failure. *Circ Res* 2003; **93**: 69–76
 80. Yamaji M, Tsutamoto T, Kawahara C, et al. Serum cortisol as a useful predictor of cardiac events in patients with chronic heart failure: the impact of oxidative stress. *Circ Heart Fail* 2009; **2**: 608–15
 81. Funder JW. Aldosterone and mineralocorticoid receptors—physiology and pathophysiology. *Int J Mol Sci* 2017; **18**: 1032
 82. Nagata K, Obata K, Xu J, et al. Mineralocorticoid receptor antagonism attenuates cardiac hypertrophy and failure in low-aldosterone hypertensive rats. *Hypertension* 2006; **47**: 656–64
 83. Funder JW. Mineralocorticoid receptor activation and oxidative stress. *Hypertension* 2007; **50**: 840–1
 84. Mihailidou AS, Mardini M, Fraser T, Knights D, Funder JW. Agonist/antagonist activity of cortisol in cardiomyocyte mineralocorticoid receptors is determined by redox state. In: *30th Annual Meeting of the International Aldosterone Conference*; 2004. New Orleans
 85. Fuller PJ, Yao Y, Yang J, Young MJ. Mechanisms of ligand specificity of the mineralocorticoid receptor. *J Endocrinol* 2012; **213**: 15–24
 86. Dahanayake JN, Kasireddy C, Karnes JP, et al. Chapter five – progress in our understanding of 19F chemical shifts. In: Webb GA, editor. *Annual Reports on NMR Spectroscopy*, **93**. Academic Press; 2018. p. 281–365. ISSN 0066-4103, ISBN 9780128149133, <https://doi.org/10.1016/bs.arnmr.2017.08.002>
 87. Murphy CD, Palmer-Brown W, Quinn L, Saccomanno M. Microbial metabolism of fluorinated drugs. In: Haufe G, Leroux FR, editors. *Fluorine in Life Sciences: Pharmaceuticals, Medicinal Diagnostics, and Agrochemicals*. Cambridge, MA: Academic Press; 2019. p. 281–99
 88. Khan MO, Lee HJ. Synthesis and pharmacology of anti-inflammatory steroidal antedugs. *Chem Rev* 2008; **108**: 5131–45
 89. Fuller PJ, Yao Y-Z, Jin R, et al. Molecular evolution of the switch for progesterone and spironolactone from mineralocorticoid receptor agonist to antagonist. *Proc Natl Acad Sci* 2019; **116**: 18578–83
 90. De Kloet ER. From receptor balance to rational glucocorticoid therapy. *Endocrinology* 2014; **155**: 2754–69
 91. Fuller PJ, Yang J, Young MJ. Mechanisms of mineralocorticoid receptor signaling. *Vitam Horm* 2019; **109**: 37–68
 92. Heegde F ter, De Rijk RH, Vinkers CH. The brain mineralocorticoid receptor and stress resilience. *Psychoneuroendocrinology* 2015; **52**: 92–110
 93. Cruz-Topete D, Oakley RH, Cidlowski JA. Glucocorticoid signaling and the aging heart. *Front Endocrinol* 2020; **11**: 347
 94. Mihailidou AS, Loan Le TY, Mardini M, Funder JW. Glucocorticoids activate cardiac mineralocorticoid receptors during experimental myocardial infarction. *Hypertension* 2009; **54**: 1306–12
 95. Edwards CRW. Renal 11-beta-hydroxysteroid dehydrogenase: a mechanism ensuring mineralocorticoid specificity. *Horm Res* 1990; **34**: 114–7
 96. Feraille E, Dizin E. Coordinated control of ENaC and Na^+ , K^+ -ATPase in renal collecting duct. *J Am Soc Nephrol* 2016; **27**: 2554–63
 97. Rosa DA de la, Li H, Canessa CM. Effects of aldosterone on biosynthesis, traffic, and functional expression of epithelial sodium channels in A6 cells. *J Gen Physiol* 2002; **119**: 427–42
 98. Chen S, Bhargava A, Mastroberardino L, et al. Epithelial sodium channel regulated by aldosterone-induced protein sgk. *Proc Natl Acad Sci* 1999; **96**: 2514–9
 99. Ozbaki-Yagan N, Liu X, Bodnar AJ, Ho J, Butterworth MB. Aldosterone-induced microRNAs act as feedback regulators of mineralocorticoid receptor signaling in kidney epithelia. *FASEB J* 2020; **34**: 11714–28
 100. Pearce D. SGK1 regulation of epithelial sodium transport. *Cell Physiol Biochem* 2003; **13**: 13–20
 101. Liu Y, Yu Y, Liu J, et al. Neuroimmune regulation in sepsis-associated encephalopathy: the interaction between the brain and peripheral immunity. *Front Neurol* 2022; **13**, 892480
 102. Hill AR, Spencer-Segal JL. Glucocorticoids and the brain after critical illness. *Endocrinology* 2021; **162**: bqaa242

103. Czempik PF, Pluta MP, Krzych LJ. Sepsis-associated brain dysfunction: a review of current literature. *Int J Environ Res Public Health* 2020; **17**: 5852
104. Meyer M, Meijer O, Hunt H, et al. Stress-induced neuroinflammation of the spinal cord is restrained by Cort113176 (Dazucorilant), a specific glucocorticoid receptor modulator. *Mol Neurobiol* 2023. <https://doi.org/10.1007/s12035-023-03554-x>. Advance Access published on August 11
105. De Nicola AF, Meyer M, Guennoun R, et al. Insights into the therapeutic potential of glucocorticoid receptor modulators for neurodegenerative diseases. *Int J Mol Sci* 2020; **21**: 2137
106. Geerling JC, Loewy AD. Aldosterone in the brain. *Am J Physiol Renal Physiol* 2009; **297**: F559–76
107. Ferrari P. The role of 11 β -hydroxysteroid dehydrogenase type 2 in human hypertension. *Biochim Biophys Acta* 2010; **1802**: 1178–87
108. Ueda K, Okamura N, Hirai M, et al. Human P-glycoprotein transports cortisol, aldosterone, and dexamethasone, but not progesterone. *J Biol Chem* 1992; **267**: 24248–52
109. De Kloet ER, Meijer OC, Nicola AF de, Rijk RH de, Joëls M. Importance of the brain corticosteroid receptor balance in metaplasticity, cognitive performance and neuroinflammation. *Front Neuroendocrinol* 2018; **49**: 124–45
110. Gomez Sanchez EP. Central mineralocorticoid receptors and cardiovascular disease. *Neuroendocrinology* 2009; **90**: 245–50
111. De Kloet ER, Meijer OC. MR/GR signaling in the brain during the stress response. In: Harvey B, Jaisser F, editors. *Aldosterone-Mineralocorticoid Receptor - Cell Biology to Translational Medicine*. IntechOpen; 2019
112. De Kloet ER, Joëls M, Holsboer F. Stress and the brain: from adaptation to disease. *Nat Rev Neurosci* 2005; **6**: 463–75
113. Joëls M, Sarabdjitsingh RA, Karst H. Unraveling the time domains of corticosteroid hormone influences on brain activity: rapid, slow, and chronic modes. *Pharmacol Rev* 2012; **64**: 901–38
114. Gomez-Sanchez EP, Venkataraman MT, Thwaites D, Fort C. ICV infusion of corticosterone antagonizes ICV-aldosterone hypertension. *Am J Physiol-Endocrinol Metab* 1990; **258**: E649–53
115. Evans LC, Ivy JR, Wyrwoll C, et al. Conditional deletion of *Hsd11b2* in the brain causes salt appetite and hypertension. *Circulation* 2016; **133**: 1360–70
116. Rossier MF. The cardiac mineralocorticoid receptor (MR): a therapeutic target against ventricular arrhythmias. *Front Endocrinol* 2021; **12**, 694758
117. Rossier MF, Python M, Maturana AD. Contribution of mineralocorticoid and glucocorticoid receptors to the chronotropic and hypertrophic actions of aldosterone in neonatal rat ventricular myocytes. *Endocrinology* 2010; **151**: 2777–87
118. Seamon K, Daly JW, Metzger H, De Souza NJ, Reden J. Structure-activity relationships for activation of adenylate cyclase by the diterpene forskolin and its derivatives. *J Med Chem* 1983; **26**: 436–9
119. Mihailidou AS, Mardini M, Funder JW. Rapid, non-genomic effects of aldosterone in the heart mediated by e protein kinase C. *Endocrinology* 2004; **145**: 773–80
120. Sato A, Liu J-P, Funder JW. Aldosterone rapidly represses protein kinase C activity in neonatal rat cardiomyocytes in vitro. *Endocrinology* 1997; **138**: 3410–6
121. Fernández-Ruiz I. Balancing stress signalling in the heart. *Nat Rev Cardiol* 2019; **16**: 384–5
122. Oakley RH, Cruz-Topete D, He B, et al. Cardiomyocyte glucocorticoid and mineralocorticoid receptors directly and antagonistically regulate heart disease in mice. *Sci Signal* 2019; **12**, eaau9685
123. Bohus B, De Kloet ER. Adrenal steroids and extinction behavior: antagonism by progesterone, deoxycorticosterone and dexamethasone of a specific effect of corticosterone. *Life Sci* 1981; **28**: 433–40
124. De Kloet ER, Versteeg DHG, Kovacs GL. Aldosterone blocks the response to corticosterone in the raphe-hippocampal serotonin system. *Brain Res* 1983; **264**: 323–7
125. Veldhuis HD, De Kloet ER. Antagonistic effects of aldosterone on corticosterone-mediated changes in exploratory behavior of adrenalectomized rats. *Horm Behav* 1983; **17**: 225–32
126. Holtorf K. The bioidentical hormone debate: are bio-identical hormones (estradiol, estriol, and progesterone) safer or more efficacious than commonly used synthetic versions in hormone replacement therapy? *Postgrad Med* 2009; **121**: 73–85
127. Santoro N, Braunstein GD, Butts CL, Martin KA, McDermott M, Pinkerton JV. Compounded bioidentical hormones in endocrinology practice: an Endocrine Society scientific statement. *J Clin Endocrinol Metab* 2016; **101**: 1318–43
128. Thorn GW, Sheppard RH, Morse WI, Reddy WJ, Beigelman PM, Renold AE. Comparative action of aldosterone and 9-alpha-fluorohydrocortisone in man. *Ann N Y Acad Sci* 1955; **61**: 609–19
129. Files JA, Ko MG, Pruthi S. Bioidentical hormone therapy. *Mayo Clin Proc* 2011; **86**: 673–80
130. Whelan AM, Jurgens TM, Trinacty M. Defining bio-identical hormones for menopause-related symptoms. *Pharm Pract Internet* 2011; **9**: 16–22
131. De Kloet ER. Why dexamethasone poorly penetrates in brain. *Stress* 1997; **2**: 13–9
132. Meijer OC, Lange ECM de, Breimer DD, Boer AG de, Workel JO, De Kloet ER. Penetration of dexamethasone into brain glucocorticoid targets is enhanced in *mdr1A* P-glycoprotein knockout mice. *Endocrinology* 1998; **139**: 1789–93
133. Bamberg K, William-Olsson L, Johansson U, Jansson-Löfmark R, Hartleib-Geschwindner J. The selective mineralocorticoid receptor modulator AZD9977 reveals differences in mineralocorticoid effects of aldosterone and fludrocortisone. *J Renin Angiotensin Aldosterone Syst* 2019; **20**, 1470320319827449
134. Tomlinson JW, Walker EA, Bujalska IJ, et al. 11 β -Hydroxysteroid dehydrogenase type 1: a tissue-specific regulator of glucocorticoid response. *Endocr Rev* 2004; **25**: 831–66
135. PubChem. Fludrocortisone. Available from: <https://pubchem.ncbi.nlm.nih.gov/compound/31378> [Accessed 21 July 2020].
136. Liu D, Ahmet A, Ward L, et al. A practical guide to the monitoring and management of the complications of systemic corticosteroid therapy. *Allergy Asthma Clin Immunol* 2013; **9**: 30
137. Robertson D, Marie Robertson R. Fludrocortisone. In: Robertson D, Biaggioni I, Burnstock G, Low PA, Paton JFR, editors. *Primer on the Autonomic Nervous System*. 3rd Edn. San Diego: Academic Press; 2012. p. 635–7

138. Tredwell M, Gouverneur V. Fluorine in medicinal chemistry: importance of chirality. In: L Carreira EM, Yamamoto H, editors. *Comprehensive Chirality*. Amsterdam: Elsevier; 2012. p. 70–85
139. Mitsky VP, Workman RJ, Nicholson WE, Vernikos J, Robertson RM, Robertson D. A sensitive radioimmunoassay for fludrocortisone in human plasma. *Steroids* 1994; **59**: 555–8
140. Florey K. Fludrocortisone acetate. In: Florey K, editor. *Analytical Profiles of Drug Substances*. Academic Press; 1974. p. 281–306. [Accessed 14 June 2020]
141. Altman LC, Hill JS, Hairfield WM, Mullarkey MF. Effects of corticosteroids on eosinophil chemotaxis and adherence. *J Clin Invest* 1981; **67**: 28–36
142. Annane D, Bellissant E, Bollaert P-E, et al. Corticosteroids in the treatment of severe sepsis and septic shock in adults: a systematic review. *JAMA* 2009; **301**: 2362
143. Furman BL. Fludrocortisone. *xPharm: the comprehensive pharmacology reference*. Elsevier; 2007. p. 1–4. <https://doi.org/10.1016/B978-008055232-3.61757-X>
144. White PC. Disorders of aldosterone biosynthesis and action. *N Engl J Med* 1994; **331**: 250–8
145. Gallo-Payet N, Martinez A, Lacroix A. Editorial: ACTH action in the adrenal cortex: from molecular biology to pathophysiology. *Front Endocrinol* 2017; **8**: 101
146. Nishimoto K, Koga M, Seki T, et al. Immunohistochemistry of aldosterone synthase leads the way to the pathogenesis of primary aldosteronism. *Mol Cell Endocrinol* 2017; **441**: 124–33
147. Vinson GP. Functional zonation of the adult mammalian adrenal cortex. *Front Neurosci* 2016; **10**: 238
148. Nishimoto K, Tomlins SA, Kuick R, et al. Aldosterone-stimulating somatic gene mutations are common in normal adrenal glands. *Proc Natl Acad Sci* 2015; **112**: E4591–9
149. Aiba M, Fujibayashi M. Alteration of subcapsular adrenocortical zonation in humans with aging: the progenitor zone predominates over the previously well-developed zona glomerulosa after 40 years of age. *J Histochem Cytochem* 2011; **59**: 557–64
150. Tezuka Y, Atsumi N, Blinder A, et al. Human adrenal cortical zone changes with aging. *J Endocr Soc* 2021; **5**: A69–A69
151. Polito A, Hamitouche N, Ribot M, et al. Pharmacokinetics of oral fludrocortisone in septic shock. *Br J Clin Pharmacol* 2016; **82**: 1509–16
152. Erlandsson F, Albayaty M, Chialda L, et al. Clinical safety, tolerability, pharmacokinetics and effects on urinary electrolyte excretion of AZD9977, a novel, selective mineralocorticoid receptor modulator. *Br J Clin Pharmacol* 2018; **84**: 1486–93
153. Hammond NE, Kumar A, Tirupakuzhi Vijayaraghavan BK, et al. Clinician preferences for prescription of corticosteroids in patients with septic shock: an international survey. *Crit Care Resusc* 2021; **23**: 234–8

Handling Editor: Jonathan Hardman