

Food Insecurity and Intimate Partner Violence Among HIV-Positive Individuals in Rural Kenya



Abigail M. Hatcher, PhD, MPhil,^{1,2} Sheri D. Weiser, MD, MPH,¹ Craig R. Cohen, MD, MPH,³ Jill Hagey, MD, MPH,⁴ Elly Weke, MA,⁵ Rachel Burger, MPH,³ Pauline Wekesa, BSc,⁵ Lila Sheira, MPH,¹ Edward A. Frongillo, PhD,⁶ Elizabeth A. Bukusi, MBChB, MMed, MPH, PhD^{3,5,7,8}

Introduction: Intimate partner violence and food insecurity are both structural drivers of HIV acquisition, care, and treatment, but little is known about how the 2 conditions intersect in the lives of those already living with HIV.

Methods: This study examined cross-sectional baseline data (collected in January 2016–December 2017) from an ongoing trial in southwestern Kenya. Trained interviewers asked enrolled participants living with HIV aged 18–60 years about household food insecurity (using the Household Food Insecurity Access Scale), intimate partner violence (using an adapted WHO multicountry study instrument), and sociodemographics. Negative binomial regression was used to examine the association between food insecurity and partner violence victimization (among women) or perpetration (among men). Secondary data were analyzed in August 2019–March 2020.

Results: Of 720 participants, more than half of women reported experiencing intimate partner violence (57.6%) and most men reported perpetrating it (58.4%). Participants reporting any partner violence had higher Household Food Insecurity Access Scale scores (21.8) compared with those reporting no violence (21.3, $p=0.02$). Each categorical change in food insecurity (mild, moderate, severe) was associated with a 41% increased risk of an additional partner violence episode. In models controlling for relationship status, wealth, season of interview (lean versus not lean), and baseline physical health, each 1-point increase in food insecurity was associated with a 6% higher risk of violence victimization among women and 4% greater risk of men perpetrating partner violence.

Conclusions: This study highlights the interconnected nature of intimate partner violence and food insecurity among women and men living with HIV. This relationship suggests that enhancing food security may be a useful intervention strategy to prevent intimate partner violence and improve HIV-related health outcomes.

Am J Prev Med 2021;60(4):563–568. © 2020 American Journal of Preventive Medicine. Published by Elsevier Inc. All rights reserved.

From the ¹Division of HIV, Infectious Diseases, and Global Medicine, Department of Medicine, University of California San Francisco, San Francisco, California; ²Faculty of Health Science, School of Public Health, University of the Witwatersrand, Johannesburg, South Africa; ³Department of Obstetrics, Gynecology and Reproductive Sciences, University of California San Francisco, San Francisco, California; ⁴Department of Medicine, Duke University, Durham, North Carolina; ⁵Center for Microbiology Research, Kenya Medical Research Institute, Nairobi, Kenya; ⁶Department of Health Promotion, Education, and Behavior, University of South Carolina, Columbia, South Carolina; ⁷Department of Global Health, University

of Washington, Seattle, Washington; and ⁸Department of Obstetrics and Gynecology, University of Washington, Seattle, Washington

Address correspondence to: Abigail M. Hatcher, PhD, Division of HIV, Infectious Diseases, and Global Medicine, Department of Medicine, University of California, San Francisco, 1001 Potrero Avenue, Building 100, Suite 335 West, San Francisco CA 94110.

E-mail: hatchera@globalhealth.ucsf.edu.

0749-3797/\$36.00

<https://doi.org/10.1016/j.amepre.2020.06.025>

INTRODUCTION

Syndemics are concurrent and mutually reinforcing social and health problems.^{1,2} The syndemic conditions of food insecurity and intimate partner violence (IPV) are associated with HIV acquisition^{3–5} and poor health among HIV-positive populations.^{6,7} In sub-Saharan Africa, all the 3 conditions are higher than elsewhere globally.^{8,9}

Food insecurity is associated with doubled odds of IPV victimization among women in the general population,^{10–14} although quantitative work among HIV-positive women is limited. Qualitative research suggests that IPV leads to food insecurity if it prompts HIV-positive women to leave violent partners or shoulder household financial needs alone.¹⁵

Among men in the general population, food insecurity is associated with higher odds of IPV perpetration.^{16,17} Little is known about this association among HIV-positive men, although IPV perpetration and food insecurity are both linked to men's HIV risk.^{5,18}

A theoretic framework underpins the relationship between food insecurity and IPV.¹⁶ Men may re-exert control through violence if unable to meet provider roles (gender resource theory),^{19,20} or deprivation could heighten acute stressors in the household (family stress theory)²¹ and deplete men's ability to attain self-control (physiology of willpower).²² A bidirectional relationship could exist if IPV exposure lowers economic earnings,²³ increases household financial burden owing to injury or missing work,²⁴ or makes relationships unstable—lowering the ability of single-earner households to obtain sufficient food.

This study examines the association between food insecurity severity and IPV among HIV-positive individuals in rural Kenya using cross-sectional data from an ongoing trial.

METHODS

This was a secondary analysis of baseline data from the Shamba Maisha cluster RCT. Shamba Maisha is a multisectoral agricultural intervention including (1) low-cost irrigation water pump, (2) group agricultural training, and (3) loan (\$150) for seeds per fertilizers.^{25,26}

A total of 8 matched pairs of health facilities in southwestern Kenya were randomized to intervention ($n=360$) or control arms ($n=362$). Inclusion criteria were being on antiretroviral treatment, being aged 18–60 years, and having access to surface water. Included participants had moderate-to-severe food insecurity, assessed using the Household Food Insecurity Access Scale (HFIAS), and were willing to save a down payment applied toward the agricultural loan. Trained researchers led structured interviews (January 2016–December 2017) and abstracted clinical data from medical records.

This study assessed IPV through 4 self-reported items on lifetime physical or sexual violence (adapted from the WHO instrument²⁷). Investigators examined the IPV experienced by female participants and perpetrated by male participants. The frequency of each violent act was assessed on a Likert-type scale (0, never; 1, once; 2, two or 3 times; 3, ≥ 4 times). The index was assessed as a dichotomous outcome (ever versus never) and continuous measure of IPV intensity (scores ranged from 0 to 12).

An HFIAS score assessed food insecurity as a continuous variable. Scores ranged from 11 to 36, and the scale had acceptable internal consistency (Cronbach's $\alpha=0.81$). Scores were categorized as mild (<14), moderate (14–21), or severe (≥ 22) food insecurity.²⁸

Sociodemographic covariates included season of interview (lean versus not lean), age, marital status (married versus widowed/separated/single), number of children, education level, baseline CD4 count, and a wealth index.²⁹

These secondary analyses were not preregistered (conducted in August 2019–May 2020) and were led using Stata, version 16. Bivariate analyses for normally distributed and categorical variables (t -test and chi-square test, respectively) and non-normally distributed variables (Wilcoxon) were conducted by IPV status.

The authors fitted negative binomial regressions by sex to account for participants with no experience of IPV (resulting in a large number of 0s). Negative binomial models assume that each individual has their own Poisson distribution, thus allowing individuals to vary in the propensity of reporting IPV. Authors adjusted for covariates of theoretic significance: season of interview, age, wealth index, marital status, education, and health status (CD4 count). Past research suggests that the similar sociodemographic covariates predict IPV victimization and IPV perpetration,^{30,31} so similar covariates were used in models for women and men. Analyses accounted for clustering by clinic using a sandwich estimator.

Written informed consent was obtained from study participants. Ethical approval for the study was granted by Kenya Medical Research Institute and the University of California San Francisco, California. Because the data are part of an ongoing trial, they are not yet publicly available.

RESULTS

Of the 720 baseline participants, 396 (55.0%) were female. The mean HFIAS score was 21 for men and 22 for women ($p=0.008$), with 569 (78.9%) participants reporting severe food insecurity (Table 1).

A majority of women (57.6%) reported IPV victimization, and a majority of men (58.4%) reported perpetrating IPV. Participants reporting any IPV had higher HFIAS scores (21.8) than those reporting no IPV (21.3, $p=0.02$), although this bivariate analysis was not significant by sex (men: $p=0.07$, women: $p=0.05$; data not shown). A scatterplot with a fitted line (Figure 1) illustrates how HFIAS was associated with IPV perpetration ($p=0.029$) and victimization ($p=0.002$).

Each 1-point increase on HFIAS was associated with a higher adjusted risk of IPV victimization among women (adjusted incident rate=1.06, 95% CI=1.01, 1.10)

Table 1. Descriptive Data and Bivariate Associations of Sociodemographics and IPV by Sex (N=720)

Sociodemographic	Total n (%)	IPV experience among women			IPV perpetration by men		
		None	Ever IPV	p-value	None	Ever IPV	p-value
Age (years), median (IQR)	40 (34–47)	43.3	42.3	0.53	38.5	37.9	0.38
Children (number), Median (IQR)	4 (2–5)	3.5	3.7	0.29	4.1	4.3	0.80
Primary school education (versus no primary school education), %	415 (57.6)	52.1	47.6	0.38	64.9	70.2	0.32
Widowed, separated, or single (versus married), %	198 (27.5)	46.1	37.9	0.10	14.2	8.0	0.07
Polygynous marriage, %	173 (29.5)	21.1	35.9	0.01	30.6	27.3	0.53
Lowest wealth quintile, %	142 (19.5)	18.9	24.1	0.54	23.5	14.0	0.13
Baseline CD4 (count), Median (IQR)	560 (397–745)	636	649	0.68	506	512	0.86
Interview season (lean versus not lean), %	165 (22.8)	22.8	22.5	0.95	17.9	27.1	0.05

IPV, intimate partner violence.

(Table 2). Among men, HFIAS was associated with a greater risk of IPV perpetration (adjusted incident rate ratio=1.04, 95% CI=1.01, 1.07). Each change in food insecurity category (mild, moderate, severe) was associated with 41% increased risk of an additional IPV episode (95% CI=1.10, 1.82, $p=0.007$) (Appendix Table 1, available online). Categorical interpretation did not reach statistical significance among women ($p=0.068$) or men ($p=0.097$).

DISCUSSION

In a sample of men and women living with HIV, food insecurity was independently associated with IPV. In

this predominately food-insecure sample, increased severity of food insecurity was linearly associated with greater IPV intensity. Each change in food insecurity category (mild, moderate, severe) was associated with a 41% increased risk of an additional IPV episode.

These findings are consistent with extant studies among the general population, where food insecurity is associated with between twofold and sixfold increased odds of any IPV perpetration^{16,17,32} or victimization.^{33–36} This is the first analysis, to the authors’ knowledge, to examine food insecurity and IPV among an HIV-positive cohort in sub-Saharan Africa. Further research may suggest that these conditions in HIV-positive samples are unique from or similar to the general population.




Figure 1. Scatterplot and fitted line of food insecurity and IPV intensity by sex. HFIAS, Household Food Insecurity Access Scale; IPV, intimate partner violence.

Table 2. Unadjusted and Adjusted Negative Binomial Regressions of IPV by Sex (N=720)

Variable	Bivariate associations		Model 1: IPV experience among women (n=396)		Model 2: IPV perpetration by men (n=324)	
	IRR (95% CI)	p-value	aIRR (95% CI)	p-value	aIRR (95% CI)	p-value
Food insecurity	1.06 (1.02, 1.09)	<0.001	1.06 (1.01, 1.10)	0.012	1.04 (1.01, 1.07)	0.043
Sociodemographics						
Age (years)	0.99 (0.98, 1.01)	0.521	0.99 (0.97, 1.01)	0.350	0.99 (0.98, 1.02)	0.616
Children	1.01 (0.67, 1.04)	0.615	0.99 (0.93, 1.05)	0.340	1.01 (0.95, 1.06)	0.819
Primary school education	0.87 (0.75, 1.02)	0.095	0.78 (0.61, 1.04)	0.097	1.17 (0.98, 1.63)	0.171
Wealth index quintiles	0.95 (0.86, 1.05)	0.306	0.91 (0.83, 1.04)	0.537	1.02 (0.92, 1.07)	0.632
Widowed, separated, or single	0.99 (0.80, 1.19)	0.834	0.98 (0.68, 1.45)	0.891	0.80 (0.43, 1.47)	0.475
Polygynous household	1.18 (0.97, 1.44)	0.079	1.41 (1.15, 1.72)	0.001	0.99 (0.60, 1.39)	0.946
Baseline CD4	1.00 (1.00, 1.00)	0.188	1.00 (1.00, 1.00)	0.589	1.00 (1.00, 1.00)	0.594
Interview season (lean versus not lean)	1.24 (0.90, 1.72)	0.184	0.08 (0.59, 1.24)	0.412	1.34 (0.96, 1.86)	0.085

Note: Boldface indicates statistical significance ($p < 0.05$). aIRR, adjusted IRR; IPV, intimate partner violence; IRR, incident risk ratio.

More than half of women (58%) reported experiencing IPV, which is higher than among HIV-positive women in Uganda (44%).³⁷ Among men, 58% reported IPV perpetration, which is higher than among HIV-positive men in Vietnam (38%).³⁸ High IPV exposure has clinical implications because it leads to incident HIV infection,³ lowers women's antiretroviral treatment adherence and viral control,⁶ and has health effects for men themselves.³⁹

These findings suggest that even when programs are unable to fully ameliorate food insecurity, slight improvements around food may be accompanied by safer relationships. Adding food security programming to clinical care could reach patients positive with HIV who experience or perpetrate IPV. By focusing on food security as an upstream determinant of health, programs may ultimately reduce IPV and improve HIV-related health.

Limitations

There are several limitations of this analysis. The small sample limits analysis of variables as categorical. Cross-sectional data limit causal conclusions, in particular, given the timing of past-month food insecurity and lifetime IPV. The study controlled for potential seasonality of food security by including season of interview as a covariate, but this does not fully ameliorate the temporality of food reporting. Self-reported measures for food insecurity or IPV may be limited by social desirability bias. Asking about women's experience and men's perpetration of IPV precludes exploring bidirectional relationship violence. Although the measure of IPV asks behaviorally specific items, there are noted limitations to IPV measures,⁴⁰ and the 4-item scale was not validated.

CONCLUSIONS

This report highlights the interconnected nature of IPV and food insecurity in an HIV-positive sample in Kenya. High prevalence of IPV in this region, alongside the declines in adherence and HIV-related health associated with IPV^{6,41} and food insecurity,⁷ makes this a crucial intersection for future work. Enhancing food security may be a useful intervention strategy to address nutrition, IPV, and HIV going forward.

ACKNOWLEDGMENTS

This study was funded by the National Institute of Mental Health (R01 MH107330 05 and K01 MH12118501).

No financial disclosures were reported by the authors of this paper.

SUPPLEMENTAL MATERIAL

Supplemental materials associated with this article can be found in the online version at <https://doi.org/10.1016/j.amepre.2020.06.025>.

REFERENCES

- Singer M, Clair S. Syndemics and public health: reconceptualizing disease in bio-social context. *Med Anthropol Q*. 2003;17(4):423–441. <https://doi.org/10.1525/maq.2003.17.4.423>.
- Singer M, Bulled N, Ostrach B, Mendenhall E. Syndemics and the bio-social conception of health. *Lancet*. 2017;389(10072):941–950. [https://doi.org/10.1016/S0140-6736\(17\)30003-X](https://doi.org/10.1016/S0140-6736(17)30003-X).
- Li Y, Marshall CM, Rees HC, Nunez A, Ezeanolue EE, Ehiri JE. Intimate partner violence and HIV infection among women: a systematic review and meta-analysis. *J Int AIDS Soc*. 2014;17(1):18845. <https://doi.org/10.7448/IAS.17.1.18845>.

4. Weiser SD, Palar K, Hatcher AM, Young SL, Frongillo EA. Food insecurity and health: a conceptual framework. In: Ivers L, editor. *Food Insecurity and Public Health*. Boca Raton, FL: CRC Press, 2015:23–50.
5. Palar K, Laraia B, Tsai AC, Johnson MO, Weiser SD. Food insecurity is associated with HIV, sexually transmitted infections and drug use among men in the United States. *AIDS*. 2016;30(9):1457–1465. <https://doi.org/10.1097/QAD.0000000000001095>.
6. Hatcher AM, Smout EM, Turan JM, Christofides N, Stöckl H. Intimate partner violence and engagement in HIV care and treatment among women: a systematic review and meta-analysis. *AIDS*. 2015;29(16):2183–2194. <https://doi.org/10.1097/QAD.0000000000000842>.
7. Weiser SD, Young SL, Cohen CR, et al. Conceptual framework for understanding the bidirectional links between food insecurity and HIV/AIDS. *Am J Clin Nutr*. 2011;94(6):1729S–1739S. <https://doi.org/10.3945/ajcn.111.012070>.
8. Devries KM, Mak JY, García-Moreno C, et al. Global health. The global prevalence of intimate partner violence against women. *Science*. 2013;340(6140):1527–1528. <https://doi.org/10.1126/science.1240937>.
9. FAO IFAD, UNICEF WFP, WHO. The State of Food Security and Nutrition in the World 2019. Safeguarding against economic slowdowns and downturns. Rome, Italy: FAO. <http://www.fao.org/3/ca5162en/ca5162en.pdf>. Published 2019. Accessed July 20, 2020.
10. Breiding MJ, Basile KC, Klevens J, Smith SG. Economic insecurity and intimate partner and sexual violence victimization. *Am J Prev Med*. 2017;53(4):457–464. <https://doi.org/10.1016/j.amepre.2017.03.021>.
11. Melchior H, Hergert A, Hofreuter-Gätgens K, et al. Predictors of treatment duration for inpatients with mental disorders—a systematic literature review [in German]. *Z Psychosom Med Psychother*. 2010;56(4):399–418. <https://doi.org/10.13109/zptm.2010.56.4.399>.
12. Ricks JL, Cochran SD, Arah OA, Williams JK, Seeman TE. Food insecurity and intimate partner violence against women: results from the California Women's Health Survey. *Public Health Nutr*. 2016;19(5):914–923. <https://doi.org/10.1017/S1368980015001986>.
13. Diamond-Smith N, Conroy AA, Tsai AC, Nekkanti M, Weiser SD. Food insecurity and intimate partner violence among married women in Nepal. *J Glob Health*. 2019;9(1):010412. <https://doi.org/10.7189/jogh.09.010412>.
14. Conroy AA, Cohen MH, Frongillo EA, et al. Food insecurity and violence in a prospective cohort of women at risk for or living with HIV in the U.S. *PLoS One*. 2019;14(3):e0213365. <https://doi.org/10.1371/journal.pone.0213365>.
15. Derose KP, Payán DD, Fulcar MA, et al. Factors contributing to food insecurity among women living with HIV in the Dominican Republic: a qualitative study. *PLoS One*. 2017;12(7):e0181568. <https://doi.org/10.1371/journal.pone.0181568>.
16. Hatcher AM, Stöckl H, McBride RS, Khumalo M, Christofides N. Pathways from food insecurity to intimate partner violence perpetration among peri-urban men in South Africa. *Am J Prev Med*. 2019;56(5):765–772. <https://doi.org/10.1016/j.amepre.2018.12.013>.
17. Chirwa ED, Sikweyiya Y, Addo-Lartey AA, et al. Prevalence and risk factors of physical or sexual intimate violence perpetration amongst men in four districts in the central region of Ghana: baseline findings from a cluster randomised controlled trial. *PLoS One*. 2018;13(3):e0191663. <https://doi.org/10.1371/journal.pone.0191663>.
18. Hatcher AM, Gibbs A, McBride RS, Rebombo D, Khumalo M, Christofides NJ. Gendered syndemic of intimate partner violence, alcohol misuse, and HIV risk among peri-urban, heterosexual men in South Africa. *Soc Sci Med*. 2019. In press Online October 22 <https://doi.org/10.1016/j.socscimed.2019.112637>.
19. Connell RW, Messerschmidt JW. Hegemonic masculinity: rethinking the concept. *Gender Soc*. 2005;19(6):829–859. <https://doi.org/10.1177/0891243205278639>.
20. Stets JE. Job autonomy and control over one's spouse: a compensatory process. *J Health Soc Behav*. 1995;36(3):244–258. <https://doi.org/10.2307/2137341>.
21. Fox GL, Benson ML, DeMaris AA, Wyk J. Economic distress and intimate violence: testing family stress and resources theories. *J Marriage Fam*. 2002;64(3):793–807. <https://doi.org/10.1111/j.1741-3737.2002.00793.x>.
22. Baumeister RF, Heatherton TF, Tice DM. *Losing Control: How and Why People Fail at Self-Regulation*. Cambridge, MA: Academic Press, 1994.
23. Lindhorst T, Oxford M, Gillmore MR. Longitudinal effects of domestic violence on employment and welfare outcomes. *J Interpers Violence*. 2007;22(7):812–828. <https://doi.org/10.1177/0886260507301477>.
24. Peterson C, Kearns MC, McIntosh WL, et al. Lifetime economic burden of intimate partner violence among U.S. adults. *Am J Prev Med*. 2018;55(4):433–444. <https://doi.org/10.1016/j.amepre.2018.04.049>.
25. Weiser SD, Bukusi EA, Steinfeld RL, et al. Shamba maisha: randomized controlled trial of an agricultural and finance intervention to improve HIV health outcomes. *AIDS*. 2015;29(14):1889–1894. <https://doi.org/10.1097/QAD.0000000000000781>.
26. Cohen CR, Steinfeld RL, Weke E, et al. Shamba Maisha: pilot agricultural intervention for food security and HIV health outcomes in Kenya: design, methods, baseline results and process evaluation of a cluster-randomized controlled trial. *Springerplus*. 2015;4:122. <https://doi.org/10.1186/s40064-015-0886-x>.
27. Garcia-Moreno C, Jansen HA, Ellsberg M, Heise L, Watts CH. Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *Lancet*. 2006;368(9543):1260–1269. [https://doi.org/10.1016/S0140-6736\(06\)69523-8](https://doi.org/10.1016/S0140-6736(06)69523-8).
28. Coates J, Swindale A, Bilinsky P. *Household Food Insecurity Access Scale (HFIAS) for Measurement of Food Access: Indicator Guide*. Washington, DC: Food and Nutrition Technical Assistance Project, Academy for Educational Development, 2006.
29. Rutstein SO, Johnson K. *DHS Comparative reports no. 6. The DHS wealth index*. Calverton, MD: ORC Macro; Published August 2004. <https://dhsprogram.com/pubs/pdf/CR6/CR6.pdf>. Accessed February 3, 2020.
30. Fulu E, Jewkes R, Roselli T, Garcia-Moreno C. UN Multi-country Cross-sectional Study on Men and Violence research team. Prevalence of and factors associated with male perpetration of intimate partner violence: findings from the UN Multi-Country Cross-sectional Study on Men and Violence in Asia and the Pacific. *Lancet Glob Health*. 2013;1(4):e187–e207. [https://doi.org/10.1016/S2214-109X\(13\)70074-3](https://doi.org/10.1016/S2214-109X(13)70074-3).
31. Costa BM, Kaestle CE, Walker A, et al. Longitudinal predictors of domestic violence perpetration and victimization: a systematic review. *Aggress Violent Behav*. 2015;24:261–272. <https://doi.org/10.1016/j.avb.2015.06.001>.
32. Gilbert L, Michalopoulos L, Ma X, et al. How do risk environment factors influence perpetration of partner violence among male migrant and non-migrant market workers in Central Asia? [published correction appears in *J Urban Health*. 2019;96(1):96]. *J Urban Health*. 2019;96(1):83–95. <https://doi.org/10.1007/s11524-018-0312-0>.
33. Field S, Onah M, van Heyningen T, Honikman S. Domestic and intimate partner violence among pregnant women in a low resource setting in South Africa: a facility-based, mixed methods study. *BMC Womens Health*. 2018;18(1):119. <https://doi.org/10.1186/s12905-018-0612-2>.
34. Gibbs A, Jewkes R, Willan S, Washington L. Associations between poverty, mental health and substance use, gender power, and intimate partner violence amongst young (18-30) women and men in urban informal settlements in South Africa: a cross-sectional study and structural equation model. *PLoS One*. 2018;13(10):e0204956. <https://doi.org/10.1371/journal.pone.0204956>.
35. Swahn MH, Dill LJ, Palmier JB, Kasirye R. Girls and young women living in the slums of Kampala: prevalence and correlates of physical and sexual violence victimization. *SAGE Open*. 2015;5(2):1–8. <https://doi.org/10.1177/2158244015580853>.
36. Willie TC, Kershaw TS, Callands TA. Examining relationships of intimate partner violence and food insecurity with HIV-related risk

- factors among young pregnant Liberian women. *AIDS Care*. 2018;30(9):1156–1160. <https://doi.org/10.1080/09540121.2018.1466983>.
37. Kabwama SN, Bukenya J, Matovu JKB, et al. Intimate partner violence among HIV positive women in care - results from a national survey, Uganda 2016. *BMC Womens Health*. 2019;19(1):130. <https://doi.org/10.1186/s12905-019-0831-1>.
 38. Hershov RB, Ha TV, Sripaipan T, et al. Perpetration of intimate partner violence among men living with HIV in Northern Vietnam. *AIDS Behav*. 2020;24:2555–2571. <https://doi.org/10.1007/s10461-020-02813-5>.
 39. Courtenay WH. Constructions of masculinity and their influence on men's well-being: a theory of gender and health. *Soc Sci Med*. 2000;50(10):1385–1401. [https://doi.org/10.1016/s0277-9536\(99\)00390-1](https://doi.org/10.1016/s0277-9536(99)00390-1).
 40. Follingstad DR, Bush HM. Measurement of intimate partner violence: a model for developing the gold standard. *Psychol Violence*. 2014;4(4):369–383. <https://doi.org/10.1037/a0037515>.
 41. Leddy AM, Weiss E, Yam E, Pulerwitz J. Gender-based violence and engagement in biomedical HIV prevention, care and treatment: a scoping review. *BMC Public Health*. 2019;19(1):897. <https://doi.org/10.1186/s12889-019-7192-4>.