





'Men are not playing their roles', maternal and child nutrition in Nanoro, Burkina Faso

Adélaïde Compaoré^{1,*}, Kadija Ouedraogo¹, Palwende R Boua¹, Daniella Watson² , Sarah H Kehoe³, Marie-Louise Newell^{4,5}, Halidou Tinto¹, Mary Barker^{3,4} and Hermann Sorgho¹  for the INPreP group

¹Clinical Research Unit of Nanoro, Institut de Recherche en Sciences de la Santé, Nanoro, Burkina Faso: ²Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK: ³MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK: ⁴School of Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK: ⁵School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Submitted 2 January 2020: Final revision received 27 July 2020: Accepted 19 August 2020: First published online 1 October 2020

Abstract

Objective: To collect context-specific insights into maternal and child health and nutrition issues, and to explore potential solutions in Nanoro, Burkina Faso.

Design: Eleven focus groups with men and women from eleven communities, facilitated by local researchers.

Setting: The study took place in the Nanoro Health district, in the West-Central part of Burkina Faso.

Participants: Eighty-six men (18–55 years) and women by age group: 18–25; 26–34 and 35–55 years, participated in the group discussions.

Results: Participants described barriers to optimal nutrition of mothers and children related to a range of community factors, with gender inequality as central. Major themes in the discussions are related to poverty and challenges generated by socially and culturally determined gender roles. Sub-themes are women lacking access to food whilst pregnant and having limited access to health care and opportunities to generate income. Although communities believe that food donations should be implemented to overcome this, they also pointed out the need for enhancing their own food production, requiring improved agricultural technologies. Given the important role that women could play in reducing malnutrition, these communities felt they needed to be empowered to do so and supported by men. They also felt that this had to be carried out in the context of an enhanced health care system.

Conclusions: Findings reported here highlight the importance of nutrition-sensitive interventions and women's empowerment in improving maternal and child nutrition. There is a need to integrate a sustainable multi-sectorial approach which goes beyond food support.

Keywords

Maternal nutrition
Child nutrition
Qualitative research
Community perceptions
Rural Burkina Faso

Burkina Faso is a landlocked and low-income country situated in West Africa. The country is among the poorest in the world with 44 % of its population below the international poverty line of US\$1.90 per day⁽¹⁾. The mortality rate of children under-five in Burkina Faso is also one of the highest in the world (85 %), 35 % of which is attributed to malnutrition⁽²⁾. The country's 2016 National Nutritional Survey revealed that 27.3 % of children under-five were stunted, 7.6 % were severely wasted and 19.2 % underweight⁽³⁾. Concurrently, overweight and obesity among women increased from 7.7 %

in 1980 to 28.8 % in 2016⁽⁴⁾. The country has implemented some important changes in nutrition policy during recent years, which has resulted in progress which includes reduced rates of stunting and wasting in the under-fives, but which do not yet put the country on track to meet the World Health Assembly targets in 2025⁽⁵⁾. Recent security issues in the northern and eastern parts of the country are jeopardising the sustainability of this progress.

Burkina Faso National Nutrition Policy suggests that child malnutrition and micronutrient deficiencies are due

*Corresponding author: Email adelhaid.compaore@gmail.com

© The Author(s), 2020. Published by Cambridge University Press on behalf of The Nutrition Society



primarily to poor breast-feeding and complementary feeding practices, accompanied by high prevalence of infectious diseases and limited health services⁽⁶⁾. The 2019 Nutrition Policy in Burkina Faso recommended that these gaps be addressed through prioritising investment in multi-sectoral programmes, which mainly benefit women and children⁽⁵⁾. Previous research in Burkina Faso and other low-and-middle income countries has shown that improvements in maternal and child nutrition are held back by deep-seated, cultural and socially defined gender inequities, which contribute to women's lack of empowerment and their limited capacity to influence household decision-making^(7–9). Women are also expected to carry heavy workloads, are subject to restrictive gender norms and to a lack of access to health-promoting resources⁽³⁾. Gender is becoming increasingly recognised as an underlying determinant of health and nutrition and an increasing number of interventions focus on women's empowerment with positive effect^(9–14). A multi-sectoral intervention, which provided dedicated land to women in Burkina Faso and trained them in agricultural production and income generation, led to significant reductions in child wasting⁽¹⁴⁾ and to improvements in mothers' nutrition and empowerment outcomes⁽¹⁵⁾.

Simultaneously, there is a growing recognition of the importance of men's involvement in ensuring adequate maternal and child nutrition^(16–18). Men play an important role in providing instrumental and emotional support to mothers and children⁽¹⁷⁾. For interventions to be effective, however, it is important to involve communities in their development; effective interventions tend to be embedded in the social life and ecology of the community which includes the men⁽¹⁹⁾.

The objective of this study was to carry out discussions with women and men in a rural community in Burkina Faso in order to collect context-specific insights into maternal and child health and nutrition issues. The intention was also to explore options for contextually appropriate interventions to improve maternal and child health and nutrition in the 'first 1000 d plus' period, this term describing the period from before conception up to a child's second birthday. This study is embedded in a larger, multicentre study entitled INPreP (Improved Nutrition Preconception, Pregnancy and Post-Delivery), which aims to understand how and where best to engage with community and relevant stakeholders to optimise nutrition in the 1000 d plus period in Burkina Faso, Ghana and South Africa⁽²⁰⁾. The qualitative findings from the other INPreP countries are presented in other papers in this special issue^(21–23).

Research methods

Study site and population

The study took place within the Nanoro Health and Demographic Surveillance System (HDSS) catchment area of the Nanoro Health district. The HDSS covers two

counties including Nanoro (fifteen villages) and Soaw (nine villages). The HDSS is located in the Center-West region of Burkina Faso, 90 km from the capital city, Ouagadougou⁽²⁴⁾. Seven peripheral health facilities and one referral hospital provide health care to the population within the HDSS. It is a poor, rural region with a population estimated in 2017 at 171 119 inhabitants⁽²⁴⁾, where approximately 55 % of people live on <0.70 US\$/d⁽²⁵⁾.

The population comprises farmers who produce food crops (sorghum, millet and maize) and cash crops (peanuts, beans, sesame and cotton) on lateritic soils. It is an area with irregular and poor distribution of rains, rudimentary cultivation techniques and soil degradation. Rearing livestock is the second economic activity in this population, particularly cattle, goats, sheep, pigs, donkeys and poultry, much of which is sold⁽²⁵⁾.

Diseases caused by inadequate potable water utilisation and poor sanitation are the main health problems⁽²⁶⁾. Malaria, acute respiratory infections and fever of unknown origin are the leading causes of mortality among adults and under-five children in this area^(27,28).

Study design and data collection

A qualitative methodology was adopted for this study because of its utility in obtaining culturally specific information about the values, opinions, behaviours, relationships and social contexts of particular populations. Focus group discussions (FGD) were chosen as the method of data collection because they allow discussion on a research topic where a specified population or group's opinions are given, and the meanings behind those perceptions are explored⁽²⁹⁾.

Discussions were conducted to explore the community's maternal and child health and nutrition priorities. Therefore, a topic guide (Appendix A and B) with standardised questions for the three INPreP sites was generated. Questions explored participants' views of the major health issues for women and children and of specific nutritional issues. Participants were invited to suggest interventions which might address these issues. Flexibility was built into the topic guide to allow for some probing where it was required for clarity.

Study participants were purposively recruited from ten of the twenty-four villages of the HDSS and from different households. Villages were selected in order to ensure that FGD were distributed across the study area (Fig. 1). Community informants facilitated the recruitment of participants for the FGD who represented age groups across the reproductive life course. Potential participants were given a Participant Information Sheet outlining the study purpose and procedures and those who agreed were invited to attend the FGD at a convenient location within the community. On the day of the FGD, participants' characteristics were confirmed and they were checked against the eligibility criteria. Written informed consent was obtained by a




Fig. 1 Map of the study area and distribution of focus group discussion per age group. W: women; M: men. •, Village; □, referral hospital; +, health facility; —, river; —, departmental road; □, Health and Demographic Surveillance System area

research assistant. Non-literate participants signed their consent form in the presence of a witness.

Participants were grouped by age and gender to enable them to freely express themselves. The research team felt that older women might inhibit younger women from speaking freely and that women might feel uncomfortable in joint meetings with men. Therefore, the groups were formed as follows: men aged 18–55 years, women aged 18–25 years; women aged 26–34 years and women aged 35–55 years.

FGD were conducted in the local language (Moore) by a moderator and an observer who took notes, these roles being taken by field assistants with a background in social research. Discussions were audio recorded.

Data analysis

Audio recordings from the FGD were simultaneously translated and transcribed verbatim from the local language into French and then into English by a social scientist. The study coordinator and the field assistant coded the transcripts independently in NVivo 12 qualitative analysis software (QSR International Pty Ltd), then inductively identified and agreed on key emerging themes through discussion. A consensus forming workshop was

held with the four INPreP research teams to thematically analyse the initial codes. Thematic analysis was used to organise and produce rich descriptions of the data following established guidelines⁽²⁹⁾. A code book was developed based on the topic guide, research objectives and the thematic grouping of the transcripts. The final set of codes was agreed by the research team, and definitions for each code were developed. Field notes taken during the data collection were used to supplement data from the transcripts.

Results

Participants characteristics

A total of eighty-six men and women participated in eleven FGD (Table 1). Of the participants, 73% were women, 57% had received no education and 27% were unemployed. Five women had never given birth (Table 2). Data are presented below as they relate to the study objectives to (i) understand context-specific issues which affected maternal and child health and nutrition and (ii) explore options for contextually appropriate interventions to improve maternal and child health and nutrition.

**Table 1** Composition of focus groups

FGD No.	Participants age (years)	Gender	Participants (n)
1	18–25	Women	8
2	35–55	Women	8
3	18–55	Men	8
4	18–55	Men	7
5	35–55	Women	8
6	35–55	Women	8
7	18–25	Women	8
8	18–55	Men	8
9	18–25	Women	8
10	26–34	Women	8
11	26–34	Women	7
Total			86

Table 2 Socio-demographic characteristics of participants

Characteristics of participants	Men		Women		Total	
	n	%	n	%	n	%
Age (years)						
18–25	4	17.4	24	30.1	28	32.6
26–55	19	82.6	39	61.9	58	67.4
Gender	23	26.7	63	73.3	86	100
Marital status						
Never married	4	17.4	7	11.1	11	12.8
Married monogamous	11	47.8	21	33.3	32	37.2
Married polygamous	7	30.4	32	50.8	39	45.3
Widowed	0	0	3	4.8	3	3.5
Religion						
Muslim	6	26.1	23	36.5	29	33.7
Catholic	10	43.5	24	38.1	34	39.5
Protestant	1	4.3	5	7.9	6	7
Animist	6	26.1	3	4.8	9	10.5
Unknown	0	0	8	12.7	8	9.3
Educational status						
No education	9	31.1	40	63.5	49	57
Primary	5	7.9	5	21.7	10	11.6
High school	5	21.7	17	27	22	25.6
Non-formal education	4	17.4	1	1.6	5	5.8
Occupation						
Unemployed	11	47.8	12	19	23	26.7
Self-employed/income-generating activities	12	52.2	51	81	63	73.3
Parity						
0	NA		5	7.9	5	5.8
1	NA		8	12.7	8	9.3
2–4	NA		28	44.4	28	32.6
5 and above	NA		22	34.9	22	25.6
Number of children						
0	5	21.7	5	7.9	10	11.6
1	2	8.7	10	15.9	12	14.0
2–4	2	8.7	29	46.0	31	36.0
5 and above	13	56.5	18	28.6	31	36.1

Obstacles to good maternal and child nutrition

Participants reported a number of barriers to optimal nutrition of mothers and children in their communities. Two major themes emerged: (1) poverty and (2) challenges generated by socially and culturally determined gender roles. Sub-themes related to access to health care and farming infrastructure to enhance food production and women lacking access to food whilst pregnant.

Themes and sub-themes are described below, supported by verbatim quotes from the FGD (Table 3).

Theme 1: The impact of poverty on women and children's health and nutritional status*Difficulties accessing health care*

Despite free universal health coverage for pregnant women and children under-five years of age implemented in Burkina Faso from 2017, poverty affected access to health care facilities. Poverty-related issues described by participants included distances between the household and the health care facility, the low quality of the free health care services and lack of availability and affordability of drugs.

After the consultation there are no drugs to buy. You can come back on next day and there is still no drug. (FGD7_Women 18–25 years)

Difficulty prioritising money for food

Lack of money also led to having to make choices between paying for health care or paying for food. Available money was in these cases often used for health care.

You can have money and you tell yourself that you are going to pay for a bag of rice, and when you get home, someone is sick and you use that money to go to the health center. There is no more bag of rice. (FGD8_Men 18–55 years)

This led to various coping mechanisms that often prioritised money spent on health care instead on nutrition.

If the child is sick, you can sell some part of your crops to pay for drugs. (FGD2_Women 35–55 years)

Climate change and food production

Both too much and too little rain reduced the crop yield for farmers and had an impact on their ability to feed themselves and their families. They described how some parts of the study area are now regularly flooded during the rainy season. This has led to displacement of some communities to places where they have to start a new life, build new houses and cultivate new, less fertile land.

Lack of rain also leads to low crop productivity and food shortages. In addition, some of the soil was poor quality and required fertilizer to grow millet, the staple food. This was challenging when farmers could not afford fertilizer or other farming inputs.

One year is not enough to solve one bad year's harvest. The year that it hadn't rained enough you manage to get through it. And the next year if you have good harvest you have to solve the old problems you get when you didn't have food to eat with the new harvest. So, if the granary becomes your box and you take grain out of there to eat and sell you are finished. (FGD3_Men 18–55 years)

Table 3 Additional illustrative quotes from research in Burkina Faso

Theme 1: The impact of poverty on women and children's health and nutritional status	
Difficulties accessing health care	<i>Woman can go into labor, and the time to find a motorcycle, put petrol and bring her to the health center she gives birth underway to the health center [...] Then we can say that the 7 km is far for us. (FGD11_Women 26–34 years)</i>
	<i>If there is no money to pay for the medicine it's another problem. If the child is sick, the mother also is sick. All of these are difficulties. (FGD3_Men 18–55 years)</i>
Climate change and food production	<i>It did not rain enough. It does not rain enough, after the rainy season some may have something but others end up empty-handed. (FGD4_Men 18–55 years)</i>
	<i>Water has overrun our houses 7 or 8 years ago, and we come here to live in the hills. Because of that we don't have enough food to eat anymore, we have few crops. It was a good field that we cultivated to have millet, and water has overrun it. (FGD6_Women 35–55 years)</i>
Water, sanitation and hygiene	<i>Stagnant dirty water and rotten food make for a lot of mosquitoes. When you throw away rotten food not far from the houses it causes a lot of mosquitoes. In addition, there are the waters where the pigs lay down, if this water is close to your house lot of mosquitoes come there. (FGD1_Women 18–25 years)</i>
	<i>There is also the problem of water [...] You can drink water from the wells thinking it is pure but it isn't. You drink and you get diarrhea, all of these are diseases. [...] We don't know what to do. (FGD3_Men 18–55 years)</i>
Theme 2: The impact of gender roles on women and children's health and nutritional status	
Women's social and domestic duties	<i>Nowadays women are more active than men; some of them travelling long distances over bicycle and falling sick. [...] When it happens that a woman is suffering from fever, no one cares. It is only when it gets worse that people are rushing to provide us with support. (FGD2_Women 35–55 years).</i>
	<i>The mother is pregnant and the child is not autonomous, so the mother suffers with the pregnancy and the child suffers too. He can't eat, his stomach becomes big. She is between two problems; she doesn't know if the child will survive, and she doesn't know if the one in her stomach will be well formed, because she can't get fatter like the others. (FGD3_Men 18–55 years)</i>
Lack of financial autonomy of women	<i>When we said here that we need warm clothes to protect children during the cold [...], it is true that it is the woman's responsibility. But the woman has nothing. The man's means are also limited so all these are difficulties for us. (FGD4_Men 18–55 years)</i>
	<i>Insufficient support from men</i>
	<i>If the children don't eat they are sick. The fathers do not take care of their family when they have money. It is outside that they spend it. (FGD7_Women 18–25 years)</i>
Theme 3: Improving maternal and child nutrition	
Involve women in income-generating activities	<i>We would like to benefit from training in making soap and dyeing, and help for the production of "soubbala *". (Women 35–55 years)</i>
	<i>We can set up a team and if we receive a financial aid we can cultivate a common field and share the crops. We can lead some income generating activities if we have financial support and we can share the earnings, and support ourselves. (FGD 6 Women 35–55 years)</i>
Facilitate access to food	<i>We cannot bring food with bags and distribute it. It is not possible. Some will have and others will not. Everyone must have an activity and be able to take care of his house and his children. (FGD11_Women 26–34 years)</i>
	<i>You must give food to women because some women can be pregnant but they don't have enough food to eat, we can distribute corn to pregnant women in order that they have enough to eat. (FGD9_women 18–25 years)</i>
Improvements in health infrastructure	<i>What we can add, we want health care workers that will sensitize people on good nutrition practices, hygiene, cleanliness, give advices to people, give them solution. We could do this through theatre; every one that follow it [the theatre] will change his behavior concerning nutrition and good hygiene practices. (FGD3_Momen 18–25 years)</i>
	<i>Sometimes women go to the Antenatal Care clinic (ANC) where they are given medications and are told to eat before taking the medication. And the woman has no money. If there could be food at the health center it will be fine. As we prepare for malnourished children. Thus, do in-patients have to eat. (FGD 8 Men 18–55 years)</i>
	<i>It's said "if you are carrying a weight more than you can, you ask for help" so we ask you for help. Even if it's not possible to do everything, you could reduce the price of medicine to allow us to pay. We don't have money. We are poor. (FGD3_Men 18–25 years)</i>
Support for agriculture	<i>If we have help to have a good harvest at the end of the season, we will have food to eat and if you get pregnant you will feed yourself well with that. (FGD10_Women 18–25 years)</i>
	<i>If we have a veterinary vaccine, it is a profitable activity: bullocks, sheep, goats. (FGD5_Women 35–55 years)</i>

*Soubbala is the fermented seeds of *Parkia biglobosa*, also known as the African locust bean.

Water, sanitation and hygiene

Lack of sanitation was also identified as a cause of poor maternal and child health. Access to potable water was limited by the presence of polluted and dirty wells. Polluted water caused nutritional problems because it is often the only source of water for drinking and cooking.

Even though the food is clean if you put it in a dirty dish it gives diseases. (FGD2_Women 35–55 years)

Men and women also reported difficulties in keeping the environment clean. Lack of facilities to dispose waste water

and trash in a hygienic way meant that households were often surrounded by rubbish, which they worried caused the spread of disease.

Theme 2: The impact of gender roles on women and children's health and nutritional status

Women's social and domestic duties

The need to fulfil gender-specific social and domestic roles was perceived as a barrier to improving maternal and child



nutrition by both men and women. Discussions reflected the fact that women were responsible for running the household, looking after the family and working the land. Participants, particularly older women, explained that prioritising such responsibilities often came at the cost of maternal and child health.

When you leave for the purchase of the goods there is no one to look after the children. While you went out of the house early [...] you did a job for which your body does not have health, and your child who stayed at home is not well taken care of. (FGD2_Women 35–55 years)

Women found it difficult to adequately fulfil their maternal role and ensure financial security.

If you're pregnant and you have a baby and you want to pedal the bike to carry out activities it's hard, and your kids will cry and people will make fun of you... (FDG11_Women 26–35 years)

Men seemed critical of women who juggled both economic activity and child care.

Some women do not take good care of children. They can put the child in the sand at the market and the child has fun with the sand while it can make him sick... (FGD4_Men 18–55 years)

Men were also critical of women's birth spacing, remarking on how pregnancy might negatively influence the mother and child's well-being. They suggested, for example, that women may not be able to properly breastfeed while pregnant. Participants said that giving birth to many children was a determinant of household food insecurity and was difficult at times of food shortages.

There are women who get children in a close way. [...] While the other one has not had enough food, he already has a little brother or sister [...] (FGD6_Men 18–55 years)

According to women, men acknowledged the burden pregnancy placed on mothers and children but were sometimes reluctant to use family planning.

There are women, if they adopt a contraceptive method, the husband can reject her... (FGD2_Women 35–55 years)

Lack of financial autonomy of women

Lack of money was a major problem for many women; they had to rely on their husband's income and lacked opportunities to generate their own money.

Women do not have any activity that generates income. If they had some, if they have a problem they can manage until the husband arrives... (FGD5_Women 35–55 years)

Insufficient support from men

The lack of male involvement in the household was widely discussed by women. Women felt overloaded with domestic

duties and saw the support they received from men in their households to be insufficient in a number of ways including nutritionally. First, women felt that men no longer took care of important needs such as paying for their children's schooling, clothing or health care.

The woman has to do her best to allow the child to wear clothes and eat. The man is going to drink his alcohol and come back late at night. He doesn't know what is going on with the woman and the children. (FGD5_Women 35–55 years)

Women felt that they were left with sole responsibility for farming and food production for their families in increasingly difficult conditions.

They have created a plough for women, and women tract themselves instead of the donkey. Women pay the price and use themselves. We call it 'I am my own donkey'. Nowadays [...] you can't even pull 10 lines and sow. (FGD6_Women 35–55 years)

Theme 3: Improving maternal and child nutrition

Participants suggested a range of solutions to the nutritional problems of women and their children in ways that were both nutrition-specific and nutrition-sensitive.

Facilitating access to food

Since poverty is a major problem and prevents access to food, participants said they would like to see food subsidies or receive these products for free.

If they could sell the food to us at 'social' prices it will be good, because nowadays you cannot expect one to come and give you free food like that. This is not possible. (FGD8_Men 18–55 years)

Participants also recognised that free or subsidised food provision was not a sustainable option nor going to solve root causes. Therefore, others felt that they needed training in skills that would help them improve their ability to grow food.

If you distribute food to everybody that will certainly bring issues; you have to train us on activities that will help us have food. Otherwise if they give you 30 bags of food you will eat all and will finish. (FGD3_Men 18–55 years)

Supporting agriculture

The community was aware of strategies for the development of resilience in agriculture and consequently wanted technical assistance and financial support to develop this sector.

If we can have ploughs and donkeys to pull them during the rainy season it will facilitate field work. If we also put fertilizer this will allow us to have good crops and our food will improve. (FGD11_Women 26–34 years)

They not only wished for help in raising livestock for breeding and for producing manure to fertilize the soil but also asked for support in keeping their animals healthy.



There is a lot to do. [. . .] If we can also breed in addition and you have natural fertilizer to enrich the ground, you will have a better harvest than someone that hasn't done the same. (FGD2_Women 35–55 years)

Involving women in income-generating activities

Women across all age groups said they would like to be employed in order to guarantee a livelihood for themselves and their families. They saw this as benefiting the whole community.

If they find us employment it will improve our living conditions and it will also develop the village. (FGD7_Women 18–25 years)

Woman asked for support to build their capacity to start businesses and acquire skills that would enable them to make their own money and take care of their families. They also talked about working their land collectively to benefit them all.

Women were clear about the connection between their ability to generate wealth and improved health.

In the other villages, women have income generating activities that they lead but here in Seguedin we do not have any. It is only us that have nothing. If we had activities with income our health would improve. (FGD10_Women 26–34 years)

Increasing men's commitment to caring for their families

Men's involvement was considered to be an essential factor in facilitating good maternal and child nutrition, especially their involvement in the decision to use family planning. Women wanted men to be made aware of the need for and benefits of family planning, even though they were not convinced that this would change men's behaviour.

One must raise men's consciousness to take care of their wives . . . But you cannot change all men. (Laughter) . . . They will not all change. There are some for whom it is not possible, you cannot change them. Some men, it is their friends who teach them to behave badly. (FGD11_Women 26–34 years)

Improving health infrastructure

The community, younger members in particular, suggested that a range of investments should be made to improve health infrastructure. These included making medications more affordable and accessible, employing more and better trained health care staff, increasing awareness of the importance of good nutrition and making food available at health care centres for in-patients.

We also want advice. When we come for consultation they can show us the food which is suitable for the children of six months, for example. It will also help us. (FGD7_Women 18–25 years)

Participants felt that making food available at health care centres would help women coming for consultation and

for in-patient stays. Currently, food for patients is only available at the market which takes place only every 3 d and the food bought there can be unsuitable. Patients have to bring food from home or go back home to eat, sometimes about 7 km from the health care centre.

They should recruit people to cook in the health centre because someone brings his patient and there is no food. [. . .] He is obliged to pay for dry bread for the patient. (FGD7_Women, 18–25 years)

Discussion

This study was carried out to explore the perspectives of men and women living in rural Burkina Faso on issues concerning maternal and child nutrition and how best to improve them. The community members in this study described factors influencing maternal and child nutrition such as income and resources, household food insecurity, lack of health services and unhealthy environments. Gender roles were widely perceived to impact nutrition; participants spoke of men limiting women's ability to control food production, and how men had become distanced from household responsibilities and care of women and children. This led to women asking for support with establishing income-generating activities to provide food for themselves and their families.

The causes of malnutrition discussed in the focus groups included changing gender roles, structural political responses and increasing vulnerability to climate change. Participants described the gender roles in their community, which they saw as affecting women and children's health and therefore their nutritional status. They spoke of the need for women to carry out their traditional domestic duties in addition to tasks outside of traditional gender roles such as farming the family's land^(30,31). In Nanoro, there is a long standing history of men migrating to Côte d'Ivoire in order to look for jobs and more financial stability⁽³²⁾. In these circumstances, women have to take on responsibilities to compensate for the loss of men's labour^(30,31). Women in this study explained, however, that they had limited financial autonomy and insufficient support from their husbands to carry out these tasks in the most effective way. Women described themselves as unable to purchase the food they needed and so consequently had to rely on men allocating them resources which were often insufficient to cover household needs. There is evidence that women's autonomy is positively linked to health care seeking behaviour for their children as well as themselves^(33,34). Likewise, women's participation in household decision-making and their ability to purchase food is significantly associated with increased diversity in the household diet⁽³⁵⁾. Participants in this study argued that improving women's financial autonomy was key to addressing malnutrition but required increased income-generating opportunities. There is evidence that empowering women improves women's health and has a positive impact on their nutritional status and that of their children^(7,12). An intervention in Burkina



Faso that gave women autonomy by providing them with land and agricultural training reported increased women's empowerment and nutritional status as well as significant reductions in child wasting^(14,15). The same intervention, however, not only had unintended consequences for women such as reduced time for child feeding and care, but also exposed them and their children to risks associated with livestock and chicken faeces⁽³⁶⁾.

The study described in this paper also reported women being too busy working at markets to take care of their children and provided more evidence that household chores and agricultural tasks can be detrimental to women's health and nutrition⁽³⁷⁾. Evidence from Ghana suggests that 70% of women had only two meals a day because needing to leave home early for work gave them too little time to eat⁽³⁸⁾. Difficult decisions regarding prioritisation of household and agricultural tasks over nutrition-related activities have to be made; time spent working on the land reduced time to gather firewood for cooking, for example, to go to the market to acquire a variety of foods and to prepare healthy and nutritious meals. All of this may affect dietary diversity of both child and mother^(39,40). Men's involvement has actually been considered an important element in supporting good maternal and child nutrition, since they have a double responsibility both as household decision-makers⁽⁴¹⁾ and as support to women and children^(44,45). Both men and women described the heavy workload that women bear and their limited financial opportunities, which were believed to damage their health and well-being and that of their children. Although research has shown the positive effect of male support on postpartum birth spacing practices⁽⁴⁴⁾, challenges remain. Further research exploring male perspectives on their roles in women's reproductive health management needs to be conducted in order to identify enablers to this and therefore helps define context and culturally specific solutions.

In addition to the debate on shifting gender roles, climate change and access to quality health care were discussed as having real impact on the quantity and quality of food for mothers and children in the communities in this study. Agriculture provides occupation for 86% of the rural population in Burkina Faso, but practices remain largely traditional and production relies on dependable climatic conditions, particularly the right amount of rainfall⁽⁴⁵⁾. Participants reported that inadequate or excessive annual rainfall was now more common. There is documentary evidence that these conditions are leading to a drop-in crop yields and a deficit of food⁽⁴⁶⁾. The effect of changes to patterns of the seasons on food intake has been described in resource-poor settings and linked with high rates of malnutrition in parts of Africa^(45–48). Because of this, the community suggested solutions to the nutrition problems that included social, economic and structural components. They would like their government to distribute free or subsidised food, in particular for the most vulnerable. Food donation or subsidisation is indeed a solution to

malnutrition. In Burkina Faso, the 'Société Nationale de Gestion du Stock de Sécurité Alimentaire' supports vulnerable people to buy cereals at subsidised prices⁽⁴⁹⁾. These initiatives have faced considerable challenges with, and therefore criticism of, processes for identifying and targeting those in need and avoiding abuse of the system by those who are not in need⁽⁵⁰⁾. Similar feelings of distrust were also expressed by community members towards health services in this study; they did not trust the quality of the free health care.

Participant also discussed causes of malnutrition which included lack of care for women and children, and household food insecurity linked to access to health services and medicines. It was perceived in the FGD that the burden of women's responsibilities is made heavier by the expectation that they will bear many children on top of their other roles already discussed. In Burkina Faso, the fertility rate remains high at 5.3 children per woman with low use of contraception (32%) among women aged 15–49 years^(51,52). Men are sometimes resistant to the use of contraception, constrained by their belief that children are bestowed by God, and by the value they place on them as a source of financial security⁽⁵³⁾. Uncontrolled fertility among women leading to inadequate birth spacing and frequent births is likely to reduce the time mothers have to provide optimum care to each child and for them to recuperate from the previous birth. This low autonomy in decision-making has been linked to sub-optimal usage of maternal health care services⁽⁵⁴⁾.

Other topics discussed by participants that affect maternal and child health and nutrition were access to the health care infrastructures and inadequate water, sanitation and hygiene. The most important contributing factor to low utilisation of health services by the poorest population groups was the expenditure it required in terms of payments for medicine, consultations and hospitalisation as well as indirect costs such as lost working time and transportation^(55,56). Therefore, short-term coping strategies were described to finance health care which, in the long term, prevented people from feeding themselves appropriately^(57,58). These sometimes included selling crops and farm animals or reducing food consumption in order to pay for catastrophic expenses. A previous study in the field of health economics conducted in 1996 in Burkina Faso showed how animals were used as 'ambulatory savings-banks' in order to pay for treatment⁽⁵⁹⁾, which was a concept indirectly described by community members. As in previous studies, participants in this study described selling crops grown for food to buy medicine, when necessary^(60,61).

Study strengths and limitations

An advantage of this study is that it was conducted in the context of multicentre research involving researchers from different backgrounds including public health, anthropology, nutrition and psychology. The thematic analysis was



conducted during a consensus forming workshop where the four INPreP research teams gather to share their data and agree themes and a set of codes from each site, thus enhancing the rigor of the analysis and interpretation. Another strength is that participants gathered in the FGD included men and women of different age groups allowing for diversity of experience and opinion. One limitation of the study, however, was that the interviews were conducted in the local languages, transcribed into French and then translated into English. Consequently, some statements may have lost their primary meanings through this process. To counteract this possibility, transcripts were shared between researchers and the representation of quotes and themes discussed and checked.

Conclusion

This paper offers context-specific insights into local maternal and child health and nutrition issues and identifies potential interventions suggested by community members as being appropriate and helpful for this population in Burkina Faso. The study highlights the importance of gender issues and of women's empowerment in addressing maternal and child nutrition in Burkina Faso. Findings reported in this paper make some fundamental points about the need for multi-sectorial interventions to address the nutrition environment in a way that is sustainable and that go beyond supplemental provision of food or nutrients.

Acknowledgements

Acknowledgements: We are most grateful to all community members for their time and efforts while participating in the study and their confidence in our research. *Financial support:* This research was funded by the National Institute for Health Research (NIHR) (17/63/154) using UK aid from the UK Government to support global health research. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care. *Conflict of interest:* The authors declare that they have no competing interests. *Authorship:* M.L.N., H.S., P.R.B., H.T., A.C. and M.B. designed the WP4 study. H.S., P.R.B., H.T. and M.B. supervised the implementation of the study and guided the manuscript conception. KO collected, transcribed and translated the data in English. A.C. checked and confirmed the accuracy of transcription and translation. A.C. and K.O. analysed the data. A.C. carried out the literature review and drafted the paper. All authors reviewed and approved the final manuscript. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki. This anthropological study was approved by

the National Health Ethics Committee in Burkina Faso, number 2018-12-156 and the University of Southampton Faculty of Medicine Ethics Committee, number 47290. All interviewees were informed before the start of the interview about project goals, the topic as well as their right to decline participation or to interrupt the conversation at any time. Anonymity was guaranteed and confidentiality of interviewees assured by assigning a unique code number to each informant. Written informed consent was obtained from all participants.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980020003365>

References

1. The World Bank (2019) World Bank country and lending groups. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups> (accessed July 2020).
2. UNICEF (2017) The State of the world's children 2017: children in a digital world. https://www.unicef.org/publications/files/SOWC_2017_ENG_WEB.pdf (accessed November 2019).
3. Ministry of Health (2016) *Enquête Nutritionnelle Nationale* [National Nutrition Survey]. Burkina Faso.
4. World Health Organisation (2020) *Nutrition Landscape Information System (NLIS) Country Profile*. Burkina Faso. <http://apps.who.int/nutrition/landscape/report.aspx?iso=BFA&rid=1620&goButton=Go> (accessed July 2020).
5. Vanderkooy A, Verstraeten R, Becquey E *et al.* (2019). *Nutrition Policy in Burkina Faso: TNWA Evidence Note 1*. Washington, DC: International Food Policy Research Institute (IFPRI).
6. Ministère de la santé Burkina Faso (2016) Politique nationale de nutrition [National nutrition policy]. Burkina Faso.
7. Shiwakoti R & Paudel R (2017) Women's empowerment and nutritional status of their children: a community-based study from villages of Bhaktapur district, Nepal. *Univer J Publ Health* **5**, 8–16.
8. Sethuraman K, Lansdown R & Sullivan K (2006) Women's empowerment and domestic violence : the role of socio-cultural determinants in maternal and child undernutrition in tribal and rural communities in South India. *Food Nutr Bull* **2**, 128–143.
9. Heise L, Greene ME, Opper N *et al.* (2019) Gender inequality and restrictive gender norms: framing the challenges to health. *Lancet* **393**, 2440–2454.
10. UNICEF (2011) Gender influences a narrative review on child survival, health and nutrition. https://www.unicef.org/Gender_Influences_on_Child_Survival_a_Narrative_review.pdf (accessed November 2019).
11. Ersino G, Zello GA, Henry CJ *et al.* (2018) Gender and household structure factors associated with maternal and child undernutrition in rural communities in Ethiopia. *PLoS One* **13**, e0203914.
12. Heckert J, Olney DK & Ruel MT (2019) Is women's empowerment a pathway to improving child nutrition outcomes in a nutrition-sensitive agriculture program? Evidence from a randomized controlled trial in Burkina Faso. *Soc Sci Med* **233**, 93–102.



13. Newton J, Verhart N & Blake A (2018) Enhancing the effectiveness of agriculture-to-nutrition pathways: key insights from a gender analysis of impact evaluation design. <http://knowledge4food.net/report-enhancing-effectiveness-agriculture-nutrition-pathways/> (accessed November 2019).
14. Olney DK, Pedehombga A, Ruel MT *et al.* (2015) A 2-year integrated agriculture and nutrition and health behavior change communication program targeted to women in Burkina Faso reduces anemia, wasting, and diarrhea in children 3–12.9 months of age at baseline: a cluster-randomized controlled trial. *J Nutr* **145**, 1317–1324.
15. Olney DK, Bliznashka L, Pedehombga A *et al.* (2016) A 2-year integrated agriculture and nutrition program targeted to mothers of young children in Burkina Faso reduces underweight among mothers and increases their empowerment: a cluster-randomized controlled trial. *J Nutr* **146**, 1109–1117.
16. Ochieng J, Afari-Sefa V, Lukumay PJ *et al.* (2017) Determinants of dietary diversity and the potential role of men in improving household nutrition in Tanzania. *PloS One* **12**, e0189022.
17. Kansime N, Atwine D, Nuwamanya S *et al.* (2017) Effect of male involvement on the nutritional status of children less than 5 years: a cross sectional study in a rural southwestern district of Uganda. *J Nutr Metab* **2017**, 3427087.
18. Barker M, Dombrowski SU, Colbourn T *et al.* (2018) Intervention strategies to improve nutrition and health behaviours before conception. *Lancet* **391**, 1853–1864.
19. McCloskey DJ, McDonald MA, Cook J *et al.* (2007) Community engagement : definitions and organizing concepts from the literature. https://www.atsdr.cdc.gov/communityengagement/pdf/PCE_Report_Chapter_1_SHEF.pdf (accessed November 2019).
20. University of Southampton Global Health Research Institute (2019) Improved Nutrition Pre- conception, Pregnancy and Post-Delivery (INPreP). <https://www.southampton.ac.uk/global-health/research/lifecourse-epidemiology/inprep/about-us.page> (accessed November 2019).
21. Erzse A, Goldstein S, Norris SA *et al.* (2020) Double-duty solutions for optimising maternal and child nutrition in urban South Africa: a qualitative study. *Public Health Nutr*, 1–11. doi: 10.1017/S1368980020002426.
22. Debpuur C, Nonterah EA, Chatio ST *et al.* (2020) Supporting maternal and child nutrition: Views from community members in rural Northern Ghana. *Public Health Nutr*. In press.
23. Watson D, Kehoe SH, Erzse A *et al.* (2020) Community perspectives of maternal and child health during nutrition and economic transition in sub-Saharan Africa. *Public Health Nutr*, 1–9. doi: 10.1017/S1368980020003018.
24. Derra K, Rouamba E, Kazienga A *et al.* (2012) Profile: Nanoro health and demographic surveillance system. *Int J Epidemiol* **41**, 1293–1301.
25. Zida Y & Kambou SH (2014) Cartographie de la pauvreté et des inégalités au Burkina Faso [Mapping poverty and inequalities in Burkina Faso]. https://www.undp.org/content/dam/burkina_faso/docs/publications/UNDP_bf_cartopauv_2014.pdf (accessed November 2019).
26. Ministry of Health, Directorate of Statistics and Planning (2016) Statistical Directory. Burkina Faso.
27. Tinto H, Valea I, Sorgho H *et al.* (2014) The impact of clinical research activities on communities in rural Africa: the development of the Clinical Research Unit of Nanoro (CRUN) in Burkina Faso. *Malar J* **13**, 113.
28. Maltha J, Guiraud I, Berenger K *et al.* (2014) Frequency of severe malaria and invasive bacterial infections among children admitted to a rural hospital in Burkina Faso. *PloS One* **9**, e89103.
29. Nyumba TO, Wilson K, Derrick CJ *et al.* (2018) The use of focus group discussion methodology: insights from two decades of application in conservation. *Methods Ecol Evol* **9**, 20–32.
30. Lantara NF (2015) The roles of woman as leader and housewife. *J Def Manag* **5**, 1–5.
31. Spurling D (1995) *Les Femmes Rurales et la Vulgarisation Agricole Dans le Sahel: Africa Reg. Find* [Rural women and agricultural vulgarisation in the Sahel Reg. Find], vol. **46**. Washington, DC: World Bank.
32. Ministère de la santé (2017) Plan d'action 2017 Du district sanitaire de Nanoro [Nanoro health district annual action plan 2017]. Burkina Faso.
33. Mkandawire E & Hendriks SL (2018) A qualitative analysis of men's involvement in maternal and child health as a policy intervention in rural Central Malawi. *BMC Pregnancy Childbirth* **18**, 37.
34. Ebot JO (2015) 'Girl power': The relationship between women's autonomy and children's immunization coverage in Ethiopia. *J Heal Popul Nutr* **33**, 18.
35. Amugsi DA, Lartey A, Kimani E *et al.* (2016) Women's participation in household decision-making and higher dietary diversity: findings from nationally representative data from Ghana. *J Heal Popul Nutr* **35**, 16.
36. Ruel MT, Quisumbing AR & Balagamwala M (2018). Nutrition-sensitive agriculture: what have we learned so far? *Glob Food Secur* **17**, 128–153.
37. Nti CA, Inkumsah D & Fleischer G (1999) Influence of women's workload on their nutritional status in selected communities in Ghana. *Int J Consum Stud* **23**, 165–170.
38. Kadiyala S, Harris J, Headey D *et al.* (2014) Agriculture and nutrition in India : mapping evidence to pathways. *Ann N Y Acad Sci* **1331**, 43–56.
39. Komatsu H, Malapit HJL & Theis S (2018) Does women's time in domestic work and agriculture affect women's and children's dietary diversity? Evidence from Bangladesh, Nepal, Cambodia. *Food Policy* **79**, 256–270.
40. The World Bank (2017) Fertility rate, total (births per woman): Burkina Faso. World Bank DataBank. =<https://data.worldbank.org/indicator/SP.DYN.TFRT.IN?locations=BF> (accessed November 2019).
41. Singh D, Lample M & Earnest J (2014) The involvement of men in maternal health care : cross-sectional, pilot case studies from Maligita. *Reprod Health* **11**, 68.
42. Thorsen D (2002) 'We Help our Husbands!' Negotiating the household budget in rural Burkina Faso. *Dev Change* **33**, 129–146.
43. Jennings L, Na M, Cherewick M *et al.* (2014) Women's empowerment and male involvement in antenatal care : analyses of Demographic and health surveys (DHS) in selected African countries. *BMC Pregnancy Childbirth* **14**, 297.
44. Daniele MAS, Sarrassat S, Cousens S *et al.* (2018) Involving male partners in maternity care in Burkina Faso: a randomized controlled trial. *Bull World Health Organ* **96**, 450–461.
45. Ibrahim B, Polcher J, Karambiri H *et al.* (2012) Characterization of the rainy season in Burkina Faso and it's representation by regional climate models. *Clim Dyn* **39**, 1287–1302.
46. Moursi M, Moura FF De & Martin-Prevel Y (2014) Seasonal differences in food and nutrient intakes among young children and their mothers in rural Burkina Faso. *J Nutr Sci* **3**, 1–9.
47. Savy M, Martin-Prevel Y, Traissac P *et al.* (2006) Dietary diversity scores and nutritional status of women change during the seasonal food shortage in rural Burkina Faso. *J Nutr* **136**, 2625–3262.
48. van Cooten MH, Bilal SM, Gebremedhin S *et al.* (2019) The association between acute malnutrition and water,



- sanitation, and hygiene among children aged 6–59 months in rural Ethiopia. *Matern Child Nutr* **15**, 1–8.
49. Société Nationale de Gestion du Stock de Sécurité Alimentaire (SONAGESS) (2019) <https://sonageSS.bf> (accessed November 2019).
 50. USAID (2012) Agriculture and Food Security Burkina Faso Fact Sheet. <https://www.usaid.gov/burkina-faso/agriculture-and-food-security> (accessed November 2019).
 51. The World Bank (2016) Contraceptive prevalence, any methods (% of women ages 15–49) Burkina Faso. World Bank DataBank. <https://data.worldbank.org/indicator/SP.DYN.CONU.ZS?view=chart> (accessed November 2019).
 52. Naik R & Smith R (2015) *Impacts of Family Planning on Nutrition*. Washington, DC: World Bank, Futures Gr. https://www.healthpolicyproject.com/pubs/690_FPandnutritionFinal.pdf (accessed November 2019).
 53. Kabagenyi A, Jennings L, Reid A *et al.* (2014) Barriers to male involvement in contraceptive uptake and reproductive health services: a qualitative study of men and women's perceptions in two rural districts in Uganda. *Reprod Health* **11**, 1–9.
 54. Tiruneh FN, Chuang KY & Chuang YC (2017) Women's autonomy and maternal healthcare service utilization in Ethiopia. *BMC Health Serv Res* **17**, 1–12.
 55. Samuelsen H (2004) Therapeutic itineraries: the medical field in rural Burkina Faso. *Anthropol Med* **11**, 27–41.
 56. Nguyen KT, Hai Khuat OT, Ma S *et al.* (2012) Coping with health care expenses among poor households: evidence from a rural commune in Vietnam. *Soc Sci Med* **74**, 724–733.
 57. Murphy A, McGowan C, McKee M *et al.* (2019) Coping with healthcare costs for chronic illness in low-and middle-income countries: a systematic literature review. *BMJ Glob Heal* **4**, e001475.
 58. Sauerborn R, Adams A & Hien M (1996) Household strategies to cope with the economic costs of illness. *Soc Sci Med* **43**, 291–301.
 59. Hausmann-Muela S (2011) The straw that breaks the camel's back. *Med Anthropol Quarterly* **25**, 103–121.
 60. Su TT, Kouyaté B & Flessa S (2006) Catastrophic household expenditure for health care in a low-income society: a study from Nouna district, Burkina Faso. *Bull World Health Organ* **84**, 21–27.
 61. Somé JW & Jones AD (2018) The influence of crop production and socioeconomic factors on seasonal household dietary diversity in Burkina Faso. *PLoS One* **13**, e0195685.